

**UC Davis**  
**Dermatology**

**Title**

Identification of psoriasis-protective IL17D variant associated with increased IL17D and FAM19A5 expression in psoriatic skin

**Permalink**

<https://escholarship.org/uc/item/3041586k>

**Authors**

Toussi, Atrin

Merleev, Alexander

Le, Stephanie

et al.

**Publication Date**

2021

**Data Availability**

The data associated with this publication are not available for this reason: N/A

# Identification of psoriasis-protective *IL17D* variant associated with increased *IL17D* and *FAM19A5* expression in psoriatic skin

Atrin Toussi,<sup>1,2</sup> Alexander Merleev,<sup>1</sup> Stephanie T. Le,<sup>1</sup> Alina Marusina,<sup>1</sup> Guillaume Luxardi,<sup>1</sup> Lam (Alex) C. Tsoi,<sup>3</sup> Johann E. Gudjonsson,<sup>4</sup> Emanuel Maverakis<sup>1</sup>

<sup>1</sup>Department of Dermatology, University of California, Davis, Sacramento, CA


<sup>2</sup>University of California, Davis, School of Medicine, Sacramento, CA

<sup>3</sup>Department of Biostatistics, Center for Statistical Genetics, University of Michigan, Ann Arbor, MI

<sup>4</sup>Department of Dermatology, University of Michigan, Ann Arbor, MI

Disclosures: None

# *IL17D* expression and variant-associated changes




Psoriasis is a chronic inflammatory skin disease with an estimated heritability of 80% (Rahman P, Elder JT 2005). The IL-17 family plays a central role in the mediation of inflammatory pathways, including psoriasis. The most well-studied of these, IL-17A, is a major pro-inflammatory cytokine (Monin L, Gaffen SL 2018; Blauvelt A, Chiricozzi A 2018).

**Methods:** RNA-Seq FASTQ files of human normal and psoriasis lesional skin were downloaded from the NCBI Sequence Read Archive (<http://www.ncbi.nlm.nih.gov/Traces/sra>). The RNA-Seq dataset used for this analysis consisted of 99 psoriasis vulgaris skin biopsy samples obtained from patients washed out of all systemic and topical therapies and 90 healthy controls.


**Results:** A, *IL17A* is highly upregulated in psoriatic skin and appears to be a driver of disease (Rembilla NC, Senra L, Boehncke, 2018). *IL17F* shares the greatest homology to *IL17A*<sup>3</sup> and is also highly upregulated in psoriasis. In contrast, RNA-Seq datasets identified that *IL17D* is highly expressed in normal skin but downregulated (Fold Change = 0.33; p = 2.5x10<sup>-14</sup>)<sup>5</sup> in psoriatic plaques.<sup>6</sup> B, an *IL17D* regulatory variant, rs9509353, was found to be protective against psoriasis (OR = 0.20, p-value = 5.9e-07), increasing expression of *IL17D* and *FAM19A5*, a chemokine-like molecule that is also normally downregulated in psoriasis.

# 2D correlation & distribution: tSNE


## Psoriasis



## Healthy



## Atopic



**Methods:** Gene expression clusters were mapped using the t-Distributed Stochastic Neighbor Embedding (t-SNE) method. Distance was calculated as  $(d) = 1 - r^2$ , where  $r$  equals Pearson's correlation coefficient, calculated with Rtsne package (Le et al. 2019).


**Results:** 2D images of the psoriasis and healthy transcriptome illustrate the close spatial relationship between *IL17D*, *FAM19A5*, and *SCARA5*, which (1) map together and (2) away from the proinflammatory cytokine cluster comprised of *IL17A*, *IL23A*, *IL36*, and *IL1B*. Notably, these proinflammatory cytokines do not cluster together in healthy skin. Correlation studies found these cytokine clusters to be independent of *IL17D* and *FAM19A5*. Combined, these results highlight a putative regulatory role for IL-17D and FAM19A5 in psoriasis.

# IL17D meta-analysis and correlations


**A. Methods:** Correlation analyses of gene expressions were performed on read counts of each identified gene normalized with DESeq2 package. Values were subsequently log transformed and winsorized when necessary. Spearman's correlation coefficients were calculated ( $r_s$ ) using the `cor.test` function in R. P values were estimated by algorithm AS 89 (Le, et al. 2019). **Results:** Correlative studies demonstrate a close relationship between *IL17D* and *FAM19A5*, *IL17D* and *SCARA5*, and an inverse relationship between *IL17D* and *NOD2* in both the healthy and psoriasis transcriptomes. Other examples of highly correlated genes include *TGF-B2*, *TGF-R3*, *BTRC*, *TP63*, and *TIMP3*.

**B. Methods:** Meta-analysis was completed using the R package "metafor." A weighted random-effects model was used to estimate a summary effect size. To estimate between-study variance, a restricted maximum-likelihood estimator was applied. A weighted estimation with inverse-variance weights was used to fit the model (Le al. 2019). **Results:** Meta-analysis across multiple psoriasis datasets confirmed close relationships between *IL17D*, *FAM19A5*, *SCARA5*, and *NOD2*, all depicted here.



A.



B.



# Single cell sequencing data



**Methods:** Raw read data was processed using the 10x Genomics Cell Ranger and results were analyzed and visualized with 10x Genomics Loupe cell Browser.

**Results:** A, single cell data demonstrating clusters of genes from different cell populations. B, putative mechanism of IL-17D regulation in psoriasis. *IL17D* is expressed by monocytes and acts in an autocrine-like fashion to increase *SCARA5* expression intracellularly, and to increase *FAM19A5* and *TGFB-2* secretion. *IL-17D* then acts on keratinocytes to reduce expression of pro-inflammatory genes, *NOD2*, *BTRC*, and *TP63*, and to increase the expression of anti-inflammatory genes, *TIMP3* and *TGFB-R3*, which encodes a *TGFB* receptor. *TGFB-2* from monocytes then interacts with its upregulated receptor on keratinocytes and separately acts on T cells to increase T regulatory cell production and to suppress T cell function, thereby reducing inflammation in psoriasis.