

UC Irvine

UC Irvine Previously Published Works

Title

Stereocontrolled Access to Quaternary Centers by Birch Reduction/Alkylation of Chiral Esters of Salicylic Acids

Permalink

<https://escholarship.org/uc/item/31z823gq>

Journal

The Journal of Organic Chemistry, 88(9)

ISSN

0022-3263

Authors

Kozlowski, Ryan A
Nguyen, Hanh T
Lehman, Michael E
[et al.](#)

Publication Date

2023-05-05

DOI

10.1021/acs.joc.3c00306

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Stereocontrolled Access to Quaternary Centers by Birch Reduction/
Alkylation of Chiral Esters of Salicylic AcidsRyan A. Kozlowski,[§] Hanh T. Nguyen,[§] Michael E. Lehman, and Christopher D. Vanderwal*Cite This: *J. Org. Chem.* 2023, 88, 6232–6236

Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: 8-Phenylmenthol esters of salicylic acid derivatives undergo efficient Birch reduction and *in situ* diastereoselective alkylations to afford methoxycyclohexadienes bearing new quaternary stereogenic centers. The use of an ester-based auxiliary is a designed improvement over the use of prolinol-derived amides, which are expensive and often very difficult to cleave.



The venerable Birch reduction, initially disclosed in 1944, has served as a key transformation in the syntheses of many complex molecules.¹ The ability to take feedstock or readily available aromatic compounds and obtain high-value-added, functional-group-rich products has proven central to the efficient synthesis of many targets.² Although recent advances include ammonia-free³ or even metal-free⁴ Birch-type reduction conditions, the classic Birch reduction using a group 1 or 2 metal dissolved in ammonia remains the most widely used method.

The presence of an anion-stabilizing group on the substrate arene—usually a carboxylic acid derivative—drives the regiochemical control of the Birch reduction and generally results in the formation of a stabilized carbanion. The enolate so formed can be used productively for C–C bond formation, generating a new quaternary carbon.⁵ The stereoselective synthesis of quaternary carbons remains a significant challenge for synthetic chemistry,⁶ and the Birch reduction/alkylation reaction is a powerful tool for directly forming quaternary carbons from aromatic rings. This Birch reduction/alkylation process of carboxylic acid derivatives was rendered diastereoselective by Schultz through the use of a proline-derived chiral auxiliary attached as an amide.^{7,8} This gives rise to enantioenriched materials following removal of the chiral auxiliary. Schultz and others have synthesized several natural products using this method;^{8b,9} however, the auxiliary removal is often nontrivial, requiring harsh or substrate-tailored conditions, or multiple steps.^{8b,9,10} Herein we disclose the first example of a nonamide based chiral auxiliary for use in a diastereoselective Birch reduction/alkylation reaction to produce highly functionalized cyclohexadienes.

During the course of efforts toward the total synthesis of a natural product, we aimed to use Schultz's auxiliary to set a critical quaternary stereogenic center starting from a substituted salicylic acid derivative (Figure 1). While we observed that the diastereoselective Birch reduction/alkylation




Figure 1. Successful Schultz-type Birch reduction/alkylation yields a product from which the auxiliary cannot be removed.

reaction proceeded well, we found that the steric bulk of the auxiliary and proximal functionality prevented subsequent desired transformations. Additionally, cleanly removing the chiral auxiliary proved impossible, as a result of poor reactivity of the amide and/or undesired side reactivity. As a result of these difficulties, we aimed to develop a more easily removable chiral auxiliary for a diastereoselective Birch reduction/alkylations of benzenoid systems.

Our attention turned toward the chiral pool in the hope of discovering an ester-based chiral auxiliary, with the assumption that hydrolytic removal would be much easier than with the amides; furthermore, a reductive option is also available that is not trivial with amides, which would likely leave the prolinol auxiliary attached via an amine linkage. Initially, and unsurprisingly, borneol, isomenthyl, and menthyl esters of 3,5-dimethoxy salicylic acid (3, Figure 2) provided essentially no diastereoselectivity (~1:1) using iodomethyl pivalate as the

Received: February 9, 2023

Published: April 11, 2023






Figure 2. Identification of a competent ester-based auxiliary.

electrophile. We attribute this lack of diastereoselection to the lack of any obvious mechanism for conformational restriction,

and thus transfer of chirality in the alkylation step. 8-Phenylmenthol esters (or esters of other chiral arene-bearing alcohols) have been found to deliver high diastereoselectivity in situations where they can engage in a π -stacking interaction with the substrate.¹¹ The most relevant work is that by Donohoe and co-workers on the diastereoselective Birch reduction of pyrroles using “cumyl” and (-)-8-phenylmenthol esters.¹² However, until recently, 8-arylmenthols were prohibitively expensive and nontrivial to synthesize. Importantly, recent work by Shenvi and co-workers has demonstrated that (-)-8-phenylmenthol and analogues bearing other aromatic substituents can easily be synthesized in two steps from pulegone,¹³ making this family of chiral auxiliaries more easily accessible than it has been previously.

In a promising initial result, we found that the (-)-8-phenylmenthyl ester of 3,0-dimethyl salicylic acid was reduced and alkylated in 46% yield and with 4:1 dr, using iodomethylpivalate as the electrophile (Figure 2). We ascribe this increase in diastereoselectivity to a potential π -stacking interaction between the phenyl ring on the chiral auxiliary and the extended enolate resulting from the reduced aromatic ring. The moderate yield and diastereoselectivity was attributed to the poor solubility of **3** in ammonia, as observed by the reaction mixture becoming a viscous, difficult-to-stir, milky-white suspension at -78 °C. Increasing the amount of THF




Figure 3. Products of Birch reduction/diastereoselective alkylation of 8-phenylmenthol esters of salicylic acid derivatives (8-PM = 8-phenylmenthyl). The relative configurations of **5** and **20** were assigned via X-ray crystallography; others assigned by analogy.

(5:1 $\text{NH}_3/\text{THF} \rightarrow 2:1$) improved solubility and increased the yield and d.r. to 66% and 7:1, respectively. Increasing the ratio further to 1:1 ammonia/THF increased the yield to 80% and kept the d.r. at a respectable 7:1. Given the ease of synthesis of various arylated menthol derivatives according to the protocol of Shenvi and co-workers,¹³ 1-naphthyl, 2-naphthyl, 4-fluorobenzenyl, and 3,5-bis(trifluoromethyl)benzenyl derivatives were investigated to see if extending the aromatic system or making the aromatic ring on the chiral auxiliary more electron-poor would increase the hypothesized π -stacking interaction with the electron-rich Birch reduction intermediate.¹⁴ Unfortunately and not unexpectedly, competitive reduction of the more electron-deficient aromatic rings of these auxiliaries prevented validation of this hypothesis. Given these results, we proceeded with (–)-8-phenylmenthol because we observed no competitive reduction of the phenyl group, good yields of the desired product, and useful levels of diastereoselectivity.

To explore the scope of this reaction, we looked to vary the substitution pattern on the aromatic ring and the identity of the electrophile (Figure 3). For otherwise unsubstituted *O*-methyl salicylic esters (products 5–9), as well as 3-methyl- (products 10–14) and 5-methyl-substituted substrates (products 15–19), benzylic, methyl, and alkyl halides were competent electrophiles, in all cases resulting in $\geq 3:1$ dr. Particularly interesting electrophiles that generate synthetically malleable products include iodomethyl pivalate (generating 8 and 13 with high selectivities) and bromoacetonitrile (18 and 19). 6-Methyl salicylate derivatives (products 20–23) performed particularly well with $\sim 20:1$ dr in all cases except for with methyl iodide. If one trend with electrophiles did emerge from these examples, it is that smaller electrophiles tend toward slightly diminished stereoselectivities. The efficiency of reaction held for the 4-methyl substrate, generating 24 with two new stereogenic centers; the configuration of the distal one at C4 is presumably not controlled by the distal auxiliary, resulting in a mixture of diastereomers.¹⁵ These results overall strongly suggest that a broad range of 8-phenylmenthyl esters of *O*-alkyl salicylates will undergo Birch reduction/alkylation with synthetically useful levels of selectivity. Furthermore, the reaction can be performed preparatively, with generation of 10 done on 2-g scale in 83% yield with the same 5:1 dr; further chromatography led to a 63% yield of diastereomerically enriched product (11:1 dr).

There were some cases wherein this protocol proved poorly effective (products 25–28). Nitrobenzyl electrophiles—designed to try to encourage crystallinity in the products—decomposed under the reaction conditions, leading to low yields. *tert*-Butyl haloacetates were poorly reactive, resulting in low conversion in the alkylation step. Attempted reaction of the intermediate enolate with chloromethyl methyl sulfide led to decomposition. Finally, and not surprisingly, β -branched electrophiles, such as the prolinol-derived reactant that might have formed 28, were completely unreactive.

Looking to expand beyond salicylates, we examined 2-methyl and 2-fluoro substrates 29 and 30 (Figure 4). Although these compounds each underwent reduction/alkylation in reasonable yields, the diastereoselectivities were well below 2:1, strongly implicating chelation as a stereocontrol element in the more selective alkylations, as we had anticipated.¹⁶ Similarly, 3-substituted substrates 31 and 32 performed efficiently in the reaction, but again with essentially no stereocontrol. Perhaps not surprisingly, 3-chloro reactant 33




Figure 4. Substrates that resulted in poor diastereoselectivity or decomposed during reduction (8-PM: 8-phenylmenthyl).

decomposed during the reduction step, presumably via Cl–C bond reduction.

We attribute the requirement of the 2-alkoxy group to its likely coordination to lithium during and after formation of the extended enolate following arene reduction. The resulting conformational control, along with likely π -stacking of the phenyl group with the cross-conjugated dienolate, leads to shielding of one enolate face, allowing for selective alkylation (Figure 5). We expect that the reactive conformation is




Figure 5. Conformational analysis of the reactive cross-conjugated dienolate.

approximated by structure 34a, in which no obvious deleterious nonbond interactions exist, and both chelation and π -stacking can be operative; the products for which we have unambiguous structural assignment from X-ray crystallography¹⁴ are consistent with this model. Rotation about the enolate C–O bond can produce conformation 34b, largely devoid of π -stacking and engendering significant nonbonded interactions between the carbinol H and R group at C6 on the arene; circumstantially, this idea is supported by the improved selectivity with R = CH_3 compared with R = H (see examples 20–23 compared to their analogues without the C6 methyl, Figure 3). Conformation 34c, while maintaining π -stacking, results in nonbonded interactions with axial C–H bonds on the phenylmenthol ring. Of course, without a C2-alkoxy group to coordinate the lithium, the substrate may form the enolate in either configuration, potentially leading to poor diastereoselectivity even if π -stacking is maintained.

One of the hallmarks of auxiliary-based diastereoselection is that separation can often lead to the isolation of stereochemically homogeneous products; unfortunately, in many of the cases described in Figure 3, significant upgrading of stereochemical purity via column chromatography was infeasible. Removal of the chiral auxiliary can be accomplished efficiently and with good recovery of (–)-8-phenylmenthol using lithium aluminum hydride (Figure 6). However, there remain some limitations of the ester-based auxiliary in that, like the Schultz amide system, simple hydrolysis is often ineffective, almost certainly owing to the substantial steric encumbrance to the approach of any nucleophile to π^* of the ester carbonyl. Lewis acid activation and dealkylation conditions were similarly ineffective.¹⁴ However, the ability to achieve simple reductive removal is an advance relative to amides in many cases, wherein reductive formation of the amine incorporating the auxiliary is generally undesired.




Figure 6. Representative examples of reductive auxiliary cleavage.

We have developed an auxiliary-controlled stereoselective Birch reduction/alkylation reaction, making use of esters of the readily available 8-phenylmenthol. The product 1,4-cyclohexadienes bearing quaternary stereogenic centers are in many cases formed in good yield and high diastereoselectivity. The ester linkage to the chiral auxiliary is easily reduced, rendering the overall process a marked improvement over previous amide-based chiral auxiliaries. Despite the requirement of 2-alkoxy substitution in almost all cases, substitution at the 3, 5, and 6 positions on the aromatic ring are tolerated, as are a variety of electrophiles. In short, this method provides chemists with a simple, scalable, and diastereoselective reaction to form all-carbon quaternary centers for use in pursuit of highly functionalized cyclohexane scaffolds.

■ ASSOCIATED CONTENT

Data Availability Statement

The data underlying this study are available in the published article and its Supporting Information.

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.3c00306>.

Experimental procedures, characterization data, NMR spectra for all new compounds, and X-ray crystallographic data (PDF)

Accession Codes

CCDC 2236575–2236577 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

Christopher D. Vanderwal – Department of Chemistry, 1102 Natural Sciences II, University of California, Irvine, California 92697-2025, United States; Department of Pharmaceutical Sciences, 101 Theory #100, University of California, Irvine, California 92617, United States; orcid.org/0000-0001-7218-4521; Email: cdv@uci.edu

Authors

Ryan A. Kozlowski – Department of Chemistry, 1102 Natural Sciences II, University of California, Irvine, California 92697-2025, United States; orcid.org/0000-0003-3860-0565

Hanh T. Nguyen – Department of Chemistry, 1102 Natural Sciences II, University of California, Irvine, California 92697-2025, United States

Michael E. Lehman – Department of Chemistry, 1102 Natural Sciences II, University of California, Irvine, California 92697-2025, United States

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.joc.3c00306>

Author Contributions

[§]R.A.K. and H.T.N. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was funded by the NSF through grant CHE-2102480, and H.T.N. was supported by an NSF Graduate Research Fellowship. We thank the Dong lab (UCI) for the use of their SFC instrument and for their assistance. SFC and HPLC analyses for enantiopurity on some samples were performed with instrumentation at the Caltech Center for Catalysis and Chemical Synthesis, a facility of the Beckman Institute at Caltech. We thank Dr. Scott Virgil for his assistance.

■ REFERENCES

- (1) (a) Rabideau, P. W.; Marcinow, Z. Birch Reduction of Aromatic Compounds. *Org. React.* **1992**, 42, 1–334. (b) Birch, A. J. The Birch Reduction in Organic Synthesis. *Pure Appl. Chem.* **1996**, 68, 553–556.
- (2) Hook, J. M.; Mander, L. N. Recent developments in the Birch reduction of aromatic compounds: applications to the synthesis of natural products. *Nat. Prod. Rep.* **1986**, 3, 35–85.
- (3) Burrows, J.; Kamo, S.; Koide, K. Scalable Birch Reduction with Lithium and Ethylenediamine in Tetrahydrofuran. *Science* **2021**, 374, 741–746.
- (4) (a) Cole, J. P.; Chen, D.-F.; Kudisch, M.; Pearson, R. M.; Lim, C.-H.; Miyake, G. M. Organocatalyzed Birch Reduction Driven by Visible Light. *J. Am. Chem. Soc.* **2020**, 142, 13573–13581. (b) Peters, B. K.; Rodriguez, K. X.; Reisberg, S. H.; Beil, S. B.; Hickey, D. P.; Kawamata, Y.; Collins, M.; Starr, J.; Chen, L.; Udyavara, S.; Klunder, K.; Gorey, T. J.; Anderson, S. L.; Neurock, M.; Minter, S. D.; Baran, P. S. Scalable and Safe Synthetic Organic Electroreduction Inspired by Li-ion Battery Chemistry. *Science* **2019**, 363, 838–845.
- (5) For a pioneering example in natural product synthesis, see: Marshall, J. A.; Wuts, P. G. M. Stereoselective Synthesis of Racemic Occidentalol and Related Cis-Fused Hexahydronaphthalenes from *m*-Toluic Acid. *J. Org. Chem.* **1977**, 42, 1794–1798.
- (6) Fuji, K. Asymmetric creation of quaternary carbon centers. *Chem. Rev.* **1993**, 93, 2037–2066.
- (7) Schultz, A. G.; Sundaraman, P. Enantioselection in the Birch Reduction–Alkylation of a Chiral Benzoic Acid Derivative. *Tetrahedron Lett.* **1984**, 25, 4591–4594.
- (8) For reviews, see: (a) Schultz, A. G. Enantioselective Methods for Chiral Cyclohexenone Synthesis. *Acc. Chem. Res.* **1990**, 23, 207–213. (b) Schultz, A. G. The asymmetric Birch reduction and reduction/alkylation strategies for synthesis of natural products. *Chem. Commun.* **1999**, 1263–1271.
- (9) For representative examples, see: (a) Zhou, Q.; Snider, B. B. Synthesis of (±)- and (–)-Vibralactone and Vibralactone C. *J. Org. Chem.* **2008**, 73, 8049–8056. (b) Jousseume, T.; Retailleau, P.; Chabaud, L.; Guillou, C. Studies on the asymmetric Birch reductive alkylation to access spiroimines. *Tetrahedron Lett.* **2012**, 53, 1370–1372.
- (10) Malachowski, W. P.; Paul, T.; Phounsavath, S. The Enantioselective Synthesis of (–)-Lycoramine with the Birch–Cope Sequence. *J. Org. Chem.* **2007**, 72, 6792–6796.

(11) 8-Phenylmenthol was introduced by Corey and Ensley and used in the form of its acrylate ester in asymmetric Diels–Alder cycloadditions: (a) Corey, E. J.; Ensley, H. E. Preparation of an Optically Active Prostaglandin Intermediate via Asymmetric Induction. *J. Am. Chem. Soc.* **1975**, *97*, 6908–6909. It has been used in many other applications, as can be found in: (b) Ort, O. (–)-Phenylmenthol. *Org. Synth.* **1987**, *65*, 203–210.

(12) Donohoe, T. J.; Guyo, P. M.; Helliwell, M. The stereoselective Birch reduction of pyrroles. *Tetrahedron Lett.* **1999**, *40*, 435–438.

(13) Crossley, S. W. M.; Martinez, R. M.; Guevara-Zuluaga, S.; Shenvi, R. A. Synthesis of the Privileged 8-Arylmenthol Class by Radical Arylation of Isopulegol. *Org. Lett.* **2016**, *18*, 2620–2623.

(14) See [Supporting Information](#) for details.

(15) Curiously, only three of the four diastereomers were observed, in a 1.8:1.1:1 ratio. It is known that a substituent in the para position can have significant impacts on the stereochemistry of protonation of dienolates resulting from Birch reductions of 4-substituted benzoic acids: Rabideau, P. W.; Sygula, A. In *The Conformational Analysis of Cyclohexenes, Cyclohexadienes, and Related Hydroaromatics*. Rabideau, P. W., Ed., VCH, New Your, 1989, Chapter 3. Owing to separation issues, we could not discern which isomers were obtained and, therefore, we cannot comment further on this unusual result.

(16) Interestingly, the 3-phenyl substrate (not shown) gave a 68% yield and a moderate 4:1 d.r. when alkylated with iodomethylpivalate. Although this result does not fit within the hypothesis of needing the 2-alkoxy group to coordinate with the lithium enolate for good diastereoselectivity, it is likely that extending the π -system allows for greater π -stacking and thus favors one conformer over the other.