

UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Evidence for height and immune function trade-offs among preadolescents in a high pathogen population

Permalink

<https://escholarship.org/uc/item/32t72045>

Journal

Evolution Medicine and Public Health, 2020(1)

ISSN

2050-6201

Authors

Garcia, Angela R
Blackwell, Aaron D
Trumble, Benjamin C
[et al.](#)

Publication Date

2020

DOI

10.1093/emph/eoaa017




Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Evidence for height and immune function trade-offs among preadolescents in a high pathogen population

Angela R. Garcia ^{1,2,*}, Aaron D. Blackwell ³, Benjamin C. Trumble ^{1,4}, Jonathan Stieglitz^{5,6}, Hillard Kaplan⁷ and Michael D. Gurven⁸

¹Center for Evolution and Medicine, Arizona State University, 427 E Tyler Mall, Tempe, AZ 85281, USA; ²School of Life Sciences, Arizona State University, 427 E Tyler Mall, Tempe, AZ 85281, USA; ³Department of Anthropology, Washington State University, Pullman, WA 99163, USA; ⁴School of Human Evolution and Social Change, Arizona State University, 900 Cady Mall, Tempe, AZ 85281, USA; ⁵Université Toulouse 1 Capitole, esplanade de l'Université 21, Allée de Brienne, Toulouse Cedex 06 31080, France; ⁶Institute for Advanced Study in Toulouse, esplanade de l'Université 21, Allée de Brienne, Toulouse Cedex 06 31080, France; ⁷Economic Science Institute, Chapman University, One University Drive, Orange, CA 92866, USA and ⁸Department of Anthropology, University of California, Santa Barbara, CA 93106, USA

*Corresponding author. School of Life Sciences, Arizona State University, 427 E Tyler Mall, Tempe, AZ 85281, USA. Tel: +1-248-830-2030; E-mail: Angela.Garcia.2@asu.edu

Received 22 November 2019; Editorial decision 24 May 2020; revised version accepted 26 May 2020

ABSTRACT

Background: In an energy-limited environment, caloric investments in one characteristic should trade-off with investments in other characteristics. In high pathogen ecologies, biasing energy allocation towards immune function over growth would be predicted, given strong selective pressures against early-life mortality.

Methodology: In the present study, we use flow cytometry to examine trade-offs between adaptive immune function (T cell subsets, B cells), innate immune function (natural killer cells), adaptive to innate ratio and height-for-age z scores (HAZ) among young children ($N = 344$; aged 2 months–8 years) in the Bolivian Amazon, using maternal BMI and child weight-for-height z scores (WHZ) as proxies for energetic status.

Results: Markers of adaptive immune function negatively associate with child HAZ, a pattern most significant in preadolescents (3+ years). In children under three, maternal BMI appears to buffer immune and HAZ associations, while child energetic status (WHZ) moderates relationships in an unexpected direction: HAZ and immune associations are greater in preadolescents with higher WHZ. Children with low WHZ maintain similar levels of adaptive immune function, but are shorter compared to high WHZ peers.

Conclusions: Reduced investment in growth in favor of immunity may be necessary for survival in high pathogen contexts, even under energetic constraints. Further, genetic and environmental factors are important considerations for understanding variation in height within this population. These findings

prompt consideration of whether there may be a threshold of investment into adaptive immunity required for survival in high pathogen environments, and thus question the universal relevance of height as a marker of health.

Lay Summary: Adaptive immune function is negatively associated with child height in this high pathogen environment. Further, low weight-for-height children are shorter but maintain similar immune levels. Findings question the relevance of height as a universal health marker, given that costs and benefits of height versus immunity may be calibrated to local ecology.

KEYWORDS: life history theory; developmental plasticity; ecological immunology; Tsimane

1. BACKGROUND AND OBJECTIVES

Energy available to organisms is limited and allocated to competing physiological demands in ways that optimize biological fitness. All else equal, energetic investment in one characteristic should trade-off with investment in other characteristics [1–3]. Natural selection can produce adaptations for managing trade-offs in species-typical ways, but it can also select for norms of reaction that manage allocations to fit local ecological contexts through developmental plasticity, allowing individuals to better approach the ‘optimal’ energy allocation strategy for a particular environment [4]. For example, given the strong selective pressure against infant mortality, early investment in immune function should be prioritized despite consequences for competing energetic requirements, like growth. This may be particularly salient in a high pathogen environment, where investment in the adaptive arm of the immune system, which confers long-lasting specific immunity, would be critical for surviving this early period [5].

Growth and immune trade-offs have been studied in humans and across vertebrate taxa [6]. In humans, numerous studies have shown that chronic immune activation [7, 8], and overall pathogen burden are associated with growth stunting [9, 10]. At young ages, while there may be early developmental origins of growth–immune associations, these relationships are often complex, and may be convoluted by changes occurring across multiple landscapes (e.g. maturational tempo, pathogen exposure, physical activity and diet) within a relatively narrow window of time. For example, in studies of infants in the Gambia [11] and Northern Kenya [12], researchers found that evidence of immune activation, as measured by (1) intestinal permeability and mucosal enteropathy and (2) immunoglobulin A (salivary), respectively, showed impaired growth, and that levels of immunoglobulins A, G and M rapidly increased with age [11]. Panter-Brick and colleagues similarly documented an age-related increase in IgG levels in a group of Nepali infants ($n = 86$), which varied by socioeconomic status, and was associated with lower weight-for-age [13]. However, the authors also found that mucosal damage, which was associated with impaired growth in height, was most severe among middle-class children, leading to the conclusion that nutritional constraints may mask or confound relationships between immune function and linear growth.

To date, many of the relevant studies have utilized either generalized markers of inflammation, or measures of immunoglobulins, and none that we are aware of include direct measurements of the leukocytes responsible for long-term adaptive immune function, including naïve and memory T cells and B cells. Numerous questions remain about how different aspects of immune function and activation relate to growth, whether these represent short- or long-term accommodations, and how factors like how nutritional environment, or energy availability, buffer or exacerbate these effects. Growth–immune trade-offs might represent short-term accommodations to temporary resource shortages, or longer-term phenotypic adjustments to expected future conditions [14]. In the short term, in addition to the direct costs of immune activation, anorexia associated with sickness behavior or gastrointestinal distress may cause sick children to eat less and absorb less nutrients [15], reducing energy available for growth. Particularly when intake is limited, fat stores may represent critical energy reserves upon which the body draws to fuel metabolic processes associated with growth and immune responses to infectious disease [16, 17]. Having larger energy intake or energy stores may allow for greater investment across multiple traits simultaneously. As a consequence, at the individual level there may be positive covariation among life history traits in individuals with ample resources [18]. In contrast to these short-term accommodations, long-term adjustments in expectation of future trade-offs may play out quite differently. For example, if short-term immune activation serves as a cue to future energy needs, we might expect individuals to shift energy into immune function and away from height, even in the absence of immediate energy shortages. We might also expect that short-term immune activation could lead to increases in energy storage, and therefore future fat accumulation [10].

One way to address the question of short- versus long-term accommodations and trade-offs is to examine multiple aspects of immunity that represent both short-term responses and long-term investments. Previous studies have found that different immune markers associate with growth on different time-scales. For example, Urlacher *et al.* [10] found that a marker of acute inflammation associated with growth over one week, and was mediated by fat reserves, while immunoglobulin levels were associated with growth over months, and with overall height-for-age z scores (HAZ).

Here, we take a similar approach in examining both innate and adaptive markers. However, in contrast to past studies, we use flow cytometry to directly characterize leukocyte populations, the cells responsible for antibody production and cellular responses. This characterization includes both cells indicative of past exposures (non-naïve cells) and cells that represent immunological resources which have not yet been activated (naïve cells). In this article, we: (i) measure multiple immune markers in peripheral leukocytes (i.e. CD4+/CD8+, naïve and non-naïve and total T cells, B cells and natural killer (NK) cells) that characterize specific components of both adaptive and innate immune function and (ii) examine whether maternal or child energetic status, as measured through weight-for-height (WHZ) (child) and body-mass-index (mother), influence potential trade-offs between the child's immune function and height (HAZ).

1.1 Measuring immune function in relation to growth-immune trade-offs

The human immune system is composed of innate and adaptive arms, which differ in their time-variant costs, time to reactivity and precision [19]. The arms of the immune system operate in an integrated way and in coordination with other organ systems (e.g. endocrine), with adaptive immune function in large part serving as an extension of, and directed by, the innate arm of the immune system [20, 21]. Innate immune functions are the body's first-line non-specific defense mechanisms which are activated immediately upon encountering a foreign pathogen. Because they are generally applicable and immediately available, in many senses the costs of innate immunity may be seen as low, but innate responses also ultimately cause more collateral damage to surrounding cells and tissue, leading to faster rates of senescence [22–24]. By comparison, slower-acting adaptive immune responses require higher initial investment into the physiological mechanisms and systems that support them, including the production, selection and maintenance of pools of diverse B- and T-cell variants, many of which may never encounter a matching antigen and thus never be used [25, 26]. However, responses of the adaptive immune system are highly specialized, and usually cause little collateral damage. The adaptive immune system is also increasingly being recognized for its importance in immunological repair of cells and tissues [27]. The thymus, a major organ of the adaptive immune system, is the primary site of T cell development and maturation [28, 29]. That functional thymic volume peaks within the first year after birth, suggests this early period is a critical window for investment in adaptive immunity [28].

The innate immune response is highly sensitized, and can be hyper-variable across days due to even minor fluctuations in environmental stimuli [30]. This sensitization makes it hard to

distinguish acute innate responses to short-term ecological stressors from longer-term calibration of the innate immune system to local ecological context. The relatively slower and less-generalized responses of the adaptive immune system suggest that, in cross-sectional studies, measures of adaptive immune function or the relative balance of adaptive to innate immune function may more reliably reflect the way the immune system is calibrated, and may thus provide a more accurate metric against which traits like growth may trade-off against.



Box 1. Components of the innate and adaptive immune system examined in this study

Immune arm	Main activity	Main target
Innate immunity		
NK cells	Kill missing-self or antigen-presenting cells	Viral; cell-mediated
Adaptive immunity		
T cells		
CD4+ (helper T)	Secrete cytokines and function to stimulate differentiation and proliferation of numerous immune cells	Aid in both cell-mediated and humoral processes
CD8+ (cytotoxic T)	Kill antigen-presenting cells	Tumor and virally-infected; cell-mediated
Naïve cells	Un-activated CD4+ or CD8+; have not yet encountered cognate antigen	Dependent on cell type
Non-naïve cells	Activated CD4+ or CD8+; antigen-specific; long-lived	Dependent on cell type
B cells	Antibody production	Humoral

1.2 Hypothesis and predictions

Immuno-development early in life is in competition with growth, given that both require substantial energetic investment. We hypothesize that in high pathogen environments, where the cost of inadequate immune function is likely higher than smaller stature, we should see trade-offs between these life history features, and these trade-offs may be buffered by adequate resources. Specifically, we hypothesize that markers of adaptive immune function will trade-off against height, visible

in this window of somatic and immunologic development, and that this trade-off will be moderated by energy availability. Data from the Tsimane, an energy-limited subsistence population in lowland Bolivia facing high pathogen loads [31], will be used to test the predictions that (P1) adaptive immune cell counts will be inversely associated with child HAZ. Furthermore, using the relative balance of investment into adaptive cellular immunity (i.e. B and T cell counts) to innate cellular immunity (i.e. NK cell counts) as a proxy for longer-term investment in costlier immune responses, we predict that (P2) a higher ratio of adaptive to innate immunity will associate with shorter stature. Finally, we predict that (P3) energetic status (proxied by child WHZ and maternal body mass index) will moderate associations between height and immune function, such that children with higher energetic status will be buffered and show a weaker trade-off (i.e. show weaker associations) between adaptive immune markers and height.

2. METHODOLOGY

2.1 Study population and data collection

2.1.1 The Tsimane of Bolivia. The Tsimane are forager-horticulturalists (population ~16 000) that live in the Beni Department of Bolivia, in the lowlands comprising the Amazon basin. Tsimane inhabit over 90 villages, most of which do not have access to electricity, running water or comprehensive waste management [32, 33]. Their diet is relatively lean, as they subsist mainly on plantains, rice, sweet manioc and corn from slash-and-burn horticulture, fish and wild game, occasionally supplemented by market goods such as refined sugar, salt and cooking oil [32, 34]. Despite evidence of child and adult stunting, wasting is uncommon; mean adult BMI is 23.6 for both sexes [31, 35]. From an early age, children are exposed to an array of pathogens, and parasitic infections among the Tsimane are common [36]. About half of Tsimane men and women have anemia, with children and adolescents showing the highest risk (56% for girls, 63% for boys) [37].

Given the high transmission rate of multiple pathogens in the Tsimane environment, investment in immune defenses is also high, with children exhibiting a concomitant early peak in humoral response (e.g. immunoglobulin-E, IgE) associated with helminthic infection compared to populations with lower transmission rates [36]. In general, Tsimane have higher levels of numerous immune components, including white blood cells (WBCs), erythrocyte sedimentation rate, B cells and NK cells than do Americans at all ages [31]. On average, 16% of WBCs are eosinophils, consistent with high levels of parasitic infection, as compared with a US reference range <5%, with approximately 85.9% of the population in eosinophilia by US standards [38]. Antibodies related to infection are also high among

Tsimane: immunoglobulin-G (IgG) levels are about twice as high, and IgE is about 100 times higher than mean US levels [36].

2.1.2 participants. Data for this study were collected as part of the Tsimane Health and Life History Project (THLHP), a longitudinal study of health and aging that began in 2002 [32]. The sample includes 344 children aged 2 months–8 years (55% female, median age 5 years) from 28 villages (see Table 1 for descriptive statistics). Participants were seen in the stationary THLHP clinic in San Borja (the closest market town to sampled villages), accompanied by a parent between mid-March and early July of 2011, whereupon participants were seen by THLHP physicians and given routine physical exams, which included patient history, symptom investigation, demographic and anthropometric information. A blood sample was taken as a part of this visit, a portion of which was used for assessing immune function.

The study was approved by the Tsimane Government (Gran Consejo Tsimane) and village leaders. Study protocols were approved by the Tsimane Government and IRBs at the



Table 1. Descriptive statistics

Variables	Sample (N = 344)	
	Mean (SD)	Range
Sex (% female)	55	
Age (years)	4.71 (2.2)	0.15–8.07
Child's BMI (kg/m ²)	16.18 (1.4)	11.19–23.11
Mother's height (cm)	150.62 (4.2)	140.7–165.0
Mother's BMI (kg/m ²)	25.00 (3.5)	17.89–39.16
Respiratory infection (% infected)	47.8	
Parasite analyses (n = 246)		
Helminthic parasites (% infected)	56.6	
Protozoal parasites (% infected)	29.2	
Leukocytes (×10 ³ per μl)	11.98 (4.3)	4.30–33.60
T cell count (×10 ³ per μl)	3.07 (1.5)	0.74–11.07
Naïve T cell count (×10 ³ per μl)	2.27 (1.3)	0.38–8.51
CD4+ cell count (×10 ³ per μl)	1.85 (1.0)	0.40–6.17
CD8+ cell count (×10 ³ per μl)	1.22 (0.7)	0.10–5.61
B cell count (×10 ³ per μl)	1.28 (0.8)	0.22–6.91
NK cell count (×10 ³ per μl)	0.42 (0.3)	0.06–2.50

University of California, Santa Barbara (#12-496) and the University of New Mexico (#07-157). Adult participants provided informed consent for their own and their children's participation. Children age six and older also provided verbal assent.

2.1.3 Anthropometric and biomarker collection. Infant and child height (or length) were measured by trained Tsimane research assistants using a portable Seca infantometer or stadiometer (Hamburg, Germany). For the purposes of analysis, length and height were treated interchangeably. Weight was measured using a Tanita digital scale to the closest tenth of a kilogram. If infants were too small to stand on the scale, their weight was calculated by subtracting the weight of the mother from the combined weight of the mother holding her infant. Maternal height and weight measures were collected using the same instruments as used for the child, either at the same visit, or within 30 days of the child's visit. Tsimane-specific z-scores for HAZ, child WHZ and maternal BMI-for-age (BMIZ) were calculated using the open-source R package *localGrowth* (<https://github.com/adblackwell/localgrowth>), which utilizes growth curves from [39]. These growth references better describe intra-population comparison by using locally-relevant growth trajectories [39]. Child WHZ and maternal BMIZ (in models for children under 3 years) were used as markers of energetic status, to model the effects of interactions between child energetic status and immune function on child HAZ. To determine the child's age, exact birthdates were verified by the mother or pulled from longitudinal THLHP medical records.

2.1.3.1 Biomarker collection

Blood was collected by a certified Bolivian biochemist into a heparin-coated vacutainer by venipuncture for children 2 years and older, and by capillary heel or finger prick for infants under 2 years of age [31]. Total WBC and hemoglobin were measured using a QBC Autoread Plus dry hematology system (QBC Diagnostics) directly after blood draw. Flow cytometry was performed in the THLHP clinic in San Borja on fresh heparinized blood within six hours of the blood draw in order to quantify lymphocyte subsets into helper T cell (CD4+CD8-), cytotoxic T cells (CD8+CD4-), NK cells (CD56+CD8-CD4-) and B cells (CD19+) using an Accuri C6 Flow Cytometer (BC Accuri Cytometers). T cells were further classified as naïve (CD45RA+) or non-naïve (CD45RA-). Absolute counts were calculated by multiplying the relative percentages determined by flow cytometer by the total lymphocyte count obtained from the QBC Autoread Plus. For full description of gating and antibodies used [31].

2.1.3.2 Parasite analyses

Fecal samples were analyzed for the presence of helminth eggs and larvae by direct identification on wet mounts or by a

modified Percoll method [9, 40]. Parasite infection was recorded as presence or absence of each species, and is based on a single fecal sample, collected and analyzed at the health clinic within one day of blood collection.

2.2 Statistical analysis

Mixed effects models fit with restricted (residual) maximum likelihood (REML), from the R package *lme4*, were used to evaluate relationships between markers of immune function and child HAZ. Maternal ID was included as a random effect, as a means to parse contribution of potential genetic clustering or clustering due to immediate environment (e.g. shared household) from individual-level factors that contribute to overall variation in HAZ [41]. Effects of community-level clustering were also considered, and initial models included community ID as a random effect. However, since the inclusion of community ID did not substantially improve fit for any model, it was not included in final analyses. Height z-scores for mothers are also included in models, for additional adjustment for potential heritable variation in height. To adjust for variation in immune function based on current infection and associated immune activation, we include covariates for respiratory illness and residual WBC counts. Additional models were ran using the subset of the sample for whom fecal parasite data were available (72% of sample), which include covariates for current protozoal and helminthic infections assessed from fecal samples. For each model, residual WBC count is WBC count minus the portion of cells that were being used as the main independent variable of interest (e.g. in a model testing the association between total T cell count and height, residual WBC = total WBC – T cell count). This adjustment is necessary to minimize collinearity. To measure relative investment in adaptive relative to innate immunity, we calculate a ratio of adaptive (B and T) cells to innate (NK) cells. Due to skewed distributions, all immune cell counts are log-transformed and standardized (z-scored) before inclusion in regression models. All analyses were performed using R statistical software version 3.4.2 [43]. All models adjust for age and sex. All results are reported as standardized betas unless otherwise noted. Both marginal and conditional R-squared values are included in tables for linear mixed effects model results. While marginal R-squared reflects the variance explained only by model fixed effects, the conditional R-squared is interpreted to reflect variance explained by both fixed and random effects (see Ref. [42] for discussion of limitations on the use of these R-squared estimates in mixed effects models).

3. RESULTS

Children in this study ranged from 2 months to 8 years of age (mean: 4.7 years, SD: 2.2) and 55% were female. Overall, 72% of participants were screened for parasites. Of this sample, 57% of

children were infected with at least one species of parasite, with 25.3% having coinfections with more than one species. The most common infections were helminths, with hookworm (*Ancylostoma duodenale* or *Necator americanus*, prevalence 50.6%) being most common, followed by roundworm (*Ascaris lumbricoides*, 16.1%), threadworm (*Strongyloides stercoralis*, 7.7%) and whipworm (*Trichuris* sp., 2.3%). Protozoan infections were also common, including *Giardia lamblia* (24.1%) and *Entamoeba histolytica* (8.4%); further, 47.8% of children were diagnosed as having a current respiratory condition. Using WHO criteria for age-specific anemia cutoffs, 45.7% of children also had at least mild anemia (<11 mg/dl of hemoglobin in children under age 5, and <11.5 mg/dl for children aged 5–11 years).

3.1 Age-related changes in lymphocyte subsets

We first examined overall differences in lymphocyte subsets by age, adjusting for sex and current infections, fit with smoothed loess curves based on generalized linear models (Fig. 1). Total T cells ($\beta = -294.89$ cells/ μL of blood/year, $P < 0.001$) and all subsets (naïve: $\beta = -264.94$, $P < 0.001$; non-naïve: $\beta = -29.96$, $P < 0.01$; CD4^+ : $\beta = -191.73$, $P < 0.001$; CD8^+ : $\beta = -103.19$, $P < 0.001$) and B cells ($\beta = -145.85$, $P < 0.001$) significantly

decline with age. Conversely, NK cells increase with age ($\beta = 30.85$, $P < 0.001$).

3.2 Associations between adaptive immune function and child height

Mixed effects linear regression models are used to evaluate relationships between child HAZ and markers of immune function, with a random effect for mother's ID (Fig. 2 and Table 2). HAZ is consistently negatively associated with adaptive immune cell counts after adjusting for age, sex, mother's height, residual WBC and current respiratory illness. Higher HAZ children have lower overall T cells ($\beta = -0.37$, $P = 0.001$), including both naïve ($\beta = -0.15$, $P = 0.009$) and non-naïve ($\beta = -0.16$, $P = 0.001$) T cell counts, CD4^+ ($\beta = -0.33$, $P = 0.002$) and CD8^+ ($\beta = -0.27$, $P = 0.006$) subsets (Fig. 2: Models 1–5) and lower B cell counts ($\beta = -0.25$, $P = 0.03$; Model 6). There is no association between innate (i.e. NK) cell counts and height ($\beta = -0.05$, $P = 0.48$; Model 7). Further, a higher ratio of adaptive to innate immune cells is not associated with child height ($\beta = -0.05$, $P = 0.35$) (Fig. 2: Model 8). Including maternal ID significantly increased model fit across all models (Table 2), suggesting that genetic and/or shared environmental factors influence variation in child HAZ.




Figure 1. (a–g) Total T cells, including naïve and non-naïve, CD4 and CD8 subsets and B cells, significantly decline with age, while NK cells do not. The trend-lines are loess fit (span = 0.75)




Figure 2. Trade-offs between child height and adaptive immunity. This figure compares parameter estimates of associations between measures of adaptive immunity and child height across all adaptive immune models from Table 2. All immune markers were logged and standardized before analysis; A:I ratio is the ratio of adaptive to innate immune cells, logged and standardized for analysis. Analyses include controls for age, mother's height, sex and WBC (logged and z-scored). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

3.3 Height and immune associations by age group

Given children younger than 3 years are likely still supplemented both immunologically and energetically by their mothers through breastfeeding, the next set of analyses examines associations between HAZ and immune cell counts including an interaction with age group. For these analyses, children are split into two groups: infants (children 2 months to 3 years; $n = 82$) and preadolescents (children 3–8 years; $n = 261$). There are no significant associations between HAZ and immune cells across cell types in infants (Table 3). In contrast, for preadolescents, HAZ and total T cells ($\beta = -0.49$, $P < 0.001$), both naïve ($\beta = -0.20$, $P = 0.002$) and non-naïve ($\beta = -0.20$, $P < 0.001$) T cell counts, CD4+ ($\beta = -0.47$, $P < 0.001$) and CD8+ ($\beta = -0.32$, $P = 0.004$) T cell subsets, and B cells ($\beta = -0.34$, $P = 0.006$), but not NK cells ($\beta = -0.03$, $P = 0.73$) or A:I ratio ($\beta = -0.10$, $P = 0.10$) (Table 3). Because children traverse through numerous developmental phases even between 3 and 8 years of age, we ran additional models which included a finer-grained age-category variable, splitting children into approximately equally weighted age groups (2 months to 3 years, $n = 82$; 3–5 years, $n = 91$; 5–6.5 years, $n = 80$; 6.5–8 years, $n = 90$), and found that

these associations were strongest among the two middle groups (3–5 and 5–6.5 years) (Supplementary Table S1).

3.4 maternal and child energetic status as a moderator of immune-height trade-offs

To test the prediction that having adequate energetic reserves may buffer a trade-off between immune function and growth in the child, we examine the effects of a three-way interaction between each immune marker, child or maternal energetic status and age group on child HAZ. In children under 3 years old, maternal energetic status (maternal BMIZ) moderates the relationship between total T ($\beta = 0.31$, $P = 0.08$) and CD4+ T cell subsets ($\beta = 0.33$, $P = 0.05$), and the A:I ratio ($\beta = 0.19$, $P = 0.04$) and height, but does not show any influence on immune cell counts and height relationships in preadolescents (Supplementary Table S2). In other words, for infants whose mothers have above the mean BMI, there is a positive association between cell count and HAZ, whereas for older children (preadolescents), there remains a negative association between cell count and HAZ, regardless of maternal BMI (Supplementary Fig. S1). Alternately, models with child




Figure 3. Graphical display of results from Table 4, model 1. For children 3 years and older, there is a negative interactive effect of child WHZ on the association between total T cell count and height. There appears to be a height-immune trade-off only in children that are at or above the mean for WHZ. Note: child height and WHZ are z-scored, and total T cell count is logged and z-scored for analyses. Model controls for age, sex, mother's height and current immune activation, and a random effect for maternal ID

are likely due to overall investment in cellular immunity and are not necessarily dependent on the causes of that investment (i.e. responsive vs. predictive). These associations suggest that it is the immune investment itself may trade-off with height, rather than direct costs of past infections.

Though in the full sample there are negative associations between adaptive immune markers and HAZ, when children are split into age-groups: infants (< 3 years old) and preadolescents (3-8 years old), cell counts and HAZ associations we find that are not statistically significant in the younger age group. The lack of an association between HAZ and immune function in infants may be due to a few statistical or biological facets. Statistically, because coefficients were consistently negative (as in the older group) after adjusting for current immune activation, the lower sample size and higher error in this group may have obscured any statistical significance in associations from emerging. Biologically, it may be that infants maintain adequate buffering by maternal energetic and immunologic resources, transmitted via breastfeeding, through this time period. In accordance with this idea, we do find a positive interaction between mother's energetic status and associations between immune cell counts and HAZ among infants whereby children with higher-BMI moms show a positive association between height and immune cell counts. There is a rich literature

showing growth faltering after weaning, which may be a consequence of increased exposure to contaminants and other environmental pathogens [45–47]. In fact, we do see that preadolescents are more likely to have helminthic parasites compared to infants, and, that protozoal infection is positively associated with greater WHZ, which in infants may be evidence of earlier weaning. Though we do not have systematic quantitative data on breastfeeding status and intensity for the present sample, a different Tsimane study found that children breastfeed until between 30-36 months of age, in concert with the introduction of complementary foods [48].

However, another possibility is that the lack of associations could be due to heterogeneity in maturational pace among individuals in this age group. For example, children who are maturationally delayed overall, would be smaller and likely have higher naïve immune cell counts. Growth is not continuous, and it is therefore difficult to confidently assess the impact of current immune profiles on growth using cross-sectional measures. It is likely that there are coordinated developmental changes across these various systems (height, weight, immune status) that vary by both chronological and maturational age which obscure our ability to tease apart trade-offs from shifts in coordination. In these data, we found that in fact the largest effect sizes for associations between immune cells and HAZ was

26. Medzhitov R. Recognition of microorganisms and activation of the immune response. *Nature* 2007;**449**:819–26.
27. Karin M, Clevers H. Reparative inflammation takes charge of tissue regeneration. *Nature* 2016;**529**:307–15.
28. George AJ, Ritter MA. Thymic involution with ageing: obsolescence or good housekeeping? *Immunol Today* 1996;**17**:267–72.
29. Thiault N, Darrigues J, Adoue V et al. Peripheral regulatory T lymphocytes recirculating to the thymus suppress the development of their precursors. *Nat Immunol* 2015;**16**:628–34.
30. Mangino M, Roederer M, Beddall MH et al. Innate and adaptive immune traits are differentially affected by genetic and environmental factors. *Nat Commun* 2017;**8**:13850.
31. Blackwell AD, Trumble BC, Maldonado Suarez I et al. Immune function in Amazonian horticulturalists. *Ann Hum Biol* 2016;**43**:382–96.
32. Gurven M, Stieglitz J, Trumble B et al. The Tsimane health and life history project: integrating anthropology and biomedicine. *Evol Anthropol* 2017;**26**:54–73.
33. Dinkel, K.A., Costa, M.E., Kraft, T.S., Stieglitz, J., Cummings, D.K., Gurven, M., Kaplan, H., Trumble, B.C., 2020. Relationship of sanitation, water boiling, and mosquito nets to health biomarkers in a rural subsistence population. *Am. J. Hum. Biol.* 32, e23356. 10.1002/ajhb.23356
34. Kraft TS, Stieglitz J, Trumble BC et al. Nutrition transition in 2 lowland Bolivian subsistence populations. *Am J Clin Nutr* 2018;**108**:1183–95.
35. Gurven M. Infant and fetal mortality among a high fertility and mortality population in the Bolivian Amazon. *Soc Sci Med* 2012;**75**: 2493–502.
36. Blackwell AD, Gurven MD, Sugiyama LS et al. Evidence for a peak shift in a humoral response to helminths: age profiles of IgE in the Shuar of Ecuador, the Tsimane of Bolivia, and the US NHANES. *PLoS Negl Trop Dis* 2011;**5**:e1218.
37. Anderson AS, Trumble BC, Hové C et al. Old friends and friendly fire: pregnancy, hookworm infection, and anemia among tropical horticulturalists. *Am J Hum Biol* 2019; 32: e23337.
38. Trumble BC, Stieglitz J, Blackwell AD et al. Apolipoprotein E4 is associated with improved cognitive function in Amazonian forager-horticulturalists with a high parasite burden. *FASEB J* 2017;**31**:1508–15.
39. Blackwell AD, Urlacher SS, Beheim B et al. Growth references for Tsimane Forager-horticulturalists of the Bolivian Amazon. *Am J Phys Anthropol* 2017;**162**:441–61.
40. Eberl M, al-Sherbiny M, Hagan P et al. A novel and sensitive method to monitor helminth infections by faecal sampling. *Acta Trop* 2002;**83**: 183–7.
41. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models Using lme4. *J Stat Softw* 2015;**67**. doi: 10.18637/jss.v067.i01.
42. Nakagawa S, Schielzeth H. A general and simple method for obtaining R² from generalized linear mixed-effects models. *Methods Ecol Evol* 2013;**4**:133–42.
43. Team R. Core. R: *A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing, 2017.
44. Soler JJ, Neve LD, Pérez-Contreras T et al. Trade-off between immunocompetence and growth in Magpies: an experimental study. *Proc R Soc Lond Ser B Biol Sci* 2003;**270**:241–8.
45. Bacha K, Mchari T, Ashenafi M. The microbial dynamics of 'borde' fermentation, a traditional Ethiopian fermented beverage. *SINET* 1998;**2**: 195–205. doi:10.4314/sinet.v2i1i2.18120.
46. Motarjemi Y, Käferstein F. Food safety, hazard analysis and critical control point and the increase in foodborne diseases: a paradox? *Food Control* 1999;**10**:325–33.
47. Padmadas SS, Hutter I, Willekens F. Weaning initiation patterns and subsequent linear growth progression among children aged 2–4 years in India. *Int J Epidemiol* 2002;**31**:855–63.
48. Martin MA, Garcia G, Kaplan HS, Gurven MD. Conflict or congruence? Maternal and infant-centric factors associated with shorter exclusive breastfeeding durations among the Tsimane. *Soc Sci Med* 2016; 170: 9-17. doi:10.1016/j.socscimed.2016.10.003.
49. Janeway CA Jr, Travers P, Walport M, Shlomchik MJ. 2001. *The Complement System and Innate Immunity*. New York, USA: Garland Science.
50. Manczinger M, Boross G, Kemény L et al. Pathogen diversity drives the evolution of generalist MHC-II alleles in human populations. *PLoS Biol* 2019;**17**:e3000131.
51. Dopico XC, Evangelou M, Ferreira RC et al. Widespread seasonal gene expression reveals annual differences in human immunity and physiology. *Nat Commun* 2015;**6**:7000.
52. Schmid-Hempel P. Variation in immune defence as a question of evolutionary ecology. *Proc Biol Sci R Soc* 2003;**270**:357–66.
53. Gurven M, Kaplan H, Winking J et al. Inflammation and Infection Do Not Promote Arterial Aging and Cardiovascular Disease Risk Factors among Lean Horticulturalists. *PLoS ONE* 2009; **4**:e6590 10.1371/journal.pone.0006590
54. Sear R, Mace R, McGregor IA. The effects of Kin on female fertility in rural Gambia. *Evol Hum Behav* 2003;**24**:25–42.
55. Moore SR, Lima AA, Conaway MR et al. Early childhood diarrhoea and helminthiasis associate with long-term linear growth faltering. *Int J Epidemiol* 2001;**30**:1457–64.
56. Motarjemi Y, Käferstein F, Moy G, Quevedo F. Contaminated weaning food: a major risk factor for diarrhoea and associated malnutrition. *Bull World Health Organ* 1993;**71**:79–92.
57. Wander K, Brindle E, O'connor KA. Sensitivity and specificity of C-reactive protein and α 1-acid glycoprotein for episodes of acute infection among children in Kilimanjaro, Tanzania. *Am J Hum Biol* 2012;**24**: 565–8.
58. West-Eberhard MJ. 2003. *Developmental Plasticity and Evolution*. Oxford University Press.