

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### Title

Inhibition of TGFbeta1 Signaling Attenuates ATM Activity in Response to Genotoxic Stress

### Permalink

<https://escholarship.org/uc/item/3476t6kx>

### Authors

Kirshner, Julia  
Jobling, Michael F.  
Pajares, Maria Jose  
et al.

### Publication Date

2006-09-15

Peer reviewed

## **Inhibition of TGF $\beta$ 1 Signaling Attenuates ATM Activity in Response to Genotoxic Stress**

Julia Kirshner.\*<sup>1</sup>, Michael F. Jobling,\*<sup>1</sup>, Maria Jose Pajares<sup>1</sup>, Shraddha A. Ravani<sup>1</sup>, Adam B. Glick<sup>2</sup>, Martin J. Lavin<sup>3</sup>, Sergei Koslov<sup>3</sup>, Yosef Shiloh<sup>4</sup> & Mary Helen Barcellos-Hoff<sup>1,5</sup>

\*Authors contributed equally.

<sup>1</sup> Life Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720

<sup>2</sup> Laboratory of Cellular Carcinogenesis and Tumor Promotion National Cancer Institute, Bethesda, MD 20892

<sup>3</sup> The Queensland Institute of Medical Research, Royal Brisbane Hospital, Herston, Queensland 4029, Australia.

<sup>4</sup> Department of Human Genetics, Sackler School of Medicine, Tel Aviv University, Ramat Aviv 69978 Israel

<sup>5</sup> Corresponding Author: Mary Helen Barcellos-Hoff  
Life Sciences Division  
1 Cyclotron Road, Bldg. 977  
Berkeley, CA 94720  
(510) 495-2535 FAX  
(510) 486-6371 PHONE  
[mhbarcellos-hoff@lbl.gov](mailto:mhbarcellos-hoff@lbl.gov)

Financial Support: Support was provided by NASA program Biomedical Research and Countermeasures Ground Research in Radiation Health to MHBH, grant number T6275-W, DOD DAMD17-01-0291 postdoctoral fellowship to MFJ, and the Low Dose Radiation Program of the DOE Office of Biological Effects Research to MHBH and the Office of Health and Environmental Research, Health Effects Division, United States Department of Energy (contract no.03-76SF00098).

Total Word Count: 6945

Abstract: 202

Text:4062

Figures: 6

Keywords: DNA damage; epithelial; cytokine

Abbreviations: Transforming growth factor  $\beta$ 1, TGF $\beta$ ; Ataxia telangiectasia mutated, ATM; ionizing radiation, IR; mammary epithelial cells, MEC; TGF $\beta$  receptor type I, T $\beta$ RI; radiation-induced foci, RIF

Running Title: TGF $\beta$  Mediates DNA Damage Responses

**ABSTRACT**

Ionizing radiation causes DNA damage that elicits a cellular program of damage control coordinated by the kinase activity of ataxia telangiectasia mutated protein (ATM). Transforming growth factor  $\beta$ 1 (TGF $\beta$ ), which is activated by radiation, is a potent and pleiotropic mediator of physiological and pathological processes. Here we show that TGF $\beta$  inhibition impedes the canonical cellular DNA damage stress response. Irradiated *Tgf $\beta$ 1* null murine epithelial cells or human epithelial cells treated with a small molecule inhibitor of TGF $\beta$  type I receptor kinase exhibit decreased phosphorylation of Chk2, Rad17 and p53, reduced  $\gamma$ H2AX radiation-induced foci, and increased radiosensitivity compared to TGF $\beta$  competent cells. We determined that loss of TGF $\beta$  signaling in epithelial cells truncated ATM autophosphorylation and significantly reduced its kinase activity, without affecting protein abundance. Addition of TGF $\beta$  restored functional ATM and downstream DNA damage responses. These data reveal a heretofore undetected critical link between the microenvironment and ATM that directs epithelial cell stress responses, cell fate and tissue integrity. Thus, TGF $\beta$ 1, in addition to its role in homeostatic growth control, plays a complex role in regulating responses to genotoxic stress, the failure of which would contribute to the development of cancer; conversely, inhibiting TGF $\beta$  may be used to advantage in cancer therapy.

## Introduction

An orchestrated response to DNA damage in multicellular organisms is important for rapid restoration of homeostasis and long-term prevention of cancer but how signaling is regulated across tissues is unknown. Little is known about the influence of extracellular signaling from growth factors on the cellular response to DNA damage, however, transforming growth factor  $\beta$ 1 (TGF $\beta$ ) is rapidly and persistently activated in tissues following IR (1). TGF $\beta$  coordinates responses to a great variety of other stimuli by regulating cell proliferation, differentiation, and apoptosis (reviewed in (2, 3)), is involved in many aspects of development, growth regulation and is known as a classical tumor suppressor (reviewed in (3)). We previously showed that epithelial tissues of *Tgf $\beta$ 1* null embryos fail to undergo either apoptosis or inhibition of cell cycle in response to IR, suggesting a surprising requirement for its activity in the responses to DNA damage (4). Neither the point at which TGF $\beta$ 1 impacts the genotoxic stress program or specific mechanisms of action have been identified.

Independent of its control of proliferation and differentiation, studies by Glick and others have implicated TGF $\beta$  in maintenance of genomic stability. *Tgf $\beta$ 1* null cells are genomically unstable (5), cannot repair alkylating damage (6) and fail to apoptose or undergo cell cycle inhibition after ionizing radiation (IR) exposure in vivo (4). *Tgf $\beta$ 1* null keratinocytes exhibit increased frequency of gene amplification as marked by N-phosphonoacetyl-L-aspartate (PALA) resistance (5). *Tgf $\beta$ 1* null keratinocytes transduced with v-ras<sup>Ha</sup> develop aneuploidy at higher frequencies, have more chromosomal abnormalities than the wild-type controls and undergo spontaneous malignant transformation more frequently and with shorter latency than wild-type counterparts (7-9). The fact that TGF $\beta$  treatment of *Tgf $\beta$ 1* null cells inhibits PALA resistance, reduces the percentage of

aneuploid metaphases, and decreases the number of spontaneous chromosome breaks indicates an ongoing process rather than selection of genomically unstable subpopulation. TGF $\beta$  is rapidly and persistently activated *in vivo* by ionizing radiation (IR) (10-14). Radiation- induced apoptosis and cell cycle arrest in epithelial cells *in vivo* are decreased in a *Tgf $\beta$ 1* gene dosage dependent fashion (4). Similarly, the mammary epithelium of *Tgf $\beta$ 1* heterozygous mice or animals treated with TGF $\beta$  neutralizing antibodies fails to undergo an apoptotic response and exhibits diminished phosphorylation of p53 in response to IR. Although these data suggest that TGF $\beta$  plays a direct role in the DNA damage response, the mechanism by which TGF $\beta$  signaling impacts the DNA damage response program has not been identified.

We postulated that TGF $\beta$  provides a microenvironment signal to ensure coordinated epithelial fate decisions and restoration of homeostasis. The primary transducer of genotoxic stress caused by IR is the nuclear protein kinase ataxia telangiectasia mutated (ATM) (15-18). ATM is a phosphoinositide 3-kinase related serine/threonine kinase that mediates DNA damage responses to initiate, recruit and activate a complex program of checkpoints for cell cycle, apoptosis and genomic integrity (see reviews (19, 20)). Mutations in human ATM lead to ataxia-telangiectasia, which is characterized by genomic instability, cellular radiation sensitivity and increased cancer. ATM is activated in response to double strand breaks caused by ionizing radiation and in turn, phosphorylates numerous substrates, thereby modulating cell fate decisions. We show here both TGF $\beta$ 1 depletion by genetic knockout in mouse cells and inhibition of TGF $\beta$  signaling in human cells compromise ATM kinase activity and autophosphorylation, leading to reduced phosphorylation of critical DNA damage transducers, abrogation of the cell cycle block and increased radiosensitivity. Linking ATM

to TGF $\beta$  ensures that cell fate decisions are functionally connected to tissue damage, which is a novel mechanism for maintaining homeostasis, but failure of this control would greatly accelerate neoplastic potential.

## Materials and methods

**Cell culture.** C57bl/129SV or Balb/c *Tgfb1*  $+/+$  and  $+/-$  primary mammary MECs were cultured in serum-free medium as described (21). Serum was removed 24 hr after culture initiation. Balb/c *Tgfb1* wildtype and null primary keratinocytes cultured from neonates gave rise to two independently derived, spontaneously immortalized keratinocyte cell lines of each genotype (5). These cell lines, H01 and H04 from *Tgfb1* heterozygote and K01 and K03 from *Tgfb1* null cultures, were used in these experiments and showed similar responses. Cells were plated in calcium-free EMEM medium containing 8% chelexed FBS and 0.2mM Ca $^{2+}$ , then passaged in serum with 0.05 mM Ca $^{2+}$ , which was changed every third day until confluence. Twelve to eighteen hours prior to irradiation, the medium was replaced with 8% serum-replacement medium (KnockoutSR, Gibco) to remove exogenous sources of TGF $\beta$ . MCF10A cells (purchased from ATCC) were cultured under serum free conditions in MEGM medium (Cambrex) supplemented with 0.1ug/ml of cholera toxin (Calbiochem).

Unless otherwise noted, confluent, growth arrested cells were used in experiments. Cells were exposed to a 5 Gy dose unless otherwise noted of 250 KVp X-ray or Co $^{60}$   $\gamma$ -radiation in air at room temperature. Control cells were sham-irradiated. In some experiments, cells were treated as indicated with 500 pg/ml recombinant TGF $\beta$ 1 (R&D Systems, Minneapolis, MN). In other

experiments, T $\beta$ RI kinase inhibitor (240nM, Calbiochem, Cat. No. 616451) was added to confluent cultures. To relieve the TGF $\beta$ RI kinase inhibitor block, medium was replaced with fresh medium and cells were cultured for an additional 48hr prior to irradiation.

**Protein analysis.** Primary cells and immortalized cell lines were isolated at indicated time points and lysed in buffer containing 50mM Tris (pH 7.5), 50mM glycerophosphate, 150mM NaCl, 10% glycerol, 1% Tween-20, 1mM of PMSF, 100  $\mu$ M DTT, 10  $\mu$ M NaVa, and 1mM NaF. Protein samples collected at times indicated post-IR were stored at -80 °C prior to separation using 4-15% SDS-PAGE gel. Proteins were transferred to Immobilon P (Millipore) PVDF membrane and incubated with primary antibodies, washed, incubated with goat anti-mouse Alexa680 (Molecular Probes cat. #A-21058) or goat anti-rabbit Dye800 (Rockland cat. #611-132-122) secondary antibodies, subsequently washed at room temperature. Membranes were scanned on the Odyssey Infrared Imaging System (LiCor). Target proteins were normalized to  $\beta$ -actin for loading and to the irradiated wildtype response for genotype comparisons; mean and standard error were determined from three or more independent experiments.

**Antibodies.** Antibodies to p53 serine 18, p53 serine 23, Rad17 serine 645, and total Rad17 were purchased from Cell Signaling Technology (Beverly, MA). Total p53 was detected using monoclonal G59-12 antibody purchased from Pharmigen. ATM was immunoprecipitated using anti-ATM antibodies (exon 53) from Bethyl (Montgomery, TX), and immunoblotted using ATM 2C1 antibodies from Genetex (San Antonio, TX).  $\beta$ -actin monoclonal clone EC-15 was from Sigma (St. Louis, MO). Monoclonal antibody clone 10H11.E12 and rabbit polyclonal antibodies to phosphorylated serine 1981 of ATM were purchased from Rockland Antibodies (Gilbertsville, PA).

Sheep anti-phospho serine 1981 ATM antibody was produced using KLH-phospho serine 1981 ATM peptide and affinity purified. Monoclonal  $\gamma$ H2AX antibody was from Upstate Cell Signaling (Lake Placid NY) and affinity purified rabbit anti-53BP1 antibody (BL181) was purchased from Bethyl Labs (Montgomery, TX).

**ATM kinase assay.** The ATM kinase assay was performed on fresh cell extracts using the GST-p53<sub>1-44</sub> substrate as described in (22). A-T human fibroblasts, purchased from Coriell Institute and cultured as recommended by supplier, were included as negative controls.

**Immunofluorescence.** Immunostaining to detect indicated target protein RIF was performed and imaged using cells cultured on LabTek 8-well chamber slides as reported (23). After treatment, cells were fixed using 2% paraformaldehyde for 5 min at room temperature followed by 100% MeOH for 20 min at -20°C. Negative controls were incubated without primary antibodies. Nuclei were counterstained with DAPI (4',6-Diamidino-2-Phenylindole) using 0.5  $\mu$ g/ml. Representative false color images are shown. In some experiments, nuclear fluorescence of 50-150 cells from 4 random fields were quantified by defining the DAPI-stained nucleus as the region of interest and integrating the mean intensity per nucleus. The mean intensity  $\pm$  S.E.M. was determined for each experimental condition.

**Flow Cytometry.** Asynchronously growing cells were pulsed for 30min with 10  $\mu$ M BrdU (Sigma) at the indicated time after irradiation. Trypsinized cells were fixed in 70% ethanol for 24hr and stained with 50  $\mu$ g/ml propidium iodide (Molecular Probes) and analyzed on Beckton Dickinson FACScan to determine cell cycle distribution.



**Colony Assay.** MCF10A cells were grown to confluence as described above, treated with 240 nM of T $\beta$ RI kinase inhibitor for 48 hr before irradiation using 250 KVp X-ray (0.61 Gy/min) . Cells were trypsinized 3 hrs later. Cells were plated in triplicate at 3 dilutions into 6-well plates and colonies were allowed to grow before fixing and staining. Colonies containing >50 cells were counted. To determine percent survival, colony forming efficiency was determined, averaged, and normalized to those of the non-irradiated control. The mean survival  $\pm$  s.e.m. was calculated for three replicate plates. The data shown are representative of three experiments.

## Results

### TGF $\beta$ dependence of the radiation response is cell-intrinsic

We have shown that p53 phosphorylation and apoptosis are significantly decreased in irradiated *Tgf $\beta$ 1* heterozygote compared to wildtype mouse mammary gland (4). To determine whether chronic TGF $\beta$  depletion in vivo had fundamentally (i.e. irreversibly) altered the radiation response of *Tgf $\beta$ 1* heterozygote epithelial cells, or if it actually mediates the execution of the radiation response, we examined primary cultures of murine mammary epithelial cells (MEC). Irradiated *Tgf $\beta$ 1* heterozygote MEC cultures showed significantly reduced phosphorylation of p53 compared to irradiated wildtype cells; p53 phosphorylation at serine 18 was reduced by 54% and at serine 23 by 63% relative to wildtype cell cultures (Figure 1A). Total MEC protein extracts were immunoblotted to confirm that p53 levels were similar between genotypes. IR-induced phosphorylations of p53 increase stability and transcriptional activity to induce downstream effector genes that mediate cell cycle delay and apoptosis, as well as initiating apoptosis directly (24, 25). Consistent with decreased p53 phosphorylation, caspase 3 cleavage, which is a marker of apoptosis, peaked at 2 hr post-IR in

wildtype MEC, but was not detected in *Tgfb1* heterozygote up to 4 hr post-IR (Figure 1B). We then asked whether this phenotype was reverted by supplementation of the serum-free culture with TGF $\beta$  (500 pg/ml). TGF $\beta$  treatment restored the ability of irradiated heterozygote MEC to phosphorylate p53 (Figure 1A) and induce caspase cleavage (Figure 1C). These data indicated that the TGF $\beta$  dependence of appropriate signaling and cell fate decisions are cell intrinsic and are compromised when TGF $\beta$  is limited.

**IR-induced phosphorylation of genotoxic stress response proteins and subsequent cell cycle arrest are severely compromised in *Tgfb1* null keratinocytes.**

To further characterize the nature and extent of the molecular defects in epithelial response to IR, we used two heterozygote and two null *Tgfb1* keratinocyte cell lines that spontaneously immortalized from primary cultures (5). *Tgfb1* heterozygote cells are competent to produce some TGF $\beta$ , albeit greatly reduced compared to wildtype (26), while null cell lines depend on, and are responsive to, exogenous sources of TGF $\beta$  such as serum. Therefore serum was eliminated by growing cell cultures to confluence before changing the media to a serum-free formulation, which also ensured that cell cycle distribution was consistent during subsequent experiments.

The prototype DNA damage response induced by IR is mobilized by the highly cytotoxic double strand break (27). The mechanism that allows this rapid dissemination of the damage alarm is based on a signal transduction pathway that begins with sensor/activator proteins that sense the damage or possibly the chromatin alterations that follow damage induction. These proteins play a major role in the activation of the transducers, which further convey the signal to multiple downstream effectors (28). Thus we examined the abundance and phosphorylation status of p53, Chk2, and Rad17 as a

function of time post-IR. Unirradiated cells of either genotype showed similar levels of total p53, Chk2, and Rad17 protein (Figure 2A). Irradiation of *Tgf $\beta$ 1* heterozygote cell lines induced prolonged phosphorylation of p53 serine 18 and Rad17 serine 645 that were maximal at 4 hr, while phosphorylation of p53 serine 23 and Chk2 threonine 68 were maximally induced at 1 hr post-IR and undetectable at later time points. In comparison, null genotype keratinocytes were hypophosphorylated in response to IR. Phosphorylation of p53 serine 18 was 30% in *Tgf $\beta$ 1* null cells relative to heterozygote cells post IR at 15 min and considerably reduced at later time points. p53 serine 23 phosphorylation was undetectable at any time in *Tgf $\beta$ 1* null keratinocyte cells. p53 serine 23 is phosphorylated by Chk2 (29), which requires phosphorylation at threonine 68 for its kinase activity (30). Compared to heterozygote cells Chk2 threonine 68 phosphorylation was also significantly reduced in null keratinocyte cells. Phosphorylation of Rad17 at serine 645, which is necessary for the DNA damage-induced activation of cell-cycle checkpoints (31), was markedly decreased in *Tgf $\beta$ 1* null cells relative to heterozygote cell lines.

A hallmark of the DNA damage response is the activation of cell cycle checkpoints, which temporarily halt the cell cycle until the damage is repaired (32). Reduced phosphorylation of Rad17, Chk2 and p53 in response to DNA damage should compromise cell cycle checkpoints in cycling cells (reviewed in (33)). Because TGF $\beta$  regulates cell cycle, and radiation response is a function of cell cycle phase, we used cells grown to fed-confluence so that cell cycle differences were not a factor. However, experiments using asynchronous cultures showed that molecular responses were also compromised in proliferating *Tgf $\beta$ 1* null cells (data not shown). Therefore, we examined cell cycle distribution in response to IR in exponentially growing cells. As expected the percent of S-phase cells in *Tgf $\beta$ 1* heterozygote cells at 5hr post irradiation was reduced from 32% to 22%

(Students t-test, p value<0.005) and the percent of cells in G2 increased nearly 3-fold. In contrast, irradiated *Tgfb1* null cells did not undergo a significant change in cell cycle distribution. The lack of cell cycle arrest in this *Tgfb1* null cell line is comparable to the absence of DNA synthesis block observed in epithelial tissues of irradiated *Tgfb1* null embryos (4).

Nuclear  $\gamma$ H2AX radiation-induced foci (RIF) are an early event elicited by DNA double strand breaks (34).  $\gamma$ H2AX RIF formed readily in *Tgfb1* heterozygote cells, but were significantly reduced in irradiated *Tgfb1* null cells (Figure 2B). In contrast, 53BP1, which binds to epitopes in methylated lysine 79 of histone H3 (35), formed RIF in both irradiated *Tgfb1* genotypes (Figure 2C). 53BP1 RIF confirm the presence of DNA damage caused by IR. The reduced phosphorylation of p53, Rad17, Chk2 and H2AX suggests that the necessary kinase is compromised.

### **Atm kinase activity and autophosphorylation are markedly reduced in *Tgfb1* null cells**

ATM is a serine/threonine protein kinase required for the rapid response to IR-induced DNA double strand breaks (36). ATM can directly phosphorylate p53 serine 18, Rad17 serine 645, Chk2 threonine 68 and H2AX in response to IR and thus is a candidate for the defective DNA damage response of *Tgfb1* null cells. To test this hypothesis, Atm kinase activity was measured in *Tgfb1* null and heterozygote keratinocyte cell lines prior to and 1 hr post-irradiation using a p53 GST-substrate *in vitro* kinase assay (Figure 3A). The level of substrate phosphorylation by Atm immunoprecipitated from irradiated *Tgfb1* null keratinocytes was 30% that of *Tgfb1* heterozygote keratinocytes (Figure 3B). We determined that Atm protein levels were unaffected by *Tgfb1* gene status as measured by immunoblotting total Atm protein in cell extracts of null versus heterozygote keratinocyte cell lines (Figure 3C), null versus wildtype primary keratinocytes or heterozygote versus wildtype mammary

primary mammary epithelial cells (not shown). These data indicate that Atm activity, rather than abundance, is affected by TGF $\beta$  depletion.

Following radiation exposure, the Atm dimer undergoes auto-phosphorylation (22, 37, 38). Bakkenist and Kastan showed that ATM autophosphorylation at serine 1981 is involved in the dissociation of inactive dimer or higher order multimers and the initiation of kinase activity and correlates with its activity (37). Atm serine 1981 phosphorylation was clearly evident in *Tgfb1* heterozygote cells from 15 min through 4 hr post-IR. In contrast, *Tgfb1* null keratinocytes showed minimal ability to undergo Atm autophosphorylation at serine 1981 immediately following irradiation, and did not recover up to 4 hr post-IR (Figure 3C). Since Bakkenist and Kastan showed that ATM autophosphorylation occurs within 2 minutes of DNA damage, these data, in conjunction with compromised substrate phosphorylation and failure of cell cycle arrest, suggest that Atm activation in these cells is absent rather than delayed.

ATM is also involved in the response to UV irradiation (39). UV causes the formation of cyclobutane pyrimidine dimers and formation of single stranded breaks as the cell attempts to repair the damage. Both UV and IR elicit p53 phosphorylation. UV irradiated *Tgfb1* heterozygote cells showed prominent p53 serine 18 phosphorylation at 1 and 3 hr post UV irradiation, but *Tgfb1* null cells did not exhibit detectable p53 phosphorylation (Figure 3D). Likewise Atm 1981P was readily observed at 1 and 3 hr post UV irradiation in *Tgfb1* heterozygote cells, although absent in *Tgfb1* null keratinocytes. Furthermore, asynchronous cultures of *Tgfb1* heterozygote cells underwent cell cycle arrest in G2 after UV (15 J/m<sup>2</sup>), while *Tgfb1* null cells did not undergo a significant change in cell cycle distribution (data not shown). Thus, TGF $\beta$ 1 depletion broadly compromises the genotoxic

stress response to physical damage caused by IR and UV.

A fraction of activated ATM binds to the DSB sites (37, 40). Many of ATM substrates are phosphorylated by the chromatin-bound fraction of this kinase (41), such as Chk2, occur at DNA double strand breaks (reviewed in (32)). Upon IR exposure, *Tgfβ1* heterozygote cells exhibited formation of Atm serine 1981P nuclear RIF (Figure 4A). Consistent with the biochemical data, neither diffuse nor punctate phosphorylated serine 1981 Atm was detectable in *Tgfβ1* null cells. The average fluorescence intensity of Atm serine 1981P did not change in irradiated null cells ( $53.5 \pm 9.1$  vs  $51.9 \pm 11.6$  s.e.m. irradiated) while the fluorescence relative to sham-irradiated heterozygote cells increased 4.6-fold at 1 hr (mean fluorescence intensity  $35.31 \pm 6.4$  vs  $164.3 \pm 22.4$  s.e.m. irradiated). The difference between genotypes did not alter up to 4 hr post-IR (data not shown). According to current models (18), recruitment of both the ATM monomer and the ATM substrates are mediated by several proteins, including the MRN complex, MDC1, and 53BP1. As shown in Figure 2C, 53BP1 RIF formation is intact. We determined Nbs-1 and Mre-11 protein levels of were unaffected by immunoblotting (data not shown). Finally, we determined that *Tgfβ1* heterozygote mammary epithelium irradiated in vivo exhibited reduced phosphorylated ATM immunoreactivity compared to tissue from wildtype mice (Figure 4B), consistent with our previous observation of reduced p53 phosphorylation. Together, the localization and biochemical data indicate that Atm activation and autophosphorylation fail to respond to IR-induced DNA damage in TGFβ compromised *murine* epithelial cells.

These data suggest Atm damage responses are a function of TGFβ availability; if so addition of TGFβ should be sufficient to restore the program. TGFβ treatment for 4 hr or more did not induce

Atm serine 1981 in unirradiated cells but restored Atm serine 1981 autophosphorylation in response to IR as determined by immunoblotting (Figure 5A). IR-induced Atm serine 1981 RIF were also restored indicating fully functional activation (Figure 5B). Restitution of autophosphorylation correlated with function as shown by restoration of p53 serine 18 phosphorylation (Figure 4A) and nuclear  $\gamma$ H2AX after IR (Figure 5C). Thus, treatment with TGF $\beta$  is sufficient to restore Atm autophosphorylation and activity in the *Tgf $\beta$ 1* null keratinocytes, indicating that TGF $\beta$  is an essential regulator of this pathway.

### **T $\beta$ RI kinase inhibitor abrogates genotoxic stress responses in human epithelial cells.**

To test whether the DNA damage response of human epithelial cells is mediated by TGF $\beta$ , MCF10A human mammary epithelial cells were treated with a small molecule inhibitor of TGF $\beta$  type I receptor kinase (42). Control experiments showed that TGF $\beta$  type I receptor (T $\beta$ RI) kinase inhibitor treatment released MCF10A cells from TGF $\beta$  growth inhibition and blocked phosphorylation of Smad 2 (data not shown). When growth arrested MCF10A cells were irradiated following a 48 hr exposure to the T $\beta$ RI kinase inhibitor, phosphorylated p53, Chk2 and Rad17 were significantly reduced compared to irradiated cells treated with vehicle (Figure 6A). Consistent with this, ATM serine 1981 phosphorylation was decreased by more than 50%. Releasing the cells from the inhibitor by re-feeding with fresh media for 48 hr prior to IR exposure, restored the DNA damage-induced phosphorylations. Furthermore, T $\beta$ RI kinase inhibitor blocked the induction of ATM serine1981 (Figure 6B) and  $\gamma$ H2AX RIF in irradiated MCF10A cells (Figure 6C). As in murine cells, 53BP1 RIF were unaffected (data not shown).

AT cells are very radiosensitive and deletion of ATM leads to radiation hypersensitivity as measured

by clonogenic survival (43). We determined that clonogenic survival following a graded IR dose response was significantly decreased by TGF $\beta$  inhibition relative to cells irradiated without inhibitor treatment (Figure 6D). The survival of MCF10A cells irradiated with 2 Gy following treatment with T $\beta$ RI kinase inhibitor decreased by 35% compared to vehicle treated controls ( $36.1 \pm 1.9$  S.E.M. vs  $56.0 \pm 2.0$  S.E.M., n=3 experiments). *Tgfb1* null keratinocytes were also more radiosensitive as measured by clonogenic survival (data not shown).

## Discussion

TGF $\beta$  is a key extracellular player for initiating and integrating multiple cellular responses to tissue response to IR and other types of damage (44). Our studies demonstrate that the activation of the ATM-mediated genotoxic stress program in mouse and human epithelial cells is severely compromised by loss of TGF $\beta$ 1 signaling. Rather than affecting kinase or substrate abundance, our data point to modulation of ATM kinase activation by one or more TGF $\beta$  transcriptional targets. Decreased Atm activity and Ser1981 autophosphorylation suggests that inhibition of TGF $\beta$  signaling affects its ability to initiate the damage response. This conclusion is supported by compromised substrate phosphorylation (i.e. p53, chk2 and Rad17) as well as by the absence of  $\gamma$ H2AX foci in irradiated *Tgfb1* heterozygote cells and human cells treated with small molecule inhibitor of the type I receptor kinase. The abrogated genotoxic stress signaling by Atm in cultured cells and compromised autophosphorylation observed in irradiated *Tgfb1* heterozygote mammary gland (Figure 5) provides a mechanism to explain our previous observation that both apoptosis and cell cycle arrest are absent in the skin or liver of irradiated *Tgfb1* null embryos (4).

Our study reveals a novel functional link between TGF $\beta$  signaling and the ATM mediated molecular



casades that dictate epithelial cell fate. Notably, TGF $\beta$  treatment of null keratinocytes prior to irradiation was essential for restoration of the DNA damage response, indicating that TGF $\beta$  signaling primes cells to respond to DNA damage either by assisting in the recruitment of ATM to the site of DNA damage or by facilitating ATM activation. This would suggest that an additional signal is required for these processes in epithelial cells or that one of the proteins normally involved in ATM activation is missing or defective. Alternatively TGF $\beta$  may directly or indirectly suppress an inhibitory function of the activation process, although, at this time, there is no known inhibitor of ATM activation. In the absence of TGF $\beta$  production or signaling, this function is dominant in both human and mouse epithelial cells.

An interesting question raised by these studies is why normal epithelial cells should require an extracellular factor in order to respond to DNA damage? The coupling of intracellular response mediated by ATM and extracellular signaling by TGF $\beta$  would ensure an integrated tissue response to damage and restoration of homeostasis, which is a novel mechanism for preventing cancer (45). Unlike fibroblasts and lymphoid cells, epithelial cells function in large part as an integrated unit, which if breached make the organism susceptible to a wide range of pathologies. By ensuring that extracellular and intracellular signaling in the response to DNA damage are intrinsically and reciprocally intertwined, then organisms can maintain homeostatic coordination of cellular events within an epithelium.

Importantly these studies suggest an additional mechanism by which early escape from TGF $\beta$  signaling could contribute to the development of cancer. Preneoplastic lesions exhibit high levels of DNA damage response proteins, which is postulated to increase the potential for genomic

instability (46). Our studies suggest that escape from TGF $\beta$ 1, in addition to releasing epithelial cells from growth control, would compromise responses to genotoxic stress, thus priming cells to become unstable. Consistent with this, keratinocytes from *Tgf $\beta$ 1* null mice exhibit a 100-1000 fold greater genomic instability measured by gene amplification than wildtype cells (5). TGF $\beta$  has been considered a canonical tumor suppressor of epithelial tissues (reviewed in (3, 47, 48); our studies indicate that TGF $\beta$  acts to maintain homeostasis in an even more comprehensive manner than previously recognized. Furthermore, current development of TGF $\beta$  inhibitors for use in cancer therapy, potentially in combination with DNA damaging agents, may well provide therapeutic advantage, as is demonstrated by increased radiosensitivity in our model systems.

### **Acknowledgments**

The authors wish to thank Ms. Yalda Afshar and Rosie Chau for experimental assistance and Mr. William Chou for preparation of figures.

**References**

1. Barcellos-Hoff, M. H., Derynck, R., Tsang, M. L.-S., and Weatherbee, J. A. Transforming growth factor- $\beta$  activation in irradiated murine mammary gland. *J Clin Invest*, 93: 892-899, 1994.
2. Siegel, P. M. and Massague, J. Cytostatic and apoptotic actions of TGF-beta in homeostasis and cancer. *Nat Rev Cancer*, 3: 807-821, 2003.
3. Derynck, R., Ackhurst, R. J., and Balmain, A. TGF- $\beta$  signaling in tumor suppression and cancer progression. *Nature Genet*, 29: 117-129, 2001.
4. Ewan, K. B., Henshall-Powell, R. L., Ravani, S. A., Pajares, M. J., Arteaga, C., Warters, R., Akhurst, R. J., and Barcellos-Hoff, M. H. Transforming Growth Factor- $\beta$ 1 Mediates Cellular Response to DNA Damage in Situ. *Cancer Res*, 62: 5627-5631, 2002.
5. Glick, A. B., Weinberg, W. C., Wu, I. H., Quan, W., and Yuspa, S. H. Transforming growth factor beta 1 suppresses genomic instability independent of a G1 arrest, p53, and Rb. *Cancer Res*, 56: 3645-3650, 1996.
6. Yamada, H., Vijayachandra, K., Penner, C., and Glick, A. Increased sensitivity of transforming growth factor (TGF) beta 1 null cells to alkylating agents reveals a novel link between TGFbeta signaling and O(6)-methylguanine methyltransferase promoter hypermethylation. *J Biol Chem*, 276: 19052-19058, 2001.
7. Glick, A., Popescu, N., Alexander, V., Ueno, H., Bottinger, E., and Yuspa, S. H. Defects in

- transforming growth factor-beta signaling cooperate with a Ras oncogene to cause rapid aneuploidy and malignant transformation of mouse keratinocytes. *Proc Natl Acad Sci USA*, *96*: 14949-14954, 1999.
8. Glick, A. B., Lee, M. M., Darwiche, N., Kulkarni, A. B., Karlsson, S., and Yuspa, S. H. Targeted deletion of the TGF- $\beta$ 1 gene causes rapid progression to squamous cell carcinoma. *Genes Dev*, *8*: 2429-2440, 1994.
  9. Glick, A. B., Kulkarni, A. B., Tennenbaum, T., Hennings, H., Flanders, K. C., O'Reilly, M., Sporn, M. B., Karlsson, S., and Yuspa, S. H. Loss of expression of transforming growth factor  $\beta$  in skin and skin tumors is associated with hyperproliferation and a high risk for malignant conversion. *Proc Natl Acad Sci USA*, *90*: 6076-6080, 1993.
  10. Barcellos-Hoff, M. H. Radiation-induced transforming growth factor  $\beta$  and subsequent extracellular matrix reorganization in murine mammary gland. *Cancer Res*, *53*: 3880-3886, 1993.
  11. Barcellos-Hoff, M. H. and Dix, T. A. Redox-mediated activation of latent transforming growth factor- $\beta$ 1. *Molec Endocrin*, *10*: 1077-1083, 1996.
  12. Martin, M., Lefaix, J.-L., Pinion, P., Crechet, C., and Daburon, F. Temporal modulation of TGF- $\beta$ 1 and  $\beta$ -actin gene expression in pig skin and muscular fibrosis after ionizing radiation. *Radiat. Res.*, *134*: 63-70, 1993.
  13. Martin, M., Vozenin, M. C., Gault, N., Crechet, F., Pfarr, C. M., and Lefaix, J. L.

- Coactivation of AP-1 activity and TGF-beta1 gene expression in the stress response of normal skin cells to ionizing radiation. *Oncogene*, *15*: 981-989, 1997.
14. Ehrhart, E. J., Carroll, A., Segarini, P., Tsang, M. L.-S., and Barcellos-Hoff, M. H. Latent transforming growth factor- $\beta$  activation in situ: Quantitative and functional evidence following low dose irradiation. *FASEB J*, *11*: 991-1002, 1997.
  15. Kurz, E. U. and Lees-Miller, S. P. DNA damage-induced activation of ATM and ATM-dependent signaling pathways. *DNA Repair (Amst)*, *3*: 889-900, 2004.
  16. Shiloh, Y. ATM and related protein kinases: safeguarding genome integrity. *Nat Rev Cancer*, *3*: 155-168, 2003.
  17. Shiloh, Y. ATM and ATR: networking cellular responses to DNA damage. *Current Opinion in Genetics & Development*, *11*: 71-77, 2001.
  18. Kastan, M. B. and Bartek, J. Cell-cycle checkpoints and cancer. *Nature*, *432*: 316-323, 2004.
  19. Abraham, R. T. Checkpoint signaling: epigenetic events sound the DNA strand-breaks alarm to the ATM protein kinase. *Bioessays*, *25*: 627-630, 2003.
  20. Kastan, M. B., Lim, D. S., Kim, S. T., Xu, B., and Canman, C. Multiple signaling pathways involving ATM. *Cold Spring Harb Symp Quant Biol*, *65*: 521-526, 2000.
  21. Barcellos-Hoff, M. H., Aggeler, J., Ram, T. G., and Bissell, M. J. Functional differentiation and alveolar morphogenesis of primary mammary epithelial cells cultures on reconstituted

- basement membrane. *Development*, 105: 223-235, 1989.
22. Kozlov, S., Gueven, N., Keating, K., Ramsay, J., and Lavin, M. F. ATP Activates Ataxia-Telangiectasia Mutated (ATM) in Vitro. Importance of Autophosphorylation. *J. Biol. Chem.*, 278: 9309-9317, 2003.
  23. Costes, S. V., Boissiere, A., Ravani, S. A., Romano, R., Parvin, B., and Barcellos-Hoff, M. H. Imaging features that discriminate between high and low LET radiation-induced foci in human fibroblasts. *Radiat Res*, 165: 505-515, 2006.
  24. Appella, E. and Anderson, C. W. Post-translational modifications and activation of p53 by genotoxic stresses. *European Journal of Biochemistry*, 268: 2764-2772, 2001.
  25. Meek, D. W. The p53 response to DNA damage. *DNA Repair (Amst)*, 3: 1049-1056, 2004.
  26. Ewan, K. B., Shyamala, G., Ravani, S. A., Tang, Y., Akhurst, R. J., Wakefield, L., and Barcellos-Hoff, M. H. Latent TGF- $\beta$  activation in mammary gland: Regulation by ovarian hormones affects ductal and alveolar proliferation. *Am J Path*, 160: 2081-2093, 2002.
  27. Bassing, C. H. and Alt, F. W. The cellular response to general and programmed DNA double strand breaks. *DNA Repair (Amst)*, 3: 781-796, 2004.
  28. Bakkenist, C. J. and Kastan, M. B. Initiating cellular stress responses. *Cell*, 118: 9-17, 2004.
  29. Keramaris, E., Hirao, A., Slack, R. S., Mak, T. W., and Park, D. S. Ataxia Telangiectasia-mutated Protein Can Regulate p53 and Neuronal Death Independent of Chk2 in Response

- to DNA Damage. *J. Biol. Chem.*, 278: 37782-37789, 2003.
30. Matsuoka, S., Rotman, G., Ogawa, A., Shiloh, Y., Tamai, K., and Elledge, S. J. Ataxia telangiectasia-mutated phosphorylates Chk2 in vivo and in vitro. *PNAS*, 97: 10389-10394, 2000.
  31. Bao, S., Tibbetts, R. S., Brumbaugh, K. M., Fang, Y., Richardson, D. A., Ali, A., Chen, S. M., Abraham, R. T., and Wang, X. F. ATR/ATM-mediated phosphorylation of human Rad17 is required for genotoxic stress responses. *Nature*, 21: 969-974, 2001.
  32. Lukas, J., Lukas, C., and Bartek, J. Mammalian cell cycle checkpoints: signalling pathways and their organization in space and time. *DNA Repair (Amst)*, 3: 997-1007, 2004.
  33. Sancar, A., Lindsey-Boltz, L. A., Unsal-Kacmaz, K., and Linn, S. Molecular Mechanisms of Mammalian DNA Repair and the DNA Damage Checkpoints. *Annual Review of Biochemistry*, 73: 39-85, 2004.
  34. Burma, S., Chen, B. P., Murphy, M., Kurimasa, A., and Chen, D. J. ATM Phosphorylates Histone H2AX in Response to DNA Double-strand Breaks. *J. Biol. Chem.*, 276: 42462-42467, 2001.
  35. Huyen, Y., Zgheib, O., Ditullio, R. A., Jr., Gorgoulis, V. G., Zacharatos, P., Petty, T. J., Sheston, E. A., Mellert, H. S., Stavridi, E. S., and Halazonetis, T. D. Methylated lysine 79 of histone H3 targets 53BP1 to DNA double-strand breaks. *Nature*, 432: 406-411, 2004.
  36. Shiloh, Y. ATM: Sounding the double-strand break alarm. *Cold Spring Harb Symp Quant*

- Biol, 65: 527-533, 2000.
37. Bakkenist, C. J. and Kastan, M. B. DNA damage activates ATM through intermolecular autophosphorylation and dimer dissociation. *Nature*, 421, 2003.
  38. Kozlov, S. V., Graham, M. E., Peng, C., Chen, P., Robinson, P. J., and Lavin, M. F. Involvement of novel autophosphorylation sites in ATM activation. *EMBO J*, 25: 3504-3514, 2006.
  39. Jazayeri, A., Falck, J., Lukas, C., Bartek, J., Smith, G. C. M., Lukas, J., and Jackson, S. P. ATM- and cell cycle-dependent regulation of ATR in response to DNA double-strand breaks. *8*: 37-45, 2006.
  40. Andegeko, Y., Moyal, L., Mittelman, L., Tsarfaty, I., Shiloh, Y., and Rotman, G. Nuclear Retention of ATM at Sites of DNA Double Strand Breaks. *J. Biol. Chem.*, 276: 38224-38230, 2001.
  41. Lukas, C., Falck, J., Bartkova, J., Bartek, J., and Lukas, J. Distinct spatiotemporal dynamics of mammalian checkpoint regulators induced by DNA damage. *Nat Cell Biol*, 5: 255-260, 2003.
  42. Sawyer, J. S., Anderson, B. D., Beight, D. W., Campbell, R. M., Jones, M. L., Herron, D. K., Lampe, J. W., McCowan, J. R., McMillen, W. T., Mort, N., Parsons, S., Smith, E. C. R., Vieth, M., Weir, L. C., Yan, L., Zhang, F., and Yingling, J. M. Synthesis and Activity of New Aryl- and Heteroaryl-Substituted Pyrazole Inhibitors of the Transforming Growth



- Factor- Type I Receptor Kinase Domain. *J Med Chem*, *46*: 3953 - 3956, 2003.
43. Xu, Y. and Baltimore, D. Dual roles of ATM in the cellular response to radiation and in cell growth control. *Genes Dev*, *10*: 2401-2410, 1996.
  44. Barcellos-Hoff, M. H. How tissues respond to damage at the cellular level: orchestration by transforming growth factor- $\beta$  (TGF- $\beta$ ). *BJR Suppl*, *27*: 123-127, 2005.
  45. Barcellos-Hoff, M. H. Integrative radiation carcinogenesis: interactions between cell and tissue responses to DNA damage. *Semin Cancer Biol*, *15*: 138-148, 2005.
  46. Bartkova, J., Horejsi, Z., Koed, K., Kramer, A., Tort, F., Zieger, K., Guldberg, P., Sehested, M., Nesland, J. M., Lukas, C., Orntoft, T., Lukas, J., and Bartek, J. DNA damage response as a candidate anti-cancer barrier in early human tumorigenesis. *Nature*, *434*: 864-870, 2005.
  47. Massague, J., Blain, S. W., and Lo, R. S. TGF- $\beta$  signaling in growth control, cancer, and heritable disorders. *Cell*, *103*: 295–309, 2000.
  48. Roberts, A. B. and Wakefield, L. M. The two faces of transforming growth factor  $\beta$  in carcinogenesis. *PNAS*, *100*: 8621-8623, 2003.

**Figures****Figure 1. p53 is hypophosphorylated and caspase cleavage are compromised in irradiated *Tgfb1* heterozygote primary MEC and are restored by TGF $\beta$  treatment.**

A. Representative immunoblots of p53 phosphorylation at serine 18 and serine 23, and total p53 in sham and irradiated wildtype (WT) and heterozygote (HT) mammary MEC in serum-free culture treated with or without TGF $\beta$  prior to IR.  $\beta$ -actin is shown as protein loading control. B. Representative immunoblots of total caspase 3 and cleavage product in wildtype and heterozygote mammary MEC following IR.  $\beta$ -actin is shown as protein loading control. C. Densitometric analysis of 17kDa cleavage fragment normalized to total caspase-3 demonstrates restoration of caspase cleavage by TGF $\beta$  treatment prior to IR.

**Figure 2. IR- induced phosphorylation of DNA damage response proteins and cell cycle arrest are compromised in *Tgfβ1* null keratinocyte cell lines.**

A. Immunoblots of p53 phosphorylation at serine 18 and serine 23, Rad17 serine 645 phosphorylation, and Chk2 threonine 685 of sham and irradiated *Tgfβ1* null and heterozygote keratinocyte cell lines as a function of time after IR. Total proteins and β actin are shown. B. Immunofluorescence detection of phosphorylated γH2AX (green) RIF formation in DAPI stained nuclei (blue) of sham and irradiated *Tgfβ1* null and heterozygote keratinocytes. γH2AX RIF formation at 30 min post-IR (2 Gy) is evident in heterozygote keratinocytes but is barely detectable in irradiated null keratinocytes. C. Immunolocalization of 53BP1 (green) in DAPI stained nuclei (blue). Sham-irradiated *Tgfβ1* heterozygote and null keratinocytes showed diffuse nuclear immunoreactivity. 53BP1 formed distinct RIF at 30 min post-IR (2 Gy) throughout the nuclei of both *Tgfβ1* heterozygote and null keratinocyte cells.

**Figure 3. ATM kinase activity and autophosphorylation are markedly reduced in TGF $\beta$ 1 compromised cells.**




A. Representative kinase assay of immortalized *Tgf $\beta$ 1* null compared to heterozygote keratinocyte cell lines. Irradiated ataxia telangiectasia (A-T) fibroblasts were included as a negative control. ATM immunoblotting after immunoprecipitation shows similar ATM loading for each genotype and treatment. Kinase activity was dramatically reduced in null versus heterozygote cells. B. Quantitation by densitometry of the kinase activity normalized to ATM protein immunoprecipitation of *Tgf $\beta$ 1* null and heterozygote cell lines (mean $\pm$ s.e.m., n=3 experiments). C. Dual immunoblot using infrared antibodies shows ATM serine 1981 autophosphorylation (green/yellow) and total ATM (red) as a function of time post-IR. Heterozygote keratinocytes show rapid and persistent ATM autophosphorylation that is lacking in null keratinocyte cells.  $\beta$ -actin is shown as protein loading control. D. Dual immunoblot analysis of *Tgf $\beta$ 1* heterozygote and knockout keratinocyte cell extracts shows that phosphorylation of p53 serine18 and ATM serine 1981 are diminished in *Tgf $\beta$ 1* null cells at 1hr post UV and do not recover by 3 hr.  $\gamma$ -tubulin is shown as protein loading control.

**Figure 4. Reduced localization of nuclear ATM serine 1981 in vitro and in vivo.** A. Immunolocalization of phospho-specific antibodies to ATM serine 1981 (green) and DAPI stained nuclei (blue) are shown. Sham-irradiated *Tgfb1* heterozygote and null keratinocytes showed little immunoreactivity. Irradiated A-T cells were negative (not shown). At 30 min post-IR (2 Gy), immunolocalized phosphorylated serine 1981 ATM was evident as distinct RIF throughout the nuclei of *Tgfb1* heterozygote keratinocyte cells but was absent from null cells. B. Immunolocalization of phospho-specific sheep anti-ATM serine 1981 (green) and DAPI stained nuclei (blue) in *Tgfb1* wildtype and heterozygote mammary gland. Sham-irradiated tissue showed little immunoreactivity in either genotype. At 1 hr post-IR (5Gy), prominent ATM serine 1981 immunostaining of nuclei was evident in the wildtype mice. Significantly less ATM serine 1981 immunostaining was present in epithelium of *Tgfb1* heterozygote mouse mammary gland.




**Figure 5. TGF $\beta$  treatment restores ATM autophosphorylation, localization, and function in *Tgfb1* null keratinocytes.**





A. Immunoblot of ATM serine 1981 phosphorylation, p53 serine 18 and  $\beta$  actin loading control for null keratinocytes. Treatment with TGF $\beta$  for 4 hr prior to IR (5Gy) significantly increased ATM serine 1981 autophosphorylation and p53 serine 18 phosphorylation. B. Immunolocalization of phospho-specific antibodies to ATM serine 1981 (green) and DAPI stained nuclei (blue) of null keratinocyte cells 30 min after irradiation (2 Gy). RIF formation was restored in TGF $\beta$  treated null keratinocyte cells. C. Histograms of the mean intensity per nucleus of  $\gamma$ H2AX immunoreactivity of *Tgfb1* null cells cultured in the presence or absence of TGF $\beta$ 1 before being irradiated.  $\gamma$ H2AX immunoreactivity was not induced by TGF $\beta$  treatment alone. TGF $\beta$ 1 exposure prior to irradiation restored  $\gamma$ H2AX in the majority of irradiated *Tgfb1* null cells.



**Figure 6. T $\beta$ R1 kinase small molecule inhibitor decreases the DNA damage response in human epithelial cells.** A. Confluent MCF10A human mammary epithelial cells cultured in serum-free medium for 96 hr were treated as follows: Control (lane 1, white bar), irradiated (lane 2, black bar), treated for the final 48 hr with T $\beta$ R1 kinase inhibitor and irradiated (lane 3, light grey bar), treated for 48 hr with T $\beta$ R1 kinase inhibitor and refed with fresh medium without inhibitor and irradiated (lane 4, dark grey bar). Cultures were harvested 1 hr after irradiation (5Gy) for immunoblotting. Quantitation of phosphorylation-specific antibodies using the LiCor Odyssey was normalized to the respective total protein. T $\beta$ R1 kinase inhibitor treatment did not change the abundance or phosphorylation status of proteins in the absence of irradiation (not shown). B, C: RIF formation in MCF10A cells treated for 24 hr with and without 240nM T $\beta$ R1 kinase inhibitor and sham or irradiated (2 Gy). Forty minutes post-IR cells were fixed and stained with phospho-specific antibodies to either  $\gamma$ H2AX (B, green fluorescence) or phosphorylated ATM serine 1981 (C, green fluorescence). Nuclei are DAPI stained (blue). Treatment with T $\beta$ R1 kinase inhibitor impedes formation of nuclear RIF of both  $\gamma$ H2AX and phosphorylated ATM serine 1981. The formation of 53BP1 nuclear RIF was unaffected (not shown). D. Colony forming efficiency of irradiated MCF10A control cells (black) and treated for 48 hr with 240nM T $\beta$ R1 kinase inhibitor (red). Inhibition of TGF $\beta$  decreased clonogenic survival compared to controls as a function of radiation dose (ANOVA,  $p < 0.0001$ ).




**a****b****c**








**a****b****c**



**a****b**

**a****b****c**

**a****b****c****d**ANOVA (tukey) test = \*\*\*,  $p < 0.0001$