Lawrence Berkeley National Laboratory

Lawrence Berkeley National Laboratory

Title

Direct Numerical Modeling of E-Cloud Driven Instability of a Bunch Train in the CERN SPS

Permalink

https://escholarship.org/uc/item/34d5c2mh

Author

Vay, J-L.

Publication Date

2011-04-29

HIFAN 1823

Direct Numerical Modeling of E-Cloud Driven Instability of a Bunch Train in the CERN SPS

by

J-L. Vay, M.A. Furman, M. Venturini

from

Lawrence Berkeley National Laboratory (on behalf of U.S. HIFS-VNL)

1 Cyclotron Road, Berkeley, CA 94720

Accelerator Fusion Research Division

University of California

Berkeley, California 94720

March 2011

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California.

This work was supported by the Director, Office of Science, Office of Fusion Energy Sciences, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231.

Direct Numerical Modeling of E-Cloud Driven Instability of a Bunch Train in the CERN SPS*

J.-L. Vay[†], M. A. Furman, M. Venturini, LBNL, USA

Abstract

The simulation package WARP-POSINST was recently upgraded for handling multiple bunches and modeling concurrently the electron cloud buildup and its effect on the beam, allowing for direct self-consistent simulation of bunch trains generating, and interacting with, electron clouds. We have used the WARP-POSINST package on massively parallel supercomputers to study the buildup and interaction of electron clouds with a proton bunch train in the CERN SPS accelerator. Results suggest that a positive feedback mechanism exists between the electron buildup and the e-cloud driven transverse instability, leading to a net increase in predicted electron density.

INTRODUCTION

Electron clouds have been shown to trigger fast growing instabilities on proton beams circulating in the SPS [1] and other accelerators [2]. So far, simulations of electron cloud buildup and their effects on beam dynamics have been performed separately. This is a consequence of the large computational cost of the combined calculation due to large space and time scale disparities between the two processes. In [3], we have presented the latest improvements of the simulation package WARP-POSINST [4] for the simulation of self-consistent ecloud effects, including mesh refinement, and generation of electrons from gas ionization and impact at the pipe walls. We also presented simulations of two consecutive bunches interacting with electrons clouds in the SPS, which included generation of secondary electrons. The distribution of electrons in front of the first beam was initialized from a dump taken from a preceding buildup calculation using the POSINST code [5, 6, 7, 8]. In this paper, we present an extension of this work where one full batch of 72 bunches is simulated in the SPS, including the entire buildup calculation and the self-consistent interaction between the bunches and the electrons.

SELF-CONSISTENT SIMULATION OF ONE BATCH IN THE SPS

One batch of 72 consecutive bunches propagating in the SPS at injection was simulated for over 1000 turns of the machine, using the parameters given in Table 1 in the Appendix. A simulated bunch-to-bunch feedback system is

Table 1: Parameters Used for Warp-POSINST Simulations

| beam energy | E_b | 26 GeV |
|--------------------------|------------------|-----------------------|
| bunch population | N_b | 1.1×10^{11} |
| rms bunch length | σ_z | 0.23 m |
| rms transverse emittance | $\epsilon_{x,y}$ | 2.8, 2.8 mm.mrad |
| rms momentum spread | δ_{rms} | 2×10^{-3} |
| beta functions | $\beta_{x,y}$ | 33.85, 71.87 m |
| betatron tunes | $Q_{x,y}$ | 26.13, 26.185 |
| chromaticities | $Q'_{x,y}$ | 0, 0 |
| Cavity voltage | V | 2 MV |
| momentum compact. factor | α | 1.92×10^{-3} |
| circumference | C | $6.911~\mathrm{km}$ |
| bucket length | δ_b | 5 ns |
| bunch spacing | Δ_b | 25 ns |
| peak SEY | δ_{SEY} | 1.2 |
| # of bunches | N_{bunch} | 72 |
| # of stations/turn | N_s | 10 |
| # of slices/bucket | N_{slices} | 64 |

used to stabilize the beam in the horizontal direction. No feedback system is used in the vertical direction. The simulation was performed on a parallel supercomputer using 2928 processors.




Figure 1: Time history of the charge density of electron cloud averaged over the pipe section at one station around the ring from a WARP-POSINST simulation.

Time histories of the electron density averaged over the pipe cross section and on axis, taken at one location around the ring, are shown in Figures 1 and 2, at turns 0 to 1000 by intervals of 200. There is an increase in average and on axis electron density between turn 200 and turn 600 for approximately the last two third of the bunch train, with an maximum enhancement of about 50% of the average electron density and 100% (i.e. doubling) of the electron

^{*}Supported by the US-DOE under Contract DE-AC02-05CH1123, the US-LHC Accelerator Research Program (LARP) and the Sci-DAC/ComPASS project. Used resources of the National Energy Research Supercomputer Center (NERSC).

[†] jlvay@lbl.gov




Figure 2: Time history of the charge density of electron cloud on axis at one station around the ring from a WARP-POSINST simulation: (top) between $0\mu s \le t \le 2\mu s$; (bottom) between $1\mu s \le t \le 1.006\mu s$.

density on axis.

The turn-by-turn history of the bunches emittance, vertical RMS size and average vertical offset, as well as the averaged spectrum of vertical bunch slices oscillations, are given in Figure 3. It shows substantial growth of emittance and vertical RMS size for bunches 25 and higher, starting to develop around turn 200 and saturating around turn 600. For bunches 40 and above, there is a net spike of average vertical offset between turn 500 and 600, followed by a steady decrease. This means that the increase that is observed in vertical emittance and RMS size is initially caused by coherent motion followed by incoherent motion due to phase mixing of the particle trajectories. For all bunches, most of the content of the vertical oscillations spectrum is below 500 MHz.

The onset of the instability around bunch 25 corresponds to a time of 25×25 ns= $6.25~\mu s$ after the passage of the first bunch at a fixed station, which matches the time at which the electron density increases in Figures 1 and 2. Buildup simulations with POSINST show in Figure 4 that doubling the vertical dimension of the bunches lead to an increase by 20-30% of the average electron density during the buildup phase. The higher increase that is observed in the fully self-consistent WARP-POSINST simulation might be due to taking into account temporal and spatial variations in the beam distributions that are not modeled in POSINST. This suggests that accurate prediction of electron buildup and




Figure 3: Bunch quantities vs bunch and turn numbers: (top-left) Bunch emittance growth $(\epsilon(t)/\epsilon(0)-1)$; (top-right) bunch vertical (Y) RMS size; (bottom-left) bunch vertical offset (averaged over all slices of absolute value of each slice offset); (bottom-right) spectrum of offset oscillations along the bunch (averaged over 1100 turns).




Figure 4: Time history of the charge density of electron cloud averaged over the pipe section at one station around the ring from POSINST simulations, using secondary electron yield peaks of 1.2 (red) and 1.18 (blue), for bunches RMS vertical sizes of 2.69 mm (solid) and 5.38 mm (dash).

their effect on the bunches necessitates the self-consistent simulation of both effects.

CONCLUSION

Direct self-consistent simulation of bunch trains generating, and interacting with, electron clouds where performed with the WARP-POSINST package on massively parallel supercomputers to study the growth rate and frequency patterns in space-time of the electron cloud driven transverse instability for a proton bunch train in the CERN SPS accel-

erator. Analysis of the turn-by-turn evolution of the electron buildup and of the bunches vertical motion shows that the vertical size increase of the bunches that is caused by its interaction with the electron cloud causes in return a net increase of up to 50% in average electron density and 100% in electron density on axis. This suggests that accurate prediction of electron buildup and their effect on the bunches necessitates the self-consistent simulation of both effects.

REFERENCES

- [1] G. Arduini et al., CERN-2005-001, 31-47 (2005)
- [2] Proc. International Workshop on Electron-Cloud Effects "ECLOUD07" (Daegu, S. Korea, April 9-12, 2007). http://chep.knu.ac.kr/ecloud07
- [3] J.-L. Vay et al., "Simulation of e-cloud driven instability and its attenuation using a feedback system in the CERN SPS", Proceedings First International Particle Accelerator Conference IPAC10, Kyoto, Japan, paper WEOBRA02 (2010).
- [4] J-L Vay et al., Update on e-cloud simulations using the package WARP/POSINST, PAC09, Proceedings, (2009). Proceedings First International Particle Accelerator Conference IPAC10, Kyoto, Japan, paper WEPEB052 (2010).
- [5] M. A. Furman and G. R. Lambertson, LBNL-41123/CBP Note-246, PEP-II AP Note AP 97.27 (Nov. 25, 1997). Proc. Intl. Workshop on Multibunch Instabilities in Future Electron and Positron Accelerators "MBI-97" (KEK, 15-18 July 1997; Y. H. Chin, ed.), KEK Proceedings 97-17, Dec. 1997, p. 170.
- [6] M. A. Furman and M. T. F. Pivi, LBNL-49771/CBP Note-415 (Nov. 6, 2002). PRST-AB 5, 124404 (2003), http://prstab.aps.org/pdf/PRSTAB/v5/i12/e124404.
- [7] M. A. Furman and M. T. F. Pivi, LBNL-52807/SLAC-PUB-9912 (June 2, 2003).
- [8] M. A. Furman, LBNL-41482/CBP Note 247/LHC Project Report 180 (May 20, 1998).