

UC Irvine

UC Irvine Previously Published Works

Title

Widespread multiple insecticide resistance in the major dengue vector *Aedes albopictus* in Hainan Province, China.

Permalink

<https://escholarship.org/uc/item/38j001tm>

Journal

Pest management science, 77(4)

ISSN

1526-498X

Authors

Li, Yiji

Zhou, Guofa

Zhong, Daibin

et al.

Publication Date

2021-04-01

DOI

10.1002/ps.6222

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

Widespread multiple insecticide resistance in the major dengue vector *Aedes albopictus* in Hainan Province, China

Yiji Li,^{a,b}  Guofa Zhou,^b Daibin Zhong,^b Xiaoming Wang,^b Elizabeth Hemming-Schroeder,^b Randy E David,^b Ming-Chieh Lee,^b Saifeng Zhong,^a Guohui Yi,^c Zhuanzhuan Liu,^d Guzhen Cui^{e*} and Guiyun Yan^{b*}



Abstract

BACKGROUND: *Aedes albopictus* is a highly invasive mosquito and has become a potential vector of dengue, chikungunya and Zika viruses. Insecticide-based mosquito interventions are the main tools for vector-borne disease control. However, mosquito resistance to insecticides is a major threat to effective prevention and control. Five *Ae. albopictus* populations across Hainan Province, China were investigated for susceptibility to multiple insecticide and resistance mechanisms.

RESULTS: Larval bioassays indicated that resistance to pyrethroids was common in all larval populations. Adult bioassays revealed all populations were either resistant or highly resistant to at least four of the six synthetic insecticides (deltamethrin, permethrin, cyfluthrin, propoxur, malathion, and DDT) tested. Pre-exposure of mosquitoes to the synergistic agent piperonyl butoxide (PBO) increased mosquito mortality by 2.4–43.3% in bioassays to DDT, malathion, and permethrin and rendered mosquito sensitive to deltamethrin, cyfluthrin, and propoxur. The frequency of knockdown resistance (*kdr*) mutations (F1534S and F1534C) ranged from 69.8% to 89.3% and from 38.1% to 87.0% in field-resistant and sensitive populations, respectively. F1534S mutation was significantly associated with pyrethroid resistance. No mutation was detected in the acetylcholinesterase (*ace-1*) gene in the two examined populations.

CONCLUSION: This study provides evidence of widespread resistance to multiple insecticides in *Ae. albopictus* in Hainan Province, China. Both *kdr* mutations and metabolic detoxification were potential causes of insecticide resistance for *Ae. albopictus*. Our findings highlight the need for insecticide resistance management and mosquito control measures that do not entirely depend on synthetic insecticides.

© 2020 The Authors. *Pest Management Science* published by John Wiley & Sons Ltd on behalf of Society of Chemical Industry. Supporting information may be found in the online version of this article.

Keywords: *Aedes albopictus*; insecticide resistance; knockdown resistance (*kdr*); piperonyl butoxide; Hainan Province

1 BACKGROUND

Aedes albopictus (Skuse) is a potential vector of at least 20 arboviruses with public health importance in tropical and subtropical regions, including yellow fever virus, dengue virus (DENV), chikungunya virus (CHIKV), and Zika virus (ZIKV).^{1–7} It is listed among the most dangerous invasive mosquito species globally.⁸ Furthermore, *Ae. albopictus* was responsible for the vast majority of dengue and chikungunya outbreaks in China during the past two decades.^{9–12} The species is widely distributed in China from the southern tropical area of Hainan Province to the temperate climate Liaoning Province in the north.^{13,14}

So far, there is no effective drug or vaccine for most *Ae. albopictus* transmitted diseases except the vaccine for yellow fever, therefore mosquito control is the most important method for preventing *Aedes* mosquito-borne infectious diseases.^{1–4} Adult *Aedes*

* Correspondence to: G Yan, School of Medicine, University of California, 3501B Hewitt Hall, UC Irvine, California, USA, E-mail: guiyun@uci.edu; or G Cui, Guizhou Medical University, No.1 Dongqing South Rd., Huaxi District, Guiyang, Guizhou 550025, China. E-mail: cui-guzhen@gmc.edu.cn

a Department of Pathogen Biology, Hainan Medical University, Haikou, China

b Program in Public Health, School of Medicine, University of California, Irvine, CA, USA

c Public Research Laboratory, Hainan Medical University, Haikou, China

d Jiangsu Key Laboratory of Immunity and Metabolism, Department of Pathogen Biology and Immunology, Xuzhou Medical University, Xuzhou, China

e Key Laboratory for Endemic and Ethnic Diseases, Ministry of Education, School of Basic Medical Science, Guizhou Medical University, Guiyang, China

mosquito control depends largely on outdoor and indoor insecticide spraying, while larval control relies on either chemical (e.g. temphos) or biological (e.g. *bti*) insecticides and community engagement for habitat management.^{15–17} In China, the primary tool for mosquito control is the use of chemical insecticides.^{18,19} However, *Ae. albopictus* resistance to all four main classes of insecticides (carbamates, organochlorines, organophosphates, and pyrethroids) has been detected in the Americas, Africa, and Asia in recent years.^{15,20,21}

Hainan Province, located in southern China, is a dengue fever epidemic area. Dengue fever outbreaks have occurred several times since the late 1970s. The 1979–1982 dengue pandemic on the island caused a total of 604 854 cases and 475 deaths.^{22,23} The most recent Dengue outbreaks occurred on this island in 2019.²⁴ As an international tourist destination and now a free-trade area, mosquito-borne infectious disease risk will likely grow due to the increasing influx of goods and human travel. This threat is underscored by the presence of dengue fever throughout the neighboring southeastern countries.²⁵ Furthermore, the tropical climate of the island makes it particularly suitable for the development and survival of vector mosquitoes. *Ae. albopictus* is widely distributed on the island, and the species is now almost the sole vector of dengue fever in Hainan.^{26,27} Previously, *Ae. aegypti* was the major dengue vector from the 1970s to the 1990s, but *Ae. aegypti* has not been found on the island during the past 10 years.^{23,28,29}

In the past 40 years, Hainan Province has used a large quantity of insecticides to help control vector-borne diseases, including malaria and dengue.²³ However, there is limited information available on the current status of insecticide resistance in *Ae. albopictus*. A previous study in Haikou city in Hainan indicated that *Ae. albopictus* had developed resistance to deltamethrin, permethrin, and beta-cypermethrin.²² Thus, monitoring the development of

insecticide resistance in *Aedes* mosquitoes has become a local key task for dengue control.⁷

Due to massive agricultural and public health use of insecticides, systematic studies on *Ae. albopictus* resistance to insecticides are critically needed for the prevention and control of mosquito-borne diseases. In the present study, we studied the insecticide resistance profiles of larval and adult *Ae. albopictus* in five areas across Hainan Province during 2017–2018. We implemented synergist and molecular assays to identify putative resistance mechanisms in *Ae. albopictus* adults.

2 MATERIALS AND METHODS

2.1 Ethics statement

No permits were required for the described field studies. Mosquito collections in breeding sites were consented orally by the owners at each location. For the insecticide usage survey, oral consent was obtained from all participating households after explanation of the study aim and design. This study did not involve collection of any human-related samples or personal information such as participants' names, addresses, phone numbers, and GPS location of their homes etc.

2.2 Study sites and mosquito collections

Immature *Ae. albopictus* mosquitoes were collected from five study sites in Hainan Province, southern China between 2017 and 2018 (Fig. 1 and Table 1). The five sites represent different ecological settings on the island (Table 1). Annual mean temperature ranges from 23.1 to 25.9 °C, and annual precipitation ranges from 1300 to 2300 mm (Table 1).

Mosquito larvae were collected from at least 50 habitats in each study site, which included both urban and suburban areas. Larvae were brought back to the insectary located at Hainan Medical




Figure 1. Map of study area in Hainan Province, China.

Table 1. Description of *Aedes albopictus* mosquito population collection sites in Hainan Province, China

| Study area | Landscape | Coordinates | | Altitude (m) |
|------------|-----------------------------|--------------|---------------|--------------|
| | | Latitude (N) | Longitude (E) | |
| Haikou | Densely populated port city | 110°20'29" | 20°04'39" | 4 |
| Wanning | Plain, agriculture area | 110°23'15" | 18°48'10" | 13 |
| Sanya | Tourist destination | 109°30'28" | 18°17'32" | 26 |
| Qiongzong | Heavily forested area | 109°50'47" | 19°03'03" | 267 |
| Danzhou | Major agricultural area | 109°33'35" | 19°31'30" | 184 |

University. Larvae were reared at 27 ± 2 °C, $70\% \pm 10\%$ relative humidity (RH) and a 12:12 h light:dark period, and fed with yeast. Emerged adults were fed with 10% sugar solution at 27 ± 2 °C, $70\% \pm 10\%$ RH.

2.3 Insecticide resistance bioassays

2.3.1 Larval bioassays

Eight insecticides were used to examine larval mortality. Seven of the insecticides were obtained from the Chinese Center for Disease Control and Prevention (China CDC). Three of them were pyrethroid insecticides: deltamethrin (technical grade 95.95%), permethrin (99%), and beta-cypermethrin (92.00%). The other chemical insecticides included were propoxur (95.56%), malathion (95%), temephos (87.40%), and DDT (94.00%). Lastly, one microbial larvicide was evaluated: *Bacillus thuringiensis israelensis* (*Bti*) (7000 ITU/mg, Wuhan Nature's Favour Bioengineering Co., Ltd, Wuhan, China). All chemical insecticides were diluted in acetone, and *Bti* was diluted in water to the required dosage following WHO guidelines.³⁰ We included adulticides for larval resistance tests because in China pyrethroid and other adulticides have been widely used for agricultural pest control and for public health use such as control mosquitoes inside rainwater drain basins. The residue of these insecticides in the water may cause larval resistance after long-term use.

The larvae of F1-2 generations were used for larval bioassays after species identification. In each test, 25 third- and fourth-instar larvae were incubated in a 100 mL solution consisting of 99 mL of distilled water and 1 mL of insecticide solution at the specified concentration using the standard 250 mL plastic cup.³⁰ Control treatments for the chemical insecticides were incubated in 99 mL of distilled water and 1 mL of acetone. Control treatments for *Bti* were incubated in 100 mL of water only. Three to five replicates were tested for each concentration. Each bioassay comprised one control group. Five to nine concentrations, providing a range of mortalities between 0% and 100%, were used to determine the 50% mortality lethal concentration (LC_{50}). Mosquitoes of the Foshan strain, which have been reared in the laboratory since 1981 without insecticide exposure, were used as a control. Larval mortality was recorded after 24 h of exposure. Bioassays were conducted at 27 ± 2 °C, $70\% \pm 10\%$ RH, and a 12:12 h light:dark period.

2.3.2 Adult resistance bioassays

The adults from F1-2 of field-collected *Ae. albopictus* larvae were tested using the standard WHO insecticide susceptibility tube test.^{31,32} The mosquitoes were subjected to insecticide susceptibility tests against pyrethroids (0.15% cyfluthrin, 0.03% deltamethrin, and 0.25% permethrin), organochlorine (4% DDT), organophosphate (0.8% malathion), and carbamate (0.1%

propoxur). The susceptible Foshan strain was used as a control. For each population of the field mosquitoes and each insecticide, eight replicates (four each in urban and suburban) of 20 non-blood-fed 3–5 day-old female mosquitoes for each replicate were subjected to WHO tube bioassays. Data analysis used pooled values of the eight replicates at each site. For control populations, two replicates were used for each insecticide. WHO test and control papers were supplied by the School of Biological Sciences, Universiti Sains Malaysia (Penang, Malaysia). Mosquitoes were exposed to test papers for 1 h. The knockdown number of females was recorded every 10 min during the 60 min exposure period. Mortality was scored after the 24 h recovery period. After the bioassay, both survivors and dead mosquitoes were stored individually in 95% alcohol at -80 °C for subsequent DNA analysis.

2.3.3 Synergist assay with piperonyl butoxide

To investigate the role of oxidase-specific metabolic resistance mechanisms, synergist assays with piperonyl butoxide (PBO) were performed together with the six insecticides tested above: deltamethrin, permethrin, cyfluthrin, malathion, propoxur, and DDT. Nonblood-fed adult mosquitoes which were 3–5-days old were pre-exposed to 4% PBO test paper for 1 h, followed by exposure to insecticide-treated test papers for 1 h. There were three replicates for each insecticide, with 25 mosquitoes per replicate. Mortality was recorded after the 24 h recovery period. Two replicates of 1 h PBO exposure were performed without subsequent exposure to an insecticide for the controls.

2.3.4 Molecular assays

Genomic DNA was extracted using the method published by Chang *et al.*³³ One adult mosquito leg for each mosquito was used for DNA extraction. Extracted DNA was stored at -20 °C or used immediately for PCR. The primers and PCR conditions used for amplification and sequencing of the voltage-gated sodium channel (VGSC) gene and the acetylcholinesterase (*ace-1*) gene of *Ae. albopictus* were according to the protocol from Su *et al* and Kasai *et al.*^{13,34} PCR products were purified and directly sequenced in both directions with the same primers. PCR products were purified with ExoSAP-IT (USB, Cleveland, OH, USA) according to the manufacturer's protocol and directly sequenced by Retrogen, Inc. (San Diego, CA, USA). The sequences were analyzed using BioEdit (<http://www.mbio.ncsu.edu/BioEdit/bioedit.html>) and Codon Code (<http://www.codoncode.com/>).

2.3.5 Community survey of insecticide usage

Questionnaire surveys were conducted in the study communities to determine insecticide usage for public health and agricultural pest control. Questionnaire surveys were conducted from September to October of 2019 in all study sites. Each survey site

included urban, suburban, and rural areas with a total of 30 questionnaires. Oral consent was obtained from all participants after explanation of the study purpose to them.

2.3.6 Statistical analysis

Resistance status of adult mosquito was classified according to WHO criteria: resistant for <90% mortality, probable resistance for 90–98% mortality, and susceptible for >98% mortality.³¹ In the larval bioassay, the median lethal concentration (LC₅₀), the 90% lethal concentration (LC₉₀), and the corresponding 95% confidence intervals were calculated based on the recorded data using Schoofs and Willhite's probit analysis program.³⁵ The degree of resistance intensity was determined by the resistance ratio (RR₅₀), calculated as the ratio of LC₅₀ value for the field population over the LC₅₀ value for the susceptible laboratory colony. By WHO definition, RR₅₀ < 5 is considered susceptible, 5 < RR₅₀ < 10 is moderately resistant, and RR > 10 is highly resistant.³¹ For the adult bioassays, χ^2 -tests were used to examine the association between *kdr* mutations and the resistance phenotype. Here, mosquitoes dead after the 24 h recovery period were classified as susceptible and surviving mosquitoes as resistant. For each population and each insecticide, the mortalities with mosquitoes exposed to PBO + insecticide were compared with those exposed with insecticide alone using *t*-tests.

3 RESULTS

3.1 Larval *Ae. albopictus* developed resistance to pyrethroid insecticides

Larval bioassays revealed that *Ae. albopictus* have developed different levels of resistance (RR₅₀ ≥ 5) to all pyrethroid insecticides at nearly all study sites, but they were all susceptible (RR₅₀ < 5) to all nonpyrethroid insecticides (Table 2 and Fig. 2). All larval populations were highly resistant to deltamethrin (LC₅₀ = 0.17–0.47 mg/L; RR₅₀ range 17–47), and most of them were moderately or highly resistant to permethrin (LC₅₀ = 0.36–1.86 mg/L, RR₅₀ range 7–37). The highest RR₅₀ was 53 for beta-cypermethrin (LC₅₀ = 1.06 mg/L) in the Haikou population, and the lowest RR₅₀ was 3 for beta-cypermethrin (LC₅₀ = 0.06 mg/L) in the Danzhou population (Fig. 2 and Table 2). In addition, all of the populations tested were susceptible to Propoxur (LC₅₀ = 0.89–4.78 mg/L, RR₅₀ range 0.8 to 4.2), malathion (LC₅₀ = 2.40–5.97 mg/L, RR₅₀ range 0.7–2.4), temephos (LC₅₀ = 0.07–0.16 mg/L, RR₅₀ range 1.8–4.0), DDT (LC₅₀ = 1.92–4.96 mg/L, RR₅₀ range

0.8–2.1), and *Bti* (LC₅₀ = 0.55–0.99 mg/L, RR₅₀ range 0.7–1.2). (Table 2 and Fig. 2).

3.2 Adult *Ae. albopictus* developed resistance to all the insecticides used

WHO tube bioassays revealed that nearly all *Ae. albopictus* populations were resistant (<90% mortality) to the six synthetic insecticides tested, as indicated by the low mortality rate after the 24 h recovery period (Fig. 3(a)). Although mortality rates varied, mosquitoes were highly resistant to DDT (mortality range 28–46%), permethrin (mortality range 47–71%), and malathion (mortality range 61–82%) at all study sites (Fig. 3(a)). Mosquito resistance levels ranged from highly resistant to suspected resistant to cyfluthrin (mortality range 69–91%), propoxur (mortality range 56–96%), and deltamethrin (mortality range 70–96%) (Fig. 3(a)).

3.3 Synergist PBO moderately increased mosquito mortality

Pre-exposure of *Ae. albopictus* mosquitoes to the PBO synergist before exposure to insecticides led to the full recovery of *Ae. albopictus* susceptibility to deltamethrin, propoxur, and cyfluthrin (except Wanning population, mortality 89%) and partial recovery of susceptibility to malathion (mortality range 72–100%) and permethrin (mortality range 82–96%) (Fig. 3(b)). However, *Ae. albopictus* mortality was still low (31–83%) against DDT after PBO treatment (Fig. 3(b)). The effect of pre-exposure to PBO on mosquito mortality varied substantially for a given insecticide among different sites and at a given site among different insecticides (Fig. 3, bottom panel), i.e. increase in mosquito mortality was observed overall, but showed strong heterogeneity (Fig. 3, bottom panel).

3.4 *kdr* mutations were associated with resistance phenotypes in adult mosquitoes

A total of 479 *Ae. albopictus* mosquitoes (234 resistant and 245 sensitive) were examined for *kdr* mutations. We detected F1534S and F1534C mutations in the VGSC gene. Overall, the total *kdr* mutation frequency was significantly higher in the resistant mosquitoes (mean 78.6%, range 69.8–89.3%) than that in the susceptible mosquitoes (mean 46.5%, range 38.1–50.9%) in four of the five populations (χ^2 -test, *P* < 0.05; Table 3). The odds ratio for the F1534S mutation ranged from 2.4 to 10.8 (but was 0.8 in Haikou) in resistant mosquitoes compared to susceptible mosquitoes, indicating the association between pyrethroid phenotypic

Table 2. Insecticide concentrations that caused 50% mortality (LC₅₀) of *Aedes albopictus* in different populations

| Insecticide | Haikou | Wanning | Sanya | Qiongzong | Danzhou | Susceptible strain |
|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|--------------------|
| Deltamethrin | 0.35 (0.25, 0.48) | 0.47 (0.38, 0.56) | 0.26 (0.20, 0.31) | 0.27 (0.15, 0.45) | 0.17 (0.09, 0.31) | 0.01 (0.01, 0.01) |
| Beta-cypermethrin | 1.06 (0.50, 2.48) | 0.15 (0.09, 0.27) | 0.17 (0.15, 0.19) | 0.09 (0.06, 0.14) | 0.06 (0.05, 0.06) | 0.02 (0.02, 0.03) |
| Permethrin | 1.86 (1.28, 2.57) | 1.24 (0.37, 2.78) | 0.51 (0.26, 0.94) | 0.86 (0.48, 1.35) | 0.36 (0.28, 0.44) | 0.05 (0.03, 0.06) |
| Malathion | 3.22 (2.90, 3.55) | 2.81 (2.21, 3.49) | 5.97 (4.45, 8.18) | 2.83 (1.97, 3.98) | 2.40 (1.88, 2.96) | 3.37 (2.70, 4.18) |
| Temephos | 0.12 (0.11, 0.13) | 0.15 (0.14, 0.17) | 0.13 (0.12, 0.14) | 0.07 (0.06, 0.07) | 0.16 (0.14, 0.17) | 0.04 (0.04, 0.04) |
| Propoxur | 4.78 (4.46, 5.12) | 2.65 (2.43, 2.88) | 3.83 (3.58, 4.09) | 0.89 (0.71, 1.10) | 1.48 (1.22, 1.78) | 1.14 (1.07, 1.23) |
| DDT | 3.72 (2.62, 5.44) | 3.42 (2.94, 3.98) | 1.92 (1.11, 3.41) | 4.96 (4.19, 5.89) | 3.46 (3.07, 3.89) | 2.41 (1.82, 3.28) |
| <i>Bti</i> | 0.55 (0.43, 0.68) | 0.60 (0.53, 0.70) | 0.82 (0.52, 1.92) | 0.99 (0.86, 1.20) | 0.95 (0.89, 1.02) | 0.81 (0.66, 1.03) |

LC₅₀ lethal concentration that kills 50% of the population (mg/L), values are LC₅₀ (95% CI).
Bti: 7000 ITU/mg.




Figure 2. The resistance of *Aedes albopictus* larvae to currently used insecticides (resistance ratio at 50% concentration). (A) Insecticide pyrethroids (cyfluthrin, deltamethrin, and permethrin); (B) organophosphate (malathion, temephos), organochlorine (DDT), carbamate (propoxur), and *Bti*. RR₅₀, resistance ratio, LC₅₀ test population/LC₅₀ laboratory-susceptible strain.




Figure 3. Mortality rates of adult *Aedes albopictus* after exposure to insecticides without (A) and with (B) pre-exposure to PBO. Error bars represent standard error of the mean. The bottom panel indicated significant level of difference in mortalities for the same insecticide between PBO and non-PBO pre-exposed populations from the same study site.

Table 3. *Kdr* mutation frequency relationship to pyrethroid resistance in different *Aedes albopictus* populations

| Study site | Phenotype | Sample size | Allele frequency (%) | | | Odds ratio (95% CI) | |
|------------|-------------|-------------|----------------------|--------|--------|------------------------------|--------------------------------|
| | | | Wildtype | F1534C | F1534S | F1534C | F1534S |
| Haikou | Resistant | 61 | 13.1 | 26.2 | 60.7 | 1.1 (0.6–2.1) ^{NS} | 0.9 (0.5–1.6) ^{NS} |
| | Susceptible | 50 | 13.0 | 24.0 | 63.0 | | |
| Qiongzong | Resistant | 14 | 10.7 | 3.6 | 85.7 | 1.5 (0.1–25.3) ^{NS} | 10.8 (3.2–37.0) ^{***} |
| | Susceptible | 21 | 61.9 | 2.4 | 35.7 | | |
| Sanya | Resistant | 36 | 16.7 | 26.4 | 56.9 | 1.5 (0.7–3.2) ^{NS} | 3.3 (1.7–6.4) ^{**} |
| | Susceptible | 42 | 52.4 | 19.0 | 28.6 | | |
| Wanning | Resistant | 48 | 30.2 | 4.2 | 65.6 | 0.6 (0.2–2.3) ^{NS} | 2.4 (1.4–4.2) ^{**} |
| | Susceptible | 56 | 49.1 | 6.4 | 44.5 | | |
| Danzhou | Resistant | 75 | 28.0 | 2.0 | 70.0 | NA | 2.4 (1.5–3.8) ^{***} |
| | Susceptible | 76 | 50.7 | 0.0 | 49.3 | | |

** $P < 0.01$.
*** $P < 0.001$.
NS, no significance.

Table 4. Summary of public health and agricultural use of pesticides based on questionnaire survey

| Site | Purpose | Environment | Insecticide | Percentage of households (%) | | | |
|-----------|---------------|-------------|-----------------------|------------------------------|--------|----------|---------|
| | | | | Never | Weekly | Biweekly | Monthly |
| Sanya | Public health | Indoor* | Pyrethroids | 3.3 | 96.7 | 0.0 | 0.0 |
| | Agricultural | Outdoor | Pyrethroids | 80.0 | 13.3 | 0.0 | 6.7 |
| | | | Organophosphate | 66.7 | 6.7 | 13.3 | 13.3 |
| Wanning | Public health | Indoor* | Microbial insecticide | 90.0 | 3.3 | 3.3 | 3.3 |
| | | | Pyrethroids | 10.0 | 36.7 | 33.3 | 20.0 |
| | Agricultural | Outdoor | Pyrethroids | 93.3 | 0.0 | 6.7 | 0.0 |
| | | | Organophosphate | 56.7 | 0.0 | 33.3 | 10.0 |
| Haikou | Public health | Indoor* | Nitromethyl | 86.7 | 0.0 | 6.7 | 6.7 |
| | | | Pyrethroids | 26.7 | 43.3 | 13.3 | 16.7 |
| | Agricultural | Outdoor | Carbamate | 93.3 | 0.0 | 3.3 | 3.3 |
| | | | Pyrethroids | 86.7 | 0.0 | 6.7 | 6.7 |
| Danzhou | Public health | Indoor* | Organophosphate | 73.3 | 0.0 | 16.7 | 10.0 |
| | | | Pyrethroids | 26.7 | 40.0 | 13.3 | 20.0 |
| | Agricultural | Outdoor | Organophosphate | 63.3 | 0.0 | 20.0 | 16.7 |
| | | | Nicotine insecticide | 86.7 | 0.0 | 10.0 | 3.3 |
| Qiongzong | Public health | Indoor* | Pyrethroids | 0.0 | 100.0 | 0.0 | 0.0 |
| | Agricultural | Outdoor | Pyrethroids | 56.7 | 0.0 | 3.3 | 40.0 |
| | | | Organophosphate | 66.7 | 0.0 | 0.0 | 33.3 |

Total number of households surveyed was 30 per site.
*Pyrethroid-based mosquito repellents.

resistance and the *kdr* mutation (Table 3). The F1534C mutation was not associated with pyrethroid resistance in any of the five populations (Table 2). The F1534L mutation was not detected at any study site. Genotyping of the *ace-1* gene in Haikou and Danzhou populations ($n = 120$), the two populations that exhibited the highest resistance to malathion, did not reveal any mutations at codon 119.

3.5 Local residents used all types of insecticides

Questionnaire surveys of local residents in the study area revealed that six classes of insecticides, i.e. pyrethroids, organophosphates, carbamates, microbial insecticide, nitromethyl, and nicotine insecticide, were used in Hainan Province (Table 4). Pyrethroids

were the most commonly used insecticide for public health and agricultural pest control, with 73.3–100.0% of the surveyed households reporting using different types of pyrethroid insecticides for mosquito control (Table 4). Organophosphates were primarily used for agricultural pest control, and carbamates were used only in the Haikou site for mosquito control among a small proportion of surveyed households (Table 4).

4 DISCUSSION

This study presents the first resistance profile of *Ae. albopictus* across Hainan Province. The results indicate that the majority of adult *Ae. albopictus* populations in Hainan are resistant or highly

resistant to the four classes of synthetic insecticides, especially the most commonly used pyrethroids. The occurrence of widespread insecticide resistance in *Ae. albopictus* adults potentially compromises the efficacy of current intervention measures in controlling *Ae. albopictus* populations on the island. While adult *Ae. albopictus* resistance to different insecticides has been reported in other places,¹⁵ *Ae. albopictus* adults in Hainan showed high resistance to all insecticides at all study sites. In Hainan province, all types of insecticides have been frequently used for mosquito control, especially for epidemic controls. These insecticides have also been widely used for agricultural pest controls, which can lead to resistance. Resistance management strategies are urgently needed to help mitigate failures in future vector control interventions.

The reemergence of dengue fever has been reported in many countries and often has been caused by invading *Ae. albopictus*.³⁶ Dengue epidemics were reported in Hainan island in the 1980s and 1990s after approximately 30 years of no reported local transmission. In 2019, Dengue fever outbreak occurred in Hainan. The high insecticide resistance in the primary vector *Ae. albopictus* adds another dimension of risk in dengue outbreak control for the near future. Currently, larval control is a key measure to control the mosquito population in Hainan. In Hainan, pyrethroids are widely used for agricultural pests and mosquito adult control but not larval control. However, we found that *Ae. albopictus* larvae are resistant to pyrethroid insecticides, which is consistent with other studies in China.^{19,37} This finding is likely caused by the residues left in the habitats from adult mosquito spraying or agricultural pest controls. However, we cannot rule out inherited resistance from adult mosquitoes, which needs further investigation. *Ae. albopictus* larvae remained sensitive to some other insecticides, including organophosphorus (malathion, temephos), propoxur, and the microbial larvicide *Bti*. However, it has been reported that *Aedes* larvae have developed high resistance to temephos in South-East Asia and South America.^{13,38–40} Therefore, *Aedes* larval resistance to insecticides must be closely monitored and integrated strategies should be developed to combat resistance.

Due to widespread multiple insecticide resistance in the dengue vector *Ae. albopictus* in Hainan Province, new strategies should be developed to enhance the efficacy of currently available insecticides and to slow down the development of insecticide resistance. The main strategies for insecticide resistance management, as recommended by WHO, include rotations, mosaics, and mixtures of insecticides among which there is no cross-resistance.⁴¹ For larval control, community-involved combinations of ecological, biological, and social strategies have been demonstrated to reduce *Aedes* mosquito abundance,^{42,43} i.e. combining larval source reduction with larvicides, especially microbial larvicides, may prevent further development of insecticide resistance in *Ae. albopictus*. Using PBO-insecticides for adult mosquito control, which combines PBO with insecticides and is widely used in the USA, is another potential option to slow down insecticide resistance.⁴⁴ However, to our knowledge PBO insecticide has not been tested in China. Our study indicated that pre-exposure to PBO could potentially fully recover the efficacy of some insecticides. In addition, field studies have indicated that insecticide-treated mosquito nets impregnated with PBO can significantly enhance the efficacy of standard net and reduces malaria infections.^{45,46} Therefore, adding PBO to insecticides may be an effective formulation for resistance management, and this strategy warrants further examination for *Ae. albopictus* control.

The *kdr* mutation in the VGSC genes is one indicator of mosquito resistance to pyrethroid insecticides.^{47,48} In malaria vector *Anopheles* mosquitoes, previous studies have found strong associations between *kdr* mutation frequency and phenotypic resistance status.^{49,50} However, this association does not necessarily hold true for all mosquito species. For example, the *kdr* mutation was not found in *An. funestus*⁵¹ nor in some populations of *An. sinensis*,⁵² even when WHO tube test mortality was very low (i.e. very high observed insecticide resistance). The *kdr* mutation, specifically the F1534 mutation, has been found in *Ae. albopictus* in China and elsewhere.^{21,22,53} Some studies found F1534 mutations to be associated with phenotypic resistance,^{54–56} but a separate study did not find this relationship.⁵⁷ Our results indicate that the F1534S mutation is associated with pyrethroid phenotypic resistance in *Ae. albopictus* in Hainan Province. However, *kdr* mutation frequency was high in both resistant and susceptible individuals, indicating that the F1534S mutation is not completely predictive of resistance status.⁵⁴ Similarly, although with very low mortality, resistant *Ae. albopictus* did not show higher F1534C mutations in comparison with susceptible mosquitoes, indicating that F1534C is also not likely to be a good indicator of resistance, which is consistent with other studies reported from China.⁵⁴ It is possible that there are regional differences in *kdr* mutations as shown in *An. sinensis*,^{54,56} but this hypothesis requires further investigation. In this study, no F1534L mutations were detected, which is different from what has been reported from nearby Guangdong province.¹³ This finding may be due to the geographic variation influenced by either environmental conditions or insecticide usage.

Ace-1 mutations have been found to confer insensitivity to organophosphates and carbamates in mosquito populations.⁴⁹ Our study did not detect any *G119* mutations in *ace-1* in *Ae. albopictus* populations, which suggests either that *Ae. albopictus* in Hainan lacks mutations at the target sites for these insecticides or that the mutation frequency was extremely low. Other studies also found that resistance to *ace-1* agonists is common in *Aedes*, but that *ace-1* mutations do not occur widely.^{13,21,58–60} Interestingly, exposure to the PBO synergist increased *Ae. albopictus* sensitivity to pyrethroids, DDT, and malathion, which suggests that metabolic resistance may play an important role in resistance. However, in high permethrin-, DDT-, and malathion-resistant *Ae. albopictus* populations, PBO exposure only partially recovered insecticide efficacy and mosquito mortality rate, which suggests that other mechanisms may also contribute to insecticide resistance in these populations.^{61,62}

Although, in general, *Ae. albopictus* populations were resistant to insecticides, it is worth noting that resistance levels varied between study sites and between different insecticides. For example, *Ae. albopictus* populations were highly resistant to permethrin and DDT in all study sites although with variations in mortality, but resistance to deltamethrin in Qiongzong was very low. These differences may be due to the differences in the usages of specific insecticide in local residents, as some insecticides may be more frequently used in certain areas but not in others. For example, in Danzhou, farmers used organophosphate twice a month. However, farmers in Qiongzong used the same type of insecticide monthly. In Qiongzong, all the surveyed households used pyrethroid for household pest controls, while only about half of the residents in Haikou did the same. These differences in insecticide usage in different sites may have led to variation in resistance levels in different *Ae. albopictus* populations. Consequently, the mechanisms of the resistance need to be further investigated.

5 CONCLUSION

Aedes mosquito resistance to insecticides is influenced by a complex interplay of factors, including environmental factors and intervention strategies. The development of insecticide resistance to multiple insecticides which has occurred in Hainan will complicate the current vector control strategies. Nonetheless, the findings from this study can help in guiding the selection of insecticides for *Aedes* mosquito control. First, high pyrethroid resistance calls for the use of alternative insecticides. Second, PBO insecticides may be an option to mitigate the negative impact of pyrethroid resistance. Finally, to achieve effective vector management, a public health response beyond routine larviciding or focal spraying is essential. Targeted vector management can make a difference in terms of reducing vector abundance and controlling mosquito-borne infectious disease outbreaks.

ACKNOWLEDGEMENTS

We wish to thank Faxing Fu, Lirong Wu, Mengyuan Lin, and Haoze Yang for their assistance with mosquito collection and resistance bioassay, and Qiang Zhang, Peilin Guo, Yitao Li, Tuquan Zheng, Xiaofang Lin, and Na Li for assistance with the insecticide use survey. This study was supported by the National Natural Science Foundation of China (81660345, 82060379 and 31760318), the Talent Introduction Fund of Hainan Medical University (2016011), the Guizhou Science and Technology Foundation ([2019]1441 and [2018]5779-17), the Jiangsu Natural Science Foundation (BK20180994), the Postdoctoral Fellowship of China (2018 M632382), and a China Scholarship Council fellowship (201808460069).

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

SUPPORTING INFORMATION

Supporting information may be found in the online version of this article.

REFERENCES

- Kraemer MU, Sinka ME, Duda KA, Mylne AQ, Shearer FM, Barker CM *et al.*, The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. *Elife* **4**:e08347 (2015).
- Gratz NG, Critical review of the vector status of *Aedes albopictus*. *Med Vet Entomol* **18**:215–227 (2014).
- Bonizzoni M, Gasperi G, Chen X and James AA, The invasive mosquito species *Aedes albopictus*: current knowledge and future perspectives. *Trends Parasitol* **29**:460–468 (2013).
- Lwande OW, Obanda V, Lindström A, Ahlm C, Evander M, Näslund J *et al.*, Globe-trotting *Aedes aegypti* and *Aedes albopictus*: risk factors for arbovirus pandemics. *Vector Borne Zoonotic Dis* **20**:71–81 (2020).
- World Health Organization, Ten threats to global health in 2019, Retrieved November 2019 (2019).
- Mousson L, Zouache K, Arias-Goeta C, Raquin V, Mavingui P and Failloux AB, The native *Wolbachia* symbionts limit transmission of dengue virus in *Aedes albopictus*. *PLoS Neglected Trop Dis* **6**:e1989 (2012).
- Zhao C, Zhu C, Kai W, Liu G, Liu Q, Lin L *et al.*, Genotypes of knockdown resistance gene and their distribution in *Aedes albopictus* in Haikou, China, in 2018. *Chin J Vector Biol Control* **30**:7–11 (2019).
- Global Invasive Species Database, 100 of the World's Worst Invasive Alien Species (2021). Available: http://www.iucngisd.org/gisd/100_worst.php. [2 July 2020].
- Li Y, Kamara F, Zhou G, Puthiyakunnon S, Li C, Liu Y *et al.*, Urbanization increases *Aedes albopictus* larval habitats and accelerates mosquito development and survivorship. *PLoS Neglected Trop Dis* **8**:e3301 (2014).
- Peng H, Lai H, Zhang Q, Xu B, Zhang H, Liu W *et al.*, A local outbreak of dengue caused by an imported case in Dongguan China. *BMC Public Health* **12**:83 (2012).
- Wu D, Wu J, Zhang Q, Zhong H, Ke C, Deng X *et al.*, Chikungunya outbreak in Guangdong Province, China, 2010. *Emerg Infect Dis* **18**:493–495 (2012).
- Wei Y, Wang J, Song Z, He Y, Zheng Z, Fan P *et al.*, Patterns of spatial genetic structures in *Aedes albopictus* (Diptera: Culicidae) populations in China. *Parasit Vectors* **12**:552 (2019).
- Su X, Guo Y, Deng J, Xu J, Zhou G, Zhou T *et al.*, Fast emerging insecticide resistance in *Aedes albopictus* in Guangzhou, China: alarm to the dengue epidemic. *PLoS Neglected Trop Dis* **13**:e0007665 (2019).
- Zheng X, Zhong D, He Y and Zhou G, Seasonality modeling of the distribution of *Aedes albopictus* in China based on climatic and environmental suitability. *Infect Dis Poverty* **8**:98 (2019).
- Moyes CL, Vontas J, Martins AJ, Ng LC, Koo SY, Dusfour I *et al.*, Contemporary status of insecticide resistance in the major *Aedes* vectors of arboviruses infecting humans. *PLoS Neglected Trop Dis* **11**:e0005625 (2017).
- Roiz D, Wilson AL, Scott TW, Fonseca DM, Jourdain F, Müller P *et al.*, Integrated *Aedes* management for the control of *Aedes*-borne diseases. *PLoS Neglected Trop Dis* **12**:e0006845 (2018).
- Marcombe S, Chonephetsarath S, Thammavong P and Brey PT, Alternative insecticides for larval control of the dengue vector *Aedes aegypti* in Lao PDR: insecticide resistance and semi-field trial study. *Parasit Vectors* **11**:616 (2018).
- Liu S, Cui F, Yan S and Qiao C, Investigation of organophosphate and pyrethroid resistance in vector mosquitoes in China. *Chin J Vector Biol Control* **22**:184–189 (2011).
- Zhao C, Zhu C, Jia Q, Yan D, Liu G, Wu H *et al.*, Resistance of *Aedes albopictus* to commonly used insecticides in different areas of China, 2017–2018. *Chin J Vector Biol Control* **30**:7–11 (2020).
- Corbel V, Fonseca DM, Weetman D, Pinto J, Achee NL, Chandre F *et al.*, International workshop on insecticide resistance in vectors of arboviruses, December 2016, Rio de Janeiro, Brazil. *Parasites Vectors* **10**:278 (2017).
- Li Y, Xu J, Zhong D, Zhang H, Yang W, Zhou G *et al.*, Evidence for multiple-insecticide resistance in urban *Aedes albopictus* populations in southern China. *Parasites Vectors* **11**:4 (2018).
- Chen H, Li K, Wang X, Yang X, Lin Y, Cai F *et al.*, First identification of *kdr* allele F1534S in VGSC gene and its association with resistance to pyrethroid insecticides in *Aedes albopictus* populations from Haikou City, Hainan Island, China. *Infect Dis Poverty* **5**:31 (2016).
- Wei S, Lin S and Yang C, The 50-year history of the development of health in Hainan Province, Haikou: *Southern Press* 396–7 (2007). [in Chinese]
- The People's Government of Hainan Province (2019). Available: <http://www.hnntv.cn/st/jrrd/2019-09-10/350737.html> [10 September 2019]
- Zhong S, Xie W, Chen J, Fu F, Peng Y, Wu L *et al.*, Knowledge, attitude and practice concerning control of mosquito vectors in residents of Haikou. *China Trop Med* **19**:731–737 (2019).
- Liu Q, Dengue fever in China: new epidemical trend, challenges and strategies for prevention and control. *Chin J Vector Biol Control* **31**:1–6 (2020).
- Gong D and Zhou H, Progress in dengue fever important vector *Aedes albopictus* in China. *Chin J Vector Biol Control* **20**:607–610 (2009).
- He C, Zhao W, Wang S, Zeng L, Li S and Ou T, Analysis of mosquito density and seasonality in urban areas of Hainan province, China in 2012. *Chin J Vector Biol Control* **25**:15–17 (2014).
- Su A, Pei Z, Fu J, Li J, Feng F, Zhu Q *et al.*, Analysis of distribution and population density changes of *Aedes aegypti* the transmission vector of dengue fever in Haikou City. *China Trop Med* **06**:1394–1395 (2005).
- World Health Organization, *Guidelines for Laboratory and Field Testing of Mosquito Larvicides*. WHO, Geneva (2005).
- World Health Organization, *Monitoring and Managing Insecticide Resistance in Aedes Mosquito Populations: Interim Guidance for Entomologists*. WHO, Geneva (2016).
- World Health Organization, *Test Procedures for Insecticide Resistance Monitoring in Malaria Vector Mosquitoes*. WHO, Geneva (2016).

- 33 Chang X, Zhong D, Li X, Huang Y, Zhu G, Wei X *et al.*, Analysis of population genetic structure of *Anopheles sinensis* based on mitochondrial DNA cytochrome oxidase subunit I gene fragment. *Nan Fang Yi Ke Da Xue Xue Bao* **35**:234–238, 247 (2015).
- 34 Kasai S, Ng LC, Lam-Phua SG, Tang CS, Itokawa K, Komagata O *et al.*, First detection of a putative knockdown resistance gene in major mosquito vector, *Aedes albopictus*. *Jpn J Infect Dis* **64**:217–221 (2011).
- 35 Finney D, *Probit Analysis*. Cambridge University Press, Cambridge (1971).
- 36 Rezza G, *Aedes albopictus* and the reemergence of dengue. *BMC Public Health* **12**:72 (2012).
- 37 Wang Y, Liu X, Li C, Su T, Jin J, Guo Y *et al.*, A survey of insecticide resistance in *Aedes albopictus* (Diptera: Culicidae) during a 2014 dengue fever outbreak in Guangzhou, China. *J Econ Entomol* **110**:239–244 (2017).
- 38 Saeung M, Ngoen-Klan R, Thanispong K, Muenworn V, Bangs MJ and Chareonviriyaphap T, Susceptibility of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) to Temephos in Thailand and surrounding countries. *J Med Entomol* **57**:1207–1220 (2020).
- 39 Valle D, Bellinato DF, Viana-Medeiros PF, Lima JBP and Martins Junior AJ, Resistance to temephos and deltamethrin in *Aedes aegypti* from Brazil between 1985 and 2017. *Mem Inst Oswaldo Cruz* **114**:e180544 (2019).
- 40 Elia-Amira NMR, Chen CD, Lau KW, Lee HL, Low VL, Norma-Rashid Y *et al.*, Organophosphate and organochlorine resistance in larval stage of *Aedes albopictus* (Diptera: Culicidae) in Sabah, Malaysia. *J Econ Entomol* **111**:2488–2492 (2018).
- 41 Dusfour I, Vontas J, David JP, Weetman D, Fonseca DM, Corbel V *et al.*, Management of insecticide resistance in the major *Aedes* vectors of arboviruses: advances and challenges. *PLoS Neglected Trop Dis* **13**:e0007615 (2019).
- 42 Quintero J, Brochero H, Manrique-Saide P, Barrera-Pérez M, Basso C, Romero S *et al.*, Ecological, biological and social dimensions of dengue vector breeding in five urban settings of Latin America: a multi-country study. *BMC Infect Dis* **14**:38 (2014).
- 43 Arunachalam N, Tana S, Espino F, Kittayapong P, Abeyewickreme W, Wai KT *et al.*, Eco-bio-social determinants of dengue vector breeding: a multicountry study in urban and periurban Asia. *Bull World Health Organ* **88**:173–184 (2010).
- 44 Dadzie SK, Chabi J, Asafu-Adjaye A, Owusu-Akrofi O, Baffoe-Wilmot A, Malm K *et al.*, Evaluation of piperonyl butoxide in enhancing the efficacy of pyrethroid insecticides against resistant *Anopheles gambiae* s.l. in Ghana. *Malar J* **16**:342 (2017).
- 45 Oumbouke WA, Rowland M, Koffi AA, Alou LPA, Camara S and N'Guessan R, Evaluation of an alpha-cypermethrin + PBO mixture long-lasting insecticidal net VEERALIN® LN against pyrethroid resistant *Anopheles gambiae* s.s.: an experimental hut trial in M'bé, central Côte d'Ivoire. *Parasit Vectors* **12**:544 (2019).
- 46 Gleave K, Lissenden N, Richardson M, Choi L and Ranson H, Piperonyl butoxide (PBO) combined with pyrethroids in insecticide-treated nets to prevent malaria in Africa. *Cochrane Database Syst Rev* **11**:CD012776 (2018).
- 47 Soderlund DM and Knipple DC, The molecular biology of knockdown resistance to pyrethroid insecticides. *Insect Biochem Mol Biol* **33**:563–577 (2003).
- 48 Fang Y, Shi W, Wu J, Li Y, Xue J, Zhang Y *et al.*, Resistance to pyrethroid and organophosphate insecticides, and the geographical distribution and polymorphisms of target-site mutations in voltage-gated sodium channel and acetylcholinesterase 1 genes in *Anopheles sinensis* populations in Shanghai, China. *Parasites Vectors* **12**:396 (2019).
- 49 Liu N, Insecticide resistance in mosquitoes: impact, mechanisms, and research directions. *Annu Rev Entomol* **60**:537–559 (2015).
- 50 Hemming-Schroeder E, Strahl S, Yang E, Nguyen A, Lo E, Zhong D *et al.*, Emerging pyrethroid resistance among *Anopheles arabiensis* in Kenya. *Am J Trop Med Hyg* **98**:704–709 (2018).
- 51 Coetzee M and Koekemoer LL, Molecular systematics and insecticide resistance in the major African malaria vector *Anopheles funestus*. *Annu Rev Entomol* **58**:393–412 (2013).
- 52 Zhong D, Chang X, Zhou G, He Z, Fu F, Yan Z *et al.*, Relationship between knockdown resistance, metabolic detoxification and organismal resistance to pyrethroids in *Anopheles sinensis*. *PLoS One* **8**:e55475 (2013).
- 53 Liu H, Liu L, Cheng P, Yang L, Chen J, Lu Y *et al.*, Bionomics and insecticide resistance of *Aedes albopictus* in Shandong, a high latitude and high-risk dengue transmission area in China. *Parasites Vectors* **13**:11 (2020).
- 54 Gao J, Chen H, Shi H, Peng H and Ma Y, Correlation between adult pyrethroid resistance and knockdown resistance (*kdr*) mutations in *Aedes albopictus* (Diptera: Culicidae) field populations in China. *Infect Dis Poverty* **7**:86 (2018).
- 55 Xu J, Bonizzoni M, Zhong D, Zhou G, Cai S, Li Y *et al.*, Multi-country survey revealed prevalent and novel F1534S mutation in voltage-gated sodium channel (VGSC) Gene in *Aedes albopictus*. *PLoS Neglected Trop Dis* **10**:e0004696 (2016).
- 56 Rath A, Mohanty I and Hazra RK, Insecticide susceptibility status of invasive *Aedes albopictus* across dengue endemic districts of Odisha, India. *Pest Manag Sci* **74**:1431–1440 (2018).
- 57 Kushwah RB, Mallick PK, Ravikumhar H, Dev V, Kapoor N, Adak TP *et al.*, Status of DDT and pyrethroid resistance in Indian *Aedes albopictus* and absence of knockdown resistance (*kdr*) mutation. *J Vector Borne Dis* **52**:95–98 (2015).
- 58 Grigoraki L, Lagnel J, Kiousoulis I, Kampouraki A, Morou E, Labbe P *et al.*, Transcriptome profiling and genetic study reveal amplified carboxylesterase genes implicated in temephos resistance, in the Asian tiger mosquito *Aedes albopictus*. *PLoS Neglected Trop Dis* **9**:e0003771 (2015).
- 59 Grisales N, Poupardin R, Gomez S, Fonseca-Gonzalez I, Ranson H and Lenhart A, Temephos resistance in *Aedes aegypti* in Colombia compromises dengue vector control. *PLoS Neglected Trop Dis* **7**:e2438 (2013).
- 60 Marcombe S, Poupardin R, Darriet F, Reynaud S, Bonnet J, Strode C *et al.*, Exploring the molecular basis of insecticide resistance in the dengue vector *Aedes aegypti*: a case study in Martinique Island (French West Indies). *BMC Genomics* **10**:494 (2009).
- 61 Kamgang B, Yougang AP, Tchoupo M, Riveron JM and Wondji C, Temporal distribution and insecticide resistance profile of two major arbovirus vectors *Aedes aegypti* and *Aedes albopictus* in Yaoundé, the capital city of Cameroon. *Parasites Vectors* **10**:469 (2017).
- 62 Chang X, Zhong D, Fang Q, Hartsel J, Zhou G, Shi L *et al.*, Multiple resistances and complex mechanisms of *Anopheles sinensis* mosquito: a major obstacle to mosquito-borne diseases control and elimination in China. *PLoS Neglected Trop Dis* **8**:e2889 (2014).