

Lawrence Berkeley National Laboratory

Recent Work

Title

Yields of Manganese in Spallation Reactions

Permalink

<https://escholarship.org/uc/item/39j9263k>

Authors

Bonner, Norman A.
Orr, William C.

Publication Date

1949-04-19

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-331
Unclassified Distribution

Copy 2

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

UNCLASSIFIED

YIELDS OF MANGANESE IN SPALLATION REACTIONS

Norman A. Bonner and William C. Orr

April 19, 1949

Berkeley, California

YIELDS OF MANGANESE IN SPALLATION REACTIONS

Norman A. Bonner* and William C. Orr
Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

* Present address: Department of Chemistry, Washington University, St. Louis, Mo.

One approach previously used in the study of high energy spallation reactions involves the bombardment of a single target isotope and the determination of the relative yields of many products.¹ In the present work a complementary approach

-
- (1) B. B. Cunningham *et al*, Phys. Rev. 72, 739 (1947); H. H. Hopkins, Jr., and B. B. Cunningham, Phys. Rev. 73, 1406 (1943).
-

has been used: the bombardment of a variety of elements and the determination of the yields of two particular product isotopes. This method has the advantage that yields can be compared more exactly, since the same radiation is always measured. It also has the disadvantage that most of the target elements have several stable isotopes, so that the reacting nucleus cannot be uniquely specified. All of the elements (excepting only rubidium and krypton) from atomic number 24 (chromium) to 38 (strontium) were bombarded with 190-Mev deuterons in the 134-inch cyclotron. The yields of 5.3-day Mn⁵² and 2.59-hr. Mn⁵⁶ were determined.

After each bombardment the target was dissolved and a known amount of inactive manganese was added, together with carriers for the other elements possibly produced. The manganese was separated chemically and the decay of its activity was followed. The chemical procedures used were varied depending on the target element, more complicated procedures being required for those far from manganese because of the low yields. The purest available materials (usually Hilger's "spectroscopically pure" grade) were used for targets.

In order to compare the yields from one bombardment with those from another, a monitor target was included in each experiment. A piece of 10 mil copper foil

cut to the exact shape and area of the target was placed immediately behind the target during exposure to the deuteron beam. Since all targets used were thin, it was assumed that each copper monitor foil received the same irradiation as the accompanying target. The yield of Cu^{64} in the monitor was taken to be a measure of the relative intensity of the deuteron beam.

The measured activities were corrected to infinite bombardment time and 100% chemical yield. In addition, the counting efficiency of Mn^{52} was assumed to be 40% relative to that of Mn^{56} and corrected accordingly. (Mn^{52} decays 35% by positron emission, 65% by K capture²; the 40% figure includes the contribution of

(2) W. M. Good, D. Peasloe, and M. Deutsch, Phys. Rev. 69, 313 (1946).

electromagnetic radiation to the counting rate.)

The data are presented in Fig. 1. The relative cross section, σ^- , is equal to the corrected manganese activity per mole of target element divided by the corrected Cu^{64} activity per mole of copper monitor. The uncertainty attached to each determination is indicated by the size of the plotted point.

It should be noted that the yield of the 21-minute isomer of Mn^{52} which decays³

(3) R. K. Osborne and M. Deutsch, Phys. Rev. 71, 467 (1947).

directly to Cr^{52} has not been included. The total yields of Mn^{52} are therefore somewhat higher than the data indicate.

The absolute cross section corresponding to $\sigma^- = 1$ is approximately 0.03 barn, according to a recent direct determination⁴ of the cross section for Cu^{64} .

(4) R. Batzel, private communication.

It is of interest that the fluctuations by which the data deviate from smooth curves can be correlated with the distribution of stable isotopes present in the

natural elements used as targets, together with the variations in excitation energy required for reaction of the different isotopes. The excitation energies were estimated with the assumption that alpha-particles are favored over other charged particles in their evaporation from excited nuclei, and the agreement obtained affords some support for this hypothesis.

The cooperation of the 184-inch cyclotron crew in performing the bombardments is gratefully acknowledged. This work was performed under the auspices of the United States Atomic Energy Commission.

Legend for Figure 1: Relative cross sections for the production of two manganese isotopes by bombardment of various elements with 190-Mev deuterons. Points with arrows attached represent upper limits.

