

Lawrence Berkeley National Laboratory

Lawrence Berkeley National Laboratory

Title

FISSION OF 238U INDUCED BY INELASTIC SCATTERING OF 120 MeV 4-PARTICLES

Permalink

<https://escholarship.org/uc/item/3bc3k6rd>

Author

Back, B.B.

Publication Date

1980-09-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Submitted to Physical Review C

FISSION OF ^{238}U INDUCED BY INELASTIC SCATTERING
OF 120 MeV α -PARTICLES

B.B. Back, A.C. Shotter, T.J.M. Symons, A. Bice,
C.K. Gelbke, T.C. Awes, and D.K. Scott

September 1980


TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782.*

RECEIVED
LAWRENCE
BERKELEY LABORATORY

JAN 8 1981

LIBRARY AND
DOCUMENTS SECTION



LBL-11852C.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Fission of ^{238}U induced by inelastic scattering
of 120 MeV α -particles

B. B. Back

Argonne National Laboratory, Argonne, Illinois 60439

A. C. Shotter,[†] T. J. M. Symons, A. Bice

Lawrence Berkeley Laboratory, Berkeley, California 94720

C. K. Gelbke, T. C. Awes, and D. K. Scott

Michigan State University, East Lansing, Michigan 48824

and

Lawrence Berkeley Laboratory, Berkeley, California 94720

PACS numbers: 25.85.Ge, 25.60.Cy, 24.30.Cz

Present address: Physics Department, University of
Edinburgh, Edinburgh EH93JZ, United Kingdom

This work was supported by the U.S. Department of Energy under Contract W-7405-ENG-48.

This manuscript was printed from originals provided by the author.

The submitted manuscript has been authored by a contractor of the U. S. Government under contract No. W-31-109-ENG-38. Accordingly, the U. S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U. S. Government purposes.

Abstract

The fission decay of ^{238}U has been measured as function of excitation energy in inelastic scattering of 120 MeV α -particles. Total kinetic energies and masses of fission fragments were measured by the double energy method. It is observed that the total kinetic energy E_K decreases and that the valley in the mass distribution is reduced when the excitation energy of the system is increased. No indication of anomalous total kinetic energy release in the region of the giant quadrupole resonance has been found. A qualitative interpretation of the data is given on the basis of a static scission point model.

I. INTRODUCTION
~~~~~

One of the long standing questions in fission research is concerned with the descent from the saddle point to scission. Whether the available energy is transferred into internal degrees of freedom or into relative motion of the two nascent fragments prior to scission is still largely unknown, since measurements of final parameters of the reaction do not directly address the question of the dynamical evolution of the fission decay.

However, some success in explaining many of the features of fission has been obtained with a static model<sup>1</sup> of the scission point configuration, which takes into account the shell structure in the nascent fragment. Most prominently, it appears that the deformed shell for  $A=138$  is responsible for the strongly asymmetric mass distributions in fission of most actinide elements. A strong spherical shell closure at  $Z=50, N=82$  is believed to be associated with the decrease of the total kinetic energy with excitation energy as observed in several light ion and neutron induced reactions on uranium and plutonium targets.<sup>2-6</sup>

To further study this latter correlation, we have measured the fission decay of  $^{238}\text{U}$  induced by the  $(\alpha, \alpha')$  reaction. We find that the total kinetic energy decreases with excitation energy at a rate of  $dE_K/dE^* = 0.38 \pm 0.07$  and that the effect is concentrated in the heavy fragment mass region of  $M_H = 125-135$ .

## II. EXPERIMENTAL PROCEDURE AND DATA ANALYSIS

A beam of 120 MeV  $\alpha$ -particles from the 88"-cyclotron at Lawrence Berkeley Laboratory was used to bombard a  $530 \mu\text{g}/\text{cm}^2$  thick, self-supporting metallic  $^{238}\text{U}$  target. Inelastically scattered  $\alpha$ -particles were identified in a triple telescope of solid state detectors located at  $\theta=16^\circ$  out of the horizontal plane. Two fission detectors were placed at  $\theta=90^\circ$  and  $\theta = -90^\circ$  in the horizontal plane, each subtending a solid angle of  $\Omega_f \approx 440 \text{ msr}$ . Coincident events between the telescope and the two fission detectors were recorded on magnetic tape for subsequent analysis.

The elements of the detector telescope were gain matched by introducing a calibrated charge pulse on the detector side of the pre-amplifiers. The energy calibration was obtained from the elastic peak in the  $\alpha$ -spectrum. The fission detectors were energy calibrated with the fission fragments from a thin  $^{252}\text{Cf}$  source.<sup>7</sup>

The data were analyzed off-line by an event-by-event reconstruction of the kinematics. Identification of  $\alpha$ -particles in the telescope was obtained by generating a particle identification spectrum from the measured pulse heights in each element in the detector telescope and selecting events in the peak corresponding to  $\alpha$ -particles. The primary energies and masses of the fission fragments were calculated by an iterative procedure taking into account the emission of neutrons from the fragments and the mass dependent pulse height defect of the detector response. The mass dependent neutron emission function measured for  $^{235}\text{U}(n,f)$

was used.<sup>8</sup> The excitation energy dependence of the neutron emission was ignored in the present analysis. The effects of this emission on the final results are discussed in a following paragraph. The main contributions to the mass resolution of the experiment came from the intrinsic detector resolution and the large solid angle of the fission detectors, which combined gave an estimated mass resolution of  $\Delta M=5$ . The experimental mass spectra have been unfolded assuming this mass resolution. The data have been corrected for accidental coincidences, which amount to  $\sim 10\%$  in the energy region above the fission threshold. Because a metallic self-supporting target was used, we are not faced with the problem of strong accidental peaks from scattering off target impurities. Due to the out-of-plane geometry used in the present experiment, there is no contribution from coincidences with recoil nuclei from light target contaminants.



### III. RESULTS AND DISCUSSION

The total kinetic energy release,  $E_K$ , as a function of excitation energy in the fissioning nucleus, and the spectrum of inelastically scattered  $\alpha$ -particles in coincidence with fission are shown in Fig. 1. We observe a strong increase in the fission coincidence rate up to the threshold for the neutron emission  $B_n$ , beyond which the competition from neutron emission introduces a sharp decrease. The increase in the fission yield due to second and third chance fission is clearly visible in the data.

In the energy region from 5-12 MeV, where only first chance fission is allowed, a strong decrease in the average total kinetic energy  $\bar{E}_K$  is observed. Rather sharp increases in  $\bar{E}_K$  are also observed above the thresholds for second and even third chance fission. These increases in  $\bar{E}_K$  most likely reflect the fact that a large fraction of the fission events above these thresholds come from near barrier fission of  $^{237}\text{U}$  and  $^{236}\text{U}$ , respectively, which have high total kinetic energies. At excitation energies above  $\sim 25$  MeV a weak increase in the total kinetic energy with excitation energy is seen.

The best fit to the total kinetic energy in the excitation energy region  $E^* = 5-12$  MeV is represented by a solid line with a slope of  $dE_K/dE^* = -0.46 \pm 0.07$  in Fig. 1. When corrected for the increase in neutron evaporation with excitation energy, this result is reduced to  $dE_K/dE^* = -0.38 \pm 0.07$ . This value is significantly lower than the result of David *et al.*<sup>5</sup> who find a

slope of  $dE_K/dE^* = -0.8 \pm 0.3$ . The origin of this discrepancy is not known at present. Decreasing total kinetic energies have been observed in several other uranium<sup>5</sup> and plutonium<sup>2-4</sup> nuclei, although there is a sizable spread in the values of  $dE_K/dE^*$  derived from various experiments. Negative slopes  $dE_K/dE^*$  have also been found in several Ac-isotopes,<sup>6</sup> however, only for the asymmetric component of the distribution.

A rather strong excitation probability of the giant quadrupole resonance has been observed in the singles  $^{238}\text{U}(\alpha, \alpha')$  reaction<sup>11</sup> under experimental conditions almost identical to the ones of the present study. Although there is still some question about the fission probability of the giant quadrupole resonance,<sup>9-12</sup> it appears that it is comparable to that of the underlying background, (which is  $P_f \sim 0.2$ ), which indicates that a substantial fraction of fission events in the energy region  $E^* \sim 8-12$  MeV can be associated with the giant quadrupole resonance. However, this does not appear to have any observable effect on the total energy release, which has a dependence on excitation energy very similar to what is observed in other reactions where the giant quadrupole resonance is not excited, e.g.,  $^{239}\text{Pu}(n, f)$ <sup>4</sup> and  $^{239}\text{Pu}(d, pf)$ .<sup>3</sup>

The slope  $dE_K/dE^*$  over the excitation energy region  $E^* = 5-12$  MeV is plotted as a function of the mass of the heavy fission fragments in Fig. 2a. A clear mass dependence of the effect is apparent, most of the effect being correlated with masses near  $M_H = 128-130$ . The total kinetic energy averaged over the same excitation energy region also displays a strong dependence on the fission fragment mass, with the maximum occurring at  $M_H \sim 130$  as

shown in Fig. 2b. However, the most abundant masses of heavy fission fragments are found at  $M_H \approx 138$ , (see Fig. 2c), which is distinctly different from the peaks in the  $dE_K/dE^*$  and  $\bar{E}_K$  data.

The dashed curve in Fig. 2b represents a calculation of the Coulomb repulsion energy between two coaxial ellipsoids each deformed to a semiaxis ratio of  $b/a = 0.6$  and separated by  $d = 2$  fm in order to simulate a neck formation between the two fragments. This comparison was performed only to show the expected dependence on fragment mass for a fixed shape of the nascent fragments. The parameters used are chosen only to give an approximately correct magnitude of the total kinetic energy and should not be considered as measured quantities.

Some qualitative insight into these shell effects may be obtained from a simple static model of the scission point configurations, which includes the effects of nuclear shells in the nascent fragments as proposed by Wilkins *et al.*<sup>1</sup> It appears that the large values of  $\bar{E}_K$  and  $d\bar{E}_K/dE^*$  found in the region  $M_H = 130$ , are correlated with the doubly magic shell for spherical shapes at  $Z=50$  and  $N=82$ . Spherical fragment shapes at scission lead to large kinetic energies due to increased Coulomb repulsion. Although this minimum in the shell energy is quite deep, it is not the main factor in determining the fission mass distribution, which peaks around  $M_H \approx 138$ . As is the case for most actinide nuclei in the region from Th to Cf, the mass distribution can be explained at least qualitatively by the occurrence of a strong deformed neutron shell at  $N=88$ . The strong Coulomb repulsion between the two nascent fission fragments at scission clearly

favors the somewhat weaker minimum in the shell energy surface at large deformation over the strong spherical minimum for the heavy fragment. Studying the  $\bar{E}_K$  dependence on the excitation energy  $E^*$ , we see that the  $A=138$  shell is very stable, whereas the  $A=132$  spherical shell rapidly disappears with excitation energy. A possible explanation of this phenomenon is that as the excitation energy is raised above the fission barrier, breaking of pairs occurs, which increases the viscous heating during the descent towards scission. Because the spherical minimum in the shell energy occurring at  $N=82$  is superimposed on a steep slope of the liquid drop potential energy surface, this minimum is rapidly destroyed with excitation energy. Contrasting this, the deformed shell at  $N=88$  has a substantial stability against excitation energy because it coincides in deformation with the minimum in the liquid drop potential energy surface (see Ref. 1).

This stability of the  $A=138$  deformed shell is also apparent in Fig. 3, where the fission fragment mass distributions are shown for several excitation energy bins. The strongly asymmetric character is preserved at least up to apparent excitation energies of 50 MeV. However, the excitation energy may not be as high as the energy of the outgoing  $\alpha$ -particle would indicate, because of 1) pre-fission neutron emission and 2) contribution from the  $(\alpha, {}^5\text{He})$  reaction, which is identified as a  $(\alpha, \alpha)$ -reaction, due to break-up of the  ${}^5\text{He}$ -ejectile.<sup>13</sup>

This interpretation is supported by the comparison with mass distribution data obtained in the  $^{238}\text{U}(\alpha, f)$  reaction,<sup>14</sup> as illustrated in Fig. 4. For the  $^{238}\text{U}(\alpha, \alpha f)$  reaction we observe an increasing abundance of symmetric mass splits with increasing excitation energy to  $E^* \sim 30$  MeV, beyond which point the peak-to-valley ratio remains almost constant. This saturation effect is not observed in fission of  $^{242}\text{Pu}$  induced by the  $^{238}\text{U}(\alpha, f)$  reaction,<sup>14</sup> where the fusion-fission process clearly dominates.

The standard deviations  $\sigma E_K$  of the total kinetic energy distributions are shown in Fig. 5 as a function of mass for different excitation energy intervals. A strong decrease in  $\sigma E_K$  is observed for increasing mass of the heavy fragment, especially in the lower excitation energy bins. The large widths of the  $E_K$  distribution at  $M_H = 120-125$  is correlated with coexistence of three competing neutron shells for  $N \sim 80$  with very different deformations of  $\beta = 0.1, 0.5, 0.85$ , respectively.<sup>1</sup> In the region of heavy fragment masses from  $M_H = 138-160$  a strong neutron shell at  $N=88$  with a deformation of  $\beta=0.65$  is dominating the scission point shapes and resulting in narrower total kinetic energy distributions, which is also observed in the data. Again we observe that these shell effects are somewhat reduced with excitation energy.

A static description of the scission point configuration as implied in the above discussion is certainly an oversimplification. However, even a dynamical description of the descent from saddle to scission must include the shell effects in the nascent fragments, and we therefore expect that such a calculation would lead to conclusions qualitatively similar to the ones presented here.

#### IV. CONCLUSIONS

~~~~~

From the present study of fission of ^{238}U induced by inelastic scattering of α -particles we conclude that the effect of nuclear shells in the nascent fragments at the scission point are of major importance for both the kinetic energy release and the mass division in fission. Although a strong deformed neutron shell at $N=88$ forces the nucleus to split preferentially into fragments of mass $A=138$ and $A=100$, we observe that the disappearance of a somewhat weaker (when imposed on the repulsive Coulomb potential) spherical shell at $A=132$ lead to a strong decrease of the total kinetic energy, when the nucleus is excited above the fission threshold.

Although it is known that a substantial fraction of the inelastic cross section at $E^* \sim 10$ MeV goes into excitation of the giant quadrupole resonance,¹² we have no evidence of anomalous kinetic energy release in this region. On the contrary, the observed slope of $d\bar{E}_K/dE^*$ is in very close agreement with measurements of neighboring nuclei using (d,p) -reactions at lower bombarding energies, where the giant quadrupole is not excited to any degree.

V. ACKNOWLEDGMENTS
~~~~~

The tireless effort of the 88"-cyclotron operating crew in providing the  $\alpha$ -beams is greatly appreciated. We would like to thank Claude Ellsworth for fabricating the high quality metallic  $^{238}\text{U}$ -targets used in the experiment.

This work was performed under the auspices of the Office of Basic Energy Sciences, Division of Nuclear Sciences, U. S. Department of Energy and in part by the National Science Foundation under grant #Phy 7822696. One of us, C.K.G., acknowledges the receipt of an Alfred P. Sloan Fellowship.

REFERENCES

1. B. D. Wilkins, E. P. Steinberg, and R. R. Chasman, Phys. Rev. C 14, 1832 (1976).
2. B. B. Back, J. M. Lebowitz and K. L. Wolf, Phys. Rev. C 20, 1819 (1979).
3. J. Lachkar, Y. Patin, and J. Sigaud, J. Phys. Lett. 36, 79 (1975).
4. N. I. Akimov, V. G. Borob'eva, V. N. Kabenin, N. P. Kolosov, B. D. Kuz'minov, A. I. Sergachev, L. D. Smirenkina, and M. Z. Tarasko, Yad. Fiz 13, 484 (1971)[Sov. J. Nucl. Phys. 13, 272, (1971)].
5. P. David, J. Debrus, F. Lübke, H. Mommsen, E. Schmitt, R. Shoenmackers, and H. Simons, Phys. Lett. 61B, 158 (1976).
6. E. Konecny, H. J. Specht, and J. Weber, in Proceedings of the Third International Atomic Energy Symposium on Physics and Chemistry of Fission, Rochester, 1973, Vol. 2, p. 3.
7. H. W. Schmitt, W. E. Kiker, C. W. Williams, Phys. Rev. 137, B837 (1965).
8. V. F. Apalin, Yu. N. Gritsyuk, I. E. Kutikov, V. I. Lebedev, and L. A. Mikaelian, Nucl. Phys. 71, 546 (1965).
9. J. van der Plicht, M. N. Harakeh, A. van der Woude, P. David and J. Debrus, Phys. Rev. Lett. 42, 1121 (1979).
10. A. C. Shotter, C. K. Gelbke, T. C. Awes, B. B. Back, J. Mahoney, T. J. M. Symons, and D. K. Scott, Phys. Rev. Lett. 43, 569 (1979).



11. J. van der Plicht, M. N. Harakeh, A. van der Woude, P. David, J. Debrus, H. Janszen and J. Schulze, preprint, 1980.
12. F. E. Bertrand, T. R. Beene, C. E. Bemis, Jr., E. E. Gross, D. J. Horen, J. R. Wu, and W. P. Jones, preprint, 1980.
13. G. Chenevert, N. S. Chant, I. Halpern, C. Glashausser, and D. L. Hendrie, Phys. Rev. Lett. 27, 434 (1971).
14. L. J. Colby, Jr., Mary LaSalle Shoaf, and J. W. Cobble, Phys. Rev. 121, 1415 (1961).

FIGURE CAPTIONS  
~~~~~

Fig. 1: The average total kinetic energy (a) and the fission coincidence spectrum (b) are shown as a function of the excitation energy. The neutron binding energy and thresholds for second and third chance fission are indicated by arrows.

Fig. 2: Fragment mass dependence of the slope dE_K/dE^* (a) and the average total kinetic energy \bar{E}_K (b) in the excitation energy region from 5-12 MeV. The mass spectrum of heavy fragments is also shown (c).

Fig. 3: Fission fragment mass distributions are shown for different excitation energy bins.

Fig. 4: Peak to valley ratios for the measured fission fragment distributions (closed circles) are compared to results obtained in the $^{238}\text{U}(\alpha, f)^*$ reaction¹⁴ (open circles).

Fig. 5: The standard deviations of the total kinetic energy distributions are shown as a function of heavy fragment mass for different excitation energy bins.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5