

# UCLA

## UCLA Previously Published Works

### Title

NF- $\kappa$ B Potentiates Caspase Independent Hydrogen Peroxide Induced Cell Death

### Permalink

<https://escholarship.org/uc/item/3bm2x69t>

### Journal

PLOS ONE, 6(2)

### ISSN

1932-6203

### Authors

Ho, Jessica Q  
Asagiri, Masataka  
Hoffmann, Alexander  
et al.

### Publication Date

2011

### DOI

10.1371/journal.pone.0016815

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

# NF- $\kappa$ B Potentiates Caspase Independent Hydrogen Peroxide Induced Cell Death

Jessica Q. Ho<sup>1</sup>, Masataka Asagiri<sup>1,2</sup>, Alexander Hoffmann<sup>1,2</sup>, Gourisankar Ghosh<sup>1\*</sup>

**1** Department of Chemistry and Biochemistry, University of California San Diego, La Jolla, California, United States of America, **2** Signaling Systems Laboratory, Department of Chemistry and Biochemistry, University of California San Diego, La Jolla, California, United States of America

## Abstract

**Background:** The pro-survival activity of NF- $\kappa$ B in response to a variety of stimuli has been extensively characterized. Although there have been a few reports addressing the pro-cell death role of NF- $\kappa$ B, the precise mechanism of NF- $\kappa$ B's pro-cell death function still remains elusive.

**Methodology/Principal Findings:** In the present study, we investigated the role of NF- $\kappa$ B in cell death induced by chronic insult with hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). Here, we show that NF- $\kappa$ B promotes H<sub>2</sub>O<sub>2</sub> induced caspase independent but PARP dependent fibroblast cell death. The pro-death activity of NF- $\kappa$ B is due to the DNA binding activity of RelA, which is induced through IKK- mediated I $\kappa$ B $\alpha$  degradation. NF- $\kappa$ B dependent pro-survival genes, Bcl-2 and XIAP, were significantly repressed, while NF- $\kappa$ B dependent pro-death genes, TNF $\alpha$  and Fas Ligand, were induced in response to H<sub>2</sub>O<sub>2</sub>.

**Conclusions/Significance:** We discovered an unexpected function of NF- $\kappa$ B, in that it potentiates chronic H<sub>2</sub>O<sub>2</sub> exposure induced cell death, and suggest that NF- $\kappa$ B mediates cell death through the repression of pro-survival genes and induction of pro-death genes. Since unremitting exposure of tissues to H<sub>2</sub>O<sub>2</sub> and other reactive oxygen species can lead to several degenerative disorders and diseases, our results have important implications for the use of NF- $\kappa$ B inhibitors in therapeutic drug design.

**Citation:** Ho JQ, Asagiri M, Hoffmann A, Ghosh G (2011) NF- $\kappa$ B Potentiates Caspase Independent Hydrogen Peroxide Induced Cell Death. PLoS ONE 6(2): e16815. doi:10.1371/journal.pone.0016815

**Editor:** Sudha Agarwal, Ohio State University, United States of America

**Received:** September 23, 2010; **Accepted:** January 11, 2011; **Published:** February 15, 2011

**Copyright:** © 2011 Ho et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by National Institutes of Health grants GM071862 and GM085490. JQH is supported by the Ruth L. Kirschstein National Research Service Award NIH/NCI T32 CA009523. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: gghosh@ucsd.edu

## Introduction

Mammalian cells are constantly exposed to reactive oxygen species (ROS), such as hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). Exogenous ROS arise from irradiation (UV, X-ray,  $\gamma$ -ray) and atmospheric pollutants, while endogenous ROS are mainly produced by the incomplete reduction of oxygen by cytochrome c during cellular respiration [1]. However, when the antioxidant capabilities of the cell are overwhelmed by ROS, a state of oxidative stress ensues, which can result in damage to DNA, proteins, and lipids [2]. Moreover, high and/or persistent levels of ROS result in aberrant cell death, which leads to aging and neurodegenerative disorders [3,4]. In particular, ROS induced fibroblast cell death can cause chronic obstructive pulmonary disease [5,6] as well as inadequate wound healing following myocardial infarction/reperfusion [7,8]. ROS induces cell death by modulating cell signaling pathways. A prominent signaling pathway involved in mediating the cell survival/cell death fate is the nuclear factor- $\kappa$ B (NF- $\kappa$ B) pathway [3].

NF- $\kappa$ B is a family of transcription factors, which are comprised of five family members: RelA/p65, RelB, c-Rel, n $\kappa$ b1/p50, and n $\kappa$ b2/p52, that form homo- or hetero-dimers in a combinatorial manner. In resting cells, the NF- $\kappa$ B dimers are retained in the cytoplasm by forming stable complexes with NF- $\kappa$ B inhibitor molecules, I $\kappa$ B ( $\alpha/\beta/\epsilon$ ). In the canonical activation pathway,

stimulation with an extracellular stimulus, such as tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), a pro-inflammatory cytokine, leads to phosphorylation of I $\kappa$ B $\alpha$  on serines 32 and 36 by IKK, the I $\kappa$ B kinase. This results in the ubiquitination of I $\kappa$ B $\alpha$ , which signals for the degradation of I $\kappa$ B $\alpha$  by the 26S proteasome. The freed NF- $\kappa$ B dimers can then translocate to the nucleus and activate transcription of their target genes [9,10,11].

Activation of NF- $\kappa$ B by exogenous H<sub>2</sub>O<sub>2</sub> has been found to be highly cell type dependent, in which NF- $\kappa$ B is activated in a variety of cell lines such as Jurkat T cells and HeLa cells [12,13], whereas NF- $\kappa$ B activation is inhibited in other cell lines such as murine neutrophils [14]. In cases where activation of NF- $\kappa$ B occurs, several mechanisms of NF- $\kappa$ B activation have been reported. While canonical activation of NF- $\kappa$ B via IKK-dependent I $\kappa$ B $\alpha$  degradation has been reported, other reports focus on an atypical mechanism of NF- $\kappa$ B activation in response to stimulation with H<sub>2</sub>O<sub>2</sub> [15,16]. This atypical mechanism involves an IKK independent mechanism and Tyr42 phosphorylation of I $\kappa$ B $\alpha$ , and only occurs in the absence of SHIP-1 [17,18]. The pathway of NF- $\kappa$ B activation in other cell lines, such as in mouse embryonic fibroblasts (MEFs), has yet to be delineated.

The anti-cell death role of NF- $\kappa$ B has been extensively characterized. RelA deficient cultured cells undergo apoptotic cell death upon treatment with TNF $\alpha$  due to deficiencies in pro-survival

and anti-oxidant gene transcription [19,20]. RelA deficiency also leads to embryonic lethality accompanied by massive apoptosis in the embryonic liver [21]. In response to a variety of other stimuli, such as ionizing radiation and chemotherapeutic drugs, RelA also appears to have an anti-apoptotic effect [22]. Finally, NF- $\kappa$ B suppression of apoptosis in cancer cells is a central event in cancer biology, as well as in chemoresistance of tumor cells [23]. However, there have also been a few scattered reports addressing the pro-cell death function of NF- $\kappa$ B in response to atypical NF- $\kappa$ B activators [24,25,26,27,28]. Yet, the mechanism by which NF- $\kappa$ B mediates a pro-cell death response remains elusive.

In the present study, we sought to define NF- $\kappa$ B's role in immortalized MEF cell death induced by chronic insult with H<sub>2</sub>O<sub>2</sub>. Here we present evidence that unremitting exposure to H<sub>2</sub>O<sub>2</sub> induces a caspase independent but PARP dependent cell death and that NF- $\kappa$ B potentiates cell death through the DNA binding activity of RelA, which is induced through the canonical activation pathway. Given that NF- $\kappa$ B dependent pro-survival genes, Bcl-2 and XIAP, were significantly repressed, while NF- $\kappa$ B dependent pro-death genes, TNF $\alpha$  and Fas Ligand, were induced in response to H<sub>2</sub>O<sub>2</sub>, we suggest that NF- $\kappa$ B mediates its pro-cell death function through the repression of pro-survival genes and induction of pro-death genes.

## Materials and Methods

### Cell culture, reagents, and antibodies

Immortalized MEFs cells were obtained from A. Hoffmann [29], cultured in humidified incubators at 37°C, 5% (v/v) CO<sub>2</sub> and were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% bovine calf serum (BCS) (Invitrogen) and 100 U/ml penicillin–streptomycin–glutamine (1xPSG) (Invitrogen). 293T were obtained from ATCC (cat # CRL-11268), and grown with DMEM supplemented with 10% Fetal Calf Serum (Invitrogen). Cells were stimulated for various periods with TNF $\alpha$  (Roche Biochemicals), glucose oxidase (Sigma), or H<sub>2</sub>O<sub>2</sub> (Sigma). Cells were labeled with CFSE (5- (and 6-) carboxyfluorescein diacetate, N-succinimidyl ester) (Sigma), H<sub>2</sub>DCFDA (Invitrogen), or propidium iodide (Sigma). Cells were treated with z-VAD-fmk (caspase inhibitor VI) and DPQ (PARP Inhibitor III), which were both purchased from Calbiochem. Antibodies against I $\kappa$ B $\alpha$  (sc371), IKK $\alpha$  (sc7184), RelA(sc372), and  $\alpha$ tubulin (sc5286) were purchased from Santa Cruz Biotechnology. Antibodies against IKK $\gamma$  (#559675), pro-caspase 3 (#611048), active caspase 3 (#559565), PARP (#556362), and PAR (#551813) were purchased from BD Pharmingen. Bcl-2 antibody (#2876) was purchased from Cell Signaling and p53 antibody (#Op03) was purchased from EMD Biosciences.

### Generation of oxidative stress

Cells were first grown to 95–98% confluency. GO was dissolved in 50 mM sodium acetate pH 5.1, and added to fresh media supplemented with 10% BCS, except in the cases where RNA was extracted, in which serum free media was used.

### Measurement of ROS production

Measurement of intracellular ROS production was carried out as described in [30]. The resulting supernatants were scanned with a Fluoromax-P instrument (J. Y. Horiba, Inc.) using a bandpass of 5 nm and ex 492 nm/em 526 nm.

### Cell Death Assay

MEF cell death was determined as the normalized value of propidium iodide (PI) incorporation =  $\left( \frac{FI_{TreatedPI} / FI_{TreatedCFSE}}{FI_{UntreatedPI} / FI_{UntreatedCFSE}} \right)$ , where FI = fluorescence intensity of PI or CFSE of GO treated or untreated

samples. A minimum of  $2 \times 10^6$  cells was used to label cells with 5  $\mu$ M CFSE. Following GO treatment, each plate was then washed twice with 1xDPBS containing 100 mg/ml CaCl<sub>2</sub> and Mg<sub>2</sub>Cl<sub>2</sub>, and treated with 10  $\mu$ g/ml PI for 15 min in the dark. To remove adherent cells, cells were incubated with Puck's buffer (5.4 mM KCl, 0.14 M NaCl, 4.2 mM NaHCO<sub>3</sub>, 5.6 mM D-glucose dextrose, 10 mM HEPES, 1 mM EDTA, pH 7.4) for 15 min in the dark. For UV treated cells, floating cells were combined with the adherent cells. The fluorescence intensity for PI (ex 535 nm/em 617 nm) and CFSE (ex 492 nm/em 517 nm) were then taken with a Fluoromax-P instrument (J. Y. Horiba, Inc.) using a bandpass of 5 nm.

### Molecular Biology

I $\kappa$ B $\alpha$  constructs were cloned into the retrovirus vector pBabe-puro between the restriction sites *Eco*RI and *Sal*I. Mutagenesis reaction was performed with the Stratagene Quickchange Mutagenesis kit. The following primers were used to clone the *R35A136A* Tg: forward: 5'- CGGGGCATGCGATTGCGCCG-CAAAATGCGAGGGGCGC-3', reverse: 5'- GCGCCCTC-GCATTTCGCGGAATCGCATGCCCG-3'.

### Retroviral transgenic system

293T cells were transfected using Lipofectamine 2000 (Invitrogen) containing 7  $\mu$ g of the retroviral vector and 3  $\mu$ g of pCI-Eco (Imgenex). Serum free DMEM containing the Lipofectamine and DNA mixture was removed 6 hrs later and replaced with DMEM supplemented with 10% FBS and 1xPSG. Cells were allowed to grow for 38–42 hrs post transfection. The media was then placed onto the *nfkb*<sup>-/-</sup> cells along with 8  $\mu$ g/ml polybrene (Sigma). These cells were then grown for another 48 hrs before selection with 2.5  $\mu$ g/ml puromycin (Calbiochem).

### Electrophoretic mobility shift assays (EMSAs)

EMSA experiments were carried out as described in [31] using a <sup>32</sup>P- labeled oligonucleotide probe (5'-GCTACAAGGGAC-TTTCGGCTGGGGACTTTCAGGGAGG -3'), which corresponds to the  $\kappa$ B site of the HIV-1 LTR.

### Western blots

Cells were lysed using RIPA buffer (20 mM Tris pH 7.5, 200 mM NaCl, 1% Triton-X 100, 2 mM DTT, 5 mM p-nitrophenyl phosphate, 2 mM sodium orthovanadate, 1X protease inhibitor cocktail [Sigma]). Membranes were developed using ECL chemiluminescence reagent (PerkinElmer) and quantitation of western blots was performed with ImageQuant TL (Amersham Biosciences).

### In vitro IKK kinase assay

Kinase assays were carried out as described in [29] using 2.0  $\mu$ g of recombinant GST-I $\kappa$ B $\alpha$  (1–54). The reaction was visualized by PhosphorImager (Molecular Dynamics) and quantitated by ImageQuant TL.

### Annexin V staining

Cells were washed with PBS and resuspended in 100  $\mu$ l of binding buffer (10 mM HEPES/NaOH, pH 7.4, 140 mM NaCl, 2.5 mM CaCl<sub>2</sub>). Alexa Fluor 647 conjugated Annexin V (BioLegend) and 7-AAD (BioLegend) were added according to the manufacturer's instructions, and the cells were incubated in the dark for 15 min at room temperature. The cells were analyzed using a Becton Dickinson FACSCalibur flow cytometer. Data analysis was performed with FlowJo software (Tree Star). Annexin

V/7-AAD double-positive cells are necrotic and excluded from analysis.

### Quantitative Real Time PCR (qPCR)

RNA was isolated using the RNeasy kit (QIAGEN) and further purified with DNase I digestion (QIAGEN). cDNA was synthesized from 1  $\mu$ g of RNA using the Superscript III First Strand synthesis kit (Invitrogen) and Oligo(dT) primers. Product accumulation was monitored by SYBR Green (Kapa Biosystems) fluorescence with Eppendorf Mastercycler ep *realplex*. The relative gene expression levels were calculated using the  $2^{-\Delta\Delta CT}$  method [32,33]. The p value was calculated using the Student's t test. The following primers were used: GAPDH Forward- AATGTGTCCTCGTGGATCT Reverse- CATCGAAGGTGGAAGAGTGG, TNF $\alpha$  Forward- CC-AGACCCCTCACACTCAGATC Reverse- CACTTGGTGGTT-TGCTACGAC, FasL (6753818a2) Forward- ACCCCCACT-CAAGGTCCAT, Reverse- CGAAGTACAACCCAGTTTCGT, Fas Receptor (6679751a3) Forward- AATCGCCTATGGTTGTT-GACC Reverse- TTGGTATGTTTTACGACTGG, Bax (6680-770a1) Forward- TGAAGACAGGGGCCTTTTTG Reverse- AA-TTCGCCGGAGCACTCG, XIAP Forward-GCAAGAGCTG-GATTTTTATGCTT, Reverse- TGCCCCCTTCATCCAATAG, Bcl-2 Forward- ATGCCCTTGTGGAAGTATATGGC, Reverse- GGTATGCACCCAGAGTGATGC, Bcl-xL (31981887a1) Forward- GACAAGGAGATGCAGGTATTGG Reverse- TCCCGT-AGAGATCCACAAAAGT, cIAP-1 (6680696a2) Forward- TCAG-TGACCTCGTTATAGCCTT Reverse- TCACACACGTCAA-ATGTTGGAA, p53 (6755881a1) Forward- GCGTAAACGCTTC-GAGATGTT Reverse- TTTTTATGGCGGGAAGTAGACTG. All primers are provided in 5' to 3' fashion. Primers obtained from the Primer Bank [34,35,36] are given with the PrimerBank ID Accession number. GAPDH primers were obtained from [37], TNF $\alpha$  primers were obtained from [38], and Bcl-2 primers were obtained from [39]. XIAP primers were designed using Primer3.

## Results

### Continuous H<sub>2</sub>O<sub>2</sub> exposure (via GO) to fibroblasts induces a caspase independent but PARP dependent cell death

In chronic obstructive pulmonary disease and inadequate wound healing following myocardial infarction/reperfusion, chronic insult of ROS to fibroblast cells can lead to aberrant cell death [6,7,8]. We attempted to simulate this condition by generating persistent oxidative stress in immortalized MEFs with glucose oxidase (GO). It has been shown that addition of GO to the media produces continuous levels of intracellular H<sub>2</sub>O<sub>2</sub>, as opposed to a bolus addition of H<sub>2</sub>O<sub>2</sub> [40], which we were also able to verify (J.Ho, data not shown). Intracellular ROS generation was verified by staining with DCFH-DA, a commonly used membrane permeable dye that fluoresces upon its oxidation to DCF by intracellular ROS [30]. Accordingly, ROS was produced in MEFs throughout the duration of treatment with 25 mU/ml GO (Fig. 1A). To roughly estimate the amount of intracellular H<sub>2</sub>O<sub>2</sub> that GO was producing, known concentrations of H<sub>2</sub>O<sub>2</sub> were also added in a bolus fashion to the cells and the intensity of fluorescence monitored. Based on the H<sub>2</sub>O<sub>2</sub> standard curve, we estimated that, within the first 3 hrs of treatment with 25 mU/ml GO, less than 50  $\mu$ M of H<sub>2</sub>O<sub>2</sub> was being produced, and following 3 hrs, roughly 50–100  $\mu$ M of H<sub>2</sub>O<sub>2</sub> was produced (Fig. 1A).

We next determined the effect of prolonged H<sub>2</sub>O<sub>2</sub> exposure (via 25 mU/ml GO) on MEFs. Significant cell death occurred within 5 hrs after GO treatment, as quantitated by the increase in propidium iodide (PI) incorporation, which correlates to a loss in


membrane integrity. PI incorporation increased even further 6 hrs following treatment (Fig. 1B). Since an increase in oxidative stress can switch the cell death mode from a caspase dependent to a caspase independent cell death mode [41], we set out to determine whether GO generated H<sub>2</sub>O<sub>2</sub> involved caspases. Treatment with irreversible general caspase inhibitor, z-VAD-fmk, was unable to prevent cell death. In agreement with these results, immunoblots against the cleaved form of caspase 3, one of the main executioners of apoptosis, revealed that cleaved caspase 3 is not present during treatment with GO (Fig. 1C). This is in contrast to UV induced MEF cell death, where caspases mediate cell death, as seen by the inhibition of cell death upon treatment with zVAD-fmk and by the production of the active form of caspase 3 (Fig. 1B, C).

Caspase independent MEF cell death induced by bolus addition of H<sub>2</sub>O<sub>2</sub> occurs in a Poly(ADP-ribose) polymerase (PARP) dependent manner [42], which we were also able to confirm (Fig. 1D). PARP, a nuclear enzyme involved in the DNA damage response and cell death, becomes activated in response to DNA damage and attaches ADP-ribose units (PAR polymer) to itself and nuclear proteins [43]. We next sought to evaluate the contribution of PARP in cell death induced by continuous exposure to H<sub>2</sub>O<sub>2</sub>. Upon GO addition, cells pretreated with PARP inhibitor, DPQ, showed significant resistance to membrane permeabilization, as opposed to cells treated without DPQ (Fig. 1E). Given that PARP protein levels remained constant while PAR formation increased, as verified by western blot against PARP and PAR, we concluded that PARP was activated within 15 min of stimulation with GO. PAR formation was then eliminated upon pretreatment with DPQ (Fig. 1E). Thus, treatment of MEFs with GO results in a caspase independent but PARP dependent cell death.

### NF- $\kappa$ B potentiates H<sub>2</sub>O<sub>2</sub> induced cell death

There have been extensive reports illustrating cooperativity between PARP and NF- $\kappa$ B. In response to DNA damaging agents such as ionizing radiation, PARP-1 has been shown to be an essential upstream mediator of NF- $\kappa$ B activation [44,45]. PARP-1 can also act as a direct co-activator of NF- $\kappa$ B [46]. Since we have shown that chronic insult with H<sub>2</sub>O<sub>2</sub> induces a PARP dependent cell death, we next wanted to determine whether NF- $\kappa$ B also plays any role in H<sub>2</sub>O<sub>2</sub> induced cell death. To examine this, we compared rates of cell death in wt and *nfkb*<sup>-/-</sup> (*nfkb1*<sup>-/-</sup> *rela*<sup>-/-</sup> *crel*<sup>-/-</sup>) MEFs treated with 25 mU/ml GO. Although we expected that cell death would be enhanced in the absence of NF- $\kappa$ B due to its well established pro-survival activity, we found, to our surprise, that MEFs which lacked p50, RelA and c-Rel were more resistant to H<sub>2</sub>O<sub>2</sub> induced cell death than wt cells (Fig. 2A).

To directly address the involvement of NF- $\kappa$ B in the delay of cell death, *nfkb*<sup>-/-</sup> cells were reconstituted, using a retroviral transgenic system, with either *rela* transgene (Tg), which is the major transactivating subunit of NF- $\kappa$ B, or a DNA defective binding mutant of *rela* (*R35A136A* Tg). Both *rela* Tg and *R35A136A* Tg reconstituted cells contained similar levels of RelA protein when compared to wt MEFs (Fig. 2B). Upon treatment with GO, *rela* Tg cells displayed increased incorporation of PI as opposed to *nfkb*<sup>-/-</sup> cells reconstituted with either empty vector (pBabe) or *R35A136A* Tg (Fig. 2B). However the amount of PI incorporation in *rela* Tg cells was decreased compared to wt cells, indicating only partial rescue of the cell death phenotype. Our attempt to study cell death in *rela* Tg reconstituted in *rela*<sup>-/-</sup> cells was unsuccessful since reconstitution with *rela* did not rescue the cell death phenotype. This could be because of the transformation of the *rela*<sup>-/-</sup> cells with viral oncoproteins, which disrupts many transcriptional regulatory pathways, and resulted in the loss of this mechanism of modulating NF- $\kappa$ B function. Regardless, the lack of




**Figure 1. Continuous H<sub>2</sub>O<sub>2</sub> exposure (via GO) to fibroblasts induces a caspase independent but PARP dependent cell death.** (A) Wild type (wt) MEFs were either untreated (-) or treated with 25 mU/ml GO for the indicated periods of time. H<sub>2</sub>O<sub>2</sub> readings for the H<sub>2</sub>O<sub>2</sub> standard curve were taken 10 min after addition. All H<sub>2</sub>O<sub>2</sub> levels are shown as normalized to untreated samples and are given in arbitrary units (a.b.u.). (B) Following pretreatment for 1 hr in the presence of 100  $\mu$ M z-VAD-fmk or DMSO, MEFs were subsequently treated with 25 mU/ml GO or 200 J/m<sup>2</sup> UV and cell death was analyzed at the indicated time points. (C) Cell lysate was analyzed by western blotting against pro-caspase3, caspase3, and  $\alpha$ -tubulin. The cell lysates were also stained by Coomassie Brilliant Blue to demonstrate equal loading. (D) Following pretreatment for 1 hr in the presence of 60  $\mu$ M of DPQ or DMSO, MEFs were subsequently treated with 25 mU/ml GO or 2 mM H<sub>2</sub>O<sub>2</sub> and cell death was analyzed at the indicated time points. (E) Cell lysate was analyzed by western blotting against PAR (poly (ADP-ribose) polymer), PARP, and  $\alpha$ -tubulin. Both z-VAD-fmk and DPQ were kept in the media following GO or UV stimulation. Cell death was determined as the normalized value of propidium iodide incorporation (see Methods). All results are presented as the average of triplicate experiments. Error bars signify  $\pm$ s.e.m (standard error of mean). doi:10.1371/journal.pone.0016815.g001

cell death in *R35A136A* Tg in *nfkb*<sup>-/-</sup> cells indicates that NF- $\kappa$ B promoted cell death is dependent on NF- $\kappa$ B, in particular, RelA DNA binding activity. Additionally, we confirmed that the differences in the rate of cell death between wt and *nfkb*<sup>-/-</sup> cells reconstituted with either pBabe or *rela* Tg were primarily dependent on NF- $\kappa$ B. This was done by determining that all cell lines contained similar basal levels of p53, a well-known and potent contributor of cell death (Fig. S1).

Next, to examine the reason for the partial rescue of the cell death phenotype in response to GO, the response of the *rela* reconstituted cells to TNF $\alpha$  induced cell death, as well as TNF $\alpha$  induced DNA binding was inspected. Since it has been shown that *de novo* synthesis prevents the occurrence of apoptotic death upon

treatment with TNF $\alpha$  [20], cells containing a transcriptionally active RelA should prevent cell death from occurring. Accordingly, upon TNF $\alpha$  stimulation, wt and *rela* Tg cells did not undergo cell death, as opposed to *R35A136A* Tg and pBabe cells (Fig. 2C). EMSA results also show that NF- $\kappa$ B was not activated in both pBabe and *R35A136A* Tg cells, in contrast to wt and, to a lower extent, *rela* Tg cells (Fig. 2D). The incomplete rescue of NF- $\kappa$ B activation in *rela* Tg cells might be due to the absence of NF- $\kappa$ B subunits, p50 and cRel. The p50:RelA heterodimer is the major NF- $\kappa$ B species, and both cRel and RelA homodimers have also been reported to play roles in the regulation of target gene expression [47]. Therefore, the absence of these dimers and the presence of only RelA homodimer alone might result in only



**Figure 2. NF-κB augments cell death in H<sub>2</sub>O<sub>2</sub> induced cell death.** Cell death assays were performed on wt, *nfkB*<sup>-/-</sup> MEFs (A), and *nfkB*<sup>-/-</sup> cells reconstituted with empty vector (pBabe), *relA* transgene (Tg), or *relA* mutant (*R35AY36A* Tg) (B) treated with 25 μM GO for the indicated periods of time. RelA was verified by western blot. Results are presented as the average of 3 independent experiments. Error bars signify ±s.e.m. \* denotes p<0.05. (C–D) wt, *relA* Tg, *R35AY36A* Tg, and pBabe *nfkB*<sup>-/-</sup> cells were analyzed for cell viability in terms of AnnexinV positive cells following 16 hrs of treatment with 10 ng/ml TNFα (C) and for nuclear localization and DNA binding by EMSA analysis following treatment with 1 ng/ml TNFα (D). Supershift analysis was performed by adding RelA antibody to the EMSA reaction. doi:10.1371/journal.pone.0016815.g002

suboptimal activation of some NF-κB target genes, and explains the partially functionally rescued *relA* reconstituted cells. Due to technical difficulties reconstituting p50 into *nfkB*<sup>-/-</sup> cells, we were unable to distinguish whether NF-κB's pro-cell death function is due to RelA homodimer or the p50:RelA heterodimer. However, from our data (Fig. 2B), it is clear that a RelA containing dimer is involved in promoting cell death.


**The canonical NF-κB activation pathway is required for the pro-cell death function of NF-κB in response to chronic exposure to H<sub>2</sub>O<sub>2</sub>**

To further determine whether NF-κB acts as a promoter of cell death in response to H<sub>2</sub>O<sub>2</sub>, we compared the rates of cell death of *ikba*<sup>-/-</sup> MEFs which were reconstituted with either wt IκBα (*wt ikba* Tg) or a mutant IκBα, where both Ser 32 and Ser 36, which are the IKK phosphorylation sites, are mutated to alanine (*aa ikba* Tg), thus preventing NF-κB activation. Interestingly, in order to achieve cell death in a timely fashion, 50 μM GO instead of

25 μM GO, was needed to induce cell death in both cell lines. This may be a reflection of cell line-specific characteristics or due to the over-expression of IκBα upon reconstitution in these cell lines (G.Ghosh, data not shown). Nevertheless, *wt ikba* Tg cells had a rapid rate of cell death compared to *aa ikba* Tg cells, where the canonical NF-κB activation pathway had been blocked (Fig. 3A). As expected, NF-κB activation was completely blocked in *aa ikba* Tg cells as demonstrated by the lack of NF-κB activation upon TNFα stimulation by EMSA analysis (Fig. 3B). These results further suggest that the resistance to death of *aa ikba* Tg cells is due to the lack of NF-κB activity. All together, these results strongly indicate that NF-κB plays a pro-cell death role in response to H<sub>2</sub>O<sub>2</sub> induced caspase independent cell death, and that the canonical activation pathway is required in mediating NF-κB's pro-cell death role.

We next examined whether NF-κB was activated in response to chronic exposure to H<sub>2</sub>O<sub>2</sub>. Indeed, as seen by EMSA analysis, NF-κB was activated in a prolonged manner upon addition of GO in





**Figure 3. The canonical activation pathway is required for the pro-cell death function of NF- $\kappa$ B.** (A) Cell viability assays were performed on *ikba*<sup>-/-</sup> MEFs reconstituted with either *wt ikba* Tg, or with *aa ikba* Tg treated with 50 mU/ml GO for the indicated periods. (B) *wt ikba* Tg and *aa ikba* Tg MEFs were treated with 1 ng/ml TNF $\alpha$  for 0.5 hr for EMSA analysis. (C) wt MEFs were treated with either 1.0 ng/ml TNF $\alpha$  or 25 mU/ml GO for the indicated times, for EMSA (C) IKK activity (D) or I $\kappa$ B $\alpha$  degradation analysis (E). Quantification of EMSA and IKK kinase assay experiments are shown. All results are presented as the average of 3 independent experiments. Error bars signify  $\pm$ s.e.m. doi:10.1371/journal.pone.0016815.g003

MEFs (Fig. 3C). Given that NF- $\kappa$ B has been reported to be activated in both an IKK independent and dependent manner in response to H<sub>2</sub>O<sub>2</sub>, we then performed IKK activity assays, in which IKK was immunoprecipitated from GO treated MEFs followed by an *in vitro* kinase assay. IKK activity assays reveal that IKK is activated following one hour of treatment with GO (Fig. 3D). Accordingly, there is also concomitant degradation of I $\kappa$ B $\alpha$  (Fig. 3E). All together these results show that NF- $\kappa$ B is activated via the canonical pathway in MEFs in response to chronic exposure to H<sub>2</sub>O<sub>2</sub>.

#### NF- $\kappa$ B-dependent survival genes are repressed whereas death promoting genes are induced by H<sub>2</sub>O<sub>2</sub>

As a transcription factor, NF- $\kappa$ B's active participation in cell death is likely to be mediated through its target genes. Therefore, we set out to examine the expression pattern of several NF- $\kappa$ B target genes that are known to impact cell death or survival

(Table 1). Using real time quantitative PCR (qPCR), we measured mRNA levels at 0 and 4 hrs after GO treatment in wt MEF and *nfkb*<sup>-/-</sup> cells reconstituted with empty vector (pBabe) or *relA* Tg. While the majority of these genes did not undergo changes in expression levels (Table 1), we identified 4 genes that underwent significant alterations: cell survival factors, Bcl-2 and XIAP, and cell death promoting factors, TNF $\alpha$  and Fas ligand (FasL). The significant reduction of Bcl-2 gene levels in both wt and *relA* Tg cells and not pBabe cells (Fig. 4) implies that Bcl-2 repression is due to the presence of RelA. Decreased Bcl-2 protein levels were also observed only in wt, and not *nfkb*<sup>-/-</sup> cells, 6 hrs following treatment with GO (Fig. S2), suggesting that changes in mRNA levels correspond to changes in protein levels. A similar trend in gene expression is also seen for X-linked inhibitor of apoptosis protein, XIAP. In contrast, TNF $\alpha$  expression was significantly induced in both wt and *relA* Tg cells, while not in pBabe cells. FasL was induced in only wt cells. These results suggest that TNF $\alpha$ , but not FasL, induction by H<sub>2</sub>O<sub>2</sub> is RelA dependent. The lack of FasL

**Table 1.** Relative gene expression levels.

	Cell Survival Genes			Cell Death Genes			
	wt	pBabe	rela Tg	wt	pBabe	rela Tg	
<b>Bcl-2</b>	0.51±0.13*	0.83±0.03*	0.71±0.08*	<b>TNF<math>\alpha</math></b>	6.5±1.00*	1.2±0.20	2.6±0.50*
<b>XIAP</b>	0.46±0.03*	0.80±0.05	0.50±0.07*	<b>Fas Ligand</b>	2.4±0.50*	0.8±0.12	1.5±0.90
<b>Bcl-xL</b>	0.79±0.12	0.91±0.11	0.67±0.20	<b>Fas Receptor</b>	1.1±0.07	0.8±0.04*	0.9±0.10
<b>clAP-1</b>	0.89±0.24	0.87±0.03*	0.69±0.07*	<b>Bax</b>	0.9±0.27	0.9±0.11	0.76±0.08*
<b>p53</b>	1.01±0.11	0.84±0.08	0.90±0.15				

Note: Relative gene expression levels are presented as the average of three triplicate experiments with the  $\pm$  standard deviation.

\*denotes  $p < 0.05$ .

doi:10.1371/journal.pone.0016815.t001


induction in *rela* Tg cells could be attributed to the partial rescue phenotype of the *rela* Tg cells (Fig. 2B).

## Discussion

This study has discovered an unexpected function of NF- $\kappa$ B in that it promotes MEF cell death in response to chronic insult with H<sub>2</sub>O<sub>2</sub>. We have shown that intracellular H<sub>2</sub>O<sub>2</sub> was continuously produced in MEFs treated with GO and that this unremitting exposure to H<sub>2</sub>O<sub>2</sub> resulted in a caspase-independent but PARP dependent cell death. We also show that the pro-death activity of NF- $\kappa$ B is dependent on the DNA binding activity of RelA, which is induced through IKK-mediated I $\kappa$ B $\alpha$  degradation. The death promoting activity of NF- $\kappa$ B might be mediated by the down regulation of a subset of NF- $\kappa$ B dependent pro-survival factors and up-regulation of NF- $\kappa$ B dependent pro-death factors, as demonstrated by the repression of Bcl-2 and XIAP, and induction of TNF $\alpha$  and FasL.

Due to the repression of pro-survival factor Bcl-2 in wt and *rela* Tg cells, we propose that the pro-cell death function of NF- $\kappa$ B is primarily due to its transcriptional down regulation of Bcl-2, since

it has been shown that in MNNG induced caspase independent but PARP-1 dependent MEF cell death, over-expression of Bcl-2 can delay cell death [42]. Bcl-2 over-expression delays cell death due to its ability to prevent translocation of apoptosis inducing factor, AIF, from the mitochondria to the nucleus. This translocation event is central in causing MNNG and H<sub>2</sub>O<sub>2</sub> induced MEF cell death [42]. Our proposed model is supported by two additional pieces of data. First, Bcl-2 protein levels decrease in wt MEFs as opposed to *nfkb*<sup>-/-</sup> MEFs after 6 hrs of treatment with 25  $\mu$ M/ml of GO (Fig. S2), indicating that the changes in Bcl-2 mRNA level translates to changes in protein levels. Secondly, the fact that Bcl-2 over-expression can only delay cell death also fits with our proposed model, since *nfkb*<sup>-/-</sup> cells, which contain higher levels of Bcl-2 than either wt or *rela* Tg cells, eventually also succumb to cell death (J. Ho, data not shown). Additionally, our result, in which Bcl-2 is down regulated in a seemingly NF- $\kappa$ B dependent manner to promote PARP dependent fibroblast cell death, suggests a novel level of cooperativity between PARP and NF- $\kappa$ B. This is because PARP and NF- $\kappa$ B cooperativity has only been shown in instances where PARP-1 can act as a direct co-activator for RelA and p50, or as an upstream



**Figure 4. NF- $\kappa$ B dependent survival genes, Bcl-2 and XIAP, are repressed, while cell death genes, TNF $\alpha$  and FasL, are induced.** Total RNA was isolated from wt and pBabe or *rela* reconstituted *nfkb*<sup>-/-</sup> MEFs without treatment (black bars) or following 4 hrs of treatment with 25  $\mu$ M/ml GO (grey bars). Relative gene expression (Rel. Gene Expr.) levels were determined using real time RT-PCR and are presented as compared to untreated samples. Results are presented as the average of three triplicate experiments. Error bars signify  $\pm$  standard deviation. \* denotes  $p < 0.05$ . doi:10.1371/journal.pone.0016815.g004



mediator of NF- $\kappa$ B activation in response to various stimuli [44,45,46].

We have convincingly demonstrated that NF- $\kappa$ B has a pro-cell death function in response to chronic insult with H<sub>2</sub>O<sub>2</sub> in MEFs. Future experiments will be performed to determine whether this effect occurs in other cell types. Aside from MEFs, this type of caspase independent but PARP dependent cell death has only been reported to occur in cortical neurons, upon *N-Methyl-D-aspartic* acid (NMDA) over-stimulation [42,48]. It is yet unknown whether NF- $\kappa$ B has any role in this NMDA induced, PARP dependent cell death in cortical neurons. Interestingly, it was shown that in HeLa cells, NF- $\kappa$ B promotes cell survival in response to a single bolus addition of low nanomolar concentration of H<sub>2</sub>O<sub>2</sub> [13]. This variation can be attributed to differences in cell type, as well as differences in the application of H<sub>2</sub>O<sub>2</sub>, and thus amount of intracellular H<sub>2</sub>O<sub>2</sub>, which, as previously stated, can switch the cell death mode. We propose that NF- $\kappa$ B can be anti-cell death in caspase dependent cell death induced by transient or low levels of H<sub>2</sub>O<sub>2</sub>, but that NF- $\kappa$ B has a pro-cell death function in caspase independent cell death induced by continuous or high levels of H<sub>2</sub>O<sub>2</sub>. Indeed, NF- $\kappa$ B's role in promoting cell survival and death is a complex event. This is demonstrated by genotoxic agent, VP16, induced cell death, in which both pro- and anti-apoptotic genes were induced in NF- $\kappa$ B dependent manner, such that the final outcome depended on a balance of the induction levels of pro- and anti-apoptotic genes [39].

Our gene expression results are strongly supported by previous studies, which showed that cell death stimulation with daunorubicin, cisplatin, or p14<sup>ARF</sup> over-expression in U2OS osteosarcoma cells resulted in RelA mediated transcriptional repression of pro-survival genes, Bcl-xL or XIAP [49]. These previous reports also showed that RelA acts in a dominantly transcriptionally repressive manner. Interestingly, we observe selective activation and repression of certain death and survival genes, in that TNF $\alpha$  was induced while Bcl-2 and XIAP were repressed in a seemingly NF- $\kappa$ B dependent manner (Fig. 4). There are currently more reports addressing NF- $\kappa$ B's singular transcriptional response in promoting cell death (either complete repression or induction of survival or death genes) [26,27,28] and significantly fewer reports describing NF- $\kappa$ B's mixed transcriptional response in promoting cell death (repression and induction of survival and death genes, respectively) [25]. This is the first report suggesting that NF- $\kappa$ B uses a mixture of transcriptional responses in promoting cell death upon ROS stimulation. It is unclear at this stage as to how NF- $\kappa$ B mediates repression of some target promoters and activation of others. However, given that oxidative stress is known to inactivate the cysteine active sites of cellular phosphatases, this would change the cellular phospho-protein state [18] and could potentially inhibit or enhance the recruitment of coactivators and corepressors. Additionally, RelA mediated repression of pro-survival genes has been reported to involve Thr505 phosphorylation of RelA, which can enhance the interaction between RelA and HDAC1 [49]. Further studies are still required to fully unravel this mechanism.

## References

- Cadenas E (1989) Biochemistry of oxygen toxicity. *Annu Rev Biochem* 58: 79–110.
- Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M (2006) Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact* 160: 1–40.
- Trachootham D, Lu W, Ogasawara MA, Nilsa RD, Huang P (2008) Redox regulation of cell survival. *Antioxid Redox Signal* 10: 1343–1374.
- Behl C, Moosmann B (2002) Oxidative nerve cell death in Alzheimer's disease and stroke: antioxidants as neuroprotective compounds. *Biol Chem* 383: 521–536.
- Nakamura Y, Romberger DJ, Tate L, Ertl RF, Kawamoto M, et al. (1995) Cigarette smoke inhibits lung fibroblast proliferation and chemotaxis. *Am J Respir Crit Care Med* 151: 1497–1503.

The induction and repression of cell death and survival genes, respectively, in response to H<sub>2</sub>O<sub>2</sub> also suggests that activation of NF- $\kappa$ B is required for mediating its pro-cell death response. Previous reports of NF- $\kappa$ B's pro-cell death function have been shown to depend on either basal [24,25] or activated NF- $\kappa$ B [49]. However, we cannot fully conclude that NF- $\kappa$ B activation is solely required to mediate the pro-cell death response, since pBabe reconstituted cells contain lower levels of basal NF- $\kappa$ B (Fig. 2D).

Our studies have also clearly shown that NF- $\kappa$ B is activated via the canonical pathway in MEFs, which is in contrast to reports where NF- $\kappa$ B activated in response to oxidative stress occurred via an atypical mechanism, involving an IKK independent mechanism and I $\kappa$ B $\alpha$  Tyr42 phosphorylation. The presence of SHIP1 has been reported to revert the mechanism of NF- $\kappa$ B activation from atypical to canonical in response to H<sub>2</sub>O<sub>2</sub> [18]. Thus, the presence of SHIP2 in MEFs, which is functionally similar to SHIP1 [50], supports our observed NF- $\kappa$ B activation mechanism.

Unremitting exposure of tissues to ROS can lead to pathological conditions, such as neurodegenerative disorders and chronic obstructive pulmonary disease [3,4,6]. Our result that NF- $\kappa$ B might play a role in promoting cell death adds another layer of complexity to therapeutic drug design and should be taken into consideration when NF- $\kappa$ B inhibitor pharmaceutical targets are used in treatment. Overall, this study shows that NF- $\kappa$ B dependent transcription is responsible for promoting H<sub>2</sub>O<sub>2</sub> induced cell death. Further experiments in future study will be done to explore the detailed molecular mechanism.

## Supporting Information

**Figure S1 wt and pBabe or *rela* Tg reconstituted *nfkb*<sup>-/-</sup> MEFs contain similar levels of basal p53.** Cell lysate of wt and *nfkb*<sup>-/-</sup> MEFs reconstituted with pBabe or *rela* Tg were analyzed by western blotting against p53 and  $\alpha$ -tubulin. (TIF)

**Figure S2 Bcl-2 protein levels significantly decrease in wt MEFs as opposed to *nfkb*<sup>-/-</sup> cells upon continuous exposure to H<sub>2</sub>O<sub>2</sub>.** wt MEFs and *nfkb*<sup>-/-</sup> MEFs were either untreated or treated with 25 mU/ml GO for 6 hrs. Cell lysate was analyzed by western blotting against Bcl-2 and  $\alpha$ -tubulin. (TIF)

## Acknowledgments

We would like to thank Erika Mathes for *wt ikba* Tg and *aa ikba* Tg cells; Dong Wang and Vincent Shih for technical assistance; Dennis Otero, Xiang-Yang Zhong and Anthony Farina for valuable discussions; and Olga Savinova, Sulakshana Mukherjee, Zhihua Tao, and Smarajit Polley for their critical reading of this manuscript.

## Author Contributions

Conceived and designed the experiments: JQH AH GG. Performed the experiments: JQH MA. Analyzed the data: JQH MA AH GG. Contributed reagents/materials/analysis tools: JQH MA AH GG. Wrote the paper: JQH GG.

6. Carnevali S, Petruzzelli S, Longoni B, Vanacore R, Barale R, et al. (2003) Cigarette smoke extract induces oxidative stress and apoptosis in human lung fibroblasts. *Am J Physiol Lung Cell Mol Physiol* 284: L955–963.
7. Takahashi A, Aoshiba K, Nagai A (2002) Apoptosis of wound fibroblasts induced by oxidative stress. *Exp Lung Res* 28: 275–284.
8. Xiao J, Lv Y, Lin S, Jin L, Zhang Y, et al. (2010) Cardiac protection by basic fibroblast growth factor from ischemia/reperfusion-induced injury in diabetic rats. *Biol Pharm Bull* 33: 444–449.
9. Karin M, Ben-Neriah Y (2000) Phosphorylation meets ubiquitination: the control of NF- $\kappa$ B activity. *Annu Rev Immunol* 18: 621–663.
10. Baldwin AS, Jr. (1996) The NF- $\kappa$ B and I $\kappa$ B proteins: new discoveries and insights. *Annu Rev Immunol* 14: 649–683.
11. Ghosh S, May MJ, Kopp EB (1998) NF- $\kappa$ B and Rel proteins: evolutionarily conserved mediators of immune responses. *Annu Rev Immunol* 16: 225–260.
12. Schreck R, Rieber P, Baeuerle PA (1991) Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF- $\kappa$ B transcription factor and HIV-1. *EMBO J* 10: 2247–2258.
13. Storz P, Toker A (2003) Protein kinase D mediates a stress-induced NF- $\kappa$ B activation and survival pathway. *EMBO J* 22: 109–120.
14. Zmijewski JW, Zhao X, Xu Z, Abraham E (2007) Exposure to hydrogen peroxide diminishes NF- $\kappa$ B activation, I $\kappa$ B $\alpha$  degradation, and proteasome activity in neutrophils. *Am J Physiol Cell Physiol* 293: C255–266.
15. Livolsi A, Busuttill V, Imbert V, Abraham RT, Peyron JF (2001) Tyrosine phosphorylation-dependent activation of NF- $\kappa$ B. Requirement for p56 LCK and ZAP-70 protein tyrosine kinases. *Eur J Biochem* 268: 1508–1515.
16. Takada Y, Mukhopadhyay A, Kundu GC, Mahabeleshwar GH, Singh S, et al. (2003) Hydrogen peroxide activates NF- $\kappa$ B through tyrosine phosphorylation of I $\kappa$ B $\alpha$  and serine phosphorylation of p65: evidence for the involvement of I $\kappa$ B $\alpha$  kinase and Syk protein-tyrosine kinase. *J Biol Chem* 278: 24233–24241.
17. Gloire G, Charlier E, Rahmouni S, Volanti C, Chariot A, et al. (2006) Restoration of SHIP-1 activity in human leukemic cells modifies NF- $\kappa$ B activation pathway and cellular survival upon oxidative stress. *Oncogene* 25: 5485–5494.
18. Gloire G, Legrand-Poels S, Piette J (2006) NF- $\kappa$ B activation by reactive oxygen species: fifteen years later. *Biochem Pharmacol* 72: 1493–1505.
19. Beg AA, Baltimore D (1996) An essential role for NF- $\kappa$ B in preventing TNF- $\alpha$ -induced cell death. *Science* 274: 782–784.
20. Papa S, Bubici C, Zazzeroni F, Pham CG, Kuntzen C, et al. (2006) The NF- $\kappa$ B-mediated control of the JNK cascade in the antagonism of programmed cell death in health and disease. *Cell Death Differ* 13: 712–729.
21. Beg AA, Sha WC, Bronson RT, Ghosh S, Baltimore D (1995) Embryonic lethality and liver degeneration in mice lacking the RelA component of NF- $\kappa$ B. *Nature* 376: 167–170.
22. Barkett M, Gilmore TD (1999) Control of apoptosis by Rel/NF- $\kappa$ B transcription factors. *Oncogene* 18: 6910–6924.
23. Kucharczak J, Simmons MJ, Fan Y, Gelinias C (2003) To be, or not to be: NF- $\kappa$ B is the answer—role of Rel/NF- $\kappa$ B in the regulation of apoptosis. *Oncogene* 22: 8961–8982.
24. Lin KI, DiDonato JA, Hoffmann A, Hardwick JM, Ratan RR (1998) Suppression of steady-state, but not stimulus-induced NF- $\kappa$ B activity inhibits alphavirus-induced apoptosis. *J Cell Biol* 141: 1479–1487.
25. Liu J, Yang D, Minemoto Y, Leitges M, Rosner MR, et al. (2006) NF- $\kappa$ B is required for UV-induced JNK activation via induction of PKC $\delta$ . *Mol Cell* 21: 467–480.
26. Ho WC, Dickson KM, Barker PA (2005) Nuclear factor- $\kappa$ B induced by doxorubicin is deficient in phosphorylation and acetylation and represses nuclear factor- $\kappa$ B-dependent transcription in cancer cells. *Cancer Res* 65: 4273–4281.
27. Campbell KJ, Rocha S, Perkins ND (2004) Active repression of antiapoptotic gene expression by RelA(p65) NF- $\kappa$ B. *Mol Cell* 13: 853–865.
28. Campbell KJ, Witty JM, Rocha S, Perkins ND (2006) Cisplatin mimics ARF tumor suppressor regulation of RelA (p65) nuclear factor- $\kappa$ B transactivation. *Cancer Res* 66: 929–935.
29. Werner SL, Barken D, Hoffmann A (2005) Stimulus specificity of gene expression programs determined by temporal control of IKK activity. *Science* 309: 1857–1861.
30. Hayakawa M, Miyashita H, Sakamoto I, Kitagawa M, Tanaka H, et al. (2003) Evidence that reactive oxygen species do not mediate NF- $\kappa$ B activation. *EMBO J* 22: 3356–3366.
31. Mathes E, O’Dea EL, Hoffmann A, Ghosh G (2008) NF- $\kappa$ B dictates the degradation pathway of I $\kappa$ B $\alpha$ . *EMBO J* 27: 1357–1367.
32. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>- $\Delta$ CT</sup> Method. *Methods* 25: 402–408.
33. Schmittgen TD, Livak KJ (2008) Analyzing real-time PCR data by the comparative C<sub>T</sub> method. *Nat Protoc* 3: 1101–1108.
34. Spandidos A, Wang X, Wang H, Dragnev S, Thurber T, et al. (2008) A comprehensive collection of experimentally validated primers for Polymerase Chain Reaction quantitation of murine transcript abundance. *BMC Genomics* 9: 633.
35. Wang X, Seed B (2003) A PCR primer bank for quantitative gene expression analysis. *Nucleic Acids Res* 31: e154.
36. Spandidos A, Wang X, Wang H, Seed B (2010) PrimerBank: a resource of human and mouse PCR primer pairs for gene expression detection and quantification. *Nucleic Acids Res* 38: D792–799.
37. Ghisletti S, Huang W, Jepsen K, Benner C, Hardiman G, et al. (2009) Cooperative NCoR/SMRT interactions establish a corepressor-based strategy for integration of inflammatory and anti-inflammatory signaling pathways. *Genes Dev* 23: 681–693.
38. Ghisletti S, Huang W, Ogawa S, Pascual G, Lin ME, et al. (2007) Parallel SUMOylation-dependent pathways mediate gene- and signal-specific transrepression by LXRs and PPAR $\gamma$ . *Mol Cell* 25: 57–70.
39. Wu ZH, Miyamoto S (2008) Induction of a pro-apoptotic ATM-NF- $\kappa$ B pathway and its repression by ATR in response to replication stress. *EMBO J* 27: 1963–1973.
40. de Oliveira-Marques V, Cyrne L, Marinho HS, Antunes F (2007) A quantitative study of NF- $\kappa$ B activation by H<sub>2</sub>O<sub>2</sub>: relevance in inflammation and synergy with TNF- $\alpha$ . *J Immunol* 178: 3893–3902.
41. Gardner AM, Xu FH, Fady C, Jacoby EJ, Duffey DC, et al. (1997) Apoptotic vs. nonapoptotic cytotoxicity induced by hydrogen peroxide. *Free Radic Biol Med* 22: 73–83.
42. Yu SW, Wang H, Poitras MF, Coombs C, Bowers WJ, et al. (2002) Mediation of poly(ADP-ribose) polymerase-1-dependent cell death by apoptosis-inducing factor. *Science* 297: 259–263.
43. Kraus WL, Lis JT (2003) PARP goes transcription. *Cell* 113: 677–683.
44. Stilmann M, Hinz M, Arslan SC, Zimmer A, Schreiber V, et al. (2009) A nuclear poly(ADP-ribose)-dependent signalosome confers DNA damage-induced I $\kappa$ B kinase activation. *Mol Cell* 36: 365–378.
45. Veuger SJ, Hunter JE, Durkacz BW (2009) Ionizing radiation-induced NF- $\kappa$ B activation requires PARP-1 function to confer radioresistance. *Oncogene* 28: 832–842.
46. Hassa PO, Covic M, Hasan S, Imhof R, Hottiger MO (2001) The enzymatic and DNA binding activity of PARP-1 are not required for NF- $\kappa$ B coactivator function. *J Biol Chem* 276: 45588–45597.
47. Gilmore TD (2006) Introduction to NF- $\kappa$ B: players, pathways, perspectives. *Oncogene* 25: 6680–6684.
48. Wang Y, Dawson VL, Dawson TM (2009) Poly(ADP-ribose) signals to mitochondrial AIF: a key event in parthanatos. *Exp Neurol* 218: 193–202.
49. Perkins ND, Gilmore TD (2006) Good cop, bad cop: the different faces of NF- $\kappa$ B. *Cell Death Differ* 13: 759–772.
50. Wang Y, Keogh RJ, Hunter MG, Mitchell CA, Frey RS, et al. (2004) SHIP2 is recruited to the cell membrane upon macrophage colony-stimulating factor (M-CSF) stimulation and regulates M-CSF-induced signaling. *J Immunol* 173: 6820–6830.