

UC Irvine

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health

Title

International Perspective from Singapore on “Methemoglobinemia and Sulfhemoglobinemia in Two Pediatric Patients after Ingestion of Hydroxylamine Sulfate”

Permalink

<https://escholarship.org/uc/item/3cc8715j>

Journal

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 10(3)

ISSN

1936-900X

Authors

Lim, Hoon Chin
Tan, Hock Heng

Publication Date

2009

Copyright Information

Copyright 2009 by the author(s). All rights reserved unless otherwise indicated. Contact the author(s) for any necessary permissions. Learn more at <https://escholarship.org/terms>

Peer reviewed

International Perspective from Singapore on “Methemoglobinemia and Sulfhemoglobinemia in Two Pediatric Patients after Ingestion of Hydroxylamine Sulfate”

Hoon Chin Lim, MBBS
Hock Heng Tan, MBBS

Changi General Hospital, Department of Emergency Medicine, Singapore

Supervising Section Editor: Mark I. Langdorf, MD, MHPE
Submission history: Submitted June 02, 2009; Accepted June 14, 2009
Reprints available through open access at www.westjem.org
[WestJEM. 2009;10:202.]

Accidental ingestional poisoning among pediatric patients is a prevalent problem. In the absence of a well-designed national injury and poisoning surveillance system, cases often go unreported. Underreporting also occurs because many pediatric ingestions are trivial and not referred to a hospital for further medical management.¹ To reduce health risks posed by common household products, emphasis and enforcement of regulations specifying child-safe packaging are required.²

The authors described two cases of methemoglobinemia (metHbemia) and sulfhemoglobinemia (sulfHbemia) following ingestion of hydroxylamine sulfate, which were treated with methylene blue and exchange transfusion. Drugs or chemicals that cause sulfHbemia can also cause metHbemia, although it is not understood why the same chemical causes metHbemia in one person and sulfHbemia in another.³ Other ingestants that may cause methHbemia and sulfHbemia are dapsona, metoclopramide, phenacetin and phenazopyridine.

Emergency physicians (EP) in some developing countries may not have access to lab equipment with advanced co-oximeter capabilities to differentiate the two types of dyshemoglobinemia. In a time-pressured situation, some bedside investigation methods may be useful in guiding decisions for specific therapy like methylene blue in a cyanosed patient with suspected poisoning. One method, the filter paper test, helps to distinguish deoxyhemoglobin from dyshemoglobin as the darkly colored blood changes to bright red after blowing some oxygen over it. No changes occur with metHb or sulfHb. To distinguish metHb from sulfHb, the addition of a few drops of potassium cyanide changes the chocolate brown of metHb to bright red as cyanometHb is formed. No reaction occurs with sulfHb.⁴ Therapeutic response to methylene blue will also aid in the diagnosis of underlying metHbemia. The response is usually fairly rapid, within 30 minutes to one hour.

EPs should consider several differentials for apparent metHbemia that does not respond to methylene blue treatment.

These include older equipment incapable of distinguishing sulfHb from metHb due to limited co-oximeter capability; Hb M disease prone to metHb formation that resists reduction; glucose-6-phosphate dehydrogenase (G6PD) deficiency; NADPH metHb reductase deficiency; poisoning with oxidizing compounds that have enterohepatic circulation, which causes prolonged elevation of metHb; and overdosing of methylene blue itself, which is an oxidizing agent.^{3,4}

Finally, G6PD deficiency, one of the most prevalent disease-causing mutations worldwide, has several variants in Asia. Patient's status will influence decision when using methylene blue as treatment for metHbemia because methylene blue itself may induce hemolysis (through development of Heinz bodies) and cause paradoxical metHbemia, especially in G6PD deficient patients.⁴

Address for Correspondence: Dr Hoon Chin Lim, Department of Emergency Medicine, Changi General Hospital, 2 Simei Street 3, Singapore 529889, Republic of Singapore. Email: hoon_chin_lim@cgh.com.sg

REFERENCES

1. Hon KL, Ho JK, Leung TF, et al. Review of children hospitalised for ingestion and poisoning at a tertiary centre. *Ann Acad Med Singapore*. 2005; 34:356-61.
2. HC Lim. Mothballs: bringing safety issues out from the closet. *Singapore Med J*. 2006; 47:1003.
3. Curry SC. Hematologic syndrome hemolysis, methemoglobinemia and sulfhemoglobiemia. In: Brent J, Wallace KL, Burkhart KK, et al., eds. *Critical Care Toxicology – Diagnosis and Management of the Critically Poisoned Patient*. Philadelphia, PA: Elsevier Mosby; 2005:335-44.
4. Wright RO, Lewander WJ, Woolf AD. Methemoglobinemia: etiology, pharmacology and clinical management. *Ann Emerg Med*. 1999; 34:646-56.