

UC Davis

UC Davis Previously Published Works

Title

Author Correction: Plk1 Regulates the Repressor Function of FoxM1b by inhibiting its Interaction with the Retinoblastoma Protein.

Permalink

<https://escholarship.org/uc/item/3fw5z42t>

Journal

Scientific Reports, 8(1)

Authors

Mukhopadhyay, Nishit
Chand, Vaibhav
Pandey, Akshay
et al.

Publication Date

2018-10-29

DOI


10.1038/s41598-018-34071-w

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

SCIENTIFIC REPORTS

**OPEN**

Author Correction: Plk1 Regulates the Repressor Function of FoxM1b by inhibiting its Interaction with the Retinoblastoma Protein

Nishit K. Mukhopadhyay¹, Vaibhav Chand¹, Akshay Pandey¹, Dragana Kopanja¹, Janai R. Carr², Yi-Ju Chen³, Xiubei Liao¹ & Pradip Raychaudhuri^{1,4}

Correction to: *Scientific Reports* <https://doi.org/10.1038/srep46017>, published online 07 April 2017

This Article contains an error in Figure 5, where the T7-FoxM1 panels have been erroneously written as 100 and not 25. The correct Figure 5 appears below as Figure 1.

¹Department of Biochemistry and Molecular Genetics (M/C 669), University of Illinois, College of Medicine, 900 S, USA Ashland Ave., Chicago, IL, 60607, USA. ²Department of Hematology/Oncology, University of California, Los Angeles, CA, USA. ³Abramson Family Cancer Research Institute, University of Pennsylvania, Philadelphia, PA, 19104, USA. ⁴Jesse Brown VA Medical Center, 820 S. Damen Ave., Chicago, IL, 60612, USA. Nishit K. Mukhopadhyay and Vaibhav Chand contributed equally. Correspondence and requests for materials should be addressed to P.R. (email: pradip@uic.edu)





Figure 1. A Plk1-site phospho-mimetic mutant of FoxM1b fails to bind Rb *in vitro*. T7-His tagged C-terminal FoxM1 (residues 508–748), Plk1-site phospho-mimetic DD mutant, and GST-Rb (residues 379–928) were all expressed separately in *E. coli*. The bacterial lysates of the wild type or DD mutant FoxM1 were mixed with the lysates containing either GST-Rb or GST-CBP-KIX and then were allowed to bind Ni-agarose column. The eluted proteins, after extensive washing of the column, were assayed for the presence of Rb and CBP by western blotting (A and B). The left lane in each of the panels indicates the absence of Rb or CBP-KIX in the column elute when GST-Rb or CBP-KIX were passed through the Ni column in absence of FoxM1.

 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018