

# **UCLA**

## **Proceedings of UCLA Health**

### **Title**

Incidental Finding of PFO After Home Sleep Apnea Test

### **Permalink**

<https://escholarship.org/uc/item/3fx8k1kh>

### **Journal**

Proceedings of UCLA Health, 27(1)

### **Authors**

Fong, Susie X.

Padilla, Alfonso J.

### **Publication Date**

2023-10-12

## CLINICAL VIGNETTE

# Incidental Finding of PFO After Home Sleep Apnea Test

Susie X. Fong, MD and Alfonso J. Padilla, MD

### *Introduction*

When a patient is referred for evaluation of a possible sleep disordered breathing disorder, a sleep study is typically ordered. The sleep study can be either a home sleep apnea test (HSAT) or an in-lab polysomnogram (PSG). Usually, the home sleep study is the first step taken in the evaluation given its low cost and convenience which increases patient access to testing.<sup>1</sup> The parameters monitored on a home sleep test, at a minimum, include respiratory effort, airflow, and oxygen saturation. The information reported from this type of study includes respiratory event index (REI), oxygen saturation nadir and baseline, and total evaluation time. The disadvantages of a HSAT include the lack of sleep stage monitoring in most kits, higher rate of technical error, and a higher false negative rate in comparison to PSGs.

Most home sleep studies have been validated against the gold standard, the in-lab polysomnogram which is an attended study with a minimum of the following: electroencephalographic activity, electrooculogram, electromyography, respiratory inductive plethysmography, airflow, snoring, oxygen saturation, body position, and videography.<sup>2</sup> While more specific and sensitive than a HSAT, this testing requires patient access to a sleep laboratory with geography and scheduling availability, and incurs a higher cost.

We present a patient presenting to sleep clinic for evaluation of snoring who completed a HSAT and subsequently found to have an anatomical heart defect, due to an abnormal result found on sleep testing. The purpose of this report is to familiarize providers with home sleep apnea testing and interpretation of findings.

### *Case Presentation*

A 69-year-old male with hypertension was referred by his primary care provider for snoring. His risk factors for sleep disordered breathing included snoring, hypertension, age greater than 50, neck circumference greater than 17 inches, and male gender. STOP BANG, a risk assessment tool for sleep apnea, was 5/8 which puts him at high risk for sleep disordered breathing. Epworth Sleepiness Scale was 0/8.

A HSAT was obtained with the following results:

Respiratory events index (REI): 12.7/hour

O2 nadir 82% from baseline 88%

Time O2 saturation less than 88%: 221 minutes

Total evaluation time: 421 minutes

The patient's REI of 12.7 per hour was diagnostic for mild obstructive sleep apnea. However, what stood out in his study was the extent of his nocturnal hypoxemia. His oxygen saturation was less than 88% for 221 minutes, which was discordant to his mild sleep apnea. The patient elected to not treat his mild sleep apnea given he was asymptomatic. He had no known pulmonary or cardiac conditions which would explain the hypoxemia. His body mass index of 35 placed him at risk for obesity hypoventilation syndrome but a recent bicarbonate of 23 made this less likely. Given the severity of his nocturnal hypoxemia, he was referred for pulmonology for further evaluation and an echocardiogram with bubble study was ordered.

The trans-thoracic echocardiogram with bubble study revealed a possible patent foramen ovale (PFO) during the injection of agitated normal saline. A subsequent trans-esophageal echocardiogram revealed an atrial septal aneurysm with a small PFO, confirmed by a positive bubble study. The patient underwent successful PFO closure by Interventional Cardiology.

### *Discussion*

While a polysomnogram is the gold standard for evaluation of sleep disordered breathing, most insurers prefer HSATs be the first step in evaluation. HSATs are more economical and provided noninferior results when compared to polysomnograms.<sup>3-5</sup> A polysomnogram should be considered in situations where other sleep conditions such as central sleep apnea, nighttime hypoventilation or narcolepsy, is suspected. Patients with increased risk for such conditions generally have GOLD stage 2-4 COPD, NYHA class III-IV heart failure, awake hypoventilation, neuromuscular weakness, opioid use, or history of stroke.<sup>6</sup> These patients should be referred to sleep medicine as the evaluation may require additional specific testing parameters besides a baseline polysomnogram. HSATs have a reported failure rate ranging from 3% to 33%.<sup>7</sup> If a HSAT has been attempted and found to be invalid due to technical error, a polysomnogram should be the next step in evaluation.

The main results obtained from HSATs include REI, oxygen saturation, and total evaluation time. The REI is calculated by the total number of apneas and hypopneas divided by the total evaluation time. Because HSATs cannot detect sleep stages, total sleep time is not used in the calculation as in polysomnograms. As total evaluation time is longer than total sleep time, the number of apneas and hypopneas in a HSAT are divided by a larger denominator leading to an underestimation of the true ratio. Therefore, false negatives are more likely in comparison to polysomnograms. If a patient has a high pretest probability for sleep disordered breathing and a negative HSAT, the next step is to evaluate further with a polysomnogram.

Our patient had a REI of 12 per hour which is diagnostic of mild obstructive sleep apnea. He was not symptomatic from his condition, and elected to not treat at this time. Usually, the oxygen saturation data is used to corroborate with the sleep apnea diagnosis. However, his condition was mild and discordant with the severity of nocturnal hypoxemia. This prompted obtaining a transthoracic echocardiogram, and referral to Pulmonology for evaluation of other etiologies.

This case summarizes the use of HSATs for evaluation of sleep disordered breathing, their advantages and disadvantages, and appropriate interpretation. It demonstrates that additional helpful information can be obtained in a HSAT, alerting to a non-sleep-related diagnosis.

## REFERENCES

1. **Kapur VK, Auckley DH, Chowdhuri S, Kuhlmann DC, Mehra R, Ramar K, Harrod CG.** Clinical Practice Guideline for Diagnostic Testing for Adult Obstructive Sleep Apnea: An American Academy of Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med.* 2017 Mar 15;13(3):479-504. doi: 10.5664/jcsm.6506. PMID: 28162150; PMCID: PMC5337595.
2. **Epstein LJ, Kristo D, Strollo PJ Jr, Friedman N, Malhotra A, Patil SP, Ramar K, Rogers R, Schwab RJ, Weaver EM, Weinstein MD; Adult Obstructive Sleep Apnea Task Force of the American Academy of Sleep Medicine.** Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *J Clin Sleep Med.* 2009 Jun 15;5(3):263-76. PMID: 19960649; PMCID: PMC2699173.
3. **Kim RD, Kapur VK, Redline-Bruch J, Rueschman M, Auckley DH, Benca RM, Foldvary-Schafer NR, Iber C, Zee PC, Rosen CL, Redline S, Ramsey SD.** An Economic Evaluation of Home Versus Laboratory-Based Diagnosis of Obstructive Sleep Apnea. *Sleep.* 2015 Jul 1;38(7):1027-37. doi: 10.5665/sleep.4804. PMID: 26118558; PMCID: PMC4481018.
4. **Mulgrew AT, Fox N, Ayas NT, Ryan CF.** Diagnosis and initial management of obstructive sleep apnea without polysomnography: a randomized validation study. *Ann Intern Med.* 2007 Feb 6;146(3):157-66. doi: 10.7326/0003-4819-146-3-200702060-00004. PMID: 17283346.
5. **Berry RB, Hill G, Thompson L, McLaurin V.** Portable monitoring and autotitration versus polysomnography for the diagnosis and treatment of sleep apnea. *Sleep.* 2008 Oct;31(10):1423-31. PMID: 18853940; PMCID: PMC2572748.
6. **Chang JL, Goldberg AN, Alt JA, Mohammed A, Ashbrook L, Auckley D, et al.** International Consensus Statement on Obstructive Sleep Apnea. *Int Forum Allergy Rhinol.* 2023 Jul;13(7):1061-1482. doi: 10.1002/alr.23079. Epub 2023 Mar 30. PMID: 36068685; PMCID: PMC10359192.
7. **Lux L, Boehlecke B, Lohr KN.** Effectiveness of Portable Monitoring Devices for Diagnosing Obstructive Sleep Apnea: Update of a Systematic Review [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2004 Sep 1. PMID: 26065047.