

Lawrence Berkeley National Laboratory

Recent Work

Title

MEASUREMENT OF ORIENTATION OF SURFACTANT MOLECULES AT A LIQUID-AIR INTERFACE BY OPTICAL SECOND-HARMONIC GENERATION

Permalink

<https://escholarship.org/uc/item/3hq8m1sw>

Author

Rasing, Th.

Publication Date

1986-12-01

LBL-22734
c2

LBL-22734

RECEIVED
LAWRENCE
BERKELEY LABORATORY

MAR 2 1987

LIBRARY AND
DOCUMENTS SECTION

To be presented at the 1987
International Congress on
Membranes and Membrane
Processes, Tokyo, Japan,
June 8-12, 1987

MEASUREMENT OF ORIENTATION OF
SURFACTANT MOLECULES AT A
LIQUID-AIR INTERFACE BY
OPTICAL SECOND-HARMONIC
GENERATION

Th. Rasing, Y.R. Shen, M.W. Kim,
P. Valint, Jr., and J. Bock

December 1986

CCM

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.*

Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Prepared for the U.S. Department of Energy
under Contract DE-AC03-76SF00098

**Center
for
Advanced
Materials**

LBL-22734
c2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

MEASUREMENT OF ORIENTATION OF SURFACTANT MOLECULES AT A LIQUID-AIR INTERFACE
BY OPTICAL SECOND-HARMONIC GENERATION

Th. Rasing,* Y. R. Shen,* Mahn Won Kim,† P. Valint, Jr.,† and J. Bock†

*Department of Physics, University of California and Center for Advanced Materials, Lawrence Berkeley Laboratory, Berkeley California 94720 USA

†Exxon Research and Engineering Company, Anandale, New Jersey 08801 USA

ABSTRACT: The average molecular orientation of a monolayer of sodium-dodecyl-naphthalene-sulfonate on a water-air interface has been determined by use of optical second-harmonic generation. The molecular orientation is shown to vary smoothly with increasing surface density of the molecules.

Insoluble molecular monolayers at gas-liquid or liquid-liquid interfaces provide an insight to the understanding of surfactants, wetting, microemulsions, and membrane structures.¹ In the interpretation of the observed properties of these systems, various assumptions about the molecular orientations are often made,² but so far, few clear experimental data are available.

In this paper we present a measurement of the molecular orientation of a monolayer of surfactant molecules at a water-air interface using the newly developed optical second-harmonic generation (SHG) technique.³ The molecules under investigation are sodium-dodecyl-naphthalene-sulfonate (SDNS) $[\text{CH}_3(\text{CH}_2)_{11}-\text{C}_{10}\text{H}_6-\text{SO}_3\text{Na}]$, where C_{10}H_6 is a double benzene structure]. Figure 1 shows the measured surface pressure π as a function of the surface area per molecule (A) for SDNS on water containing 2% NaCl. The π - A diagram does not exhibit any discontinuous phase transition usually observed in this pressure range for this type of molecular monolayer.²

That optical SHG is an effective surface probe has been demonstrated recently in a number of cases.³⁻⁶ It is based on the idea that SHG is forbidden in centrosymmetric media but allowed at the interfaces where the inversion symmetry is necessarily broken. The surface nonlinear susceptibility $\chi_S^{(2)}$ which is responsible for the SHG at an interface generally reflects the properties of the surface layer. If $\chi_S^{(2)}$ arises mainly from a monolayer of molecular adsorbates, and takes the form

$$\chi_S^{(2)} = N_S \langle \bar{\alpha}^{(2)} \rangle, \quad (1)$$

where N_S is the surface density of the molecules, and $\langle \bar{\alpha}^{(2)} \rangle$ is the nonlinear polarizability averaged over the molecular orientational distribution, then a measurement of $\chi_S^{(2)}$ should enable us to obtain information about the orientation of the adsorbates. This is particularly true for rodlike molecules whose $\bar{\alpha}^{(2)}$ is dominated by a single element $\alpha_{\xi\xi\xi}^{(2)}$ along the molecular axis $\hat{\xi}$. For example, when the orientational distribution of the molecules in the azimuthal plane is random, the nonvanishing components of $\chi_S^{(2)}$ can be written as⁴

$$\begin{aligned} \chi_{S,111}^{(2)} &= N_S \langle \cos^3 \theta \rangle \alpha_{\xi\xi\xi}^{(2)}, \\ \chi_{S,1\bar{1}\bar{1}}^{(2)} &= \chi_{S,111}^{(2)} = \chi_{S,\bar{1}\bar{1}\bar{1}}^{(2)} = \frac{1}{2} N_S \langle \cos \theta \sin^2 \theta \rangle \alpha_{\xi\xi\xi}^{(2)}. \end{aligned} \quad (2)$$

where θ is the polar angle between the molecular axis and the surface normal, and the subindices $\bar{1}$ and 1 refer to directions perpendicular and parallel to the surface, respectively. We see from Eq. (2) that a measurement of the ratio of any two linear combinations of $\chi_{S,111}^{(2)}$ and $\chi_{S,\bar{1}\bar{1}\bar{1}}^{(2)}$ can yield a weighted average of θ . The average value or the most probable value of θ can then be deduced by assuming a certain orientational distribution function.

In analyzing the data, we assumed a δ function for the orientational distribution in θ . This means that the orientation of SDNS would be specified by a single value of θ . Figure 2 then gives the result on how θ of SDNS on water varies with the surface pressure π . It appears that at high pressures towards a saturated monolayer, the molecules tilted at $\sim 30^\circ$ from the surface normal, while at low pressures, they inclined more towards the surface plane (see also Ref. 7).

In conclusion, we have shown for the first time how optical second-harmonic generation can be used to measure the average molecular orientation of a molecular monolayer at a water-air interface. The

orientation of the SDNS molecules appears to vary continuously and tilt more towards the surface normal with increasing surface pressure. It shows no discontinuity in the variation and approaches a limiting inclination angle - 30° .

This work was supported by the Director, Office of Energy Research, Office of Energy Research, Office of Basic Energy Sciences, Materials Sciences Division of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.




Fig. 1 Surface pressure π of SDNS as a function of the area per molecule A on a water surface containing 2% NaCl.




Fig. 2 Tilt angle θ between the molecular axis and the surface normal as a function of the surface pressure π for SDNS on water containing 2% NaCl.

REFERENCES

1. Micellization, Solubilization and Microemulsions, ed. K. L. Mittal (Plenum, New York, 1977); J. S. Huang and M. W. Kim, *Phys. Rev. Lett.* **47**, 1463 (1981).
2. For a review, see, e.g., G. M. Bell, L. L. Coombs, and L. J. Dunne, *Chem. Rev.* **81**, 15 (1981), and references therein.
3. C. K. Chen, T. F. Heinz, D. Ricard, and Y. R. Shen, *Phys. Rev. Lett.* **46**, 15 (1981).
4. T. F. Heinz, H. W. K. Tom, and Y. R. Shen, *Phys. Rev. A* **28**, 1883 (1983).
5. H. W. K. Tom, T. F. Heinz, and Y. R. Shen, *Phys. Rev. Lett.* **51**, 1983 (1983).
6. H. W. K. Tom, C. M. Made, X. D. Zhu, J. E. Crowell, T. F. Heinz, G. A. Somorjai, and Y. R. Shen, *Phys. Rev. Lett.* **52**, 348 (1984).
7. Th. Rasing, Y. R. Shen, M. W. Kim, P. Valint, Jr., and J. Bock, *Phys. Rev. A* **31**, 537 (1985).

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*