

UC Irvine

UC Irvine Previously Published Works

Title

Breast Cancer Polygenic Risk Score and Contralateral Breast Cancer Risk

Permalink

<https://escholarship.org/uc/item/3hw3g9q1>

Journal

American Journal of Human Genetics, 107(5)

ISSN

0002-9297

Authors

Kramer, Iris
Hooning, Maartje J
Mavaddat, Nasim
et al.

Publication Date

2020-11-01

DOI

10.1016/j.ajhg.2020.09.001

Peer reviewed

Breast Cancer Polygenic Risk Score and Contralateral Breast Cancer Risk

Iris Kramer,¹ Maartje J. Hooning,² Nasim Mavaddat,³ Michael Hauptmann,^{4,5} Renske Keeman,¹ Ewout W. Steyerberg,^{6,7} Daniele Giardiello,^{1,6} Antonis C. Antoniou,³ Paul D.P. Pharoah,^{3,8} Sander Canisius,^{1,9} Zumuruda Abu-Ful,¹⁰ Irene L. Andrulis,^{11,12} Hoda Anton-Culver,¹³ Kristan J. Aronson,¹⁴ Annelie Augustinsson,¹⁵ Heiko Becher,^{16,17} Matthias W. Beckmann,¹⁸ Sabine Behrens,¹⁹ Javier Benitez,^{20,21} Marina Bermisheva,²² Natalia V. Bogdanova,^{23,24,25} Stig E. Bojesen,^{26,27,28} Manjeet K. Bolla,³ Bernardo Bonanni,²⁹ Hiltrud Brauch,^{30,31,32} Michael Bremer,²³ Sara Y. Brucker,³³ Barbara Burwinkel,^{34,35} Jose E. Castelao,³⁶ Tsun L. Chan,^{37,38} Jenny Chang-Claude,^{19,39} Stephen J. Chanock,⁴⁰ Georgia Chenevix-Trench,⁴¹ Ji-Yeob Choi,^{42,43}

(Author list continued on next page)

Summary

Previous research has shown that polygenic risk scores (PRSs) can be used to stratify women according to their risk of developing primary invasive breast cancer. This study aimed to evaluate the association between a recently validated PRS of 313 germline variants (PRS₃₁₃) and contralateral breast cancer (CBC) risk. We included 56,068 women of European ancestry diagnosed with first invasive breast cancer from 1990 onward with follow-up from the Breast Cancer Association Consortium. Metachronous CBC risk (N = 1,027) according to the distribution of PRS₃₁₃ was quantified using Cox regression analyses. We assessed PRS₃₁₃ interaction with age at first diagnosis, family history, morphology, ER status, PR status, and HER2 status, and (neo)adjuvant therapy. In studies of Asian women, with limited follow-up, CBC risk associated with PRS₃₁₃ was assessed using logistic regression for 340 women with CBC compared with 12,133 women with unilateral breast cancer. Higher PRS₃₁₃ was associated with increased CBC risk: hazard ratio per standard deviation (SD) = 1.25 (95%CI = 1.18–1.33) for Europeans, and an OR per SD = 1.15 (95%CI = 1.02–1.29) for Asians. The absolute lifetime risks of CBC, accounting for death as competing risk, were 12.4% for European women at the 10th percentile and 20.5% at the 90th percentile of PRS₃₁₃. We found no evidence of confounding by or interaction with individual characteristics, characteristics of the primary tumor, or treatment. The C-index for the PRS₃₁₃ alone was 0.563 (95%CI = 0.547–0.586). In conclusion, PRS₃₁₃ is an independent factor associated with CBC risk and can be incorporated into CBC risk prediction models to help improve stratification and optimize surveillance and treatment strategies.

Introduction

Due to the high incidence of breast cancer and improving survival, an increasing number of breast cancer survivors are at risk of developing contralateral breast cancer

(CBC). The 10-year cumulative incidence of CBC is ~4%,^{1,2} but estimates vary widely depending on factors such as germline genetics, family history, and (neo)adjuvant systemic therapy for the first breast cancer.³ The risk of developing CBC is particularly high in women with

¹The Netherlands Cancer Institute - Antoni van Leeuwenhoek Hospital, Division of Molecular Pathology, Amsterdam 1066 CX, the Netherlands; ²Erasmus MC Cancer Institute, Department of Medical Oncology, Rotterdam 3015 CN, the Netherlands; ³University of Cambridge, Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, Cambridge CB1 8RN, UK; ⁴The Netherlands Cancer Institute - Antoni van Leeuwenhoek Hospital, Department of Epidemiology and Biostatistics, Amsterdam 1066 CX, the Netherlands; ⁵Brandenburg Medical School Theodor Fontane, Institute of Biostatistics and Registry Research, Neuruppin 16816, Germany; ⁶Leiden University Medical Center, Department of Biomedical Data Sciences, Leiden 2333 ZA, the Netherlands; ⁷Erasmus MC, Department of Public Health, Rotterdam 3015 GD, the Netherlands; ⁸University of Cambridge, Centre for Cancer Genetic Epidemiology, Department of Oncology, Cambridge CB1 8RN, UK; ⁹The Netherlands Cancer Institute - Antoni van Leeuwenhoek Hospital, Division of Molecular Carcinogenesis, Amsterdam 1066 CX, the Netherlands; ¹⁰Carmel Medical Center and Technion Faculty of Medicine, Clalit National Cancer Control Center, Haifa 35254, Israel; ¹¹Lunenfeld-Tanenbaum Research Institute of Mount Sinai Hospital, Fred A. Litwin Center for Cancer Genetics, Toronto, ON M5G 1X5, Canada; ¹²University of Toronto, Department of Molecular Genetics, Toronto, ON M5S 1A8, Canada; ¹³University of California Irvine, Department of Epidemiology, Genetic Epidemiology Research Institute, Irvine, CA 92617, USA; ¹⁴Queen's University, Department of Public Health Sciences, and Cancer Research Institute, Kingston, ON K7L 3N6, Canada; ¹⁵Lund University, Department of Cancer Epidemiology, Clinical Sciences, Lund 222 42, Sweden; ¹⁶University Medical Center Hamburg-Eppendorf, Institute of Medical Biometry and Epidemiology, Hamburg 20246, Germany; ¹⁷Charité – Universitätsmedizin Berlin, Institute of Biometry and Clinical Epidemiology, Berlin 10117, Germany; ¹⁸University Hospital Erlangen, Friedrich-Alexander-University Erlangen-Nuremberg, Department of Gynecology and Obstetrics, Comprehensive Cancer Center ER-EMN, Erlangen 91054, Germany; ¹⁹German Cancer Research Center (DKFZ), Division of Cancer Epidemiology, Heidelberg 69120, Germany; ²⁰Centro de Investigación en Red de Enfermedades Raras (CIBERER), Madrid 28029, Spain; ²¹Spanish National Cancer Research Centre (CNIO), Human Cancer Genetics Programme, Madrid 28029, Spain; ²²Ufa Federal Research Centre of the Russian Academy of Sciences, Institute of Biochemistry and Genetics, Ufa 450054, Russia; ²³Hannover Medical School, Department of Radiation Oncology, Hannover 30625, Germany; ²⁴Hannover Medical School, Gynaecology Research Unit, Hannover 30625, Germany; ²⁵N.N. Alexandrov Research Institute of Oncology and Medical Radiology, Minsk 223040, Belarus; ²⁶Copenhagen University Hospital, Copenhagen General Population Study, Herlev and Gentofte Hospital, Herlev 2730, Denmark; ²⁷Copenhagen University Hospital, Department of Clinical Biochemistry,

(Affiliations continued on next page)



Christine L. Clarke,⁴⁴ NBCS Collaborators,^{45,46,47,48,49,50,51,52,53,54,55,56} J. Margriet Collée,⁵⁷ Fergus J. Couch,⁵⁸ Angela Cox,⁵⁹ Simon S. Cross,⁶⁰ Kamila Czene,⁶¹ Mary B. Daly,⁶² Peter Devilee,^{63,64} Thilo Dörk,²⁴ Isabel dos-Santos-Silva,⁶⁵ Alison M. Dunning,⁸ Miriam Dwek,⁶⁶ Diana M. Eccles,⁶⁷ D. Gareth Evans,^{68,69} Peter A. Fasching,^{18,70} Henrik Flyger,⁷¹ Manuela Gago-Dominguez,^{72,73} Montserrat García-Closas,⁴⁰ José A. García-Sáenz,⁷⁴ Graham G. Giles,^{75,76,77} David E. Goldgar,⁷⁸ Anna González-Neira,²¹ Christopher A. Haiman,⁷⁹ Niclas Håkansson,⁸⁰ Ute Hamann,⁸¹ Mikael Hartman,^{82,83} Bernadette A.M. Heemskerk-Gerritsen,² Antoinette Hollestelle,² John L. Hopper,⁷⁶ Ming-Feng Hou,⁸⁴ Anthony Howell,⁸⁵ ABCTB Investigators,⁸⁶ kConFab Investigators^{87,88} Hidemi Ito,^{89,90} Milena Jakimovska,⁹¹ Anna Jakubowska,^{92,93} Wolfgang Janni,⁹⁴ Esther M. John,⁹⁵ Audrey Jung,¹⁹ Daehee Kang,^{42,43,96} C. Marleen Kets,⁹⁷ Elza Khusnutdinova,^{22,98} Yon-Dschun Ko,⁹⁹ Vessela N. Kristensen,^{45,56} Allison W. Kurian,^{95,100} Ava Kwong,^{37,101,102} Diether Lambrechts,^{103,104} Loic Le Marchand,¹⁰⁵ Jingmei Li,¹⁰⁶ Annika Lindblom,^{107,108} Jan Lubiński,⁹² Arto Mannermaa,^{109,110,111} Mehdi Manoochehri,⁸¹ Sara Margolin,^{112,113} Keitaro Matsuo,^{89,90} Dimitrios Mavroudis,¹¹⁴ Alfons Meindl,¹¹⁵ Roger L. Milne,^{75,76,77} Anna Marie Mulligan,^{116,117} Taru A. Murañen,¹¹⁸ Susan L. Neuhausen,¹¹⁹ Heli Nevanlinna,¹¹⁸ William G. Newman,^{68,69} Andrew F. Olshan,¹²⁰ Janet E. Olson,¹²¹ Håkan Olsson,¹⁵

(Author list continued on next page)

Herlev and Gentofte Hospital, Herlev 2730, Denmark; ²⁸University of Copenhagen, Faculty of Health and Medical Sciences, Copenhagen 2200, Denmark; ²⁹IEO, European Institute of Oncology IRCCS, Division of Cancer Prevention and Genetics, Milan 20141, Italy; ³⁰Dr. Margarete Fischer-Bosch-Institute of Clinical Pharmacology, Stuttgart 70376, Germany; ³¹University of Tübingen, iFIT-Cluster of Excellence, Tübingen 72074, Germany; ³²German Cancer Research Center (DKFZ), German Cancer Consortium (DKTK), Partner Site Tübingen, Tübingen 72074, Germany; ³³University of Tübingen, Department of Gynecology and Obstetrics, Tübingen 72076, Germany; ³⁴German Cancer Research Center (DKFZ), Molecular Epidemiology Group, C080, Heidelberg 69120, Germany; ³⁵University of Heidelberg, Molecular Biology of Breast Cancer, University Womens Clinic Heidelberg, Heidelberg 69120, Germany; ³⁶Instituto de Investigación Sanitaria Galicia Sur (IISGS), Xerencia de Xestión Integrada de Vigo-SERGAS, Oncology and Genetics Unit, Vigo 36312, Spain; ³⁷Hong Kong Hereditary Breast Cancer Family Registry, Hong Kong; ³⁸Hong Kong Sanatorium and Hospital, Department of Pathology, Happy Valley, Hong Kong; ³⁹University Medical Center Hamburg-Eppendorf, Cancer Epidemiology Group, University Cancer Center Hamburg (UCCH), Hamburg 20246, Germany; ⁴⁰National Cancer Institute, National Institutes of Health, Department of Health and Human Services, Division of Cancer Epidemiology and Genetics, Bethesda, MD 20850, USA; ⁴¹QIMR Berghofer Medical Research Institute, Department of Genetics and Computational Biology, Brisbane, QLD 4006, Australia; ⁴²Seoul National University Graduate School, Department of Biomedical Sciences, Seoul 03080, Korea; ⁴³Seoul National University, Cancer Research Institute, Seoul 03080, Korea; ⁴⁴University of Sydney, Westmead Institute for Medical Research, Sydney, NSW 2145, Australia; ⁴⁵Oslo University Hospital-Radiumhospitalet, Department of Cancer Genetics, Institute for Cancer Research, Oslo 0379, Norway; ⁴⁶University of Oslo, Institute of Clinical Medicine, Faculty of Medicine, Oslo 0450, Norway; ⁴⁷Vestre Viken Hospital, Department of Research, Drammen 3019, Norway; ⁴⁸Oslo University Hospital-Ullevål, Section for Breast and Endocrine Surgery, Department of Cancer, Division of Surgery, Cancer and Transplantation Medicine, Oslo 0450, Norway; ⁴⁹Oslo University Hospital, Department of Radiology and Nuclear Medicine, Oslo 0379, Norway; ⁵⁰Akershus University Hospital, Department of Pathology, Lørenskog 1478, Norway; ⁵¹Oslo University Hospital, Department of Tumor Biology, Institute for Cancer Research, Oslo 0379, Norway; ⁵²Oslo University Hospital-Radiumhospitalet, Department of Oncology, Division of Surgery, Cancer and Transplantation Medicine, Oslo 0379, Norway; ⁵³Oslo University Hospital-Radiumhospitalet, National Advisory Unit on Late Effects after Cancer Treatment, Department of Oncology, Oslo 0379, Norway; ⁵⁴Akershus University Hospital, Department of Oncology, Lørenskog 1478, Norway; ⁵⁵Oslo University Hospital, Oslo Breast Cancer Research Consortium, Oslo 0379, Norway; ⁵⁶Oslo University Hospital and University of Oslo, Department of Medical Genetics, Oslo 0379, Norway; ⁵⁷Erasmus University Medical Center, Department of Clinical Genetics, Rotterdam 3015 CN, the Netherlands; ⁵⁸Mayo Clinic, Department of Laboratory Medicine and Pathology, Rochester, MN 55905, USA; ⁵⁹University of Sheffield, Sheffield Institute for Nucleic Acids (SiNFoNiA), Department of Oncology and Metabolism, Sheffield S10 2TN, UK; ⁶⁰University of Sheffield, Academic Unit of Pathology, Department of Neuroscience, Sheffield S10 2TN, UK; ⁶¹Karolinska Institutet, Department of Medical Epidemiology and Biostatistics, Stockholm 171 65, Sweden; ⁶²Fox Chase Cancer Center, Department of Clinical Genetics, Philadelphia, PA 19111, USA; ⁶³Leiden University Medical Center, Department of Pathology, Leiden 2333 ZA, the Netherlands; ⁶⁴Leiden University Medical Center, Department of Human Genetics, Leiden 2333 ZA, the Netherlands; ⁶⁵London School of Hygiene and Tropical Medicine, Department of Non-Communicable Disease Epidemiology, London WC1E 7HT, UK; ⁶⁶University of Westminster, School of Life Sciences, London W1B 2HW, UK; ⁶⁷University of Southampton, Faculty of Medicine, Southampton SO17 1BJ, UK; ⁶⁸University of Manchester, Manchester Academic Health Science Centre, Division of Evolution and Genomic Sciences, School of Biological Sciences, Faculty of Biology, Medicine and Health, Manchester M13 9WL, UK; ⁶⁹St Mary's Hospital, Manchester University NHS Foundation Trust, Manchester Academic Health Science Centre, North West Genomics Laboratory Hub, Manchester Centre for Genomic Medicine, Manchester M13 9WL, UK; ⁷⁰University of California at Los Angeles, David Geffen School of Medicine, Department of Medicine Division of Hematology and Oncology, Los Angeles, CA 90095, USA; ⁷¹Copenhagen University Hospital, Department of Breast Surgery, Herlev and Gentofte Hospital, Herlev 2730, Denmark; ⁷²Grupo de Medicina Xenómica, Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS), Fundación Pública Galega de Medicina Xenómica, Santiago de Compostela 15706, Spain; ⁷³University of California San Diego, Moores Cancer Center, La Jolla, CA 92037, USA; ⁷⁴Instituto de Investigación Sanitaria San Carlos (IdISSC), Centro Investigación Biomédica en Red de Cáncer (CIBERONC), Medical Oncology Department, Hospital Clínico San Carlos, Madrid 28040, Spain; ⁷⁵Cancer Council Victoria, Cancer Epidemiology Division, Melbourne, VIC 3004, Australia; ⁷⁶The University of Melbourne, Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, Melbourne, VIC 3010, Australia; ⁷⁷Monash University, Precision Medicine, School of Clinical Sciences at Monash Health, Clayton, VIC 3168, Australia; ⁷⁸Huntsman Cancer Institute, University of Utah School of Medicine, Department of Dermatology, Salt Lake City, UT 84112, USA; ⁷⁹University of Southern California, Department of Preventive Medicine, Keck School of Medicine, Los Angeles, CA 90033, USA; ⁸⁰Karolinska Institutet, Institute of Environmental Medicine, Stockholm 171 77, Sweden; ⁸¹German Cancer Research Center (DKFZ), Molecular Genetics of Breast Cancer, Heidelberg 69120, Germany; ⁸²National University of Singapore and National University Health System, Saw Swee Hock School of Public Health, Singapore 119077, Singapore; ⁸³National University Health System, Department of Surgery, Singapore 119228, Singapore; ⁸⁴Kaohsiung Medical University, Chung-Ho Memorial Hospital, Kaohsiung 807, Taiwan; ⁸⁵University of Manchester, Division of Cancer Sciences, Manchester M13 9PL, UK; ⁸⁶University of Sydney, Australian Breast Cancer Tissue Bank, Westmead Institute for Medical Research, Sydney, NSW 2145, Australia; ⁸⁷Peter MacCallum Cancer Center, Melbourne, VIC 3000, Australia; ⁸⁸The University of Melbourne, Sir Peter MacCallum Department of Oncology, Melbourne, VIC 3000, Australia; ⁸⁹Aichi Cancer Center Research Institute, Division of Cancer Epidemiology and Prevention, Nagoya 464-8681, Japan; ⁹⁰Nagoya University Graduate School of Medicine, Division of Cancer Epidemiology, Nagoya 466-8550, Japan; ⁹¹MASA, Research Centre for Genetic Engineering and Biotechnology 'Georgi D. Efremov', Skopje 1000, Republic of North Macedonia; ⁹²Pomeranian Medical

(Affiliations continued on next page)

Tjong-Won Park-Simon,²⁴ Julian Peto,⁶⁵ Christos Petridis,¹²² Dijana Plaseska-Karanfilska,⁹¹ Nadege Presneau,⁶⁶ Katri Pylkäs,^{123,124} Paolo Radice,¹²⁵ Gad Rennert,¹⁰ Atocha Romero,¹²⁶ Rebecca Roylance,¹²⁷ Emmanouil Saloustros,¹²⁸ Elinor J. Sawyer,¹²⁹ Rita K. Schmutzler,^{130,131,132} Lukas Schwentner,⁹⁴ Christopher Scott,¹²¹ Mee-Hoong See,¹³³ Mitul Shah,⁸ Chen-Yang Shen,^{134,135} Xiao-Ou Shu,¹³⁶ Sabine Siesling,^{137,138} Susan Slager,¹²¹ Christof Sohn,¹³⁹ Melissa C. Southey,^{75,77,140} John J. Spinelli,^{141,142} Jennifer Stone,^{76,143} William J. Tapper,⁶⁷ Maria Tengström,^{109,144,145} Soo Hwang Teo,^{146,147} Mary Beth Terry,¹⁴⁸ Rob A.E.M. Tollenaar,¹⁴⁹ Ian Tomlinson,^{150,151} Melissa A. Troester,¹²⁰ Celine M. Vachon,¹⁵² Chantal van Ongeval,¹⁵³ Elke M. van Veen,^{68,69} Robert Winqvist,^{123,124} Alicja Wolk,^{80,154} Wei Zheng,¹³⁶ Argyrios Ziogas,¹³ Douglas F. Easton,^{3,8} Per Hall,^{61,112} and Marjanka K. Schmidt^{1,155,*}

rare mutations in certain genes including *BRCA1*, *BRCA2*, and *CHEK2*, with approximately 2- to 4-fold higher risks reported compared with women without these mutations.³

Recently, genome-wide association studies (GWASs) have identified multiple common germline variants that

are associated with first primary breast cancer risk.^{4,5} These are associated with small differences in risk individually, but their combined effects can be summarized in a polygenic risk score (PRS), which has been shown to stratify women according to their risk of developing breast cancer.^{6–9} Using a large GWAS dataset from the Breast Cancer

University, Department of Genetics and Pathology, Szczecin 71-252, Poland; ⁹³Pomeranian Medical University, Independent Laboratory of Molecular Biology and Genetic Diagnostics, Szczecin 71-252, Poland; ⁹⁴University Hospital Ulm, Department of Gynaecology and Obstetrics, Ulm 89075, Germany; ⁹⁵Stanford Cancer Institute, Stanford University School of Medicine, Department of Epidemiology & Population Health, Stanford, CA 94304, USA; ⁹⁶Seoul National University College of Medicine, Department of Preventive Medicine, Seoul 03080, Korea; ⁹⁷the Netherlands Cancer Institute - Antoni van Leeuwenhoek Hospital, Department of Clinical Genetics, Amsterdam 1066 CX, the Netherlands; ⁹⁸Bashkir State University, Department of Genetics and Fundamental Medicine, Ufa 450000, Russia; ⁹⁹Johanniter Krankenhaus, Department of Internal Medicine, Evangelische Kliniken Bonn gGmbH, Bonn 53177, Germany; ¹⁰⁰Stanford University School of Medicine, Department of Health Research and Policy, Stanford, CA 94305, USA; ¹⁰¹The University of Hong Kong, Department of Surgery, Pok Fu Lam, Hong Kong; ¹⁰²Hong Kong Sanatorium and Hospital, Cancer Genetics Center and Department of Surgery, Happy Valley, Hong Kong; ¹⁰³VIB Center for Cancer Biology, Leuven 3001, Belgium; ¹⁰⁴University of Leuven, Laboratory for Translational Genetics, Department of Human Genetics, Leuven 3000, Belgium; ¹⁰⁵University of Hawaii Cancer Center, Epidemiology Program, Honolulu, HI 96813, USA; ¹⁰⁶Genome Institute of Singapore, Human Genetics Division, Singapore 138672, Singapore; ¹⁰⁷Karolinska Institutet, Department of Molecular Medicine and Surgery, Stockholm 171 76, Sweden; ¹⁰⁸Karolinska University Hospital, Department of Clinical Genetics, Stockholm 171 76, Sweden; ¹⁰⁹University of Eastern Finland, Translational Cancer Research Area, Kuopio 70210, Finland; ¹¹⁰University of Eastern Finland, Institute of Clinical Medicine, Pathology and Forensic Medicine, Kuopio 70210, Finland; ¹¹¹Kuopio University Hospital, Biobank of Eastern Finland, Kuopio 70210, Finland; ¹¹²Södersjukhuset, Department of Oncology, Stockholm 118 83, Sweden; ¹¹³Karolinska Institutet, Department of Clinical Science and Education, Södersjukhuset, Stockholm 118 83, Sweden; ¹¹⁴University Hospital of Heraklion, Department of Medical Oncology, Heraklion 711 10, Greece; ¹¹⁵University of Munich, Campus Großhadern, Department of Gynecology and Obstetrics, Munich 81377, Germany; ¹¹⁶University of Toronto, Department of Laboratory Medicine and Pathobiology, Toronto, ON M5S 1A8, Canada; ¹¹⁷University Health Network, Laboratory Medicine Program, Toronto, ON M5G 2C4, Canada; ¹¹⁸Helsinki University Hospital, Department of Obstetrics and Gynecology, University of Helsinki, Helsinki 00290, Finland; ¹¹⁹Beckman Research Institute of City of Hope, Department of Population Sciences, Duarte, CA 91010, USA; ¹²⁰University of North Carolina at Chapel Hill, Department of Epidemiology, Gillings School of Global Public Health and UNC Lineberger Comprehensive Cancer Center, Chapel Hill, NC 27599, USA; ¹²¹Mayo Clinic, Department of Health Sciences Research, Rochester, MN 55905, USA; ¹²²King's College London, Research Oncology, Guy's Hospital, London SE1 9RT, UK; ¹²³University of Oulu, Laboratory of Cancer Genetics and Tumor Biology, Cancer and Translational Medicine Research Unit, Biocenter Oulu, Oulu 90220, Finland; ¹²⁴Northern Finland Laboratory Centre Oulu, Laboratory of Cancer Genetics and Tumor Biology, Oulu 90220, Finland; ¹²⁵Fondazione IRCCS Istituto Nazionale dei Tumori (INT), Unit of Molecular Bases of Genetic Risk and Genetic Testing, Department of Research, Milan 20133, Italy; ¹²⁶Hospital Universitario Puerta de Hierro, Medical Oncology Department, Madrid 28222, Spain; ¹²⁷UCLH Foundation Trust, Department of Oncology, London NW1 2PG, UK; ¹²⁸University Hospital of Larissa, Department of Oncology, Larissa 411 10, Greece; ¹²⁹King's College London, School of Cancer & Pharmaceutical Sciences, Comprehensive Cancer Centre, Guy's Campus, London SE1 1UL, UK; ¹³⁰Faculty of Medicine and University Hospital Cologne, University of Cologne, Center for Familial Breast and Ovarian Cancer, Cologne 50937, Germany; ¹³¹Faculty of Medicine and University Hospital Cologne, University of Cologne, Center for Integrated Oncology (CIO), Cologne 50937, Germany; ¹³²Faculty of Medicine and University Hospital Cologne, University of Cologne, Center for Molecular Medicine Cologne (CMMC), Cologne 50931, Germany; ¹³³University of Malaya, Breast Cancer Research Unit, University Malaya Cancer Research Institute, Faculty of Medicine, Kuala Lumpur 50603, Malaysia; ¹³⁴Academia Sinica, Institute of Biomedical Sciences, Taipei 115, Taiwan; ¹³⁵China Medical University, School of Public Health, Taichung 40402, Taiwan; ¹³⁶Vanderbilt University School of Medicine, Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Nashville, TN 37232, USA; ¹³⁷Netherlands Comprehensive Cancer Organisation (IKNL), Department of Research, Utrecht 3511 DT, the Netherlands; ¹³⁸University of Twente, Department of Health Technology and Service Research, Technical Medical Center, Enschede 7522 NB, the Netherlands; ¹³⁹University Hospital and German Cancer Research Center, National Center for Tumor Diseases, Heidelberg 69120, Germany; ¹⁴⁰The University of Melbourne, Department of Clinical Pathology, Melbourne, VIC 3010, Australia; ¹⁴¹BC Cancer, Population Oncology, Vancouver, BC V5Z 1G1, Canada; ¹⁴²University of British Columbia, School of Population and Public Health, Vancouver, BC V6T 1Z4, Canada; ¹⁴³Curtin University and University of Western Australia, The Curtin UWA Centre for Genetic Origins of Health and Disease, Perth, WA 6000, Australia; ¹⁴⁴Kuopio University Hospital, Department of Oncology, Cancer Center, Kuopio 70210, Finland; ¹⁴⁵University of Eastern Finland, Institute of Clinical Medicine, Oncology, Kuopio 70210, Finland; ¹⁴⁶Cancer Research Malaysia, Breast Cancer Research Programme, Subang Jaya, Selangor 47500, Malaysia; ¹⁴⁷University of Malaya, Department of Surgery, Faculty of Medicine, Kuala Lumpur 50603, Malaysia; ¹⁴⁸Columbia University, Department of Epidemiology, Mailman School of Public Health, New York, NY 10032, USA; ¹⁴⁹Leiden University Medical Center, Department of Surgery, Leiden 2333 ZA, the Netherlands; ¹⁵⁰University of Birmingham, Institute of Cancer and Genomic Sciences, Birmingham B15 2TT, UK; ¹⁵¹University of Oxford, Wellcome Trust Centre for Human Genetics and Oxford NIHR Biomedical Research Centre, Oxford OX3 7BN, UK; ¹⁵²Mayo Clinic, Department of Health Science Research, Division of Epidemiology, Rochester, MN 55905, USA; ¹⁵³Leuven Cancer Institute, University Hospitals Leuven, Leuven Multidisciplinary Breast Center, Department of Radiology, Leuven 3000, Belgium; ¹⁵⁴Uppsala University, Department of Surgical Sciences, Uppsala 751 05, Sweden; ¹⁵⁵The Netherlands Cancer Institute - Antoni van Leeuwenhoek hospital, Division of Psychosocial Research and Epidemiology, Amsterdam 1066 CX, the Netherlands

*Correspondence: mk.schmidt@nki.nl
<https://doi.org/10.1016/j.ajhg.2020.09.001>

Association Consortium (BCAC), we previously developed and validated a 313-variant PRS (PRS₃₁₃) among women of European descent. In independent prospective studies, this PRS₃₁₃ predicted the risk of primary invasive breast cancer with an odds ratio (OR) per standard deviation (SD) of 1.61 (95% confidence interval (95%CI) = 1.57–1.65).⁷ The PRS₃₁₃ has also been externally validated using the UK Biobank cohort.

The aim of the current study was to evaluate the association between PRS₃₁₃ and CBC risk, using data from BCAC. Other studies have shown associations between risk of CBC and both a 67-variant PRS¹⁰ and individual variants,¹¹ but not yet with PRS₃₁₃, the most extensively validated PRS. Further, the dataset currently evaluated is larger than those previously tested. We carried out two types of analyses. We conducted a cohort study among studies of European ancestry women with follow-up data available and performed Cox regression analyses to estimate hazard ratios (HRs) for CBC. Potential confounding and interaction with characteristics of the individual, characteristics of the primary tumor, or treatment were tested. In addition, to directly compare with the OR reported for PRS₃₁₃ and first breast cancer, we selected case-case series and performed logistic regression analyses comparing the PRS₃₁₃ distribution in women with CBC versus those with unilateral breast cancer. These analyses were conducted separately in European and Asian women (follow-up was too limited to perform a cohort study for the Asian population). Use of PRS₃₁₃ may lead to more accurate CBC risk prediction to support decision making for women who may or may not benefit from additional surveillance and risk-reducing treatment strategies.

Material and Methods

Study Subjects

Case-Case Series

We selected women who were diagnosed with breast cancer and women without any diagnosis of breast cancer from the BCAC including all women of European ancestry, based on genotyping data, and selecting only those studies which reported on CBC (62 studies) (Figure S1A, Table S1 and S2). BCAC database version freeze 12 was used. All women diagnosed with invasive breast cancer as a first cancer were included in the analysis; the small number of tumors with unknown invasiveness were considered invasive (Table S2). In the case-case series, a CBC was defined as a breast cancer (*in situ* or invasive) in the contralateral breast irrespective of the time since the first breast cancer. The case-case series comprised 81,000 women with unilateral breast cancer, 3,607 women with CBC, and 62,830 women without any diagnosis of breast cancer (Figure S1A). We also compared women with unilateral breast cancer to women without any diagnosis of breast cancer to reproduce the estimate that was previously reported for first breast cancer risk⁷ in our study selection.

We selected for a separate analysis women of Asian ancestry from the BCAC data, comprising 12,133 women with unilateral breast cancer, 340 women with CBC, and 13,398 women without

any diagnosis of breast cancer from eight studies (Figure S1B, Table S2).

European Cohort

In the European cohort, we used metachronous CBC as the outcome, defined as a breast cancer in the contralateral breast (*in situ* or invasive) diagnosed at least 3 months after the first breast cancer. We used a cut-off of 3 months to reduce the likelihood that these CBCs represent metastases rather than true second primary tumors. We selected all women diagnosed with breast cancer from the European case-case series and excluded four studies that did not provide follow-up information on vital status (Figure S1A). We did not include Asian women since follow-up was too limited in these studies. We additionally excluded 6,207 women with no follow-up and 2,208 women who developed synchronous CBC, distant metastasis, or who died or were last known to be alive within 3 months after the first breast cancer diagnosis. Since BCAC also included prevalent cases, we excluded 3,796 women who developed CBC or were censored before study entry. The case-case series included women diagnosed between 1947 and 2018. In the European cohort, we excluded 2,235 women who were diagnosed with their first breast cancer before 1990 or who had missing year of first diagnosis. We restricted to women diagnosed from 1990 onward so that diagnostic procedures and treatment would be more representative of current practice. Moreover, clinico-pathological, treatment, and follow-up data were more complete after 1990. In addition, we excluded 16 studies (9,783 women) without information about metachronous CBC events (Figure S1A). After these exclusions, the cohort for this analysis comprised data from 42 studies, including 56,068 women with invasive breast cancer among whom 1,027 metachronous CBC occurred (Table S2).

All individuals provided written informed consent, and all studies were approved by the relevant institutional review boards. BCAC data were centrally harmonized and cleaned in communication with the study data managers and principal investigators. Data collection for individual studies is described in Table S1.

Genotyping and PRSs

DNA samples from participants were genotyped using the iCOGS array^{12,13} or the OncoArray,^{4,14} with genotypes for variants not on the arrays estimated by imputation.^{4,13} The PRS₃₁₃ was calculated as a weighted sum of the minor allele dosages; the variant selection and weights are as given by Mavaddat et al.⁷ We also calculated estimates for a previously published PRS₇₇⁶ and for estrogen receptor (ER)-specific PRSs (ER-positive PRS₃₁₃ and ER-negative PRS₃₁₃).⁷ The ER-specific PRSs were constructed by defining subtype-specific weights for the 313 variants using a hybrid approach.⁷ Variants and corresponding coefficients used to construct the PRS are shown in Table S3. We standardized the PRS in our analyses by dividing it by the SD of the PRS of the control subjects (PRS₇₇ SD = 0.45; PRS₃₁₃ SD = 0.61; ER-positive PRS₃₁₃ SD = 0.65; ER-negative PRS₃₁₃ SD = 0.59) exactly as was done in the analyses of the PRS and first breast cancer risk.^{6,7} This allows a direct comparison of the magnitude of the CBC relative risk estimation to that of the first breast cancer.

For samples genotyped with both OncoArray and iCOGS array (9,071 samples), OncoArray data were used in preference as the imputation quality was generally higher. The intraclass correlation coefficient (ICC) between the PRSs derived from the two platforms was 0.99 (95%CI = 0.99–0.99) for the PRS₇₇ and 0.96 (95%CI = 0.95–0.96) for PRS₃₁₃ (Figure S2). Given the high correlation

between the two platforms, PRS measures from both platforms were used in the analyses without adjustment.

Statistical Analysis

European Cohort

The primary outcome in the European cohort was the development of metachronous CBC. Cox proportional hazards models were used to estimate HRs for metachronous CBC risk by PRS, stratified by country. Since previous studies have shown that age at first breast cancer diagnosis is an important predictor of CBC,³ the analyses were performed with attained age as the timescale. Time at risk started 3 months after the first breast cancer diagnosis and ended at the age of CBC diagnosis, distant metastasis (where available), death, or end of follow-up, whichever came first. For women who had a study entry more than 3 months after first breast cancer diagnosis, follow-up started at the age of study entry. We also performed a fixed-effect meta-analysis of country-specific effects using the STATA command *metan*. We performed a fixed-effect meta-analysis over a random-effect meta-analysis since there was no evidence for heterogeneity in effect sizes between countries (I-squared = 0%, Figure S3). For some analyses, only invasive CBC was used as the outcome; in these analyses we censored on *in situ* CBC. Separate analyses were conducted for ER-positive CBC (censored on ER-negative and ER-unknown CBC) and ER-negative CBC (censored on ER-positive and ER-unknown CBC).

We evaluated the linearity of the association between PRS₃₁₃ per unit SD and CBC risk using restricted cubic splines with three knots. There was no evidence for violation of the linearity assumption. Therefore, in the main analysis, the PRS₃₁₃ was treated as a continuous covariate, and estimated the HR per unit SD of the PRS₃₁₃. Violation of the proportional hazard assumption was assessed by inspection of the Schoenfeld residuals.¹⁵ As a second analysis, we used the per SD log HR of the PRS₃₁₃ to calculate the predicted HR at different percentiles of the PRS₃₁₃, compared to the 50th percentile. Third, the PRS₃₁₃ was categorized into percentile groups (0th to 10th, 10th to 20th, 20th to 40th, 40th to 60th, 60th to 80th, 80th to 90th, 90th to 100th) to illustrate the differences between PRS₃₁₃ subgroups, with the middle quintile (40th to 60th) as the reference.

We also performed multivariable Cox regression analyses to determine whether the log HR of CBC risk by PRS changed when adjusting for year of first breast cancer diagnosis, family history of breast cancer in a first degree relative, and several clinical characteristics of the first breast cancer such as nodal status, tumor size, morphology, ER status, progesterone receptor (PR) status, human epidermal growth factor receptor 2 (HER2) status, (neo) adjuvant chemotherapy, adjuvant endocrine therapy, and radiotherapy. These analyses were performed in all women, a complete case set (excluding those with unknown values for the covariates), and in a set excluding studies oversampling case subjects with family history. Potential effect modification of the PRS₃₁₃ effect by the same variables was evaluated by fitting interaction terms in different models using complete case sets, including the standardized PRS₃₁₃, modifier, and interaction.

The discriminative ability of different models (model 1: PRS₃₁₃ alone; model 2: other risk factors [the adjustment variables from the multivariable Cox regression analyses]; model 3: PRS₃₁₃ + other risk factors) was calculated using Harrell's C-index.¹⁶ Since no standard performance measures are currently available to account for left-truncated follow-up time (i.e., to start analyses at

age at study entry), we used time since first breast cancer as the timescale to calculate the C-index.

Absolute Risks

Absolute risks of developing CBC at PRS₃₁₃ percentiles were calculated using the estimated log HRs per SD from the breast cancer cohort (BCAC) under the log-linear model, assuming the PRS is normally distributed. The PRS₃₁₃ and age-specific incidences were constrained to the age-specific CBC incidences from women diagnosed with a first invasive breast cancer in the period 2003–2010 from the Netherlands Cancer Registry (NCR).¹ The procedure for constraining the incidences has been previously described.¹⁷ The age-specific CBC incidences were calculated overall and for age-specific groups, censoring on death and distant metastasis. We used data from the NCR since this registry has complete coverage of all newly diagnosed cancers in the Netherlands. The NCR cohort included all females aged ≥ 18 years and follow-up for second cancers was complete until February 1, 2016.¹ We then applied the competing risk of dying on the absolute CBC risks. The absolute CBC risk (AR_g) by age t in PRS₃₁₃ category g , taking into account the competing risk of dying was calculated by:

$$AR_g(t) = \sum_{u=0}^{t-1} \mu_g(u) S_g(u) S_m(u)$$

where $\mu_g(t)$ is the CBC incidence associated with PRS₃₁₃ category g , $S_g(t)$ the probability of being free of CBC to age t , and $S_m(t)$ the probability of surviving to age t .

Case-Case Series

For the case-case series (European and Asian), logistic regression models were used to estimate the ORs for CBC risk (comparing with unilateral breast cancer) and for unilateral breast cancer risk (comparing with women without any diagnosis of breast cancer) associated with PRS₃₁₃. All analyses were adjusted for age and country (Table S1). For all unilateral- and contralateral breast cancer patients, we used age at first breast cancer diagnosis, and for women without any diagnosis of breast cancer we used age at baseline questionnaire.

For direct comparison with the estimate reported for PRS₃₁₃ and first breast cancer, we also performed logistic regression analyses in the same BCAC study participants included in the validation of the association between PRS₃₁₃ and first breast cancer.⁷ This validation set comprised a subsample from 24 studies and included 3,781 women with unilateral breast cancer, 94 women with CBC, and 3,753 women without any diagnosis of breast cancer (Table S2). For this analysis, we adjusted for 10 principal components, in line with Mavaddat et al.⁷

For European women who had follow-up time available more than 3 months after the first breast cancer diagnosis, a sensitivity analysis was performed for metachronous CBC (1,702 CBCs). We also did a separate analysis for invasive CBC (N = 3,246), by excluding CBC *in situ*.

All p values are two sided; tests with $p < .05$ are referred to as statistically significant. Analyses were performed using STATA, v.13.1 (StataCorp) and R v.3.3.2.

Results

European (Cohort) Cox Regression Analyses

The European cohort included 56,068 women diagnosed with first invasive breast cancer with 1,027 metachronous CBC events. Median follow-up was 8.4 years. Patient, tumor, and treatment characteristics are summarized in Table S4.

Table 1. Association between PRSs and Contralateral Breast Cancer Risk in the European Cohort (N = 56,068)

Polygenic Risk Score (PRS)	No. of CBC	HR per Unit SD ^a	95%CI	p Value
PRS₇₇^b				
All CBC	1,027	1.21	1.14–1.29	<.001
Invasive CBC	923	1.21	1.13–1.29	<.001
PRS₃₁₃^b				
All CBC	1,027	1.25	1.18–1.33	<.001
Invasive CBC	923	1.24	1.16–1.32	<.001
ER-positive invasive CBC ^d	275	1.38	1.23–1.55	<.001
ER-negative invasive CBC ^d	97	0.92	0.75–1.12	.39
ER-Positive PRS₃₁₃^{b,c}				
All CBC	1,027	1.23	1.16–1.31	<.001
Invasive CBC	923	1.22	1.15–1.30	<.001
ER-positive invasive CBC ^d	275	1.37	1.22–1.54	<.001
ER-Negative PRS₃₁₃^{b,c}				
All CBC	1,027	1.25	1.17–1.33	<.001
Invasive CBC	923	1.24	1.16–1.33	<.001
ER-negative invasive CBC ^d	97	1.06	0.86–1.30	.58

Abbreviations: PRS, polygenic risk score; No., number; CBC, contralateral breast cancer; HR, hazard ratio; CI, confidence interval; ER, estrogen receptor; SD, standard deviation.

^aAll analyses were performed with attained age as timescale.


^bCoefficients to construct the PRSs are shown in Table S3. All PRSs were standardized by the same SD as was used by Mavaddat et al.⁷ The SD was 0.45 for overall breast cancer PRS₇₇, 0.61 for overall breast cancer PRS₃₁₃, 0.65 for ER-positive PRS₃₁₃, and 0.59 for ER-negative PRS₃₁₃.

^cER-specific PRSs were constructed using a hybrid method, as described by Mavaddat et al.⁷

^dWomen with ER-unknown CBC (N = 551) were censored in these analyses.

The associations between the different PRSs and CBC risk are shown in Table 1. The HR for CBC per SD of PRS₃₁₃ was 1.25 (95%CI = 1.18–1.33). For comparison, the HR per SD for PRS₇₇ was 1.21 (95%CI = 1.14–1.29). Women within the 0th to 10th and the 90th to 100th percentile of the PRS₃₁₃ had 0.59-fold (95%CI = 0.45–0.78) and 1.38-fold (95%CI = 1.13–1.69) risk of CBC, respectively, compared with women within the 40th to 60th percentile (Figure 1, Table S5). The predicted HRs of CBC for women at the 10th and 90th percentile of the PRS₃₁₃ were 0.75 and 1.33, respectively, compared to the 50th percentile (Figure 1). Since we observed evidence of departure from the proportional hazards assumption ($p = 0.02$),¹⁵ we also calculated HRs stratified for follow-up duration (<5 and ≥5 years). The HR by SD of the PRS₃₁₃ was 1.21 (95%CI = 1.10–1.32) for CBC diagnosed ≤5 years after first breast cancer diagnosis (CBC N = 428) and 1.28 (95%CI = 1.18–1.38) for CBC diagnosed >5 years after first diagnosis (CBC N = 599).

The HR per SD of PRS₃₁₃ for ER-positive invasive CBC was 1.38 (95%CI = 1.23–1.55) compared to a HR per SD

**Figure 1. Estimates for Contralateral Breast Cancer Risk by Percentile Categories of the 313-Variant PRS (PRS₃₁₃)**

The figure shows the hazard ratios per SD and 95% confidence intervals for percentiles of the PRS₃₁₃ relative to the middle quintile (underlying table can be found in Table S5). The solid line denotes the estimates for contralateral breast cancer risk with the PRS₃₁₃ fitted as a continuous covariate. Coefficients to construct the PRS₃₁₃ are shown in Table S3. The PRS₃₁₃ was standardized by SD = 0.61, in line with Mavaddat et al.⁷ The analyses were performed with attained age as timescale. PRS, polygenic risk score; SD, standard deviation.

of the ER-positive PRS₃₁₃ of 1.37 (95%CI = 1.22–1.54) (Table 1). For ER-negative invasive CBC, the HR per SD was 0.92 (95%CI = 0.75–1.12) for PRS₃₁₃ and 1.06 (95%CI = 0.86–1.30) for the ER-negative PRS₃₁₃.

Sensitivity analysis using the overall PRS₃₁₃ showed a HR per SD of 1.24 (95%CI = 1.16–1.32) for invasive CBC risk. When we used time since first breast cancer as the timescale, we found similar results (HR per SD = 1.25, 95%CI = 1.18–1.33). Meta-analysis of country-specific effects showed a HR per SD of 1.25 (95%CI = 1.18–1.33) for CBC risk by PRS₃₁₃ (Figure S3).

The association between the PRS₃₁₃ and CBC risk did not change when adjusting for characteristics of the individual, tumor, or treatment, nor when excluding studies oversampling case subjects with a family history (Table S6). When considering potential modifiers of the effect of the PRS₃₁₃ on CBC risk (Table 2), we found that the HR was the lowest in women aged <40 years at first breast cancer diagnosis (HR per SD = 1.13; 95%CI = 0.98–1.31) and tended to increase with age, although these effects were not statistically significant ($P_{\text{heterogeneity}} = 0.26$; $P_{\text{trend}} = 0.05$). We found no indication for effect modification by family history ($P_{\text{heterogeneity}} = 0.63$), morphology ($P_{\text{heterogeneity}} = 0.14$), ER status ($P_{\text{heterogeneity}} = 0.13$), PR status ($p = 0.26$), HER2 status ($P_{\text{heterogeneity}} = 0.42$), chemotherapy ($P_{\text{heterogeneity}} = 0.60$), endocrine therapy ($P_{\text{heterogeneity}} = 0.79$), or radiotherapy ($P_{\text{heterogeneity}} = 0.40$) (Table 2).

The C-index was 0.563 (95%CI = 0.547–0.586) for the model only including PRS₃₁₃, 0.605 (95%CI = 0.591–0.629)

Table 2. Association between the 313-Variant PRS (PRS₃₁₃) and Contralateral Breast Cancer Risk for Subgroups

Subgroups	No. of Patients	No. of CBC	HR per Unit SD ^{a,b}	95%CI	p Value	P _{heterogeneity} ^{c,d}	P _{trend} ^{c,e}
All patients	56,068	1,027	1.25	1.18–1.33	<.001	-	-
Age at First Breast Cancer Diagnosis (Years)						.26	.05
<40	5,877	171	1.13	0.98–1.31	.09		
40–49	11,928	265	1.25	1.11–1.41	<.001		
50–59	16,882	320	1.22	1.09–1.36	<.001		
60+	21,381	271	1.36	1.21–1.52	<.001		
Family History (First Degree Relative)						.63	-
No	33,623	618	1.26	1.16–1.36	<.001		
Yes	10,369	302	1.22	1.09–1.36	<.001		
Morphology						.14	-
Ductal	37,324	621	1.21	1.12–1.31	<.001		
Lobular	5,878	118	1.32	1.10–1.59	.002		
Mixed (ductal and lobular)	2,174	46	1.52	1.15–2.02	.004		
Other	3,344	70	1.20	0.96–1.50	.11		
ER Status						.13	-
Negative	9,527	194	1.13	0.98–1.30	.08		
Positive	38,090	670	1.28	1.19–1.38	<.001		
PR Status						.26	-
Negative	13,098	244	1.16	1.03–1.32	.02		
Positive	27,044	554	1.27	1.17–1.38	<.001		
HER2 Status						.42	-
Negative	23,787	352	1.29	1.17–1.44	<.001		
Positive	4,969	60	1.45	1.13–1.85	.004		
(Neo)adjuvant Chemotherapy						.60	-
No	18,110	361	1.28	1.16–1.42	<.001		
Yes	18,559	363	1.24	1.12–1.37	<.001		
(Neo)adjuvant Endocrine Therapy						.79	-
No	10,781	242	1.28	1.13–1.44	<.001		
Yes	27,322	460	1.30	1.19–1.43	<.001		
Radiotherapy						.40	-
No	11,023	188	1.33	1.15–1.53	<.001		
Yes	29,142	617	1.24	1.15–1.34	<.001		

Abbreviations: PRS, polygenic risk score; No., number; CBC, contralateral breast cancer; HR, hazard ratio; CI, confidence interval; ER, estrogen receptor; PR, progesterone receptor; HER2, human epidermal growth factor receptor 2.

^aHR for CBC risk by unit SD of PRS₃₁₃. All analyses were performed with attained age as timescale.

^bCoefficients to construct the PRS₃₁₃ are shown in Table S3. The PRS₃₁₃ was standardized by standard deviation = 0.61, in line with Mavaddat et al.⁷

^cThe interaction between the PRS₃₁₃ and each subgroup was tested in different models including the standardized PRS₃₁₃, modifier, and interaction. Patients with unknown values were excluded from these analyses. Since attained age was used as timescale in all models, the model with age at first breast cancer only included the PRS₃₁₃ and interaction.

^dP for interaction based on test for heterogeneity across categories.

^eP for interaction based on a trend test with age as continuous variable.

for the model only including other risk factors, and 0.623 (95%CI = 0.608–0.645) for the complete model (Table 3).

Absolute Risks

Based on the HR estimates for PRS₃₁₃, the predicted CBC risk by age 80 years was 12.4% at the 10th percentile of

Table 3. Discriminatory Ability (C-Index) of the 313-Variant PRS (PRS₃₁₃) and Other Risk Factors for Contralateral Breast Cancer Risk in the European Cohort

	C-Index (95%CI) ^{a,b}
Model 1: PRS ₃₁₃ ^c alone	0.563 (0.547–0.586)
Model 2: Other risk factors ^d	0.605 (0.591–0.629)
Model 3: PRS ₃₁₃ ^c + other risk factors ^d	0.623 (0.608–0.645)

Abbreviations: PRS, polygenic risk score; CI, confidence interval.

^aThe Harrell's C-index was obtained by the STATA `stcox` postestimation command "estat concordance," using time since first breast cancer on the time-scale without taking delayed entry (prevalent cases) into account. We did not consider delayed entry since no standard performance measures are currently available in the statistical literature to account for left-truncated follow-up time. The median of delayed entry was 0.4 years (standard deviation = 2.7) in our study.

^bThe 95% CIs were obtained by use of the `somersd` package in STATA.

^cCoefficients to construct the PRS₃₁₃ are shown in Table S3. The PRS₃₁₃ was standardized by SD = 0.61, in line with Mavaddat et al.⁷

^dIncluding age at first diagnosis, year of first diagnosis, family history for breast cancer in a first degree relative, and clinical characteristics of the first breast cancer (nodal status, tumor size, differentiation grade, morphology, estrogen receptor status, human epidermal growth factor receptor 2 status, chemotherapy, endocrine therapy, radiotherapy).

the PRS₃₁₃, compared with 20.5% at the 90th percentile of the PRS₃₁₃ (Figure 2), accounting for death as competing risk. When death was not taken into account as competing risk, the corresponding predicted risks by age 80 were 17.0% at the 10th percentile and 27.9% at the 90th percentile of the PRS₃₁₃ (Figure S4). Table 4 shows the 5- and 10-year cumulative CBC risks by PRS₃₁₃ for different age groups, accounting for death as competing risk (Table S7 shows results without competing risks).

European and Asian (Case-Case Series) Logistic Regression Analyses

Figure 3 shows the distribution of the PRS₃₁₃ per SD in the European case-case series. Median PRS₃₁₃ was -0.4 (interquartile range [IQR] = 1.35) for control women without any diagnosis of breast cancer (N = 81,000), 0.2 (IQR = 1.36) for women with unilateral breast cancer (N = 62,830), and 0.5 (IQR = 1.40) for women with CBC (N = 3,607). The OR for unilateral breast cancer per SD of the PRS₃₁₃, compared to control women, was 1.82 (95%CI = 1.80–1.84) (Table S8). The OR for CBC per SD of PRS₃₁₃, compared to unilateral breast cancer, was 1.30 (95%CI = 1.26–1.35).

In sensitivity analyses, the OR per SD of PRS₃₁₃ was 1.27 (95%CI = 1.21–1.33) for metachronous CBC and the OR per SD was 1.29 (95%CI = 1.24–1.33) for invasive CBC, compared to unilateral breast cancer. When analyses were restricted to the validation set of Mavaddat et al.,⁷ the OR for unilateral breast cancer per SD of the PRS₃₁₃ was 1.67 (95%CI = 1.59–1.76) compared to control women, and the OR for CBC per SD of PRS₃₁₃ was 1.39 (95%CI = 1.13–1.70) compared to unilateral breast cancer (Table S8).

For women of Asian descent, the OR for unilateral breast cancer per SD of the PRS₃₁₃ was 1.56 (95%CI = 1.52–1.60)




Figure 2. Predicted Contralateral Breast Cancer Risk by Percentile of the 313-Variant PRS (PRS₃₁₃) with Death as Competing Risk Coefficients to construct the PRS₃₁₃ are shown in Table S3. The PRS₃₁₃ was standardized by SD = 0.61, in line with Mavaddat et al.⁷ The CBC incidences were calculated based on incidence data from the Netherlands Cancer Registry¹ and relative risks estimated as described in the Material and Methods. PRS, polygenic risk score; CBC, contralateral breast cancer.

compared to control women, and the OR for CBC per SD of PRS₃₁₃ was 1.15 (95%CI = 1.02–1.29) compared to women with unilateral breast cancer (Table S8).

Discussion

Previous studies have shown that a PRS, summarizing the effects of common germline variants, can be used to stratify women with respect to their risk to develop a primary breast cancer.^{6–9} In this study, we observed a clear association between the PRS₃₁₃ and CBC risk in women of both European and Asian ancestry. The association was observed in both the case-case series and the European cohort. The HRs per SD of CBC for women at the 10th and 90th percentile of the continuous predicted PRS₃₁₃ were 0.75 and 1.33, respectively, compared to the 50th percentile. This translates to absolute risks at the 10th and the 90th percentile of the PRS₃₁₃ of 12.4% and 20.5%, respectively, by age 80 years. We estimated a C-index for the PRS₃₁₃, summarizing its discriminatory ability, of 0.563 in the European cohort.

One previous study has investigated the effect of a PRS, including 67 variants, and CBC risk.¹⁰ This study found a risk ratio of 1.75 (95%CI = 1.41–2.18) for women in the upper quartile of the PRS compared with women in the lowest quartile. To facilitate comparison, we performed a similar analysis in our case-case series, showing an OR of 1.98 (95%CI = 1.79–2.18), adjusted for country and age at first diagnosis, for women in the upper quartile of the PRS₃₁₃. This indicates that the PRS₃₁₃ improves stratification relative to PRSs including fewer variants. Moreover, in our European cohort, the C-index for the PRS alone

Table 4. 5- and 10-Year Cumulative Risks of Contralateral Breast Cancer by the 313-Variant PRS (PRS₃₁₃) for Different Age Groups with Death as Competing Risk

Age at First Breast Cancer Diagnosis (years)	5-Year Cumulative CBC Risks (%) Range by Age					10-Year Cumulative CBC risks (%) Range by Age				
	5 th Percentile PRS ₃₁₃	10 th Percentile PRS ₃₁₃	50 th Percentile PRS ₃₁₃	90 th Percentile PRS ₃₁₃	95 th Percentile PRS ₃₁₃	5 th Percentile PRS ₃₁₃	10 th Percentile PRS ₃₁₃	50 th Percentile PRS ₃₁₃	90 th Percentile PRS ₃₁₃	95 th Percentile PRS ₃₁₃
30–34	1.9–3.1	2.1–3.4	2.7–4.5	3.6–5.9	4.0–6.5	3.1–4.1	3.4–4.5	4.5–5.9	5.9–7.7	6.5–8.5
35–39	0.8–2.1	0.9–2.3	1.2–3.0	1.5–3.9	1.7–4.3	2.1–3.5	2.3–3.8	3.0–5.0	3.9–6.6	4.3–7.2
40–44	1.5–2.8	1.7–3.1	2.2–4.1	2.9–5.3	3.2–5.9	2.8–4.6	3.1–5.0	4.1–6.6	5.3–8.6	5.9–9.4
45–49	1.4–2.5	1.5–2.7	2.0–3.6	2.6–4.7	2.9–5.2	2.5–3.9	2.7–4.3	3.6–5.6	4.7–7.4	5.2–8.1
50–54	1.4–2.8	1.5–3.0	1.9–4.0	2.6–5.2	2.8–5.8	2.8–4.5	3.0–4.9	4.0–6.4	5.2–8.4	5.8–9.3
55–59	1.6–3.1	1.8–3.4	2.3–4.5	3.1–5.9	3.4–6.5	3.1–4.8	3.4–5.2	4.5–6.9	5.9–9.0	6.5–9.9
60–64	1.7–3.3	1.9–3.6	2.5–4.7	3.3–6.2	3.6–6.8	3.3–5.0	3.6–5.4	4.7–7.1	6.2–9.3	6.8–10.2
65–70	1.5–3.2	1.6–3.5	2.1–4.6	2.8–6.1	3.1–6.7	3.2–4.1	3.5–4.5	4.6–5.9	6.1–7.7	6.7–8.5

Abbreviations: PRS, polygenic risk score; CBC, contralateral breast cancer. Coefficients to construct the PRS₃₁₃ are shown in Table S3. The PRS₃₁₃ was standardized by SD = 0.61, in line with Mavaddat et al.⁷ The CBC incidences for each age group were calculated based on incidence data from the Netherlands Cancer Registry¹ and relative risks estimated as described in the Material and Methods. Death was taken into account as competing risk.

improved from 0.547 (95%CI = 0.536–0.575) for the previously reported PRS₇₇⁶ to 0.563 (95%CI = 0.547–0.586) for the PRS₃₁₃.

We found no evidence that the association between the PRS₃₁₃ and CBC risk was confounded by family history, adjuvant therapy, morphology, age, or tumor receptor status of the first breast cancer, nor that there was effect modification by those factors. The absence of notable effect modification is in line with the abovementioned study of a 67-variant PRS and CBC risk; no heterogeneity in association was found by age, family history, morphology, ER status, and adjuvant treatment.¹⁰

To provide an external validation of our findings, we examined data from UK Biobank, which includes many

women diagnosed with breast cancer with data available on the PRS₃₁₃ (Supplemental Note). Unfortunately, UK Biobank has no information available on the laterality of the tumor, and it is, therefore, not possible to distinguish between contralateral and ipsilateral breast cancers. We therefore performed analyses using any second breast cancer as the endpoint. This secondary analysis did confirm the association between the PRS₃₁₃ and second breast cancer risk (HR per SD = 1.13, 95%CI = 1.01–1.27), but with a lower estimate than in our European cohort. The lower estimate may be explained by the inclusion of the ipsilateral breast cancers, which may be more likely to be recurrences than new primary breast cancers compared to CBCs. Indeed, when we used ipsilateral breast cancer as the outcome in our European cohort, we found no association with the PRS₃₁₃ (HR = 1.02, 95%CI = 0.90–1.15).

The association between the PRS₃₁₃ and CBC risk (OR per SD = 1.30; 95%CI = 1.26–1.35) in the BCAC database was weaker (expressed in terms of an OR) than was found for first breast cancer among independent prospective studies (OR per SD = 1.61; 95%CI = 1.57–1.65). Under a simple polygenic model, the relative risk would be expected to be similar for the second breast cancer. The attenuated estimate for CBC might however be explained by several factors. Some attenuation of the estimate might have been due to dilution in the end-point definition, i.e., if some of the CBCs were metastases. Previous studies investigating the clonal relatedness of first breast cancers and CBCs using tumor sequencing have shown that 6%–12% of CBCs represent metastases.^{18,19} This hypothesis would be consistent with our finding of a slightly stronger association between the PRS₃₁₃ and late CBCs, diagnosed >5 years after the first breast cancer, than for early CBCs, diagnosed ≥5 years after the first cancer, since the latter are more likely to be metastases. In addition, 3%–5% of




Figure 3. Distribution of the 313-Variant PRS (PRS₃₁₃) in 62,830 Control Women without Any Diagnosis of Breast Cancer, 81,000 Women with Unilateral Breast Cancer, and 3,607 Women with Contralateral Breast Cancer

Coefficients to construct the PRS₃₁₃ are shown in Table S3. The PRS₃₁₃ was standardized by SD = 0.61, in line with Mavaddat et al.⁷ PRS, polygenic risk score; BC, breast cancer; CBC, contralateral breast cancer; SD, standard deviation.

the women with breast cancer will have a mutation in *BRCA1* or *BRCA2*,^{20,21} who have high CBC risks. It has been shown that the relative risk associated with PRS is lower (for the first breast cancer) for women with a *BRCA1* and *BRCA2* mutation than in the general population,²² diluting the overall relative risk for CBC. More generally, it is possible that the CBC association may be attenuated due to the effect of other, unmeasured, genetic or other risk factors. If the risks are high, case subjects with higher PRS₃₁₃ will have, on average, lower values of other risk factors, due to elimination of the highest risk individuals, again attenuating the CBC association. Finally, given the limited information on family history in our dataset, the estimate could have been biased due to a family history effect not detected in our data.

There was some suggestion that the relative risk associated with PRS₃₁₃ decreased with younger age ($P_{\text{trend}} = 0.05$) and, specifically, was lower for women aged <40 years (HR per SD = 1.13; 95%CI = 0.98–1.31). Interestingly, Mavaddat et al.⁷ also found a lower relative risk below age 40 for first breast cancer. This effect may reflect the different characteristics of breast cancers at young ages, both in terms of germline susceptibility and pathology.^{23,24} For example, the proportion of ER-negative breast cancers is higher at young ages, and the PRS is less predictive for ER-negative disease.^{6,7,24}

In the logistic regression analyses in Asian women, the association between the PRS₃₁₃ and CBC risk was slightly weaker than in European women. This finding is consistent with a recent analysis investigating the association between a 287-variant PRS and first breast cancer risk in the Asian population,²⁵ which showed an attenuated OR in Asian women (OR = 1.52, 95%CI = 1.49–1.56) compared to European women (OR = 1.61, 95%CI = 1.57–1.66). The lower estimate for Asian women might reflect the fact the PRS₃₁₃ was developed in European populations, and the different LD structure in Asians may attenuate the association since the variants in the PRS are likely to be surrogates for the causal variants. Other explanations for the attenuated estimate may be the slightly younger age at first breast cancer diagnosis and the higher proportion ER-negative CBCs in Asian women compared to European women in our study. Finally, the imputation quality for variants was somewhat lower, on average, for the Asian than for the European dataset, with three variants on OncoArray and four variants on ICOGs with an imputation quality score < 0.3 (Table S3). Nevertheless, we included those variants in the PRS for both European and Asian women, to keep the PRS comparable between ethnicities and studies. Future studies including larger numbers of Asian women, and women of other ethnicities, are needed to generate population-specific PRSs and to validate our findings in these groups.

A major strength of this study is the very large sample size in the BCAC dataset, including genotype information for ~150,000 women and a large number of CBC

events. A limitation of this study is missing data on the patient, tumor, and treatment characteristics, which reduces the power of the multivariable Cox regression analyses and interaction analyses. In addition, registration of CBC was not complete; the 10-year cumulative CBC incidence was 2.2% in the BCAC dataset, compared to 3.8% using complete data from the Netherlands Cancer Registry.¹ For this reason, we estimated relative risk estimates using the BCAC data and applied these to external registry data to obtain absolute risk estimates. The under-reporting of CBC should not bias our HR estimates, given that the event rate is low and reporting of CBC is unlikely to be related to the PRS₃₁₃. Moreover, we reran the cohort analysis in the subset of countries with a 10-year cumulative CBC incidence $\geq 3.0\%$ in the BCAC dataset, and the estimates were very similar to the main analyses (HR per SD = 1.23, 95%CI = 1.14–1.33) (Figure S3).

In conclusion, the PRS₃₁₃ is predictive for the development of CBC. We found no evidence for confounding or effect modification by other previously established CBC risk factors. The PRS₃₁₃ is therefore likely to be an independent risk factor for CBC. Since the predictive ability of the PRS on its own is modest, it should be combined with other breast cancer risk factors to provide more useful CBC risk prediction models. More accurate risk prediction will help identify women at high CBC risk who will benefit from additional surveillance and/or risk reducing mastectomy, and equally important, to identify those women at low risk in order to avoid unnecessary surgeries.

Data and Code Availability

Data used in this manuscript may be requested through the original providers. Data of the Breast Cancer Association Consortium may be requested for non-profit research through an application procedure with the Breast Cancer Association Consortium. Data of the UK Biobank needs to be requested through UK Biobank.

Supplemental Data

Supplemental Data can be found online at <https://doi.org/10.1016/j.ajhg.2020.09.001>.

Declaration of Interests

M.W.B. conducts research funded by Amgen, Novartis, and Pfizer, outside the submitted work. P.A.F. conducts research funded by Amgen, Novartis, and Pfizer, outside the submitted work. He received honoraria from Roche, Novartis, and Pfizer. H.N. received honorarium from Astra Zeneca outside the submitted work.

Received: May 15, 2020

Accepted: September 2, 2020

Published: October 5, 2020

Web Resources

Breast Cancer Association Consortium, <http://bcac.ccge.medschl.cam.ac.uk/bcacdata/>
UK Biobank, <https://www.ukbiobank.ac.uk/researchers/>

References

1. Kramer, I., Schaapveld, M., Oldenburg, H.S.A., Sonke, G.S., McCool, D., van Leeuwen, F.E., Van de Vijver, K.K., Russell, N.S., Linn, S.C., Siesling, S., et al. (2019). The influence of adjuvant systemic regimens on contralateral breast cancer risk and receptor subtype. *J. Natl. Cancer Inst.* *111*, 709–718.
2. Xiong, Z., Yang, L., Deng, G., Huang, X., Li, X., Xie, X., Wang, J., Shuang, Z., and Wang, X. (2018). Patterns of Occurrence and Outcomes of Contralateral Breast Cancer: Analysis of SEER Data. *J. Clin. Med.* *7*, 7.
3. Akdeniz, D., Schmidt, M.K., Seynaeve, C.M., McCool, D., Giardiello, D., van den Broek, A.J., Hauptmann, M., Steyerberg, E.W., and Hooning, M.J. (2019). Risk factors for metachronous contralateral breast cancer: A systematic review and meta-analysis. *Breast* *44*, 1–14.
4. Michailidou, K., Lindström, S., Dennis, J., Beesley, J., Hui, S., Kar, S., Lemaçon, A., Soucy, P., Glubb, D., Rostamianfar, A., et al.; NBCS Collaborators; ABCTB Investigators; and ConFab/AOCS Investigators (2017). Association analysis identifies 65 new breast cancer risk loci. *Nature* *551*, 92–94.
5. Milne, R.L., Kuchenbaecker, K.B., Michailidou, K., Beesley, J., Kar, S., Lindström, S., Hui, S., Lemaçon, A., Soucy, P., Dennis, J., et al.; ABCTB Investigators; EMBRACE; GEMO Study Collaborators; HEBON; kConFab/AOCS Investigators; and NBCS Collaborators (2017). Identification of ten variants associated with risk of estrogen-receptor-negative breast cancer. *Nat. Genet.* *49*, 1767–1778.
6. Mavaddat, N., Pharoah, P.D., Michailidou, K., Tyrer, J., Brook, M.N., Bolla, M.K., Wang, Q., Dennis, J., Dunning, A.M., Shah, M., et al. (2015). Prediction of breast cancer risk based on profiling with common genetic variants. *J. Natl. Cancer Inst.* *107*, 107.
7. Mavaddat, N., Michailidou, K., Dennis, J., Lush, M., Fachal, L., Lee, A., Tyrer, J.P., Chen, T.H., Wang, Q., Bolla, M.K., et al.; ABCTB Investigators; kConFab/AOCS Investigators; and NBCS Collaborators (2019). Polygenic Risk Scores for Prediction of Breast Cancer and Breast Cancer Subtypes. *Am. J. Hum. Genet.* *104*, 21–34.
8. Brentnall, A.R., van Veen, E.M., Harkness, E.F., Rafiq, S., Byers, H., Astley, S.M., Sampson, S., Howell, A., Newman, W.G., Cuzick, J., and Evans, D.G.R. (2020). A case-control evaluation of 143 single nucleotide polymorphisms for breast cancer risk stratification with classical factors and mammographic density. *Int. J. Cancer* *146*, 2122–2129.
9. Shieh, Y., Hu, D., Ma, L., Huntsman, S., Gard, C.C., Leung, J.W., Tice, J.A., Vachon, C.M., Cummings, S.R., Kerlikowske, K., and Ziv, E. (2016). Breast cancer risk prediction using a clinical risk model and polygenic risk score. *Breast Cancer Res. Treat.* *159*, 513–525.
10. Robson, M.E., Reiner, A.S., Brooks, J.D., Concannon, P.J., John, E.M., Mellemejaer, L., Bernstein, L., Malone, K.E., Knight, J.A., Lynch, C.F., et al. (2017). Association of Common Genetic Variants With Contralateral Breast Cancer Risk in the WECARE Study. *J. Natl. Cancer Inst.* *J. Natl. Cancer Inst.* *109*, dxj051, dxj051.
11. Teraoka, S.N., Bernstein, J.L., Reiner, A.S., Haile, R.W., Bernstein, L., Lynch, C.F., Malone, K.E., Stovall, M., Capanu, M., Liang, X., et al.; WECARE Study Collaborative Group (2011). Single nucleotide polymorphisms associated with risk for contralateral breast cancer in the Women's Environment, Cancer, and Radiation Epidemiology (WECARE) Study. *Breast Cancer Res.* *13*, R114.
12. Michailidou, K., Beesley, J., Lindstrom, S., Canisius, S., Dennis, J., Lush, M.J., Maranian, M.J., Bolla, M.K., Wang, Q., Shah, M., et al.; BOCS; kConFab Investigators; AOCS Group; NBCS; and GENICA Network (2015). Genome-wide association analysis of more than 120,000 individuals identifies 15 new susceptibility loci for breast cancer. *Nat. Genet.* *47*, 373–380.
13. Michailidou, K., Hall, P., Gonzalez-Neira, A., Ghoussaini, M., Dennis, J., Milne, R.L., Schmidt, M.K., Chang-Claude, J., Bojesen, S.E., Bolla, M.K., et al.; Breast and Ovarian Cancer Susceptibility Collaboration; Hereditary Breast and Ovarian Cancer Research Group Netherlands (HEBON); kConFab Investigators; Australian Ovarian Cancer Study Group; and GENICA (Gene Environment Interaction and Breast Cancer in Germany) Network (2013). Large-scale genotyping identifies 41 new loci associated with breast cancer risk. *Nat. Genet.* *45*, 353–361, e1–e2.
14. Amos, C.I., Dennis, J., Wang, Z., Byun, J., Schumacher, F.R., Gayther, S.A., Casey, G., Hunter, D.J., Sellers, T.A., Gruber, S.B., et al. (2017). The OncoArray Consortium: A Network for Understanding the Genetic Architecture of Common Cancers. *Cancer Epidemiol. Biomarkers Prev.* *26*, 126–135.
15. Schoenfeld, D.A. (1983). Sample-size formula for the proportional-hazards regression model. *Biometrics* *39*, 499–503.
16. Harrell, F.E., Jr., Califf, R.M., Pryor, D.B., Lee, K.L., and Rosati, R.A. (1982). Evaluating the yield of medical tests. *JAMA* *247*, 2543–2546.
17. Antoniou, A.C., Beesley, J., McGuffog, L., Sinilnikova, O.M., Healey, S., Neuhausen, S.L., Ding, Y.C., Rebbeck, T.R., Weitzel, J.N., Lynch, H.T., et al.; Ontario Cancer Genetics Network; SWE-BRCA; HEBON; EMBRACE; GEMO; GEMO; Breast Cancer Family Registry; kConFab; and CIMBA (2010). Common breast cancer susceptibility alleles and the risk of breast cancer for BRCA1 and BRCA2 mutation carriers: implications for risk prediction. *Cancer Res.* *70*, 9742–9754.
18. Klevebring, D., Lindberg, J., Rockberg, J., Hilliges, C., Hall, P., Sandberg, M., and Czene, K. (2015). Exome sequencing of contralateral breast cancer identifies metastatic disease. *Breast Cancer Res. Treat.* *151*, 319–324.
19. Begg, C.B., Ostrovnaya, I., Geyer, F.C., Papanastasiou, A.D., Ng, C.K.Y., Sakr, R.A., Bernstein, J.L., Burke, K.A., King, T.A., Piscuoglio, S., et al. (2018). Contralateral breast cancers: Independent cancers or metastases? *Int. J. Cancer* *142*, 347–356.
20. Thompson, D., and Easton, D. (2004). The genetic epidemiology of breast cancer genes. *J. Mammary Gland Biol. Neoplasia* *9*, 221–236.
21. van den Broek, A.J., van 't Veer, L.J., Hooning, M.J., Cornelissen, S., Broeks, A., Rutgers, E.J., Smit, V.T., Cornelisse, C.J., van Beek, M., Janssen-Heijnen, M.L., et al. (2016). Impact of Age at Primary Breast Cancer on Contralateral Breast Cancer Risk in BRCA1/2 Mutation Carriers. *J. Clin. Oncol.* *34*, 409–418.
22. Kuchenbaecker, K.B., McGuffog, L., Barrowdale, D., Lee, A., Soucy, P., Dennis, J., Domchek, S.M., Robson, M., Spurdle, A.B., Ramus, S.J., et al. (2017). Evaluation of Polygenic Risk Scores for Breast and Ovarian Cancer Risk Prediction in

- BRCA1 and BRCA2 Mutation Carriers. *J. Natl. Cancer Inst.* *109*, 109.
23. Azim, H.A., Jr., Michiels, S., Bedard, P.L., Singhal, S.K., Crischiello, C., Ignatiadis, M., Haibe-Kains, B., Piccart, M.J., Sotiriou, C., and Loi, S. (2012). Elucidating prognosis and biology of breast cancer arising in young women using gene expression profiling. *Clin. Cancer Res.* *18*, 1341–1351.
24. Anders, C.K., Hsu, D.S., Broadwater, G., Acharya, C.R., Foekens, J.A., Zhang, Y., Wang, Y., Marcom, P.K., Marks, J.R., Febbo, P.G., et al. (2008). Young age at diagnosis correlates with worse prognosis and defines a subset of breast cancers with shared patterns of gene expression. *J. Clin. Oncol.* *26*, 3324–3330.
25. Ho, W.K., Tan, M.M., Mavaddat, N., Tai, M.C., Mariapun, S., Li, J., Ho, P.J., Dennis, J., Tyrer, J.P., Bolla, M.K., et al. (2020). European polygenic risk score for prediction of breast cancer shows similar performance in Asian women. *Nat. Commun.* *11*, 3833.