

UCSF

UC San Francisco Previously Published Works

Title

The Mediating Roles of Coping, Sleep, and Anxiety Motives in Cannabis Use and Problems Among Returning Veterans With PTSD and MDD

Permalink

<https://escholarship.org/uc/item/3j1936sw>

Journal

Psychology of Addictive Behaviors, 30(7)

ISSN

0893-164X

Authors

Metrik, Jane
Jackson, Kristina
Bassett, Shayna S
et al.

Publication Date

2016-11-01

DOI

10.1037/adb0000210

Peer reviewed

The Mediating Roles of Coping, Sleep, and Anxiety Motives in Cannabis Use and Problems Among Returning Veterans With PTSD and MDD

Jane Metrik

Providence VA Medical Center, Providence, Rhode Island, and
Brown University School of Public Health

Kristina Jackson

Brown University School of Public Health

Shayna S. Bassett

Providence VA Medical Center, Providence, Rhode Island, and
Alpert Medical School, Brown University

Michael J. Zvolensky

University of Houston and University of Texas MD Anderson
Cancer Center

Karen Seal and Brian Borsari

San Francisco VA Medical Center, San Francisco, California, and University of California, San Francisco

Veterans with posttraumatic stress disorder (PTSD) and major depressive disorder (MDD), the 2 most prevalent mental health disorders in the Iraq and Afghanistan veterans, are at increased risk for cannabis use and problems including cannabis use disorder (CUD). The present study examined the relationship of PTSD and MDD with cannabis use frequency, cannabis problems, and CUD as well as the role of 3 coping-oriented cannabis use motives (coping with negative affect, situational anxiety, and sleep) that might underlie this relationship. Participants were veterans ($N = 301$) deployed post-9/11/2001 recruited from a Veterans Health Administration facility in the Northeast United States based on self-reported lifetime cannabis use. There were strong unique associations between PTSD and MDD and cannabis use frequency, cannabis problems, and CUD. Mediation analyses revealed the 3 motives accounted, in part, for the relationship between PTSD and MDD with 3 outcomes in all cases but for PTSD with cannabis problems. When modeled concurrently, sleep motives, but not situational anxiety or coping with negative affect motives, significantly mediated the association between PTSD and MDD with use. Together with coping motives, sleep motives also fully mediated the effects of PTSD and MDD on CUD and in part the effect of MDD on cannabis problems. Findings indicate the important role of certain motives for better understanding the relation between PTSD and MDD with cannabis use and misuse. Future work is needed to explore the clinical utility in targeting specific cannabis use motives in the context of clinical care for mental health and CUD.

Keywords: cannabis, motives, PTSD, depression, sleep

Prevalence rates of cannabis use and cannabis use disorder (CUD) have more than doubled in the past decade in the general population of United States (U.S.) adults (Hasin et al., 2015) and among the U.S. military veterans (Bonn-Miller, Harris, & Trafton, 2012). A growing number of veterans are also diagnosed with posttraumatic stress disorder (PTSD), with PTSD prevalence rates of 23% among those returning from Operation Enduring Freedom,

Operation Iraqi Freedom, and Operation New Dawn (OEF/OIF/OND; Fulton et al., 2015). Although PTSD is the most common mental health diagnosis among OEF/OIF/OND veterans, more than 17% of returning veterans also meet criteria for a depressive disorder (Seal et al., 2009). The prevalence rates of these mental health disorders are disproportionately higher than the rates of PTSD and MDD in the general population (3.5% and 6.7%,

This article was published Online First October 27, 2016.

Jane Metrik, Providence VA Medical Center, Providence, Rhode Island, and Center for Alcohol and Addiction Studies, Brown University School of Public Health; Kristina Jackson, Center for Alcohol and Addiction Studies, Brown University School of Public Health; Shayna S. Bassett, Providence VA Medical Center, and Department of Psychiatry and Human Behavior, Alpert Medical School, Brown University; Michael J. Zvolensky, Department of Behavioral Science, University of Houston, and University of Texas MD Anderson Cancer Center; Karen Seal and Brian Borsari, San Francisco VA Medical Center, San Francisco, California, and Department of Psychiatry, University of California, San Francisco.

All authors contributed in a significant way to the manuscript and have all read and approved the final manuscript. This study was funded

by a National Institute on Drug Abuse grant (R01 DA033425), awarded to Jane Metrik and Brian Borsari. Kristina Jackson's work on this project was supported by K02 AA13938. The funding sources had no other role other than financial support. The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs. The authors have no financial relationship with the study sponsor. The authors gratefully acknowledge Cassandra Delapaix, Rebecca Swagger, Madeline Benz, Hannah Wheeler, Suzanne Sales, and Julie Evon for their contribution to the project.

Correspondence concerning this article should be addressed to Jane Metrik, Center for Alcohol and Addiction Studies, Brown University, Box G-S121-4, Providence, RI 02912. E-mail: Jane_Metrik@brown.edu

respectively; Kessler, Chiu, Demler, Merikangas, & Walters, 2005). Many service members with PTSD are also dually or multiply diagnosed (Cohen et al., 2010; Seal, Bertenthal, Miner, Sen, & Marmar, 2007), with major depressive disorder (MDD) being the second most current co-occurring psychiatric disorder (30% comorbidity with PTSD; Seal et al., 2008), followed by substance use disorders (SUDs; Golub, Vazan, Bennett, & Liberty, 2013; Seal et al., 2008, 2011). In fact, having either a PTSD or MDD diagnosis increased the odds of having any drug use disorder more than threefold in this veteran population (Seal et al., 2011).

Notably, population comorbidity estimates specific to cannabis use and CUD are not well-established among OEF/OIF/OND veterans because there is no routine screening or assessment for cannabis use or CUD in the Veterans Health Administration (VHA; Department of Veterans Affairs, 2009). Relative to many other comorbid psychiatric disorders (Agosti, Nunes, & Levin, 2002; Chen, Wagner, & Anthony, 2002; Conway, Compton, Stinson, & Grant, 2006), one study found that PTSD is the most prevalent co-occurring psychiatric disorder among veterans with CUD presenting to VHA, at 29% (Bonn-Miller et al., 2012). Similar to other SUDs, co-occurrence of CUD and PTSD is associated with greater PTSD symptom severity, decreased likelihood of CUD cessation, and worse PTSD and CUD clinical outcomes (Bonn-Miller et al., 2015; Bonn-Miller, Vujanovic, & Drescher, 2011) as well as greater health services use relative to no comorbidity (Ouimette, Finney, & Moos, 1999; Saladin, Brady, Dansky, & Kilpatrick, 1995; Watkins, Burnam, Kung, & Paddock, 2001).

Although the literature has been primarily focused on the comorbidity between PTSD and cannabis use and CUD (Cogle, Bonn-Miller, Vujanovic, Zvolensky, & Hawkins, 2011; Kevorkian et al., 2015), depression appears to also co-occur with cannabis use problems and dependence in both the general population (Chen et al., 2002; Degenhardt, Hall, & Lynskey, 2003; Feingold, Fox, Rehm, & Lev-Ran, 2015; Grant et al., 2004) and in veterans (Farris, Zvolensky, Boden, & Bonn-Miller, 2014; Goldman et al., 2010). However, the relative roles of MDD and PTSD with respect to cannabis use and problems have not been examined. Given that MDD is more prevalent than other mental health disorders in terms of its co-occurrence with PTSD among returning veterans (Seal et al., 2011), it is presently unclear whether increased risk of cannabis misuse is unique to PTSD symptomatology relative to MDD and whether the mechanisms that explain the pathways from PTSD and MDD to cannabis-related problems and CUD are similar.

Mechanisms Underlying Associations Between PTSD, MDD, and Cannabis Outcomes

Individuals with affective vulnerabilities, such as PTSD and MDD, are especially likely to use cannabis as a means of coping (Boden, Babson, Vujanovic, Short, & Bonn-Miller, 2013; Bonn-Miller, Vujanovic, Feldner, Bernstein, & Zvolensky, 2007; Buckner & Zvolensky, 2014; Bujarski, Norberg, & Copeland, 2012; Johnson, Mullin, Marshall, Bonn-Miller, & Zvolensky, 2010; Mitchell, Zvolensky, Marshall, Bonn-Miller, & Vujanovic, 2007; Simons, Gaher, Correia, Hansen, & Christopher, 2005). In fact, there is broad-based evidence that cannabis may be used by some persons with clinical disorders or individual differences in affective vulnerability factors as an (short-term) emotion regulatory strategy to reduce or manage perceived aversive psychological and

mood states (Metrik, Kahler, McGeary, Monti, & Rohsenow, 2011). For example, emotionally vulnerable cannabis users may rely on cannabis to decrease distress (Potter, Vujanovic, Marshall-Berenz, Bernstein, & Bonn-Miller, 2011), or to cope with symptoms of anxiety and PTSD (Bonn-Miller, Vujanovic, & Zvolensky, 2008; Bonn-Miller, Vujanovic, et al., 2007; Bonn-Miller, Zvolensky, & Bernstein, 2007; Buckner, Bonn-Miller, Zvolensky, & Schmidt, 2007), which, may in turn theoretically be related to greater problematic cannabis use and CUD (Bonn-Miller & Zvolensky, 2009; Farris, Metrik, Bonn-Miller, Kahler, & Zvolensky, in press; Moitra, Christopher, Anderson, & Stein, 2015). Indeed, cannabis coping motives mediate the relation between different indices of affective vulnerability (e.g., anxiety sensitivity, distress intolerance, social anxiety) implicated in the etiology of mood and anxiety disorders and a range of cannabis outcomes, including cannabis use frequency (Johnson, Bonn-Miller, Leyro, & Zvolensky, 2009), severity of cannabis problems (Buckner & Zvolensky, 2014; Bujarski et al., 2012), and cannabis dependence (Johnson et al., 2010) in nonveteran samples.

Importantly, coping to relieve PTSD-related negative affect and distress has been established as a powerful motivator for cannabis use among nonveteran users (Bonn-Miller, Vujanovic, et al., 2007; Bonn-Miller, Vujanovic, Boden, & Gross, 2011; Bujarski et al., 2012; Potter et al., 2011) and veterans (Boden et al., 2013; Bremner, Southwick, Darnell, & Charney, 1996). Two veteran studies have shown that cannabis expectancies about anticipated benefits of cannabis use, a construct conceptually related to motives (Cox & Klinger, 1988), mediated the relation between symptoms of PTSD and cannabis use in an Internet survey of combat veterans (Earleywine & Bolles, 2014), and the relation between depressive symptoms and cannabis use among cannabis-dependent military veterans (Farris et al., 2014). Other work suggests the remission of PTSD symptoms is associated with a 75% or greater reduction in drug and alcohol use days, but reduction in SUD symptoms does not improve PTSD symptoms (Hien et al., 2010). Conversely, lower levels of change in PTSD symptom severity after PTSD treatment was associated with increased use of cannabis postdischarge (Bonn-Miller et al., 2011).

In contrast to other cannabis use motives (e.g., enhancement, social) that are associated with nonproblem patterns of cannabis use, motives broadly related to coping with negative affect are specifically linked with heightened cannabis use, including cannabis-related problems and CUD (Bonn-Miller & Zvolensky, 2009; Bonn-Miller, Boden, Bucossi, & Babson, 2014; Moitra et al., 2015; Simons et al., 2005). Cognitive-motivational factors related to coping and avoidance of social situations have also been linked with cannabis use and related problems (Buckner & Zvolensky, 2014). However, using cannabis specifically as a means of managing sleep problems has received far less attention; yet, sleep disturbance is a prominent symptom of both PTSD and MDD (American Psychiatric Association, 2013). In particular, veterans with PTSD suffer from chronic sleep disturbance (Lavie, Katz, Pillar, & Zinger, 1998; Woodward, Arsenault, Murray, & Bliwise, 2000), and thus, may be using cannabis to help cope with sleep problems in the short-term, particularly because cannabis can produce sedation (Lamarque & De Koninck, 2007; Stewart, Pihl, Conrad, & Dongier, 1998; Vandrey, Babson, Herrmann, & Bonn-Miller, 2014). Although not yet examined among veterans, Bonn-Miller and colleagues (Bonn-Miller, Babson, & Vandrey, 2014;

Bonn-Miller, Boden, et al., 2014) found that medical cannabis users with elevated PTSD symptoms, relative to those with lower PTSD scores, were more likely to use cannabis to improve sleep and used cannabis more frequently. Specifically, sleep but not coping motives interacted with PTSD symptoms to predict frequency of cannabis use. In the overall sample, coping and other motives (boredom, alcohol, enjoyment, celebration, altered perception, social anxiety, availability, and experimentation) but not sleep motives were associated with cannabis problems. Given the dearth of studies on sleep motives in veterans but well-established sleep dysfunction in those with PTSD and MDD, we would expect sleep to be a prominent motive for using cannabis among this population.

Present Study

Past work has not evaluated the affective-motivational model, which emphasizes the central role of negative affect in motivating drug use (Baker, Piper, McCarthy, Majeskie, & Fiore, 2004; Simons et al., 2005), as applied to cannabis use motives as mediators of the relation between PTSD and MDD with cannabis use problems or CUD among veterans. Furthermore, extant studies linking PTSD or MDD with cannabis use severity lack specificity in terms of using cannabis to cope with negative affect versus using to cope with sleep problems that are highly prevalent among veterans with PTSD and MDD. The present study addresses these critical gaps in knowledge in an OEF/OIF/OND sample of lifetime cannabis users with a full range of cannabis involvement (frequency of use ranging from lifetime to current weekly/daily use). The first novel aim was to concurrently examine the relative roles of PTSD and MDD on multiple indicators of cannabis use including cannabis use frequency, cannabis-related problems, and diagnosis with past 12-month *DSM-5* CUD. Based on prior work (e.g., Boden et al., 2013; Earleywine & Bolles, 2014; Farris et al., 2014), we hypothesized that each affective disorder will be significantly associated with each of the three cannabis use indices. The hypotheses pertaining to the strength of the association of PTSD relative to MDD with the cannabis outcomes were exploratory given these predictors have not been previously examined concurrently in one comprehensive study.

The second novel aim was to specifically examine the mediating role of coping motives relative to sleep and anxiety motives as pathways between PTSD and MDD and cannabis use outcomes in one comprehensive multivariate model. Based on findings from research that investigated individual components of our multivariate model (e.g., Boden et al., 2013; Bonn-Miller, Vujanovic, et al., 2007, 2008; Earleywine & Bolles, 2014), we hypothesized that the three motives would significantly mediate the relations of PTSD and MDD with the three dependent variables. Furthermore, sleep and coping motives were expected to have a stronger influence than situational anxiety motives in explaining the relation between MDD and cannabis indices.

Method

Sample and Procedure

Participants were OEF/OIF/OND veterans deployed post-9/11/2001 eligible for the study if they met the following inclusion

criteria: (a) at least 18 years old; (b) an OEF/OIF/OND veteran as confirmed by the Providence VHA Computerized Patient Record System (CPRS); and (c) used cannabis at least once in his or her lifetime. Exclusion criteria were: (a) suicidal risk in the past 2 weeks ascertained with items from the Suicidality Scale of the Inventory of Depression and Anxiety Symptoms (Watson et al., 2007) and clinical interview; (b) psychotic symptoms in the past month assessed by the Structured Clinical Interview for *DSM-IV* Non-Patient Edition (SCID-IV-NP; First, Spitzer, Gibbon, & Williams, 2002); (c) score ≤ 23 on the Mini-Mental Status Exam (Folstein, Folstein, & McHugh, 1975); or (d) active duty at the time of the baseline assessment (due to increased likelihood of study dropout due to deployment). The study was approved by the Institutional Review Boards of Brown University and the Providence VHA.

Participants were recruited from the Veterans Health Administration (VHA) facility in Providence, RI between February 2013 and December 2015. In addition to study advertisements at the Providence VHA, veterans were recruited by utilizing the VHA OEF/OIF/OND Roster, an accruing database of combat veterans who have recently returned from military service in Iraq and Afghanistan and enrolled in VHA. Using the Roster, study information was mailed to potential participants in RI, MA, and CT. Veterans could indicate lack of interest in being contacted for the screening by mailing back a preaddressed, stamped do-not-contact postcard (4.6% of all mailings). Veterans were screened for eligibility by telephone (53% of all outreach calls made) and were invited to complete a baseline interview, at which time they signed informed consent, completed the in-person screening survey, and completed assessments for the baseline portion for this ongoing prospective study. Of the 830 individuals screened for the study, 35% ($n = 292$) did not meet preliminary inclusion/exclusion criteria and nearly 26% ($n = 213$) were eligible but did not show for the baseline assessment ($n = 154$), declined to participate in the study ($n = 32$), or asked to be contacted in the future ($n = 27$). Twenty-one eligible screeners were scheduled for a future study appointment. Of the 304 enrolled in the study, three were deemed ineligible at baseline. Thus, results are based on 301 participants.

The baseline assessment consisted of a battery of interview and self-report assessments and a urine toxicology screen for cannabis. Zero breath alcohol concentration was verified with an Alco-Sensor IV (Intoximeters, Inc., St Louis, MO) to ensure validity of the assessment. All participants were compensated \$50 upon completion of the study session.

Measures

Demographic information. Demographic and background information such as sex, ethnicity, marital status, employment, branch of service, location and number of deployments (OEF/OIF/OND) was collected at baseline and verified through CPRS.

The Clinician Administered PTSD Scale for *DSM-IV*. CAPS (Blake et al., 1995) is a semistructured interview used to assess for lifetime and past month *DSM-IV* PTSD diagnosis. Participants were scored for frequency (score ≥ 1) and intensity (score ≥ 2) of symptoms using established diagnostic guidelines (Weathers, Keane, & Davidson, 2001). Depending on whether they met diagnostic criteria for PTSD, participants were then

assigned a lifetime and past month dichotomous diagnosis score (1 = PTSD, 0 = No PTSD).

Structured Clinical Interview for DSM-IV, Non-Patient Edition. SCID-NP was used to determine diagnosis of lifetime and current (past month) MDD (First et al., 2002). No modifications were necessary to ascertain DSM-5 MDD diagnoses following the release of DSM-5 (American Psychiatric Association, 2013) as the criteria for MDD did not change.

DSM-5 diagnosis of lifetime and current (past year) CUD. This was also determined with the SCID-NP. With the release of DSM-5 (American Psychiatric Association, 2013), legal problems diagnostic criterion was excluded and items assessing cannabis withdrawal and craving were included. Participants endorsing two or more of the 11 symptoms in the past year met criteria for current DSM-5 CUD.

Cannabis-related problems. Cannabis-related problems were assessed with the Marijuana Problems Scale (MPS; Stephens, Roffman, & Curtin, 2000), a self-report 22-item questionnaire that evaluates problems experienced in the past 90 days related to cannabis use. A total count of combined minor and serious problems was used rather than a severity score (only 4.6% endorsed major problems). The MPS has strong internal consistency (Peters, Nich, & Carroll, 2011; Stephens et al., 2000); in this sample it was excellent ($\alpha = .92$).

Cannabis use. The Time-Line Follow-Back Interview (TLFB; Dennis, Funk, Godley, Godley, & Waldron, 2004; Sobell & Sobell, 1992) was conducted at baseline for the 6 months prior to the visit, and used to characterize the sample in terms of percent days cannabis use. In addition to TLFB, all participants were assessed for lifetime cannabis use (study eligibility criterion) and use in the past year. Based on their TLFB interview and endorsement of past-year cannabis use, participants were classified into the following three use frequency categories: lifetime user (no past year use, $n = 181$), past-year user (using less than two times a week, $n = 60$), and frequent user (at least two or more times a week, $n = 60$); with the latter group reflective of regular heavy patterns of use. Frequent users reported using cannabis on an average of 74.5 ($SD = 33.7$) percent of days on the TLFB relative to past-year users, 11.8 ($SD = 17.2$) percent of days.

The Comprehensive Marijuana Motives Questionnaire (CMMQ). Participants rated on a 1 = *almost never/never* to 5 = *almost always/always* scale how often they used cannabis for each of 36 reasons; three items per subscale were used to derive a mean composite score for 12 different motives (Lee, Neighbors, Hendershot, & Grossbard, 2009). Consistent with the affective-motivational model in the present study, only the coping (e.g., “because you were depressed”, “to forget your problems”), social (henceforth situational) anxiety (e.g., “because it relaxes you when you are in an insecure situation”), and sleep (e.g., “because you are having problems sleeping”) cannabis use motives were studied. Internal consistencies for the three factors were good ($\alpha = .84$ to $.91$).

The Pittsburgh Sleep Quality Index (PSQI). The PSQI (Buysse, Reynolds, Monk, Berman, & Kupfer, 1989) is a psychometrically sound self-report questionnaire for assessing sleep quality and disturbances over a 1-month period (Carpenter & Andrykowski, 1998). A global PSQI score greater than 5 indicates poor sleep quality (Buysse et al., 1989). PSQI was used descriptively to characterize sleep patterns in this sample.

Data Analytic Strategy

First, preliminary descriptive analyses were conducted including examining cannabis use motives as a function of diagnosis of PTSD, MDD, and CUD as well as correlations among all predictor and outcome variables. Regression analyses were performed to test the effects of PTSD and MDD on: (a) cannabis use (b) number of cannabis-related problems, and (c) CUD. Ordinal logistic regression was used for the three-level cannabis use variable, Poisson regression was used to examine predictors of the count variable (cannabis problems), and logistic regression was used for CUD. The two predictors were simultaneously entered, covarying for age, as age has been shown to be an important factor related to both drug use and PTSD diagnosis among veterans (Seal et al., 2007, 2011). Sex and marital status were initially also covaried, but were subsequently dropped from all analyses because they were not associated with any outcomes. We also tested whether the effects of the two predictors significantly differed from each other using the Wald test of parameter constraints. Finally, we tested the mediating role of cannabis use motives in the association between diagnoses of PTSD and MDD and the three cannabis outcome variables. Each mediator was first tested in a separate model, for each of the three cannabis outcomes (9 models); then, mediators were entered simultaneously in one model for each of the three criterion variables (3 models) to test whether the associations remained with the other motives in one model. Additionally, within this final model we tested whether the three indirect effects significantly differed from one another. To conduct the tests, the indirect effect for each mediator was computed by taking the product of the “a” path (predictor to mediator) and the “b” path (mediator to outcome); a Wald test of parameter constraints tested whether these product terms were significantly different from each other. Thus, a total of 12 mediational models were tested. All models included the direct effect of the diagnosis and the outcome variable. Models were conducted using Mplus Version 7.3 (Muthén & Muthén, 1998–2012). Mediation was tested using the delta method for the indirect effect (MacKinnon, 2008).

Results

Descriptive Statistics

The sample was predominantly male with a mean age of 33.4 ($SD = 9.5$) years (see Table 1). The median annual family income bracket of participants was \$40,000–49,000; 20.3% were unemployed; and 46.8% were married or cohabiting. The average percent of drinking days was 25.6 ($SD = 29.7$) with participants consuming an average of 4.6 ($SD = 3.6$) drinks per drinking day. Tobacco smokers ($n = 136$) reported cigarette use an average of 78.4% ($SD = 33.8$) of days, and averaged 11.3 ($SD = 7.7$) cigarettes per day on smoking days.

The three cannabis use frequency groups did not significantly differ by sex ($p = .19$), race ($p = .69$), or ethnicity ($p = .82$). Lifetime users were significantly older ($M = 35.66$, $SD = 10.23$) than past-year ($M = 30.23$, $SD = 7.42$) and frequent users ($M = 29.83$, $SD = 6.56$; $F(2, 298) = 13.87$, $p < .001$). Table 2 and Table 3 display sample distribution by PTSD diagnosis (No PTSD,

Table 1
Sample Demographics, Diagnostic, and Military Service-Related Characteristics ($N = 301$)

Variable	<i>n</i>	%
Gender		
Male	285	94.7
Race		
White	244	81.1
Black/African American	11	3.7
Asian	5	1.7
Native Hawaiian/ Pacific Islander	2	.7
Multiracial/Other	38	12.6
Ethnicity		
Hispanic/Latino(a)	38	12.6
Marital Status		
Single/Never married	95	31.6
Married/Living with partner	141	46.8
Divorced/Separated	65	21.6
Employment status ^a		
Employed	233	77.4
Unemployed/Homemaker	61	20.3
Student	71	23.6
Military service	83	27.6
Combat operation(s) served in ^a		
Operation Enduring Freedom (OEF)	225	74.8
Operation Iraqi Freedom (OIF)	160	53.2
Operation New Dawn (OND)	61	20.3
Most recent branch of service		
Army	220	73.1
Marines	35	11.6
Air Force	24	8.0
Navy	21	7.0
Coast Guard	1	.3
THC positive toxicology screen	65	21.6
Major depressive disorder, current	46	15.3
Major depressive disorder, lifetime	150	49.8
Posttraumatic stress disorder, current	41	13.6
Posttraumatic stress disorder, lifetime	78	25.9
Cannabis use disorder, current	47	15.6
Cannabis use disorder, lifetime	119	39.5
	<i>M</i>	<i>SD</i>
Age	33.4	9.5
Years of Education Completed	13.7	2.0
Number of deployments post-9/11/2001	1.8	1.1
Years since last deployment	3.7	2.5
Global PSQI score	8.9	4.1

^a Multiple responses permitted.

PTSD) and MDD diagnosis (No MDD, MDD) within CUD (Table 2) and each of the cannabis use frequency groups (Table 3). Levels of cannabis use motives and outcomes as a function of PTSD and MDD are shown in Table 4.

Association Between PTSD, MDD, and the Cannabis-Dependent Variables

Bivariate correlations are presented in Table 5. The PTSD and MDD variables were significantly positively associated with each other. Both PTSD and MDD were positively significantly associated with cannabis use motives. The three cannabis use motives were significantly positively associated with all three criterion variables. The criterion variables were significantly intercorrelated (medium to large sized associations). Importantly, there were

correlations in the small-to-medium size range among the predictors (PTSD and MDD) and outcomes, including significant but small-sized correlation between PTSD and problems.

Next, we conducted regression analyses to examine the effects of PTSD and MDD on the three dependent variables, controlling for age. Findings indicated that PTSD and MDD each were significantly associated with greater cannabis use ($OR = 1.75$, 95% CI [1.22, 2.52], $p < .01$; $OR = 1.65$, 95% CI [1.17, 2.33], $p < .01$, respectively); and the two predictors did not significantly differ from each other, Wald $\chi^2(1, N = 301) = 0.04$, $p = .83$. Similarly, PTSD and MDD both uniquely predicted the likelihood of being diagnosed with a CUD ($OR = 1.71$, 95% CI [1.14, 2.59], $p \leq .01$; $OR = 1.78$, 95% CI [1.19, 2.66], $p < .01$, respectively); and the two predictors did not significantly differ from each other, Wald $\chi^2(1, N = 301) = 0.01$, $p = .91$. In the model predicting cannabis problems, there was a significant main effect for MDD but not PTSD, such that diagnosis of MDD was significantly associated with greater number of cannabis problems ($b = .59$, $SE = .06$, $\beta = .84$; $p < .001$); these effects significantly differed from each other, Wald $\chi^2(1, N = 299) = 21.66$, $p < .001$.

Mediation Results

Finally, we conducted a set of mediation models for each of the three outcomes (see Figure 1). Mediation of the association between PTSD and cannabis problems was not tested due to lack of significant association between PTSD and the criterion in the multivariate model. Results are presented in Table 6.

Cannabis use frequency. In separate single mediator models (top panel of Table 6), there was a significant indirect effect of PTSD and MDD through each use motive on cannabis use frequency; inclusion of each mediator accounted for a sizable portion of the effects of PTSD and MDD on the dependent variables (between 26% and 57% for PTSD; between 26% and 43% for MDD). In the multiple mediator model (bottom panel of Table 6), the indirect effect of PTSD and MDD through sleep motives on cannabis use was significant and accounted for 57% and 43% of the total effect, respectively. The indirect effects of PTSD and MDD through coping and situational anxiety motives were non-significant (and did not differ from each other), indicating that these motives did not mediate the effects of the affective disorders when sleep motives were also in the model. Furthermore, with cannabis use motives in the model predicting cannabis use frequency, the direct effects of PTSD and MDD were nonsignificant, indicating full mediation.

Cannabis problems. In separate single mediator models, there was a significant indirect effect of MDD through each use

Table 2
Posttraumatic Stress Disorder (PTSD) by Major Depressive Disorder (MDD) by Cannabis Use Disorder (CUD)

	CUD		No CUD		Total <i>n</i>
	No MDD <i>n</i> (%)	MDD <i>n</i> (%)	No MDD <i>n</i> (%)	MDD <i>n</i> (%)	
PTSD	8 (26.7)	8 (47.1)	13 (5.8)	12 (41.4)	41
No PTSD	22 (73.3)	9 (52.9)	212 (94.2)	17 (58.6)	260
Total	30 (100)	17 (100)	225 (100)	29 (100)	301

Table 3
 Posttraumatic Stress Disorder (PTSD) by Major Depressive Disorder (MDD) by Cannabis Use Frequency

	Frequent use		Past-year use		Lifetime use		Total
	No MDD n (%)	MDD n (%)	No MDD n (%)	MDD n (%)	No MDD n (%)	MDD n (%)	n (%)
PTSD	9 (23.1)	12 (57.1)	2 (3.8)	3 (42.9)	10 (6.1)	5 (27.8)	41
No PTSD	30 (76.9)	9 (42.9)	51 (96.2)	4 (57.1)	153 (93.9)	13 (72.2)	260
Total	39 (100)	21 (100)	53 (100)	7 (100)	163 (100)	18 (100)	301

motive on cannabis problems, indicating that each motive was a partial mediator of MDD; the indirect effect through each motives variable accounted for between 16% and 28% of the total effect. In the multiple mediator model, the indirect effects of MDD through sleep and coping motives on cannabis problems were significant (percent mediated was 22% and 18%, respectively); these indirect effects were similar in magnitude and did not significantly differ from each other. In contrast, the indirect effects of MDD through situational anxiety motive was nonsignificant, indicating that this motive did not mediate the effects of MDD when the other two motives were also in the model. There was still a significant direct effect between MDD and problems, even after accounting for mediators in the model, thus suggesting that there was only partial mediation.

CUD. In separate single mediator models, there were significant indirect effects of PTSD and MDD through each motive on CUD, with each mediator accounting for between 17% and 51% of the total effect. In the multiple mediator model the indirect effects of MDD through sleep and coping motives on CUD were significant and accounted for 26% and 21% of the variance, respectively. The indirect effect of PTSD through sleep motive also was significant, accounting for 42% of the variance, and through coping was at trend level ($p = .054$), accounting for 20% of the total effect. The indirect effects of PTSD and MDD through situational anxiety motive were nonsignificant, indicating that this motive did not mediate the effects of disorders when the other two motives

were also in the model. The indirect effect for situational anxiety motives significantly differed from those of coping and sleep motives, which did not significantly differ from each other. With cannabis use motives in the model predicting CUD, the direct effects of PTSD and MDD were nonsignificant, indicating full mediation.

Discussion

This is the first study to evaluate the affective-motivational model of substance use with coping-oriented cannabis use motives including coping with negative affect, situational anxiety, and sleep problems as mediators of the relation between PTSD and MDD with cannabis use, related problems, and CUD. Results indicated that MDD was significantly associated with all three cannabis use indices; PTSD was significantly associated with cannabis use and CUD. Both mental health disorders were significantly associated with all three cannabis use motives. In support of our hypotheses, there was evidence of coping-oriented reasons for use explicating these relations of the MDD and PTSD with greater frequency of cannabis use or with CUD. Furthermore, specific cannabis use motives related to managing sleep and coping with negative affect had stronger influence than situational anxiety motives on veterans' risk for cannabis misuse when tested simultaneously. Finally, comparison of the relative pathways of PTSD and MDD with the indices of cannabis use revealed the

Table 4
 Comparisons of Cannabis Use Motives and Outcomes as a Function of Diagnosis of Posttraumatic Stress Disorder (PTSD) and Major Depressive Disorder (MDD)

	No PTSD M (SD)	PTSD M (SD)	t-test	d	No MDD M (SD)	MDD M (SD)	t-test	d
Coping motives	1.51 (.93)	2.25 (1.19)	4.55***	.69	1.50 (.90)	2.25 (1.28)	4.84***	.68
Sleep motives	1.80 (1.19)	2.87 (1.49)	5.15***	.79	1.82 (1.23)	2.65 (1.40)	4.15***	.63
Situational anxiety motives	1.54 (.88)	2.24 (1.22)	4.41***	.66	1.54 (.88)	2.17 (1.23)	4.16***	.59
Cannabis-related problems	.98 (2.63)	1.98 (3.56)	2.14*	.32	.81 (2.36)	2.76 (4.14)	4.49***	.58
	%	%	χ^2	OR	%	%	χ^2	OR
Cannabis use			28.9***				22.47***	
Lifetime users ^a	64.1	36.6			64.2	39.1		
Past-year users	20.8	12.2		1.02	20.5	15.2		1.22
Frequent users	15.1	51.2		5.96	15.4	45.7		4.88
Cannabis use disorder	11.9	39.0	19.74***	4.73	11.8	37.0	18.77***	4.40

Note. d = Cohen's d; ^a = reference group; OR = odds ratio; Results are based on t-tests and chi-square tests.

* $p < .053$. *** $p < .001$.

Table 5
Correlations Among Variables

Variable	2	3	4	5	6	7	8	9
1. Posttraumatic stress disorder	.37**	.25**	.29**	.25**	.27**	.12*	.26**	-.07
2. Major depressive disorder		.27**	.23**	.23**	.25**	.25**	.25**	-.01
3. Coping motives			.56**	.61**	.37**	.32**	.38**	-.09
4. Sleep motives				.60**	.61**	.32**	.44**	-.28**
5. Situational anxiety motives					.37**	.22**	.27**	-.11
6. Cannabis use						.33**	.62**	-.27**
7. Cannabis-related problems							.40**	-.08
8. Cannabis use disorder								-.14*
9. Age								—

Note. Correlations among continuous variables are Pearson correlations; correlations among categorical variables are phi coefficients, and correlations between continuous and categorical variables are point-biserial correlations. For the purposes of the correlation matrix, cannabis use was treated as a continuous variable, although it is treated as an ordered categorical variable in regression analyses.

* $p \leq .05$. ** $p \leq .01$.

unique and overall equivalent predictive roles of the two disorders in increasing likelihood of cannabis use and CUD.

The present study replicates the findings from prior studies that support the central role of cannabis coping motives in PTSD (Boden et al., 2013; Bonn-Miller et al., 2011; Bremner et al., 1996; Bujarski et al., 2012; Potter et al., 2011) and more broadly in emotional dysregulation (Bonn-Miller, Vujanovic, et al., 2007, 2008; Zvolensky et al., 2009) among cannabis users. We extended these findings on coping with negative affect to the other two more specific coping-oriented reasons: using cannabis to manage sleep problems and to avoid situational anxiety. Each of these motives appears to directly map onto the clinical presentation of PTSD and MDD. The situational anxiety and sleep relations are consistent with prior research demonstrating that lower levels of subjective improvement in PTSD symptoms of avoidance and hyperarousal are associated with increased risk of cannabis use in veterans (Bonn-Miller et al., 2011). The relationship between MDD and cannabis use via cannabis use motives observed here differ from past work focused on depressive symptoms among nonveteran samples of young adult cannabis users (Bonn-Miller, Zvolensky, Bernstein, & Stickle, 2008; Johnson et al., 2009), potentially due to the sample type (e.g., severity). In both of these studies, self-reported depressive symptoms were significantly associated with cannabis coping motives but not with past month cannabis use frequency.

In the multiple mediator models, sleep motives emerged as a dominant factor in explicating the relations between each of the two affective disorders with cannabis use. Specifically, using cannabis to manage sleep disturbance fully mediated the effect of both PTSD and MDD on frequency of cannabis use. Together with motives broadly related to coping with negative affect, sleep motives also fully mediated the effects of PTSD and MDD on increased risk of CUD and partially mediated the effect of MDD on cannabis-related problems. These findings are consistent with one prior study with medical marijuana users that showed sleep, but not coping motives, were associated with symptoms of PTSD and increased cannabis use (Bonn-Miller, Babson, et al., 2014). The novel aspect of the current study, however, is extending these findings to the diagnosis of MDD and to cannabis use outcomes across the full spectrum from frequency of use to cannabis-related problems to CUD.

Situational anxiety motive did not mediate the association between the affective disorders and the three indices of cannabis use in the multiple mediator models. Using cannabis as a means of managing situational or social anxiety may be closely related to the PTSD's avoidance symptoms, as demonstrated in the single mediator models. However, in line with our hypothesis, sleep and coping with negative affect appear to be more salient motives to use cannabis in veterans with PTSD and MDD. Future studies should consider examining the construct validity of the situational (social) anxiety motives of the CMMQ in relation to social and other anxiety disorders. Alternative motivational processes underlying the relation between cannabis use and affective disorders could also be explored.

The analysis of different types of motives for using cannabis has several significant clinical implications. For instance, omitting sleep when discussing cannabis use in clinical samples, especially in veterans, may result in missing an important underlying reason for why individuals are using cannabis and why they may be ambivalent about quitting use. Pharmacological properties of cannabis are conducive to sedation and relaxation (Bonn-Miller, Babson, et al., 2014), which could translate to subjective perception of improved sleep onset by many cannabis users (Conroy & Arnedt, 2014) and may also explain its widespread use among veterans with PTSD, MDD, and sleep disturbances. Nevertheless, cannabis, particularly at high doses, may not be beneficial to sleep as it disrupts sleep architecture including decreased REM and slow wave sleep, and increased sleep onset latency (Gates, Albertella, & Copeland, 2014; Garcia & Salloum, 2015). Furthermore, withdrawal from cannabis results in significantly disturbed sleep in the majority of heavy cannabis users (Budney, Hughes, Moore, & Vandrey, 2004), which subsequently places abstinent users at greater risk for relapse (Vandrey et al., 2014). Engaging veterans who use cannabis in evidence-based treatments that address sleep, such as cognitive-behavioral therapy for insomnia (CBT-I; Morin et al., 2006; Perlis, Jungquist, Smith, & Posner, 2006), may have high impact on clinical outcomes for PTSD and CUD (Babson, Ramo, Baldini, Vandrey, & Bonn-Miller, 2015). The present findings also suggest targeting coping motives in the context of integrated behavioral treatments for CUD and PTSD as well as MDD may be useful. Comorbid treatments that specifically focus on coping skills for mood and stress management (Buckner et al.,




Figure 1. Mediation analysis of the relation between PTSD/MDD and cannabis use (Panel A), cannabis-related problems (Panel B), and cannabis use disorder (Panel C).

2014) or increase an individual's tolerance for emotional distress and decrease avoidance of distress via cannabis use (Bonn-Miller, Vujanovic, Twohig, Medina, & Huggins, 2010) may be particularly effective in veterans with CUD and co-occurring PTSD/MDD.

Identifying theory-driven mechanisms that maintain the association between the two most prevalent affective disorders in veterans and cannabis use will also help improve the screening procedures currently in place in clinical settings to detect cannabis misuse as early as possible, which has the potential to increase early intervention. A short screener that assesses cannabis consumption, problems, dependence, and perhaps even psychological sequelae might be warranted (e.g., the Cannabis Use Disorders Test; Adamson et al., 2010). Cannabis users may

develop problems all along a continuum ranging in severity of negative consequences (e.g., procrastination, low energy, memory problems, sleep problems on the lower end of the spectrum) to more severe problems, consistent with CUD (e.g., loss of control over drug use; Piomelli, Haney, Budney, & Piazza, 2016). Clinically, this distinction has great utility as screening for cannabis-related problems (in addition to CUD) may facilitate a therapeutic intervention by tapping into problem areas that help elicit motivation for changing cannabis use. Naturally, effective screening and discussion will not occur if providers are hesitant or unsure of how to assess cannabis use for reasons such as current political and legal climate regarding medical marijuana, use of cannabis to alleviate mental health conditions that have been caused or exacerbated by military service,

Table 6

Indirect and Direct Effects for the Mediation of the Association Between Posttraumatic Stress Disorder (PTSD) and Major Depressive Disorder (MDD) With Cannabis Use, Cannabis-Related Problems, and Cannabis Use Disorder (CUD), by Three Cannabis Use Motives

Cannabis use motives	Cannabis use		Cannabis-related problems	CUD	
	PTSD	MDD	MDD	PTSD	MDD
Single mediator					
Coping	.14 (26%)*	.16 (31%)**	.16 (27%***)	.18 (31%)*	.20 (32%)**
Situational anxiety	.18 (26%)*	.13 (26%)*	.09 (16%***)	.12 (22%)*	.10 (17%)*
Sleep	.39 (57%***)	.24 (43%)*	.17 (28%***)	.31 (51%)**	.19 (29%)*
Multiple mediators					
Total indirect effect	.39 (57%***)	.24 (43%)*	.21 (35%***)	.33 (52%***)	.25 (39%)**
Coping	.004 (1%) ^a	.004 (1%) ^a	.11 (18%) ^{a***}	.13 (20%) ^a $p = .054$.14 (21%) ^{a*}
Situational anxiety	-.003 (0%) ^a	-.003 (0%) ^a	-.04 (0%) ^b	-.06 (0%) ^b	-.05 (0%) ^b
Sleep	.39 (57%) ^{b***}	.24 (43%) ^{b*}	.13 (22%) ^{a***}	.27 (42%) ^{a***}	.16 (26%) ^{a*}
Direct effect	.30	.32 ^{p = .095}	.39***	.31	.39 ^{p = .09}

Note. Parameters are unstandardized. All models control for age. Mediation of the association between PTSD and cannabis-related problems was not tested due to lack of significant association between the predictor and the criterion. Indirect effects sharing common superscripts did not significantly differ from each other.

* $p \leq .05$. ** $p \leq .01$. *** $p \leq .001$.

stigma related to military regulations of substance use, or feeling that one lacks the expertise or education on cannabis use and treatments (Bujarski et al., 2016).

The strengths of the current study include a large veteran sample and selecting on lifetime cannabis use and therefore permitting the examination of the full spectrum from lifetime to problem/dependent end. Consistent with prior reports on the comorbidity of CUD with PTSD (Bonn-Miller et al., 2012), 34% of veterans with CUD in the current sample had co-occurring PTSD and similarly 36% had co-occurring MDD, while 17% were comorbid for CUD-PTSD-MDD. Although CUD prevalence estimates are not well-established among OEF/OIF/OND veterans, the current sample rate of 15.6% exceeds the CUD estimate of 10% in the general nonveteran population of lifetime cannabis users (Hasin et al., 2015). This sample was also comparable in terms of PTSD and MDD rates (Fulton et al., 2015; Seal et al., 2009) and age, gender, and other demographic characteristics to other OEF/OIF samples (e.g., Cohen et al., 2010; Seal et al., 2007, 2011) but younger than studies based on veterans from other eras, treatment-engaged (e.g., PTSD in Bonn-Miller et al., 2012), or selected based on presence of a substance use disorder (e.g., CUD in Boden et al., 2013). Another strength was the use of interview-based “gold standard” diagnostic measures for PTSD, MDD, and CUD. Cannabis use status was biochemically verified with the urine toxicology screen. Furthermore, prior research linking PTSD or depression with cannabis use severity lacked specificity in terms of using cannabis to cope with negative affect and other symptoms versus using to specifically cope with sleep problems highly prevalent among veterans with PTSD and MDD. This study found specific motives of coping with negative affect, situational anxiety and sleep problems to underlie the PTSD and MDD relationship with cannabis use and misuse, especially sleep motives.

However, the findings of the study must be considered in the context of some limitations and may not generalize to all OEF/OIF/OND veterans who are using the VHA for health care services. In this initial examination, the cross-sectional design limits causal inferences about the association between the affective dis-

orders and cannabis use outcomes. Nevertheless, similar to the temporal relation between PTSD and CUD (e.g., Hien et al., 2010), MDD is typically found to precede CUD and contribute to its etiology (Agosti et al., 2002; Conway et al., 2006). Our continuing longitudinal study, which was designed to assess these variables at independent time points, will help clarify the temporal sequencing of these relations and examine cannabis use motives as mediators. Longitudinal models will also help test specific hypotheses for current versus lifetime users with respect to potentially different mechanisms underlying the relationship of PTSD and MDD with CUD and cannabis problems. Next, a small number of female veterans in our sample limited the generalizability of our findings and resulted in low power to detect any possible gender differences in motivational processes linking affective disorders with cannabis use. For example, previous research with nonveterans have found gender differences in coping motives mediating the relationship between distress tolerance and cannabis use (Bujarski et al., 2012). Multiple statistical tests were conducted in the present analyses, which may have inflated the risk of Type I error. While the risk of Type I error was likely mitigated to a large degree by our theory-based approach and a priori hypotheses, replication is needed to further substantiate these findings. Next, similar to other recent studies (e.g., Bonn-Miller et al., 2015) we employed *DSM-IV* criteria for PTSD diagnosis due to the timing of the study relative to the transition to *DSM-5* version. Replication studies using *DSM-5* criteria for PTSD may be needed. Finally, although interviewers were trained in the reliable use of the SCID and the CAPS diagnostic instruments and were required to demonstrate competence and adherence, interrater reliability ratings of the diagnoses were not conducted in the present study.

In conclusion, despite the coping-oriented premise for cannabis use steering a number of U.S. states to legalize medical cannabis as treatment for individuals with PTSD (the only psychological condition among other medical conditions sanctioned for medical marijuana use), the science behind this legislative action is lacking. Controlled studies on its effectiveness and safety in use to cope with PTSD symptoms are needed to substantiate the perceived

self-reported benefits of cannabis in this patient population and to test whether benefits of using plant-based cannabis outweigh the risks (Korem, Zer-Aviv, Ganon-Elazar, Abush, & Akirav, 2016). Findings from the present study provided further evidence that individuals with PTSD and MDD, the two most prevalent mental health disorders among Iraq and Afghanistan veterans, are at increased risk for cannabis use and related problems including CUD. Future studies should consider examining other potential mediators that may also explain the link between PTSD and MDD with cannabis use (e.g., difficulty in emotion regulation; Bonn-Miller et al., 2011). Findings highlight the benefit of identifying specific motives for cannabis use, ones that are relevant to symptoms of PTSD and MDD such as sleep and coping with negative affect, as targets in empirically supported treatments that can alleviate these symptoms.

References

- Adamson, S. J., Kay-Lambkin, F. J., Baker, A. L., Lewin, T. J., Thornton, L., Kelly, B. J., & Sellman, J. D. (2010). An improved brief measure of cannabis misuse: The Cannabis Use Disorders Identification Test-Revised (CUDIT-R). *Drug and Alcohol Dependence, 110*, 137–143. <http://dx.doi.org/10.1016/j.drugalcdep.2010.02.017>
- Agosti, V., Nunes, E., & Levin, F. (2002). Rates of psychiatric comorbidity among U.S. residents with lifetime cannabis dependence. *The American Journal of Drug and Alcohol Abuse, 28*, 643–652. <http://dx.doi.org/10.1081/ADA-120015873>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Babson, K. A., Ramo, D. E., Baldini, L., Vandrey, R., & Bonn-Miller, M. O. (2015). Mobile app-delivered cognitive behavioral therapy for insomnia: Feasibility and initial efficacy among veterans with cannabis use disorders. *JMIR Research Protocols, 4*(3), e87. <http://dx.doi.org/10.2196/resprot.3852>
- Baker, T. B., Piper, M. E., McCarthy, D. E., Majeskie, M. R., & Fiore, M. C. (2004). Addiction motivation reformulated: An affective processing model of negative reinforcement. *Psychological Review, 111*, 33–51. <http://dx.doi.org/10.1037/0033-295X.111.1.33>
- Blake, D. D., Weathers, F. W., Nagy, L. M., Kaloupek, D. G., Gusman, F. D., Charney, D. S., & Keane, T. M. (1995). The development of a Clinician-Administered PTSD Scale. *Journal of Traumatic Stress, 8*, 75–90. <http://dx.doi.org/10.1002/jts.2490080106>
- Boden, M. T., Babson, K. A., Vujanovic, A. A., Short, N. A., & Bonn-Miller, M. O. (2013). Posttraumatic stress disorder and cannabis use characteristics among military veterans with cannabis dependence. *The American Journal on Addictions, 22*, 277–284. <http://dx.doi.org/10.1111/j.1521-0391.2012.12018.x>
- Bonn-Miller, M. O., Babson, K. A., & Vandrey, R. (2014). Using cannabis to help you sleep: Heightened frequency of medical cannabis use among those with PTSD. *Drug and Alcohol Dependence, 136*, 162–165. <http://dx.doi.org/10.1016/j.drugalcdep.2013.12.008>
- Bonn-Miller, M. O., Boden, M. T., Bucossi, M. M., & Babson, K. A. (2014). Self-reported cannabis use characteristics, patterns and helpfulness among medical cannabis users. *The American Journal of Drug and Alcohol Abuse, 40*, 23–30. <http://dx.doi.org/10.3109/00952990.2013.821477>
- Bonn-Miller, M. O., Harris, A. H., & Trafton, J. A. (2012). Prevalence of cannabis use disorder diagnoses among veterans in 2002, 2008, and 2009. *Psychological Services, 9*, 404–416. <http://dx.doi.org/10.1037/a0027622>
- Bonn-Miller, M. O., Moos, R. H., Boden, M. T., Long, W. R., Kimerling, R., & Trafton, J. A. (2015). The impact of posttraumatic stress disorder on cannabis quit success. *The American Journal of Drug and Alcohol Abuse, 41*, 339–344. <http://dx.doi.org/10.3109/00952990.2015.1043209>
- Bonn-Miller, M. O., Vujanovic, A. A., Boden, M. T., & Gross, J. J. (2011). Posttraumatic stress, difficulties in emotion regulation, and coping-oriented marijuana use. *Cognitive Behaviour Therapy, 40*, 34–44. <http://dx.doi.org/10.1080/16506073.2010.525253>
- Bonn-Miller, M. O., Vujanovic, A. A., & Drescher, K. D. (2011). Cannabis use among military veterans after residential treatment for posttraumatic stress disorder. *Psychology of Addictive Behaviors, 25*, 485–491. <http://dx.doi.org/10.1037/a0021945>
- Bonn-Miller, M. O., Vujanovic, A. A., Feldner, M. T., Bernstein, A., & Zvolensky, M. J. (2007). Posttraumatic stress symptom severity predicts marijuana use coping motives among traumatic event-exposed marijuana users. *Journal of Traumatic Stress, 20*, 577–586. <http://dx.doi.org/10.1002/jts.20243>
- Bonn-Miller, M. O., Vujanovic, A. A., Twohig, M. P., Medina, J. L., & Huggins, J. L. (2010). Posttraumatic stress symptom severity and marijuana use coping motives: A test of the mediating role of non-judgmental acceptance within a trauma-exposed community sample. *Mindfulness, 1*, 98–106. <http://dx.doi.org/10.1007/s12671-010-0013-6>
- Bonn-Miller, M. O., Vujanovic, A. A., & Zvolensky, M. J. (2008). Emotional dysregulation: Association with coping-oriented marijuana use motives among current marijuana users. *Substance Use & Misuse, 43*, 1653–1665. <http://dx.doi.org/10.1080/10826080802241292>
- Bonn-Miller, M. O., & Zvolensky, M. J. (2009). An evaluation of the nature of marijuana use and its motives among young adult active users. *The American Journal on Addictions, 18*, 409–416. <http://dx.doi.org/10.3109/10550490903077705>
- Bonn-Miller, M. O., Zvolensky, M. J., & Bernstein, A. (2007). Marijuana use motives: Concurrent relations to frequency of past 30-day use and anxiety sensitivity among young adult marijuana smokers. *Addictive Behaviors, 32*, 49–62. <http://dx.doi.org/10.1016/j.addbeh.2006.03.018>
- Bonn-Miller, M. O., Zvolensky, M. J., Bernstein, A., & Stickle, T. R. (2008). Marijuana coping motives interact with marijuana use frequency to predict anxious arousal, panic-related catastrophic thinking, and worry among current marijuana users. *Depression and Anxiety, 25*, 862–873. <http://dx.doi.org/10.1002/da.20370>
- Bremner, J. D., Southwick, S. M., Darnell, A., & Charney, D. S. (1996). Chronic PTSD in Vietnam combat veterans: Course of illness and substance abuse. *The American Journal of Psychiatry, 153*, 369–375. <http://dx.doi.org/10.1176/ajp.153.3.369>
- Buckner, J. D., Bonn-Miller, M. O., Zvolensky, M. J., & Schmidt, N. B. (2007). Marijuana use motives and social anxiety among marijuana-using young adults. *Addictive Behaviors, 32*, 2238–2252. <http://dx.doi.org/10.1016/j.addbeh.2007.04.004>
- Buckner, J. D., & Zvolensky, M. J. (2014). Cannabis and related impairment: The unique roles of cannabis use to cope with social anxiety and social avoidance. *The American Journal on Addictions, 23*, 598–603. <http://dx.doi.org/10.1111/j.1521-0391.2014.12150.x>
- Buckner, J. D., Zvolensky, M. J., Schmidt, N. B., Carroll, K. M., Schatschneider, C., & Crapanzano, K. (2014). Integrated cognitive behavioral therapy for cannabis use and anxiety disorders: Rationale and development. *Addictive Behaviors, 39*, 495–496. <http://dx.doi.org/10.1016/j.addbeh.2013.10.023>
- Budney, A. J., Hughes, J. R., Moore, B. A., & Vandrey, R. (2004). Review of the validity and significance of cannabis withdrawal syndrome. *The American Journal of Psychiatry, 161*, 1967–1977. <http://dx.doi.org/10.1176/appi.ajp.161.11.1967>
- Bujarski, S. J., Galang, J. N., Short, N. A., Trafton, J. A., Gifford, E. V., Kimerling, R., . . . Bonn-Miller, M. O. (2016). Cannabis use disorder treatment barriers and facilitators among veterans with PTSD. *Psychology of Addictive Behaviors, 30*, 73–81. <http://dx.doi.org/10.1037/adb0000131>

- Bujarski, S. J., Norberg, M. M., & Copeland, J. (2012). The association between distress tolerance and cannabis use-related problems: The mediating and moderating roles of coping motives and gender. *Addictive Behaviors*, *37*, 1181–1184. <http://dx.doi.org/10.1016/j.addbeh.2012.05.014>
- Buysse, D. J., Reynolds, C. F., III, Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Research*, *28*, 193–213. [http://dx.doi.org/10.1016/0165-1781\(89\)90047-4](http://dx.doi.org/10.1016/0165-1781(89)90047-4)
- Carpenter, J. S., & Andrykowski, M. A. (1998). Psychometric evaluation of the Pittsburgh Sleep Quality Index. *Journal of Psychosomatic Research*, *45*, 5–13. [http://dx.doi.org/10.1016/S0022-3999\(97\)00298-5](http://dx.doi.org/10.1016/S0022-3999(97)00298-5)
- Chen, C. Y., Wagner, F. A., & Anthony, J. C. (2002). Marijuana use and the risk of major depressive episode: Epidemiological evidence from the United States National Comorbidity Survey. *Social Psychiatry and Psychiatric Epidemiology*, *37*, 199–206. <http://dx.doi.org/10.1007/s00127-002-0541-z>
- Cohen, B. E., Gima, K., Bertenthal, D., Kim, S., Marmar, C. R., & Seal, K. H. (2010). Mental health diagnoses and utilization of VA non-mental health medical services among returning Iraq and Afghanistan veterans. *Journal of General Internal Medicine*, *25*, 18–24. <http://dx.doi.org/10.1007/s11606-009-1117-3>
- Conroy, D. A., & Arnedt, J. T. (2014). Sleep and substance use disorders: An update. *Current Psychiatry Reports*, *16*, 487. <http://dx.doi.org/10.1007/s11920-014-0487-3>
- Conway, K. P., Compton, W., Stinson, F. S., & Grant, B. F. (2006). Lifetime comorbidity of DSM-IV mood and anxiety disorders and specific drug use disorders: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *The Journal of Clinical Psychiatry*, *67*, 247–258. <http://dx.doi.org/10.4088/JCP.v67n0211>
- Cogle, J. R., Bonn-Miller, M. O., Vujanovic, A. A., Zvolensky, M. J., & Hawkins, K. A. (2011). Posttraumatic stress disorder and cannabis use in a nationally representative sample. *Psychology of Addictive Behaviors*, *25*, 554–558. <http://dx.doi.org/10.1037/a0023076>
- Cox, W. M., & Klinger, E. (1988). A motivational model of alcohol use. *Journal of Abnormal Psychology*, *97*, 168–180. <http://dx.doi.org/10.1037/0021-843X.97.2.168>
- Degenhardt, L., Hall, W., & Lynskey, M. (2003). Exploring the association between cannabis use and depression. *Addiction*, *98*, 1493–1504. <http://dx.doi.org/10.1046/j.1360-0443.2003.00437.x>
- Dennis, M. L., Funk, R., Godley, S. H., Godley, M. D., & Waldron, H. (2004). Cross-validation of the alcohol and cannabis use measures in the Global Appraisal of Individual Needs (GAIN) and Timeline Followback (TLFB; Form 90) among adolescents in substance abuse treatment. *Addiction*, *99*(Suppl. 2), 120–128. <http://dx.doi.org/10.1111/j.1360-0443.2004.00859.x>
- Department of Veterans Affairs. (2009). *VA/DoD clinical practice guideline for management of substance use disorders*. Retrieved from http://www.healthquality.va.gov/guidelines/MH/sud/sud_full_601f.pdf
- Earleywine, M., & Bolles, J. R. (2014). Marijuana, expectancies, and posttraumatic stress symptoms: A preliminary investigation. *Journal of Psychoactive Drugs*, *46*, 171–177. <http://dx.doi.org/10.1080/02791072.2014.920118>
- Farris, S. G., Metrik, J., Bonn-Miller, M. O., Kahler, C. W., & Zvolensky, M. J. (in press). Anxiety sensitivity and distress intolerance in predicting cannabis dependence symptoms, problems, and craving: The mediating role of coping motives. *Journal of Studies on Alcohol and Drugs*.
- Farris, S. G., Zvolensky, M. J., Boden, M. T., & Bonn-Miller, M. O. (2014). Cannabis use expectancies mediate the relation between depressive symptoms and cannabis use among cannabis-dependent veterans. *Journal of Addiction Medicine*, *8*, 130–136. <http://dx.doi.org/10.1097/ADM.0000000000000010>
- Feingold, D., Fox, J., Rehm, J., & Lev-Ran, S. (2015). Natural outcome of cannabis use disorder: A 3-year longitudinal follow-up. *Addiction*, *110*, 1963–1974. <http://dx.doi.org/10.1111/add.13071>
- First, M. B., Spitzer, R. L., Gibbon, M., & Williams, J. B. W. (2002). *Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Non-patient Edition* (SCID-I/NP). New York, NY: Biometrics Research, New York State Psychiatric Institute.
- Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). Mini-mental state: A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, *12*, 189–198. [http://dx.doi.org/10.1016/0022-3956\(75\)90026-6](http://dx.doi.org/10.1016/0022-3956(75)90026-6)
- Fulton, J. J., Calhoun, P. S., Wagner, H. R., Schry, A. R., Hair, L. P., Feeling, N., . . . Beckham, J. C. (2015). The prevalence of posttraumatic stress disorder in Operation Enduring Freedom/Operation Iraqi Freedom (OEF/OIF) veterans: A meta-analysis. *Journal of Anxiety Disorders*, *31*, 98–107. <http://dx.doi.org/10.1016/j.janxdis.2015.02.003>
- Garcia, A. N., & Salloum, I. M. (2015). Polysomnographic sleep disturbances in nicotine, caffeine, alcohol, cocaine, opioid, and cannabis use: A focused review. *The American Journal on Addictions*, *24*, 590–598. <http://dx.doi.org/10.1111/ajad.12291>
- Gates, P. J., Albertella, L., & Copeland, J. (2014). The effects of cannabinoid administration on sleep: A systematic review of human studies. *Sleep Medicine Reviews*, *18*, 477–487. <http://dx.doi.org/10.1016/j.smrv.2014.02.005>
- Goldman, M., Suh, J. J., Lynch, K. G., Szucs, R., Ross, J., Xie, H., . . . Oslin, D. W. (2010). Identifying risk factors for marijuana use among Veterans Affairs patients. *Journal of Addiction Medicine*, *4*, 47–51. <http://dx.doi.org/10.1097/ADM.0b013e3181b18782>
- Golub, A., Vazan, P., Bennett, A. S., & Liberty, H. J. (2013). Unmet need for treatment of substance use disorders and serious psychological distress among veterans: A nationwide analysis using the NSDUH. *Military Medicine*, *178*, 107–114. <http://dx.doi.org/10.7205/MILMED-D-12-00131>
- Grant, B. F., Stinson, F. S., Dawson, D. A., Chou, S. P., Dufour, M. C., Compton, W., . . . Kaplan, K. (2004). Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Archives of General Psychiatry*, *61*, 807–816. <http://dx.doi.org/10.1001/archpsyc.61.8.807>
- Hasin, D. S., Saha, T. D., Kerridge, B. T., Goldstein, R. B., Chou, S. P., Zhang, H., . . . Grant, B. F. (2015). Prevalence of marijuana use disorders in the United States between 2001–2002 and 2012–2013. *Journal of the American Medical Association Psychiatry*, *72*, 1235–1242. <http://dx.doi.org/10.1001/jamapsychiatry.2015.1858>
- Hien, D. A., Jiang, H., Campbell, A. N., Hu, M. C., Miele, G. M., Cohen, L. R., . . . Nunes, E. V. (2010). Do treatment improvements in PTSD severity affect substance use outcomes? A secondary analysis from a randomized clinical trial in NIDA's Clinical Trials Network. *The American Journal of Psychiatry*, *167*, 95–101. <http://dx.doi.org/10.1176/appi.ajp.2009.09091261>
- Johnson, K. A., Bonn-Miller, M. O., Leyro, T. M., & Zvolensky, M. J. (2009). Anxious arousal and anhedonic depression symptoms and the frequency of current marijuana use: Testing the mediating role of marijuana-use coping motives among active users. *Journal of Studies on Alcohol and Drugs*, *70*, 543–550. <http://dx.doi.org/10.15288/jsad.2009.70.543>
- Johnson, K., Mullin, J. L., Marshall, E. C., Bonn-Miller, M. O., & Zvolensky, M. (2010). Exploring the mediational role of coping motives for marijuana use in terms of the relation between anxiety sensitivity and marijuana dependence. *The American Journal on Addictions*, *19*, 277–282. <http://dx.doi.org/10.1111/j.1521-0391.2010.00041.x>
- Kessler, R. C., Chiu, W. T., Demler, O., Merikangas, K. R., & Walters, E. E. (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, *62*, 617–627. <http://dx.doi.org/10.1001/archpsyc.62.6.617>
- Kevorkian, S., Bonn-Miller, M., Belendiuk, K., Carney, D., Roberson-Nay, R., & Berenz, E. (2015). Associations among trauma, posttraumatic stress disorder, cannabis use, and cannabis use disorder in a nationally

- representative epidemiologic sample. *Psychology of Addictive Behaviors*, 29, 633–638. <http://dx.doi.org/10.1037/ad0000110>
- Korem, N., Zer-Aviv, T. M., Ganon-Elazar, E., Abush, H., & Akirav, I. (2016). Targeting the endocannabinoid system to treat anxiety-related disorders. *Journal of Basic and Clinical Physiology and Pharmacology*, 27, 193–202. <http://dx.doi.org/10.1515/jbcpp-2015-0058>
- Lamarche, L. J., & De Koninck, J. (2007). Sleep disturbance in adults with posttraumatic stress disorder: A review. *The Journal of Clinical Psychiatry*, 68, 1257–1270. <http://dx.doi.org/10.4088/JCP.v68n0813>
- Lavie, P., Katz, N., Pillar, G., & Zinger, Y. (1998). Elevated awaking thresholds during sleep: Characteristics of chronic war-related posttraumatic stress disorder patients. *Biological Psychiatry*, 44, 1060–1065.
- Lee, C. M., Neighbors, C., Hendershot, C. S., & Grossbard, J. R. (2009). Development and preliminary validation of a comprehensive Marijuana Motives Questionnaire. *Journal of Studies on Alcohol and Drugs*, 70, 279–287. <http://dx.doi.org/10.15288/jsad.2009.70.279>
- MacKinnon, D. P. (2008). *Introduction to statistical mediation analysis*. New York, NY: Erlbaum and Taylor Francis Group.
- Metrik, J., Kahler, C. W., McGeary, J. E., Monti, P. M., & Rohsenow, D. J. (2011). Acute effects of marijuana smoking on negative and positive affect. *Journal of Cognitive Psychotherapy*, 25, 31–46. <http://dx.doi.org/10.1891/0889-8391.25.1.31>
- Mitchell, H., Zvolensky, M. J., Marshall, E. C., Bonn-Miller, M. O., & Vujanovic, A. A. (2007). Incremental validity of coping-oriented marijuana use motives in the prediction of affect-based psychological vulnerability. *Journal of Psychopathology and Behavioral Assessment*, 29, 277–288. <http://dx.doi.org/10.1007/s10862-007-9047-z>
- Moitra, E., Christopher, P. P., Anderson, B. J., & Stein, M. D. (2015). Coping-motivated marijuana use correlates with DSM-5 cannabis use disorder and psychological distress among emerging adults. *Psychology of Addictive Behaviors*, 29, 627–632. <http://dx.doi.org/10.1037/ad0000083>
- Morin, C. M., Bootzin, R. R., Buysse, D. J., Edinger, J. D., Espie, C. A., & Lichstein, K. L. (2006). Psychological and behavioral treatment of insomnia: Update of the recent evidence (1998–2004). *Sleep*, 29, 1398–1414.
- Muthén, L. K., & Muthén, B. O. (1998–2012). *Mplus user's guide* (7th ed.). Los Angeles, CA: Author.
- Ouimette, P., Finney, J. W., & Moos, R. H. (1999). Two-year posttreatment functioning and coping of substance abuse patients with posttraumatic stress disorder. *Psychology of Addictive Behaviors*, 13, 105–114. <http://dx.doi.org/10.1037/0893-164X.13.2.105>
- Perlis, M. L., Jungquist, C., Smith, M. T., & Posner, D. (2006). *Cognitive behavioral treatment of insomnia: A session-by-session guide* (Vol. 1). New York, NY: Springer Science & Business Media.
- Peters, E. N., Nich, C., & Carroll, K. M. (2011). Primary outcomes in two randomized controlled trials of treatments for cannabis use disorders. *Drug and Alcohol Dependence*, 118, 408–416. <http://dx.doi.org/10.1016/j.drugalcdep.2011.04.021>
- Piomelli, D., Haney, M., Budney, A. J., & Piazza, P. V. (2016). Roundtable discussion: Legal or illegal, cannabis is still addictive. *Cannabis and Cannabinoid Research*, 1, 47–53. <http://dx.doi.org/10.1089/can.2015.29004.rtd>
- Potter, C. M., Vujanovic, A. A., Marshall-Berenz, E. C., Bernstein, A., & Bonn-Miller, M. O. (2011). Posttraumatic stress and marijuana use coping motives: The mediating role of distress tolerance. *Journal of Anxiety Disorders*, 25, 437–443. <http://dx.doi.org/10.1016/j.janxdis.2010.11.007>
- Saladin, M. E., Brady, K. T., Dansky, B. S., & Kilpatrick, D. G. (1995). Understanding comorbidity between PTSD and substance use disorders: Two preliminary investigations. *Addictive Behaviors*, 20, 643–655. [http://dx.doi.org/10.1016/0306-4603\(95\)00024-7](http://dx.doi.org/10.1016/0306-4603(95)00024-7)
- Seal, K. H., Bertenthal, D., Maguen, S., Gima, K., Chu, A., & Marmar, C. R. (2008). Getting beyond “Don’t ask; don’t tell”: An evaluation of U.S. Veterans Administration postdeployment mental health screening of veterans returning from Iraq and Afghanistan. *American Journal of Public Health*, 98, 714–720. <http://dx.doi.org/10.2105/AJPH.2007.115519>
- Seal, K. H., Bertenthal, D., Miner, C. R., Sen, S., & Marmar, C. (2007). Bringing the war back home: Mental health disorders among 103,788 US veterans returning from Iraq and Afghanistan seen at Department of Veterans Affairs facilities. *Archives of Internal Medicine*, 167, 476–482. <http://dx.doi.org/10.1001/archinte.167.5.476>
- Seal, K. H., Cohen, G., Waldrop, A., Cohen, B. E., Maguen, S., & Ren, L. (2011). Substance use disorders in Iraq and Afghanistan veterans in VA healthcare, 2001–2010: Implications for screening, diagnosis and treatment. *Drug and Alcohol Dependence*, 116, 93–101. <http://dx.doi.org/10.1016/j.drugalcdep.2010.11.027>
- Seal, K. H., Metzler, T. J., Gima, K. S., Bertenthal, D., Maguen, S., & Marmar, C. R. (2009). Trends and risk factors for mental health diagnoses among Iraq and Afghanistan veterans using Department of Veterans Affairs health care, 2002–2008. *American Journal of Public Health*, 99, 1651–1658. <http://dx.doi.org/10.2105/AJPH.2008.150284>
- Simons, J. S., Gaher, R. M., Correia, C. J., Hansen, C. L., & Christopher, M. S. (2005). An affective-motivational model of marijuana and alcohol problems among college students. *Psychology of Addictive Behaviors*, 19, 326–334. <http://dx.doi.org/10.1037/0893-164X.19.3.326>
- Sobell, L. C., & Sobell, M. B. (1992). Timeline followback: A technique for assessing self-reported ethanol consumption. In R. Z. Litten & J. P. Allen (Eds.), *Measuring alcohol consumption: Psychosocial and biological methods* (pp. 41–72). Totowa, NJ: Humana Press.
- Stephens, R. S., Roffman, R. A., & Curtin, L. (2000). Comparison of extended versus brief treatments for marijuana use. *Journal of Consulting and Clinical Psychology*, 68, 898–908. <http://dx.doi.org/10.1037/0022-006X.68.5.898>
- Stewart, S. H., Pihl, R. O., Conrod, P. J., & Dongier, M. (1998). Functional associations among trauma, PTSD, and substance-related disorders. *Addictive Behaviors*, 23, 797–812. [http://dx.doi.org/10.1016/S0306-4603\(98\)00070-7](http://dx.doi.org/10.1016/S0306-4603(98)00070-7)
- Vandrey, R., Babson, K. A., Herrmann, E. S., & Bonn-Miller, M. O. (2014). Interactions between disordered sleep, post-traumatic stress disorder, and substance use disorders. *International Review of Psychiatry*, 26, 237–247. <http://dx.doi.org/10.3109/09540261.2014.901300>
- Watkins, K. E., Burnam, A., Kung, F. Y., & Paddock, S. (2001). A national survey of care for persons with co-occurring mental and substance use disorders. *Psychiatric Services*, 52, 1062–1068. <http://dx.doi.org/10.1176/appi.ps.52.8.1062>
- Watson, D., O’Hara, M. W., Simms, L. J., Kotov, R., Chmielewski, M., McDade-Montez, E. A., . . . Stuart, S. (2007). Development and validation of the Inventory of Depression and Anxiety Symptoms (IDAS). *Psychological Assessment*, 19, 253–268. <http://dx.doi.org/10.1037/1040-3590.19.3.253>
- Weathers, F. W., Keane, T. M., & Davidson, J. R. (2001). Clinician-administered PTSD Scale: A review of the first ten years of research. *Depression and Anxiety*, 13, 132–156. <http://dx.doi.org/10.1002/da.1029>
- Woodward, S. H., Arsenault, N. J., Murray, C., & Bliwise, D. L. (2000). Laboratory sleep correlates of nightmare complaint in PTSD inpatients. *Biological Psychiatry*, 48, 1081–1087. [http://dx.doi.org/10.1016/S0006-3223\(00\)00917-3](http://dx.doi.org/10.1016/S0006-3223(00)00917-3)
- Zvolensky, M. J., Marshall, E. C., Johnson, K., Hogan, J., Bernstein, A., & Bonn-Miller, M. O. (2009). Relations between anxiety sensitivity, distress tolerance, and fear reactivity to bodily sensations to coping and conformity marijuana use motives among young adult marijuana users. *Experimental and Clinical Psychopharmacology*, 17, 31–42. <http://dx.doi.org/10.1037/a0014961>

Received March 15, 2016

Revision received July 18, 2016

Accepted July 22, 2016 ■