

UCSF

UC San Francisco Previously Published Works

Title

Mental and Physical Health Needs of Lesbian, Gay, and Bisexual Clients in Substance Abuse Treatment

Permalink

<https://escholarship.org/uc/item/3k4657nf>

Authors

Flentje, Annesa

Livingston, Nicholas A

Roley, Jason

et al.

Publication Date

2015-11-01

DOI

10.1016/j.jsat.2015.06.022

Peer reviewed



Mental and Physical Health Needs of Lesbian, Gay, and Bisexual Clients in Substance Abuse Treatment



Annesa Flentje, Ph.D.^{a,*}, Nicholas A. Livingston, M.A.^b, Jason Roley, B.A.^{c,d}, James L. Sorensen, Ph.D.^c

^a Department of Community Health Systems, University of California, San Francisco, 2 Koret Way, San Francisco, California, 94143

^b Department of Psychology, The University of Montana, 32 Campus Drive Missoula, Montana, 59812

^c Department of Psychiatry, University of California, San Francisco, 1001 Potrero Avenue, San Francisco, California, 94110

^d School of Medicine, Drexel University, 3300 Henry Avenue, Philadelphia, Pennsylvania, 19129

ARTICLE INFO

Article history:

Received 2 March 2015

Received in revised form 28 May 2015

Accepted 29 June 2015

Keywords:

Lesbian, gay, bisexual
Substance abuse treatment
Physical health disparities
Mental health disparities

ABSTRACT

Objective: Lesbian, gay, and bisexual (LGB) orientation predicts greater substance use, treatment utilization, and poorer mental and physical health, but health needs of LGB individuals in substance abuse treatment remain largely unknown. The purpose of this study was to identify differences in mental and physical health needs of LGB individuals in substance abuse treatment.

Methods: Substance abuse treatment admissions data from the County of San Francisco were used in this investigation of differences in mental and physical health problems and service utilization between LGB ($n = 1,441$) and heterosexual individuals ($n = 11,770$).

Results: LGB individuals were more likely to have mental health diagnoses ($_{adj}ORs$ ranging from 1.86 to 4.00) and current mental health prescription medications ($_{adj}ORs$ from 1.79 to 4.99) than heterosexual counterparts. Gay and bisexual men and bisexual women but not lesbian women, were more likely to be receiving mental health treatment. Gay men and bisexual women were more likely than heterosexual counterparts to report physical health problems. Gay and bisexual men and bisexual women but not lesbian women were more likely to be receiving health care. There were no differences between LGB individuals and heterosexual counterparts in the number of emergency room visits or hospital overnight stays.

Discussion: This study found that LGB individuals entering substance abuse treatment have greater mental and physical health needs than heterosexual counterparts. Implications for healthcare integration, research, and practice are discussed.

© 2015 Elsevier Inc. All rights reserved.

1. Introduction

Alcohol and illicit drug use remain significant public health concerns among the general population (Substance Abuse and Mental Health Services Administration, 2012). Additionally, rates of use and substance use disorder diagnoses (SUDs) among lesbian, gay, and bisexual (LGB) individuals remain disproportionately high relative to heterosexuals (Green & Feinstein, 2012; McCabe, Hughes, Bostwick, West, & Boyd, 2009). As outlined in Meyer's Minority Stress Model (Meyer, 2003), this disparity has been linked to higher rates of minority stress—often stemming from LGB-based institutional oppression and interpersonal mistreatment (e.g., discrimination and victimization)—which has been shown to confer risk for substance use and comorbidity through heightened emotional regulation demands (Weber, 2008). Consistent with elevated rates of use and SUD diagnoses, LGB individuals are also more likely to seek substance abuse treatment compared to heterosexual individuals (McCabe, West, Hughes, & Boyd, 2013). Further, LGB

individuals evidence elevated rates of physical and mental health concerns relative to heterosexual individuals (Dilley, Simmons, Boysun, Pizacani, & Stark, 2010; Fredriksen-Goldsen et al., 2013), consistent with the minority stress hypothesis (Meyer, 2003). Such comorbidity is a salient concern among substance use treatment seekers, as comorbid mental health conditions may precede substance use disorders (Swendsen et al., 2010) and co-occurring health complications may negatively impact substance use treatment outcomes (Grella, Hser, Joshi, & Rounds-Bryant, 2001). Furthermore, treatment for SUD is an important tactic for reducing the both substance use-related deaths and health problems (U.S. Department of Health and Human Services, 2014). At present, it is unclear whether programs and providers are sufficiently aware of the specific mental and physical health needs of LGB individuals seeking substance use treatment, an important precursor to providing appropriate services.

Historically, LGB individuals have also experienced health disparities with regard to receiving adequate physical, mental, and substance use treatment services. For instance, LGB individuals experience greater barriers to securing appropriate medical and mental health services due, in part, to provider or institutional bias (Cochran, Peavy, & Robohm, 2007) and lower rates of adequate health care coverage relative to

* Corresponding author at: Department of Community Health Systems, University of California, San Francisco, 2 Koret Way Room N505, San Francisco, California, 94143.

E-mail address: annesaf.flentje@ucsf.edu (A. Flentje).

heterosexuals (Buchmueller & Carpenter, 2010; Dilley et al., 2010). A potential consequence is that, for many, health care services may be less available, especially in the absence of publicly-funded treatment options. Adequate care is also compromised by the lack of research on the specific physical and health care needs of LGB individuals. Despite reduced access to health care coverage, LGB individuals have been shown to utilize emergency care services at greater rates than their heterosexual counterparts (Sánchez, Hailpern, Lowe, & Calderon, 2007), which is consistent with previously mentioned reports of worse general physical health among LGB individuals relative to heterosexual individuals and greater unmet medical needs.

Special mental and physical health considerations have also been documented among LGB individuals seeking substance use treatment services. For instance, researchers have shown that LGB individuals seeking substance use treatment present with elevated rates of comorbid mental health diagnoses (Lipsky et al., 2012), substance use severity, and past-year medical service utilization (Cochran & Cauce, 2006), relative to heterosexual individuals. Nonetheless, the research literature remains limited, as the aforementioned research (Cochran & Cauce, 2006; Lipsky et al., 2012) took place within the only US State (Washington) that, at the time, asked treatment applicants to identify their sexual orientation and gender identity.

The goal of this study was to examine how mental and physical health needs and treatment utilization of LGB individuals differed compared to heterosexual counterparts among those seeking substance abuse treatment within a publicly-funded system. Based on previous research documenting higher rates of mental and physical health problems among LGB individuals, and the minority stress hypothesis (Meyer, 2003), we hypothesized that LGB individuals would have higher rates of mental and physical health problems and service utilization relative to heterosexual individuals. This study can inform substance abuse treatment implementation and integration of psychiatric and medical health care to improve screening and service delivery.

2. Method

The methods for this study were similar to the methods used by Flentje, Heck, and Sorensen (2015), which utilized the same treatment database and sample for a study examining the primary substance of abuse, route of administration, age of initiation of that substance, and the frequency at which that substance was used among LGB individuals entering substance abuse treatment. The methods and sample description are described briefly here and also in Flentje, Heck, et al. (2015).

De-identified data were obtained from the Department of Public Health in the County of San Francisco, which collected client admission and discharge information from all substance abuse treatment programs in the county that received any government funding. For individuals who sought treatment between July 2007 and December 2009, the treatment record(s) for that timeframe were included as well as all other existing treatment records. This resulted in 107,470 total treatment episodes attended by 14,015 different individuals. A treatment episode was defined as contact with any treatment program which initiated a billing entry into the San Francisco billing information system. These entries were associated with an admission record collected by the treatment agency. As such, treatment episodes could include a range of services including residential, detox, or outpatient services. A single treatment episode was selected for each individual based on the criterion that it was the most recent treatment episode, and information provided at admission was used for this study. Individuals with mental and physical health data were included in this study if they: (1) reported a male or female sex, (2) reported a lesbian, gay, bisexual, or heterosexual sexual orientation, and (3) did not endorse a transgender identity. Information regarding the admission characteristics of transgender individuals within the same substance abuse treatment database was documented in a separate study (Flentje, Heck, & Sorensen, 2014).

2.1. Measures

The questions asked of clients at treatment and discharge were from the California Outcomes Measurement System (CALOMS), a procedure created to monitor substance abuse treatment outcomes within California which has been used in peer-reviewed research (e.g. Brecht & Urada, 2011; Conner, Hampton, Hunter, & Urada, 2011; Evans, Jaffe, Urada, & Anglin, 2012; Flentje, Heck, et al., 2015; Flentje et al., 2014; Gonzales, Brecht, Mooney, & Rawson, 2011; Swartz, 2010). This measurement system was used across programs in California, but localities could also add questions of interest. At the time of data collection, the County of San Francisco had elected to add questions about sexual orientation and gender identity, data that were not collected in other areas. Sexual orientation was queried with the following response options: “Lesbian: Female/Female,” “Gay: Male/Male,” “Bisexual: Both Male & Female,” “Heterosexual,” “Decline to Answer,” and “Unsure.”

Clients entering substance abuse treatment were required to answer multiple questions regarding demographics, substance use, mental health, and physical health. Substance abuse treatment programs provided client admission data to the County of San Francisco where it was compiled. Some questions that were used in this study queried for the time period of 30 days prior to admission, specifically, questions which asked if an individual had: taken prescribed medication for mental health, been in a hospital or psychiatric facility for mental health, experienced physical problems, gone to the emergency room (ER), or stayed in the hospital overnight for a physical health problem. Participants were also asked if they: had a prior mental health diagnosis, were receiving mental health treatment, had a recent mental health assessment, were receiving physical health care, or had a recent physical health assessment. Questions querying recent mental and physical health assessments and whether individuals were receiving mental health treatment were only queried in a specific iteration of the data collection system (which underwent adjustments during the time of data collection for this study), thus only a portion of the sample was queried with these questions.

2.2. Analyses

Analyses were conducted separately by sex. Demographics and demographic differences by sexual orientation are reported elsewhere (Flentje, Heck, et al., 2015). Logistic regression was used to predict outcomes of interest in this study, with gay (for men) or lesbian (for women) and bisexual orientation entered (heterosexual was the reference group), and age, race (dichotomized as White and non-White), and ethnicity (dichotomized as Hispanic or not Hispanic) entered as covariates. To control for type I error, the alpha level was set to .01, and 99% confidence intervals were reported accordingly.

3. Results

3.1. Participants

Participants who did not meet sexual orientation and gender identity inclusion criteria were as follows: transgender ($n = 199$), sexual orientation criterion not met ($n = 210$, with $n = 135$ who declined answering and $n = 75$ who responded “unsure”), sexual orientation did not match with reported sex (e.g., endorsed female sex and gay male sexual orientation, $n = 23$). A total of 13,211 participants met inclusion criteria for the study. Demographic information for participants is reported by sex and sexual orientation in Table 1 (similarly reported in Flentje, Heck, et al., 2015). Participants were an average of 38.10 years old ($SD = 13.48$), with over 90% of participants being age 18 or older ($n = 12,012$, 90.9%). Overall, among males, there were differences by sexual orientation in race and level of education, but not

Table 1
Demographic information by sex and sexual orientation for individuals seeking substance abuse treatment in San Francisco ($N = 13,211$).

	Overall Sample ($N = 13,211$)	Male participants ($n = 9330$)			Female participants ($n = 3881$)		
		Heterosexual ($n = 8318$)	Gay ($n = 797$)	Bisexual ($n = 215$)	Heterosexual ($n = 3452$)	Lesbian ($n = 156$)	Bisexual ($n = 273$)
Age (M, SD)	38.10 (13.48)	39.24 (13.42)	39.81 (10.78)	39.89 (11.19)	35.70 (13.81)	36.12 (11.17)	33.44 (12.16)
Race, n (%)							
White	4,705 (35.6%)	2,779 (33.4%)	535 (67.1%)	137 (63.7%)	1093 (31.7%)	59 (37.8%)	102 (37.4%)
Black	4,844 (36.7%)	3,201 (38.5%)	81 (10.2%)	40 (18.6%)	1385 (40.1%)	56 (35.9%)	81 (29.7%)
Native American/Alaska Native	174 (1.3%)	86 (1.0%)	10 (1.3%)	5 (2.3%)	68 (2.0%)	2 (1.3%)	3 (1.1%)
Asian American/Pacific Islander	738 (5.6%)	512 (6.2%)	24 (3.0%)	4 (1.9%)	178 (5.2%)	11 (7.1%)	9 (3.3%)
Multi-racial	678 (5.1%)	332 (4.0%)	63 (7.9%)	14 (6.5%)	213 (6.2%)	12 (7.7%)	44 (16.1%)
Other race	2071 (15.7%)	1,407 (16.9%)	84 (10.5%)	15 (7.0%)	515 (14.9%)	16 (10.3%)	34 (12.5%)
Ethnicity, n (%)							
Not Hispanic	10,778 (81.6%)	6,756 (81.2%)	666 (83.6%)	194 (90.2%)	2815 (81.5%)	122 (78.2%)	225 (82.4%)
Mexican/Mexican American	1,048 (7.9%)	682 (8.2%)	52 (6.5%)	6 (2.8%)	274 (7.9%)	14 (9.0%)	20 (7.3%)
Cuban	59 (0.4%)	43 (0.5%)	5 (0.6%)	4 (1.9%)	7 (0.2%)	0 (0.0%)	0 (0.0%)
Puerto Rican	185 (1.4%)	106 (1.3%)	8 (1.0%)	2 (0.9%)	56 (1.6%)	5 (3.2%)	8 (2.9%)
Other Hispanic/Latino	1,141 (8.6%)	731 (8.8%)	66 (8.3%)	9 (4.2%)	300 (8.7%)	15 (9.6%)	20 (7.3%)
Education in years (M, SD)	11.92 (2.53)	11.81 (2.41)	14.07 (2.64)	12.78 (2.22)	11.65 (2.51)	12.54 (2.17)	12.21 (2.61)

ethnicity or age. Percentages indicate that gay and bisexual men were more likely to be of White race, and reported higher levels of education. For females there were differences by sexual orientation in level of education, but not race, ethnicity, or age (specific analyses on differences in demographic characteristics by sexual orientation reported in Flentje, Heck, et al. (2015)).

3.2. Mental health problems and service utilization

Mental and physical health problems, care, and service utilization by sex and sexual orientation of participants are reported by sex and sexual orientation in Table 2. LGB status was predictive of higher rates of mental health diagnoses for both men (gay men, adjusted odds ratio [$_{adj}OR$]: 4.00, 99% confidence interval [CI]: 3.23, 4.94; bisexual men, $_{adj}OR$: 3.56, 99% CI: 2.42, 5.23) and women (lesbian women, $_{adj}OR$: 1.86, 99% CI: 1.22, 2.83; bisexual women, $_{adj}OR$: 2.26, 99% CI: 1.59, 3.20). LGB status predicted higher rates of current mental health prescription medication for both men (gay men, $_{adj}OR$: 4.99, 99% CI: 4.04, 6.16; bisexual men, $_{adj}OR$: 2.95, 99% CI: 1.99, 4.37) and women (lesbian women, $_{adj}OR$: 1.87, 99% CI: 1.20, 2.91; bisexual women, $_{adj}OR$: 1.79, 99% CI: 1.24, 2.60). Gay ($_{adj}OR$: 3.38, 99% CI: 2.43, 4.71) and bisexual men ($_{adj}OR$: 2.59, 99% CI: 1.37, 4.90), and bisexual women ($_{adj}OR$: 1.97, 99% CI: 1.13, 3.45) were more likely to be receiving mental health treatment, but there were no differences between lesbian women and heterosexual women regarding the likelihood of receiving mental health treatment. Gay men and bisexual women were more likely than heterosexual men and women, respectively, to have undergone a recent mental health assessment (gay men, $_{adj}OR$: 1.96, 99% CI: 1.42, 2.71; bisexual women, $_{adj}OR$: 2.10, 99% CI: 1.22, 3.61). Lastly,

gay men were more likely to have recently been in a psychiatric hospital or facility ($_{adj}OR$: 2.21, 99% CI: 1.47, 3.35) than heterosexual men, but no differences were observed in the other three groups when compared to heterosexual counterparts. Complete results of logistic regression analyses are reported in Table 3.

3.3. Physical health problems and service utilization

Among men, gay men were more likely than heterosexual men to report physical health problems in the previous 30 days ($_{adj}OR$: 1.42, 99% CI: 1.14, 1.77), but this difference was not evident for bisexual men. Among women, bisexual women were more likely to report physical health problems ($_{adj}OR$: 1.70, 99% CI: 1.18, 2.47), but there was no difference between lesbian and heterosexual women. Gay ($_{adj}OR$: 4.25, 99% CI: 3.00, 6.04) and bisexual men ($_{adj}OR$: 2.61, 99% CI: 1.38, 4.96), and bisexual women ($_{adj}OR$: 1.83, 99% CI: 1.07, 3.13) had greater odds than heterosexual counterparts of receiving health care, but there were no differences observed between lesbian women and heterosexual women. Gay men were more likely to report a recent physical health assessment ($_{adj}OR$: 2.10, 99% CI: 1.49, 2.97), but there were no differences between bisexual men, lesbian women, nor bisexual women when compared to heterosexual counterparts. LGB status was not predictive of ER visits nor hospital stays among males or females.

4. Discussion

Consistent with hypotheses, and Meyer's minority stress hypothesis (2003), our study found that sexual orientation is a predictor of mental and physical health status, and that important mental and physical

Table 2
Physical and mental health problems, risks, care, and service utilization by sex and sexual orientation for individuals seeking treatment in San Francisco ($N = 13,211$).

	Male participants			Female participants		
	Heterosexual	Gay	Bisexual	Heterosexual	Lesbian	Bisexual
Mental health problems and service utilization						
Prior mental health diagnosis	2153 (27.0%)	499 (64.6%)	127 (61.4%)	1229 (38.1%)	75 (50.7%)	142 (55.9%)
Taken prescribed medication for mental health in past 30 days	1108 (13.7%)	380 (48.7%)	74 (35.7%)	743 (22.2%)	50 (33.3%)	84 (31.5%)
Receiving mental health treatment ^a	576 (19.1%)	147 (48.4%)	30 (41.1%)	392 (27.4%)	24 (35.8%)	41 (39.0%)
Recent mental health assessment ^a	965 (31.9%)	155 (51.0%)	36 (49.3%)	452 (31.6%)	25 (37.3%)	48 (45.7%)
Psychiatric hospital or facility	236 (2.9%)	55 (7.1%)	11 (5.3%)	120 (3.6%)	8 (5.3%)	12 (4.5%)
Physical health problems and service utilization						
Physical health problems in previous 30 days	1761 (21.7%)	253 (32.4%)	65 (31.4%)	785 (23.5%)	39 (26.0%)	83 (31.1%)
Receiving physical health care	1037 (34.3%)	211 (69.4%)	42 (57.5%)	608 (42.5%)	31 (46.3%)	57 (54.3%)
Recent physical health assessment ^a	1517 (50.2%)	212 (69.7%)	44 (60.3%)	765 (53.5%)	34 (50.7%)	58 (55.2%)
ER visits in previous 30 days	863 (10.6%)	85 (10.9%)	30 (14.5%)	352 (10.5%)	15 (10.0%)	37 (13.9%)
Hospital overnight stays	433 (5.3%)	47 (6.0%)	16 (7.7%)	179 (5.4%)	5 (3.3%)	14 (5.2%)

^a For these descriptives, only a subset of participants were available ($n = 5000$) due to changes in data collection practices.

Table 3

Results of logistic regression analyses using sexual orientation to predict mental and physical health problems and service utilization (separate analyses by sex) among individuals seeking substance abuse treatment in San Francisco ($N = 13,211$), adjusted for age, race, and ethnicity with heterosexual as reference ($N = 13,211$).

	Male participants		Female participants	
	Gay	Bisexual	Lesbian	Bisexual
	adjOR (99% CI)	adjOR (99% CI)	adjOR (99% CI)	adjOR (99% CI)
Mental health problems and service utilization				
Prior mental health diagnosis	4.00 (3.23, 4.94)*	3.56 (2.42, 5.23)*	1.86 (1.22, 2.83)*	2.26 (1.59, 3.20)*
Taken prescribed medication for mental health in past 30 days	4.99 (4.04, 6.16)*	2.95 (1.99, 4.37)*	1.87 (1.20, 2.91)*	1.79 (1.24, 2.60)*
Receiving mental health treatment ^a	3.38 (2.43, 4.71)*	2.59 (1.37, 4.90)*	1.64 (0.86, 3.13)	1.97 (1.13, 3.45)*
Recent mental health assessment ^a	1.96 (1.42, 2.71)*	1.82 (0.97, 3.39)	1.42 (0.75, 2.67)	2.10 (1.22, 3.61)*
Psychiatric hospital or facility	2.21 (1.47, 3.35)*	1.66 (0.73, 3.79)	1.67 (0.69, 4.00)	1.36 (0.61, 3.04)
Physical health problems and service utilization				
Physical health problems in previous 30 days	1.42 (1.14, 1.77)*	1.41 (0.94, 2.11)	1.24 (0.78, 1.99)	1.70 (1.18, 2.47)*
Receiving health care	4.25 (3.00, 6.04)*	2.61 (1.38, 4.96)*	1.22 (0.66, 2.27)	1.83 (1.07, 3.13)*
Recent physical health assessment ^a	2.10 (1.49, 2.97)*	1.40 (0.74, 2.66)	0.90 (0.48, 1.66)	1.19 (0.70, 2.04)
ER visits in previous 30 days	0.81 (0.59, 1.11)	1.18 (0.69, 1.99)	1.18 (0.63, 2.20)	1.52 (0.93, 2.47)
Hospital overnight stays	0.92 (0.61, 1.40)	1.29 (0.64, 2.57)	0.76 (0.27, 2.10)	1.11 (0.53, 2.33)

* Indicates an analysis where $p < .01$.

^a For these analyses, only a subset of participants were available ($n = 3397$ for males and $n = 1612$ for females) due to changes in data collection practices.

health status disparities exist among LGB substance abuse treatment-seeking individuals. A consistent trend across LGB individuals was that there were higher rates of previous mental health diagnoses. Similarly, all LGB groups were more likely to be taking psychiatric medications. Also, gay and bisexual men and bisexual women were more likely to be receiving mental health treatment. This suggests that these groups are entering treatment with a need for continuity of care for co-occurring disorders.

Within our study, lesbian and bisexual women had about 2 times greater odds of previous mental health diagnoses than heterosexual women, while gay and bisexual men had around 3.5–4 times greater odds, when compared to heterosexual men. Of note is the observed difference in rates of mental health diagnoses between the heterosexual comparison groups (38.1% for heterosexual women, and 27.0% for heterosexual men), which may reflect greater rates of psychiatric complications associated with substance use for women than men, consistent with other studies (Denier, Thevos, Latham, & Randall, 1991; Hernandez-Avila, Rounsaville, & Kranzler, 2004). Even given the higher base rate for women, in our study we still detected greater odds of mental health diagnoses and mental health medication use for lesbian and bisexual women, and mental health treatment for bisexual women. The higher base rates of co-occurring mental health disorders observed among heterosexual women seeking treatment likely was reflected in the overall smaller observed odds ratios for lesbian and bisexual women (relative to gay and bisexual men compared to heterosexual men). More than half of all lesbian and bisexual women entered treatment with a mental health diagnosis, highlighting the importance of consideration of co-occurring mental health need within this population. Similarly, the overall lower rates of mental health service utilization among heterosexual men likely attributed to the large odds ratios observed among gay men (ranging from 3.38 for mental health treatment utilization to 4.99 for mental health medication usage). The absolute percentages among gay and bisexual men were also notable, as nearly two-thirds of gay and bisexual men entered treatment with a mental health diagnosis. The need for treatment addressing both substance use and co-occurring disorders is therefore extremely relevant for one-half to two-thirds of LGB people seeking substance abuse treatment services and is an imperative and not a complementary service among this population.

The null effects for mental health treatment utilization among lesbian women could be attributed to unmet mental health treatment need among lesbian clients. This may be consistent with prior research documenting compounded stress for sexual minority women (e.g., contending with both sexism and heterosexism; see Drabble & Eliason, 2012), but inconsistent with reports of greater utilization among sexual minority women in general (Drabble & Eliason, 2012).

That being said, this effect is most likely attributable to insufficient power to detect meaningful effects due to a relatively small number of lesbian women (our smallest observed LGB group) who reported recent mental health treatment or assessment, thus these results should be interpreted with caution.

Gay men and bisexual women were at greater risk for experiencing recent physical health problems compared to heterosexual individuals. Gay and bisexual men and bisexual women were more likely to be receiving health care, but only gay men were more likely to have had a recent physical health assessment. This suggests that health care continuity needs are an area for future investigation for this treatment population.

In contrast to previous research (Cochran & Cauce, 2006; Sánchez et al., 2007), LGB individuals seeking substance abuse treatment did not report higher rates of recent ER visits or hospital overnight stays. Whereas Sánchez et al. (2007) analyzed data from a convenience sample of LGBT individuals in New York City in 2004, Cochran and Cauce's (2006) results were based on data from individuals who received publicly-funded substance use treatment in the state of Washington between July 1, 2001 and December 31, 2002. The discrepancy between the results presented here and those presented elsewhere may be attributable to differing operationalization of emergency room utilization across studies (i.e., self-reported ER visits in last 12 months versus 30 days in the current study), low response variability given our retrospective 30-day measurement window, or actual regional differences with regard to ER visits across studies. For instance, the null effects for ER visits reported here could reflect that substance use treatment-seeking individuals in San Francisco, irrespective of sexual orientation, are at higher risk for ER visits (which ranged from 10 to 14.5% of participants across all sexual orientations) and hospital overnight stays (which ranged from 3.3% to 7.7% of participants across sexual orientations) overall. It could also reflect that within San Francisco it may be easier for LGB individuals to access other healthcare services, thus circumventing the need for emergency department use to fill the need of regular access to healthcare.

We found higher mental health treatment utilization among all LGB groups except for lesbian women. This is similar to data published from the population-based California Quality of Life Survey (Grella, Greenwell, Mays, & Cochran, 2009) which found higher mental health treatment utilization among lesbian and bisexual women. It is unknown if access to mental health care is easier for LGB individuals within California, and these results should be replicated in other locations. Research has shown that lesbian and bisexual women who disclose their sexual minority status report greater satisfaction with their providers (Mosack, Brouwer, & Petroll, 2013) and are more likely utilize health care services (Bergeron & Senn, 2003) compared to those for whom sexual identity remains undisclosed. Decisions to disclose are linked to openness and

individuals' comfort level with their providers (Bergeron & Senn, 2003; Polek, Hardie, & Crowley, 2008). It is possible that within San Francisco, an area with a track record for moving forward the gay rights movement (Armstrong, 2002), disclosure and barriers to access to care are removed or greatly reduced for LGB individuals.

4.1. Implications for care

These observed mental and physical health disparities suggest that additional screening, outreach, provider training, and service delivery integration are needed to best coordinate care among members of this at-risk population. It also may suggest that substance abuse treatment settings are well suited for linkage to psychiatric and primary care services. Our finding that sexual minority status predicts important mental and physical health problems suggests that providers should query about client sexual minority status (in a way that facilitates healthy and supporting provider–client dialog), and perform careful screening of mental and physical health care needs among LGB individuals seeking services. Efforts to improve screening and outreach to LGB individuals would be further improved to the degree that providers are trained to meet the specific needs of LGB individuals seeking services.

Another suggestion includes the integration and coordination of care for LGB clients seeking substance use treatment. The benefits of integrating mental and physical health care into substance use treatment (Drainoni et al., 2014; Grella & Stein, 2006; Hart, Tulloch, & O'Cleirigh, 2014; Hides et al., 2010; Zaller, Gillani, & Rich, 2007) as well as LGB-specific interventions into existing substance use treatment settings (Reback, Veniegas, & Shoptaw, 2014; Senreich, 2010; Shoptaw et al., 2008), have been recognized. Our study also indicates that LGB clients are more likely to enter treatment on psychiatric medications, and therefore may benefit from integrated psychiatric care. The dearth of research on fully integrated and/or coordinated LGB-specific services necessitates further inquiry. Thus far, research has shown integrated services to be an efficacious alternative among difficult to reach clinical populations with unmet treatment needs (Drainoni et al., 2014). Expected advantages of health care integration to LGB clients include not having to repeatedly disclose sexual orientation to new providers, coordination of care, concurrent treatment of comorbid mental and physical health conditions that might otherwise complicate substance use treatment, greater support in terms of facilitating client treatment adherence and follow through, and increased accountability regarding health care service delivery. These suspected benefits represent plausible and testable hypotheses for future public health promotion and intervention research. In the meantime, it is incumbent on providers to consider ways in which they may enhance continuity of care for LGB individuals seeking substance abuse treatment.

Lastly, this study supports the importance of collecting sexual orientation data in electronic health records and research, as sexual orientation can be a predictor of disparities in research and in health care settings. Despite this need, sexual orientation is often not queried or reported in research literature (Flentje, Bacca, & Cochran, 2015). Querying about sexual orientation and including it in electronic health records may facilitate future LGB health disparity research, which is needed to more fully understand the specific needs of LGB clients in substance use treatment and in other settings. Further, having this information available in electronic health records may promote open dialog between clients and ongoing/future providers, inform provider case conceptualizations, and improve therapeutic and medical referrals and recommendations.

4.2. Limitations

Data for the current study were collected from an urban area of San Francisco, potentially limiting the degree to which these results generalize broadly. Although minority stress hypotheses were largely supported, these effects might be less pronounced in this region, relative

to more socially or politically conservative regions of the United States. Thus, replication is needed. The self-report nature of these data represents another potential limitation. Another consideration is that these data may only generalize to LGB substance use treatment-seeking individuals rather than LGB individuals in general. Furthermore, as this study was done with data collected for other purposes, sexual orientation categories were predetermined and limited to LGB individuals, thus did not allow for individuals to report other sexual orientations (e.g., queer). Similarly, because data were released by a county health system, some variables which may have differed by sexual orientation were not available, including the level of care that participants were seeking and received while in treatment. Regardless, this population represents an important and at-risk demographic, for which there remains a need for additional research regarding best-practices in substance use treatment settings.

4.3. Conclusions

This study is among the first to document the mental and physical health care needs of LGB individuals seeking substance use treatment, and it is the first in recent years to document ongoing mental and physical health disparities among LGB individuals seeking substance use treatment. Consistent with hypotheses, the effects reported here suggest that substance abuse treatment-seeking LGB individuals are at elevated risk for mental and physical health problems compared to heterosexuals. Additional research is needed to address how to best meet mental and physical health care needs among substance use treatment-seeking LGB individuals.

Acknowledgements

We would like to acknowledge that our work on this study was completed with support from the National Institute on Drug Abuse under award numbers T32DA007250, P50DA09253, and U10DA015815. The authors would also like to thank the San Francisco Department of Public Health, Community Behavioral Health Service for making this research possible (in particular Tom Bleecker and Alice Gleghorn in helping us to obtain these data). The content is solely the responsibility of the authors, and the views expressed herein do not necessarily reflect the official policies or views of the City and County of San Francisco or the National Institutes of Health; nor does mention of the San Francisco Department of Public Health or the National Institutes of Health imply its endorsement.

References

- Armstrong, E. A. (2002). *Forging gay identities: Organizing sexuality in San Francisco, 1950–1994*. University of Chicago Press.
- Bergeron, S., & Senn, C. Y. (2003). Health care utilization in a sample of Canadian lesbian women: Predictors of risk and resilience. *Women & Health, 37*(3), 19–35.
- Brecht, M. L., & Urada, D. (2011). Treatment outcomes for methamphetamine users: California Proposition 36 and comparison clients. *Journal of Psychoactive Drugs, 43*(Suppl. 1), 68–76.
- Buchmueller, T., & Carpenter, C. S. (2010). Disparities in health insurance coverage, access, and outcomes for individuals in same-sex versus different-sex relationships, 2000–2007. *American Journal of Public Health, 100*(3), 489–495.
- Cochran, B. N., & Cauce, A. M. (2006). Characteristics of lesbian, gay, bisexual, and transgender individuals entering substance abuse treatment. *Journal of Substance Abuse Treatment, 30*(2), 135–146.
- Cochran, B. N., Peavy, K. M., & Robohm, J. S. (2007). Do specialized services exist for LGBT individuals seeking treatment for substance misuse? A study of available treatment programs. *Substance Use & Misuse, 42*(1), 161–176.
- Conner, B. T., Hampton, A. S., Hunter, J., & Urada, D. (2011). Treating opioid use under California's Proposition 36: Differential outcomes by treatment modality. *Journal of Psychoactive Drugs, 43*(Suppl. 1), 77–83.
- Denier, C. A., Thevos, A. K., Latham, P. K., & Randall, C. L. (1991). Psychosocial and psychopathology differences in hospitalized male and female cocaine abusers: A retrospective chart review. *Addictive Behaviors, 16*(6), 489–496.
- Dille, J. A., Simmons, K. W., Boysun, M. J., Pizacani, B. A., & Stark, M. J. (2010). Demonstrating the importance and feasibility of including sexual orientation in public health surveys: Health disparities in the Pacific Northwest. *American Journal of Public Health, 100*(3), 460–467.

- Drabble, L., & Eliason, M. J. (2012). Substance use disorders treatment for sexual minority women. *Journal of LGBT Issues in Counseling*, 6(4), 274–292. <http://dx.doi.org/10.1080/15538605.2012.726150>.
- Drainoni, M. L., Farrell, C., Sorensen-Alawad, A., Palmisano, J. N., Chaisson, C., & Walley, A. Y. (2014). Patient perspectives of an integrated program of medical care and substance use treatment. *AIDS Patient Care and STDs*, 28(2), 71–81.
- Evans, E., Jaffe, A., Urada, D., & Anglin, M. D. (2012). Differential outcomes of court-supervised substance abuse treatment among California parolees and probationers. *International Journal of Offender Therapy and Comparative Criminology*, 56(4), 539–556.
- Flentje, A., Bacca, C. L., & Cochran, B. N. (2015). Missing data in substance abuse research? Researchers' reporting practices of sexual orientation and gender identity. *Drug and Alcohol Dependence*, 147, 280–284.
- Flentje, A., Heck, N. C., & Sorensen, J. L. (2014). Characteristics of transgender individuals entering substance abuse treatment. *Addictive Behaviors*, 39(5), 969–975.
- Flentje, A., Heck, N. C., & Sorensen, J. L. (2015). Substance use among lesbian, gay, and bisexual clients entering substance abuse treatment: Comparisons to heterosexual clients. *Journal of Consulting and Clinical Psychology*. <http://dx.doi.org/10.1037/a0038724>.
- Fredriksen-Goldsen, K. I., Emlert, C. A., Kim, H. J., Muraco, A., Erosheva, E. A., Goldsen, J., et al. (2013). The physical and mental health of lesbian, gay male, and bisexual (LGB) older adults: The role of key health indicators and risk and protective factors. *The Gerontologist*, 53(4), 664–675.
- Gonzales, R., Brecht, M. L., Mooney, L., & Rawson, R. A. (2011). Prescription and over-the-counter drug treatment admissions to the California public treatment system. *Journal of Substance Abuse Treatment*, 40(3), 224–229.
- Green, K. E., & Feinstein, B. A. (2012). Substance use in lesbian, gay, and bisexual populations: An update on empirical research and implications for treatment. *Psychology of Addictive Behaviors*, 26(2), 265–278.
- Grella, C. E., Greenwell, L., Mays, V. M., & Cochran, S. D. (2009). Influence of gender, sexual orientation, and need on treatment utilization for substance use and mental disorders: Findings from the California Quality of Life Survey. *BMC Psychiatry*, 9(1), 52.
- Grella, C. E., Hser, Y. I., Joshi, V., & Rounds-Bryant, J. (2001). Drug treatment outcomes for adolescents with comorbid mental and substance use disorders. *The Journal of Nervous and Mental Disease*, 189(6), 384–392.
- Grella, C. E., & Stein, J. A. (2006). Impact of program services on treatment outcomes of patients with comorbid mental and substance use disorders. *Psychiatric Services*, 57(7), 1007–1015.
- Hart, T. A., Tulloch, T. G., & O'Cleirigh, C. (2014). Integrated cognitive behavioral therapy for social anxiety and HIV prevention for gay and bisexual men. *Cognitive and Behavioral Practice*, 21(2), 149–160.
- Hernandez-Avila, C. A., Rounsaville, B. J., & Kranzler, H. R. (2004). Opioid-, cannabis- and alcohol-dependent women show more rapid progression to substance abuse treatment. *Drug and Alcohol Dependence*, 74(3), 265–272.
- Hides, L., Carroll, S., Catania, L., Cotton, S. M., Baker, A., Scaffidi, A., et al. (2010). Outcomes of an integrated cognitive behaviour therapy (CBT) treatment program for co-occurring depression and substance misuse in young people. *Journal of Affective Disorders*, 121(1), 169–174.
- Lipsky, S., Krupski, A., Roy-Byrne, P., Huber, A., Lucenko, B. A., & Mancuso, D. (2012). Impact of sexual orientation and co-occurring disorders on chemical dependency treatment outcomes. *Journal of Studies on Alcohol and Drugs*, 73(3), 401–412.
- McCabe, S. E., Hughes, T. L., Bostwick, W. B., West, B. T., & Boyd, C. J. (2009). Sexual orientation, substance use behaviors and substance dependence in the United States. *Addiction*, 104(8), 1333–1345.
- McCabe, S. E., West, B. T., Hughes, T. L., & Boyd, C. J. (2013). Sexual orientation and substance abuse treatment utilization in the United States: Results from a national survey. *Journal of Substance Abuse Treatment*, 44(1), 4–12.
- Meyer, I. H. (2003). Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: Conceptual issues and research evidence. *Psychological Bulletin*, 129(5), 674–697.
- Mosack, K. E., Brouwer, A. M., & Petroll, A. E. (2013). Sexual identity, identity disclosure, and health care experiences: Is there evidence for differential homophobia in primary care practice? *Women's Health Issues*, 23(6), e341–e346.
- Polek, C. A., Hardie, T. L., & Crowley, E. M. (2008). Lesbians' disclosure of sexual orientation and satisfaction with care. *Journal of Transcultural Nursing*, 19(3), 243–249.
- Reback, C. J., Veniegas, R., & Shoptaw, S. (2014). Getting off: development of a model program for gay and bisexual male methamphetamine users. *Journal of Homosexuality*, 61(4), 540–553.
- Sánchez, J. P., Hailpern, S., Lowe, C., & Calderon, Y. (2007). Factors associated with emergency department utilization by urban lesbian, gay, and bisexual individuals. *Journal of Community Health*, 32(2), 149–156.
- Senreich, E. (2010). Are specialized LGBT program components helpful for gay and bisexual men in substance abuse treatment? *Substance Use & Misuse*, 45(7–8), 1077–1096.
- Shoptaw, S., Reback, C. J., Larkins, S., Wang, P. C., Rotheram-Fuller, E., Dang, J., et al. (2008). Outcomes using two tailored behavioral treatments for substance abuse in urban gay and bisexual men. *Journal of Substance Abuse Treatment*, 35(3), 285–293.
- Substance Abuse and Mental Health Services Administration (2012). State estimates of substance use and mental disorders from the 2009–2010. *National Surveys on Drug Use and Health, NSDUH Series H-43, HHS Publication No. (SMA) 12-4703* (Retrieved from: <http://samhsa.gov/data/NSDUH/2k10State/NSDUHsae2010/FullReport/NSDUHsaeMainReport2010.htm>).
- Swartz, R. (2010). Medical marijuana users in substance abuse treatment. *Harm Reduction Journal*, 7(3).
- Swendsen, J., Conway, K. P., Degenhardt, L., Glantz, M., Jin, R., Merikangas, K. R., et al. (2010). Mental disorders as risk factors for substance use, abuse and dependence: Results from the 10-year follow-up of the National Comorbidity Survey. *Addiction*, 105(6), 1117–1128.
- U.S. Department of Health and Human Services (2014). Healthy people 2020, substance abuse. Retrieved from [HealthyPeople.gov https://www.healthypeople.gov/2020/topics-objectives/topic/substance-abuse](https://www.healthypeople.gov/2020/topics-objectives/topic/substance-abuse).
- Weber, G. N. (2008). Using to numb the pain: Substance use and abuse among lesbian, gay and bisexual individuals. *Journal of Mental Health Counseling*, 30(1), 31–48. <http://dx.doi.org/10.1521/jscp.2009.28.5.597>.
- Zaller, N., Gillani, F. S., & Rich, J. D. (2007). A model of integrated primary care for HIV-positive patients with underlying substance use and mental illness. *AIDS Care*, 19(9), 1128–1133.