

UCSF

UC San Francisco Previously Published Works

Title

Cardiac magnetic resonance in the diagnosis of the unusually detected acute myocarditis in the young people: a case report.

Permalink

<https://escholarship.org/uc/item/3kb0p94f>

Authors

de la Guía-Galipienso, Fernando
García-González, Pilar
Fabregat-Andrés, Oscar
et al.

Publication Date

2021

DOI

10.21037/acr-21-24

Peer reviewed



Cardiac magnetic resonance in the diagnosis of the unusually detected acute myocarditis in the young people: a case report

Fernando de la Guía-Galipienso^{1,2,3}, Pilar García-González⁴, Oscar Fabregat-Andrés⁵, Aurelio Quesada-Dorador^{6,7}, Christoph Meyer-Josten^{1,2}, Carl J. Lavie⁸, Daniel P. Morin⁸, Fabian Sanchis-Gomar^{9^}

¹Glorieta Policlínica, Denia, Alicante, Spain; ²REMA Sports Cardiology Clinic, Denia, Alicante, Spain; ³Cardiology Service, Hospital Clínica Benidorm, Alicante, Spain; ⁴Cardiovascular Unit ASCIRES, Valencia, Spain; ⁵Cardiology Service, Hospital IMED Valencia, Valencia, Spain; ⁶Arrhythmia Unit, Cardiology Service, General University Hospital Consortium of Valencia, Valencia, Spain; ⁷School of Medicine, Catholic University of Valencia San Vicente Mártir, Valencia, Spain; ⁸John Ochsner Heart and Vascular Institute, Ochsner Clinical School - The University of Queensland School of Medicine, New Orleans, LA, USA; ⁹Department of Physiology, Faculty of Medicine, University of Valencia and INCLIVA Biomedical Research Institute, Valencia, Spain

Correspondence to: Fabian Sanchis-Gomar, MD, PhD. Department of Physiology, Faculty of Medicine, University of Valencia, Av. Blasco Ibañez, 15, 46010 Valencia, Spain. Email: fabian.sanchis@uv.es.

Abstract: Myocarditis is among the causes of arrhythmic sudden cardiac death (SCD) in young athletes, with viral infection being the most common cause worldwide. Myocarditis recently has been reported as one of the cardiac complications of coronavirus disease 2019 (COVID-19) in athletes. Here we present a case of a 20-year-old male recreational soccer player with an episode of loss of consciousness in the context of respiratory infection. The patient reports having woken up with symptoms of an upper respiratory tract infection, and after playing a soccer match, he developed dizziness and a headache. He then suffered vasovagal syncope without loss of sphincter control. Physical examination, heart auscultation, peripheral and carotid pulses, and blood, microbiological/serological tests result on admission were normal. Moreover, no jugular engorgement at 45°, malleolar edema, or other heart failure signs were found. The 12-lead electrocardiogram (ECG), echocardiogram, 24-hour Holter-ECG did not reveal any significant finding. A cardiac magnetic resonance (CMR) was finally performed, revealing an abnormal signal increase was observed at the apical level in the short-tau inversion-recovery (STIR) and 4-chamber sequences. In addition, a pattern of apical fibrosis was observed in 4- and 2-chamber and short-axis late enhancement sequences for assessment of myocardial viability confirming the diagnosis of myocarditis. In athletes with suspected myocarditis, CMR seems to be a useful diagnostic tool, with excellent sensitivity for detecting inflammation, myocardial edema, and/or focal scarring.

Keywords: Myocarditis; sudden cardiac death (SCD); athlete; cardiac magnetic resonance; case report

Received: 06 April 2021; Accepted: 02 July 2021; Published: 25 October 2021.

doi: 10.21037/acr-21-24

View this article at: <https://dx.doi.org/10.21037/acr-21-24>

[^] ORCID: 0000-0003-0424-4208.

Introduction

Myocarditis is a major cause of arrhythmic sudden cardiac death (SCD) in young athletes (1-3). Between 7% to 20% of SCD in young athletes is due to myocarditis. The proportion of SCD caused by myocarditis (SCD-myocarditis) is highly variable (between 1% and 14% among young people) due to the populations' homogeneity and differences in the definition of SCD and post-mortem myocarditis. The etiology of myocarditis is heterogeneous, with viral infection being the most common cause worldwide, especially enterovirus, coxsackievirus B, parvovirus B-19, and human herpesvirus 6. Importantly, myocarditis recently has been reported as one of the cardiac complications of coronavirus disease 2019 (COVID-19) in athletes (4). This fact is particularly relevant since physical exercise during the acute phase of viral myocarditis may precipitate malignant ventricular arrhythmias. The diagnosis is based on a detailed medical history, physical examination, and complementary tests, such as appropriate blood work (e.g., troponin), 12-lead electrocardiogram (ECG), echocardiography, and eventually, cardiac magnetic resonance (CMR) (5).

We present the following article in accordance with the CARE reporting checklist (available at <https://dx.doi.org/10.21037/acr-21-24>).

Case presentation

A 20-year-old male recreational soccer player was admitted to the cardiology unit after presenting with an episode of loss of consciousness in the context of respiratory infection without associated fever. The patient reports having woken up with symptoms of an upper respiratory tract infection, and after playing a soccer match, he developed dizziness and a headache. He then suffered vasovagal syncope without loss of sphincter control. No chest pain, nausea, vomiting, diarrhea, or other associated symptoms were reported.

The patient had no relevant family history of disease, with no cases of SCD. He had not previously reported palpitations, syncopal episodes, or dizziness associated with sports participation. No previous interventions were reported.

The physical examination on admission included a heart rate of 77 bpm and resting blood pressure of 134/76 mmHg; temperature: 36.7 °C; SpO₂ 98% on room air. Heart auscultation was normal. Symmetrical peripheral pulses were palpable. The patient had symmetrical carotid pulses,

with no jugular engorgement at 45°, malleolar edema, or other heart failure signs. We also performed a blood test (*Table 1*). We also requested microbiological/serological tests.

The 12-lead ECG showed sinus rhythm with a right bundle branch block (RBBB), with pronounced T-wave inversions (TWI) in inferolateral leads (*Figure 1*). The echocardiogram revealed non-dilated ventricular and atrial cavities, non-hypertrophic ventricular septum, and normal systolic and diastolic function. There was no pericardial effusion. Finally, a 24-hour Holter-ECG study was performed, which did not reveal any significant arrhythmia.

Viral serological test results were negative (i.e., adenovirus, enterovirus, toxoplasma *Gondi*, influenza virus A and B). A CMR was finally performed, revealing the absence of significant ventricular hypertrophy in a short-axis cine sequence of the left ventricle (LV) and right ventricle (RV) (*Figure 2*). However, an abnormal signal increase was observed at the apical level in the short-tau inversion-recovery (STIR) and 4-chamber sequences (*Figure 3A,B*). In addition, a pattern of apical fibrosis was observed in 4- and 2-chamber and short-axis late enhancement sequences for assessment of myocardial viability (*Figure 3C*), confirming the diagnosis of myocarditis.

The patient had a favorable evolution of symptoms, with abatement of the fever, an absence of heart failure signs or symptoms, arrhythmias or hemodynamic instability, and a good response to the symptomatic treatment administered (1 gram of acetaminophen every 8 hours). Given the good clinical evolution, the patient was discharged.

Sports practice was discouraged for six months, during which regular medical checkups were performed. In this regard, it is worth highlighting the last examination carried out two years after the initial presentation. The 12-lead ECG showed persistent T wave inversion in the inferolateral wall (*Figure 4*). The patient underwent an exercise stress test, showing anterolateral T-wave pseudonormalization with maximum effort (*Figure 5*), without induction of ventricular arrhythmias and negative for myocardial ischemia (ST elevation/depression). No arrhythmic episodes were found on a 24-hour Holter monitor. Finally, a control CMR showed apical fibrosis persistence in the 4- and 2-chamber late gadolinium enhancement (LGE) sequences without edema (*Figure 6*). The patient returned to exercise, playing soccer recreationally, and practicing light-to-moderate-intensity physical activity with high dynamic components.

All procedures performed in this case were in accordance

Table 1 Blood test results

Variables	Patient's values	Normal values
Leukocytes/L	24,200	4,500–11,000
Neutrophils, %	90.8	40–60
Lymphocytes, %	3.9	20–40
Monocytes, %	4.6	2–8
Eosinophils, %	0.4	1–4
Basophils, %	0.3	0.5–1
Platelets, $\times 10^9/L$	229	150–450
Hemoglobin, mg/dL	14.8	13.5–17.5
Hematocrit, %	43.7	40–50
Fibrinogen, mg/dL	412	200–400
Na, mEq/L	135	135–145
K, mEq/L	3.4	3.5–5
hs-cTnI, ng/mL	0.11	0.060
CRP, mg/dL	1.9	<0.3

hs-cTnI, high-sensitivity cardiac troponin I; CRP, C-reactive protein.

with the ethical standards of the institutional and national research committees and with the Helsinki Declaration (as revised in 2013). Written informed consent was obtained from the patient.

Discussion

A complete evaluation is recommended when myocarditis is suspected, including CMR (6). The case we have presented here is especially relevant in the COVID-19 era since severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is among the potential causes provoking acute myocarditis in young athletes and other patients. In effect, SARS-CoV-2 infection has been associated with myocarditis. Current evidence indicates that approximately 20% of hospitalized patients suffered cardiac injury, with elevated levels of high-sensitivity troponins (hs-cTnI) (7). A meta-summary of cases reflected that CMR established the diagnosis of myocarditis in many patients, with the presence of edema and cardiac injury, highlighting the valuable role of CMR in these patients (8). In the COVID-19 era, CMR imaging will be a very useful and effective tool to assess

cardiac involvement. LGE is the best predictor of mortality regardless of symptoms, functional class, ejection fraction, or LV end-diastolic volume (9). Ventricular scarring/fibrosis (as evidenced by LGE on the CMR), which may be missed by echocardiography, can cause SCD due to ventricular arrhythmias (10).

The finding of unexplained TWI on athletes' 12-lead ECG should be thoroughly investigated by CMR imaging (11). In a study carried out by Schnell *et al.* (12), heart disease was established in 44.5% of 155 athletes with deep TWI (80% in lateral leads), with hypertrophic cardiomyopathy the most diagnosis (81%). Echocardiography was abnormal in 37 (53.6%) cases, while CMR revealed another 24 (16.6%) athletes with heart disease. Thus, CMR significantly improves the accuracy of diagnosing cardiomyopathy in athletes with marked repolarization alterations in the ECG and normal echocardiogram, as was also evidenced in the present clinical case.

CMR is currently the most widely used noninvasive imaging technique to confirm acute myocarditis, especially in hemodynamically stable patients during the first week of symptoms. CMR allows optimal myocardial tissue characterization, detecting inflammatory damage as necrosis and fibrosis in the form of LGE. Although the value of repeating the CMR at six months is unknown, as are its clinical and prognostic implications, it may be advisable to monitor the evolution of myocardial damage.

The optimal treatment of patients with acute myocarditis, and their prognosis, remain controversial. It is essential to highlight the need for prolonged monitoring with Holter-ECG or other devices regardless of ventricular function.

Due to the risk of exercise-induced arrhythmia, athletes with acute myocarditis should not return to competitive activity for 3–6 months after recovery from the acute process (13). The current sports cardiology guidelines of the European Cardiology Society (14) advise that athletes diagnosed with recent myocarditis should refrain from physical or sports activities while active inflammation persists, regardless of other factors (age, sex, LV systolic dysfunction). Likewise, it reinforces the previous recommendation of abstinence from moderate-to-high intensity exercise for a period of 3 to 6 months (5), highlighting the essential role of the CMR in the monitoring and evolution of this entity, together with a comprehensive evaluation after a complete recovery to




Figure 1 Baseline ECG at admission. ECG, electrocardiogram.




Figure 2 LV and RV short-axis cine sequence ruling out significant ventricular hypertrophy. LV, left ventricle; RV, right ventricle.




Figure 3 CMR on admission. (A) 4-chamber STIR sequence showing signal increase at apical level. Regions of interest used to measure T2 value at the apical (n=214) and lateral (n=88) levels. Ratio T2 value apical/lateral level >2. (B) LV and RV short-axis STIR sequence showing increased signal at the apical level. (C) Viability sequence in 2 and 4 chambers and short-axis planes showing increased signal at the apical level. Pattern compatible with fibrosis (red arrows) and confirming the diagnosis of apical myocarditis. CMR, cardiac magnetic resonance; STIR, short-tau inversion-recovery; LV, left ventricle; RV, right ventricle.




Figure 4 Follow-up ECG 2 years later. ECG, electrocardiogram.




Figure 5 Control stress test 2 years later. Anterolateral TWI with maximal effort. Clinically and electrically negative for ischemia. TWI, T-wave inversion.




Figure 6 Control CMR. (A,B) Viability sequence in 2 and 4 chamber planes in which the persistence of apical fibrosis is shown. (C) Sequence without signal increase at the apical level in 4-chamber STIR sequence. CMR, cardiac magnetic resonance; STIR, short-tau inversion-recovery.

assess the risk of exercise-related SCD.

Conclusions

CMR seems to be a useful diagnostic tool in athletes with suspected myocarditis, with excellent sensitivity for detecting inflammation, myocardial edema, and/or focal scarring. The distribution and extent of LGE with a non-ischemic pattern is an independent predictor of adverse cardiac events during follow-up.

Acknowledgments

Funding: None.

Footnote

Reporting Checklist: The authors have completed the CARE reporting checklist. Available at <https://dx.doi.org/10.21037/acr-21-24>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://dx.doi.org/10.21037/acr-21-24>). FSG serves as an unpaid editorial board member of *AME Case Reports* from Apr 2021 to Mar 2023. The other authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All procedures

performed were in accordance with the ethical standards of the institutional and national research committees and with the Helsinki Declaration (as revised in 2013). Written informed consent was obtained from the patient.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Thiene G. Sudden cardiac death in the young: a genetic destiny? *Clin Med (Lond)* 2018;18:s17-23.
2. Lynge TH, Nielsen TS, Gregers Winkel B, et al. Sudden cardiac death caused by myocarditis in persons aged 1-49 years: a nationwide study of 14294 deaths in Denmark. *Forensic Sci Res* 2019;4:247-56.
3. Bagnall RD, Weintraub RG, Ingles J, et al. A Prospective Study of Sudden Cardiac Death among Children and Young Adults. *N Engl J Med* 2016;374:2441-52.
4. Phelan D, Kim JH, Elliott MD, et al. Screening of Potential Cardiac Involvement in Competitive Athletes Recovering From COVID-19: An Expert Consensus Statement. *JACC Cardiovasc Imaging* 2020;13:2635-52.
5. Maron BJ, Udelson JE, Bonow RO, et al. Eligibility and Disqualification Recommendations for Competitive

- Athletes With Cardiovascular Abnormalities: Task Force 3: Hypertrophic Cardiomyopathy, Arrhythmogenic Right Ventricular Cardiomyopathy and Other Cardiomyopathies, and Myocarditis: A Scientific Statement From the American Heart Association and American College of Cardiology. *J Am Coll Cardiol* 2015;66:2362-71.
6. Sinagra G, Anzini M, Pereira NL, et al. Myocarditis in Clinical Practice. *Mayo Clin Proc* 2016;91:1256-66.
 7. Shi S, Qin M, Shen B, et al. Association of Cardiac Injury With Mortality in Hospitalized Patients With COVID-19 in Wuhan, China. *JAMA Cardiol* 2020;5:802-10.
 8. Ho JS, Sia CH, Chan MY, et al. Coronavirus-induced myocarditis: A meta-summary of cases. *Heart Lung* 2020;49:681-5.
 9. Gräni C, Eichhorn C, Bière L, et al. Prognostic Value of Cardiac Magnetic Resonance Tissue Characterization in Risk Stratifying Patients With Suspected Myocarditis. *J Am Coll Cardiol* 2017;70:1964-76.
 10. Zorzi A, Perazzolo Marra M, Rigato I, et al. Nonischemic Left Ventricular Scar as a Substrate of Life-Threatening Ventricular Arrhythmias and Sudden Cardiac Death in Competitive Athletes. *Circ Arrhythm Electrophysiol* 2016;9:e004229.
 11. Sharma S, Drezner JA, Baggish A, et al. International Recommendations for Electrocardiographic Interpretation in Athletes. *J Am Coll Cardiol* 2017;69:1057-75.
 12. Schnell F, Riding N, O'Hanlon R, et al. Recognition and significance of pathological T-wave inversions in athletes. *Circulation* 2015;131:165-73.
 13. Hurwitz B, Issa O. Management and Treatment of Myocarditis in Athletes. *Curr Treat Options Cardiovasc Med* 2020;22:65.
 14. Pelliccia A, Sharma S, Gati S, et al. 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease. *Eur Heart J* 2021;42:17-96. Erratum in: *Eur Heart J* 2021;42:548-9.

doi: 10.21037/acr-21-24

Cite this article as: de la Guía-Galipienso F, García-González P, Fabregat-Andrés O, Quesada-Dorador A, Meyer-Josten C, Lavie CJ, Morin DP, Sanchis-Gomar F. Cardiac magnetic resonance in the diagnosis of the unusually detected acute myocarditis in the young people: a case report. *AME Case Rep* 2021;5:35.