

UC Davis

UC Davis Previously Published Works

Title

Dispositional Negativity: An Integrative Psychological and Neurobiological Perspective

Permalink

<https://escholarship.org/uc/item/3p2339r7>

Journal

Psychological Bulletin, 142(12)

ISSN

0033-2909

Authors

Shackman, Alexander J
Tromp, Do PM
Stockbridge, Melissa D
[et al.](#)

Publication Date

2016-12-01

DOI

10.1037/bul0000073

Peer reviewed

Dispositional negativity: An integrative psychological and neurobiological perspective

Alexander J. Shackman^{a,c,d}

Do P. M. Tromp^{e,g,i,j}

Melissa D. Stockbridge^b, Claire M. Kaplan^{a*}, Rachael M. Tillman^{a*}

Andrew S. Fox^{e,f}

Departments of ^aPsychology and ^bHearing and Speech Sciences, ^cNeuroscience and Cognitive Science Program, ^dMaryland Neuroimaging Center, University of Maryland, College Park, MD 20742 USA

^eDepartment of Psychology and ^fCalifornia National Primate Research Center, University of California, Davis, CA 95616 USA

* contributed equally

ACKNOWLEDGEMENTS

Authors acknowledge assistance from L. Friedman, L. Pessoa, and J. Smith and constructive feedback from M. Barstead, J. Curtin, M. Gamer, D. Grupe, E. Lemay, K. Rubin, J. Stern, three anonymous reviewers, and the Editor. This work was supported by the University of California, University of Maryland, National Institute of Drug Abuse (DA040717), National Institute of Mental Health (MH107444), and a National Science Foundation Graduate Research Fellowship. Authors declare no conflicts of interest.

PLEASE CITE THIS MANUSCRIPT AS:

Shackman, A. J., Tromp, D. M. P., Stockbridge, M. D., Kaplan, C. M., Tillman, R. M. & Fox, A. S. (*in press*). Dispositional negativity: An integrative psychological and neurobiological perspective. *Psychological Bulletin*.

ADDRESS CORRESPONDENCE TO

Alexander J. Shackman (shackman@umd.edu)
Department of Psychology
3123G Biology-Psychology Building
University of Maryland
College Park, Maryland 20742 USA

ABSTRACT

Dispositional negativity—the propensity to experience and express more frequent, intense, or enduring negative affect—is a fundamental dimension of childhood temperament and adult personality. Elevated levels of dispositional negativity can have profound consequences for health, wealth, and happiness, drawing the attention of clinicians, researchers, and policy makers. Here, we highlight recent advances in our understanding of the psychological and neurobiological processes linking stable individual differences in dispositional negativity to momentary emotional states. Self-report data suggest that three key pathways—increased stressor reactivity, tonic increases in negative affect, and increased stressor exposure—explain most of the heightened negative affect that characterizes individuals with a more negative disposition. Of these three pathways, tonically elevated, indiscriminate negative affect appears to be most central to daily life and most relevant to the development of psychopathology. New behavioral and biological data provide insights into the neural systems underlying these three pathways and motivate the hypothesis that seemingly ‘tonic’ increases in negative affect may actually reflect increased reactivity to stressors that are remote, uncertain, or diffuse. Research focused on humans, monkeys, and rodents suggests that this indiscriminate negative affect reflects trait-like variation in the activity and connectivity of several key brain regions, including the central extended amygdala and parts of the prefrontal cortex. Collectively, these observations provide an integrative psychobiological framework for understanding the dynamic cascade of processes that bind emotional traits to emotional states and, ultimately, to emotional disorders and other kinds of adverse outcomes.

Humans, monkeys, and other animals show marked individual differences in temperament, the tendency to experience fear, anger, disgust, joy, and other fleeting emotional states (Gosling, 2008)¹. This distinction between emotional traits and emotional states has its origins in antiquity. More than 2,000 years ago, the Roman intellectual Cicero drew a sharp distinction between anxious temperament (*anxietas*) and anxiety (*angor*) (Eysenck, 1983). Like many contemporary researchers, he characterized traits as the proneness to experience particular emotional states (*proclivitas*). Our own working definition is that traits represent enduring emotional and cognitive biases that first emerge early in life, but continue to evolve and grow in complexity across the lifespan (Shiner, *in press-b*). Emotional traits account, in a probabilistic manner, for consistency in thoughts, feelings, physiology, and actions across time and situations (Caspi, Roberts, & Shiner, 2005; Fleeson, 2001; Shiner, *in press-a*; Shiner et al., 2012). Like other psychological constructs that vary across individuals, emotional traits reflect the combined influence of genes and experience on brain structure and function (Polderman et al., *in press*).

The study of temperament has proven theoretically informative and practically important. Individual differences in temperament and personality have profound consequences for health, wealth, and happiness. Accordingly, temperament has increasingly drawn the attention of educators, social scientists, neurobiologists, clinicians, economists, and public policy makers (Duckworth & Allred, 2012; Ferguson, Heckman, & Corr, 2011; Lahey, 2009; Moffitt, Poulton, & Caspi, 2013; Roberts, Kuncel, Shiner, Caspi, & Goldberg, 2007). Despite this growing interest, fundamental questions about the nature and origins of temperament have remained unresolved. One of the most basic questions concerns the nature of the relations between trait-like differences in temperament and more transient emotional experiences and behaviors (Barlow, Sauer-Zavala, Carl, Bullis, & Ellard, 2013; Epstein, 1994). As the pioneering

¹ 'Emotion' is a fuzzy, contentious category that conventionally includes valenced processes (e.g., action tendencies, attention, motivated learning, overt behavior, subjective feelings, and alterations in peripheral physiology) that are triggered by specific external or internal antecedents, such as actual or remembered threat in the case of fear (Anderson & Adolphs, 2014; Beck, 2015; Gendron & Barrett, 2009; Izard, 2010; Lapate & Shackman, *in press*).

psychologist David Funder noted, traits “describe patterns and consistencies in behavior, but they don't explain where those patterns and consistencies come from” (Funder, 1994).

Here, we highlight recent advances in our understanding of the psychological and neurobiological processes linking emotional traits to emotional states, focusing on dispositional negativity, one of the most intensively studied dimensions of temperament and personality. We begin by describing the nature of dispositional negativity and surveying its association with well-being and disease. Next, we review self-report and behavioral data suggesting that three key pathways—increased stressor reactivity, tonic increases in negative affect, and increased stressor exposure—explain most of the heightened negative affect characteristic of individuals with a negative disposition (**Figure 1**). In the third section of the review, we explore recent advances in our understanding of the neural systems underlying these three pathways, focusing especially on studies of fear and anxiety. This work motivates the hypothesis that seemingly ‘tonic’ increases in negative affect may actually reflect increased reactivity to stressors that are uncertain, temporally remote, or perceptually diffuse (e.g., an unfamiliar experimental context, a pitch-black room; **Figure 2**). These observations raise a number of interesting new questions. We conclude by outlining several strategies for addressing them and for developing a deeper understanding of the pathways linking emotional traits to momentary emotional states and, ultimately, to psychopathology and other kinds of adverse outcomes.

Method

The goal of our review was to integrate psychological and biological perspectives on dispositional negativity into a coherent theoretical framework. Accordingly, it includes 579 citations covering the period between 1966 and 2016, with the major emphasis on recent research (median year of publication = 2011, *SD* = 8.1; ‘in press’ publications were coded as 2016). This body of research encompasses

laboratory, field (e.g., daily diary, ecological momentary assessments), and epidemiological research. It includes published studies of mice, rats, monkeys, and humans, including studies of children, adolescents, adults, and elders. Although most of the work is focused on unselected individuals, we also highlight relevant evidence gleaned from studies of psychiatric and neurological patients. Of necessity, we draw on published articles and reviews from a broad spectrum of scholarly disciplines, from psychology and psychiatry to genetics and neuroscience. For all articles, backward citation checks were used to increase coverage. Although our aims and scope precluded the use of formal meta-analytic techniques, in order to maximize reproducibility and generalizability we place special emphasis on evidence gleaned from meta-analyses² and large-scale studies, including prospective longitudinal studies and nationally representative samples.

The Nature and Consequences of Elevated Dispositional Negativity

Dispositional negativity or ‘negative emotionality’—the propensity to experience and express more frequent, intense, or enduring negative affect—is a fundamental dimension of childhood temperament and adult personality. Individuals with a more negative disposition tend to be anxious, guilt-prone, insecure, moody, critical, angry, and dissatisfied. They tend to perceive the world as dangerous and threatening and themselves as inadequate (Barlow et al., 2013; Caspi et al., 2005; L. A. Clark & Watson, 2008; Lahey, 2009). Dispositional negativity is sometimes parsed into two mid-level dimensions: *anxious*

² Our review incorporates evidence derived from 41 more narrowly focused meta-analyses and systematic reviews (Avery, Clauss, & Blackford, 2016; Bastiaansen et al., 2014; Buhle et al., 2014; Calder, Ewbank, & Passamonti, 2011; Cavanagh & Shackman, 2015; Chase, Eickhoff, Laird, & Hogarth, 2011; Clauss & Blackford, 2012; Connor-Smith & Flachsbart, 2007; Duits et al., 2015; Etkin & Wager, 2007; A. S. Fox & Kalin, 2014; A. S. Fox, Oler, Tromp, Fudge, & Kalin, 2015; Hakulinen, Elovainio, et al., 2015; Hakulinen, Hintsanen, et al., 2015; Hamilton et al., 2012; Houben, Van Den Noortgate, & Kuppens, 2015; Jokela, Pulkki-Raback, Elovainio, & Kivimaki, 2014; Karney & Bradbury, 1995; Kotov, Gamez, Schmidt, & Watson, 2010; Kuhn & Gallinat, 2011; Lahey, 2009; Linetzky, Pergamin-Hight, Pine, & Bar-Haim, 2015; Malouff, Thorsteinsson, Rooke, & Schutte, 2007; Malouff, Thorsteinsson, Schutte, Bhullar, & Rooke, 2010; Matthews, Deary, & Whiteman, 2009; Ng, Eby, Sorensen, & Feldman, 2005; Ormel et al., 2013; Polderman et al., *in press*; Prinzie, Stams, Dekovic, Reijntjes, & Belsky, 2009; Roberts & DelVecchio, 2000; Roberts, Walton, & Viechtbauer, 2006; Shackman et al., 2011; Steel, Schmidt, & Shultz, 2008; Tang, Fellows, Small, & Dagher, 2012; Thibodeau, Jorgensen, & Kim, 2006; Turkheimer, Pettersson, & Horn, 2014; Vukasovic & Bratko, 2015; Wacker, Chavanon, & Stemmler, 2010; Watson & Clark, 1984; Watson & Naragon-Gainey, 2014).

distress, encompassing feelings of anxiety, fear, and depression; and *irritable distress*, encompassing feelings of anger, frustration, and hostility (Caspi et al., 2005). At present, there is greater theoretical consensus about the nature and significance of anxious distress (Caspi et al., 2005; Ormel et al., 2013; Soto & John, *in press*) and it has received more empirical attention (Barlow et al., 2013). Anxious distress, in turn, subsumes a variety of narrower facet traits, including anxious temperament, anxiety sensitivity, behavioral inhibition, harm avoidance, neuroticism, and trait anxiety (Barlow et al., 2013; Caspi et al., 2005; Markon, Krueger, & Watson, 2005; van den Berg et al., 2014; Widiger, 2009)³.

We conceptualize dispositional negativity as an extended family of closely related phenotypes that first emerge early in development, persist into adulthood, and reflect a combination of heritable and non-heritable factors (A. S. Fox & Kalin, 2014; Lake, Eaves, Maes, Heath, & Martin, 2000; Ormel et al., 2013; Power & Pluess, 2015; Smith et al., 2015; Soto & John, 2014; Turkheimer et al., 2014; Vukasovic & Bratko, 2015). Among adults, concordance between self- and informant-reported (e.g., friends, family members, co-workers) dispositional negativity is substantial (Connolly, Kavanagh, & Viswesvaran, 2007), particularly when multiple informants are employed (McCrae & Costa, 1987), suggesting that it is more than just a negative response bias. Core features of this phenotypic family—including increased behavioral inhibition, heightened vigilance, and other signs of fear and anxiety—are expressed similarly across mammalian species, enabling mechanistic studies to be performed in rodents and monkeys (Boissy, 1995; Mobbs & Kim, 2015; Oler, Fox, Shackman, & Kalin, 2016). Although the molecular underpinnings of dispositional negativity and its neural substrates remain poorly understood (Bastiaansen et al., 2014; Bogdan, Pagliaccio, Baranger, & Hariri, 2016; Christian et al., 2009), some

³ While there are potential benefits to focusing on narrowly defined phenotypic traits (Block, 1995, 2010; Clifford, Lemery-Chalfant, & Goldsmith, 2015; K. H. Rubin & Asendorpf, 1993), adopting a narrow perspective substantially reduces the size and scope of the relevant evidentiary record. At present, very little is known about some facets of dispositional negativity—including anger, guilt, and shame—contribute to adverse outcomes (Kopala-Sibley et al., *in press*), making this an important challenge for future research. From a public health perspective, focusing on broad phenotypes that confer elevated risk for a range of adverse outcomes maximizes the opportunity to develop broad-spectrum interventions (Barlow, Ellard, Sauer-Zavala, Bullis, & Carl, 2014; Chronis-Tuscano et al., 2015; Moffitt et al., 2013).

promising candidates have recently been identified in humans (Buckholtz et al., 2008; Okbay et al., *in press*), monkeys (Alisch et al., 2014; A. S. Fox et al., 2012; Kalin et al., *in press*; Oler et al., 2009; Rogers et al., 2013; Roseboom et al., 2014), and rodents (Turner, Clinton, Thompson, Watson, & Akil, 2011).

Dispositional negativity is stable, but not immutable, and like other emotional traits continues to develop and change across the lifespan (Fraley & Roberts, 2005; Roberts & DeVecchio, 2000; Roberts & Mroczek, 2008; Roberts et al., 2006). In fact, mean levels of dispositional negativity show substantial fluctuations—equivalent to *T*-scores of 2 in males and 5 in females—between the ages of 10 and 65, peaking in adolescence (Soto, John, Gosling, & Potter, 2011). Several large ($n = 4,850 - 1,267,218$) international studies indicate that, from about age 14 on, women tend to report substantially higher levels of dispositional negativity than men (De Bolle et al., 2015; Schmitt, Realo, Voracek, & Allik, 2008; Soto et al., 2011).

A range of evidence indicates that dispositional negativity can be increased by stress, trauma, and negative life events (e.g., death of a spouse, birth of a child, chronic disease; Barlow et al., 2013; Hutteman, Bleidorn, Kerestes, et al., 2014; Jeronimus, Riese, Sanderman, & Ormel, 2014; Jokela, Hakulinen, Singh-Manoux, & Kivimaki, 2014; Jokela, Kivimaki, Elovainio, & Keltikangas-Jarvinen, 2009; Laceulle, Nederhof, Karreman, Ormel, & Van Aken, 2011; Ludtke, Roberts, Trautwein, & Nagy, 2011; Parker, Ludtke, Trautwein, & Roberts, 2012; Roberts, Caspi, & Moffitt, 2003; Robins, Caspi, & Moffitt, 2002). But importantly it can also be decreased by cognitive-behavioral (Barlow et al., 2013; Bennett et al., 2015; Mihalopoulos et al., 2015) and pharmacological interventions for anxiety and depression (Barlow et al., 2013; Soskin, Carl, Alpert, & Fava, 2012), raising the possibility of developing strategies for identifying high-risk individuals and preventing the onset of more severe sequelae. Identifying the psychological and neurobiological mechanisms governing the malleability of temperament is a

particularly important avenue for future research, one that promises to provide new targets for intervention (A. S. Fox et al., 2012).

There is clear evidence that dispositionally negative individuals tend to experience heightened levels of momentary negative affect. Self-report measures of dispositional negativity (trait) and negative affect (state) are strongly correlated (Lieberman et al., *in press*; Matthews et al., 2009; Watson & Clark, 1984, 1992). In fact, a recent meta-analysis incorporating data from more than 30,000 individuals showed that dispositional negativity explains 30-50% of the variance in negative affect (Steel et al., 2008). Ratings obtained from other informants, such as clinicians and spouses, yield similar conclusions, indicating that these trait-state relations are not an artifact of response biases (Lieberman et al., *in press*; McCrae & Costa, 1991; Soto & John, *in press*; Steel et al., 2008; Watson & Clark, 1984).

It merits comment that the impact of dispositional negativity on momentary emotional experience is not limited to negative affect. Individuals with a more negative disposition are also prone to lower levels of positive affect (Aldinger et al., 2014; Gable, Reis, & Elliot, 2000; Jacobs et al., 2011; Soto & John, *in press*; Watson & Clark, 1984; Zautra, Affleck, Tennen, Reich, & Davis, 2005) and subjective wellbeing (Steel et al., 2008), perhaps reflecting a suppressive consequence of negative affect on reward and appetitive motivation (i.e., sometimes termed 'stress-induced' anhedonia; Pizzagalli, 2014).

Dispositional negativity predicts a multitude of practically important outcomes, from satisfaction and wealth to marital stability and disease. Increased dispositional negativity is associated with lower levels of educational attainment (Damian, Su, Shanahan, Trautwein, & Roberts, 2015; Hengartner, Kawohl, Haker, Rossler, & Ajdacic-Gross, 2016) and occupational success (Heineck, 2011; Hengartner, Kawohl, et al., 2016; Ng et al., 2005; Shanahan, Bauldry, Roberts, Macmillan, & Russo, 2014; Soldz & Vaillant, 1999; Sutin, Costa, R., & Eaton, 2009; Uysal & Pohlmeier, 2011; Viinikainen, Kokko, Pulkkinen, & Pehkonen,

2010). In a nationally representative sample of 81,000 high school students, individuals who were one standard-deviation above the mean lost the equivalent of half an academic year in educational attainment and \$3,628 in annual income by the time they reached mid-life compared to those one standard-deviation below the mean (in 2014 dollars; Damian et al., 2015).

Individuals with elevated dispositional negativity report reduced satisfaction with their lives (Dyrenforth, Kashy, Donnellan, & Lucas, 2010; Soto & Luhmann, 2013), jobs (Wayne, Musisca, & Fleeson, 2004), friends (R. E. Wilson, Harris, & Vazire, 2015), and spouses (Dyrenforth et al., 2010; Solomon & Jackson, 2014). In fact, elevated levels of dispositional negativity in adolescence have been shown to predict lower levels of job satisfaction 50 years later (Staw, Bell, & Clausen, 1986) and lower levels of psychological well-being 36 years later (Abbott et al., 2008). Heightened dispositional negativity is also a strong predictor of future loneliness (Pressman et al., 2005; Stokes, 1985) and divorce (Karney & Bradbury, 1995; Kurdek, 1993). In one particularly compelling example, 278 married couples were longitudinally assessed between 1936 and 1981 (Kelly & Conley, 1987). Of these, 50 divorced. Self- and acquaintance-ratings of husbands' and wives' dispositional negativity at the initial assessment were among the strongest prenuptial predictors of divorce across the 45-year follow-up period. Similar results have been reported for spouses with pre-existing anxiety disorders—for example, the odds of getting divorced among patients with a pre-marital diagnosis of generalized anxiety disorder is 1.7-fold greater than psychiatrically healthy controls (Kessler, Walters, & Forthofer, 1998). More frequent marital dissolution is, in turn, associated with reduced reproductive success. In a prospective study of more than 1,500 Finnish adults, a one standard-deviation increase in dispositional negativity decreased the odds of having a second and third child by 11% and 15%, respectively, largely due to the higher frequency of marital dissolution and divorce (Jokela et al., 2009).

From the perspective of physical health, dispositional negativity is associated with sleep problems (Hintsanen et al., 2014), metabolic syndrome (Phillips et al., 2010), elevated cholesterol levels (Hengartner, Kawohl, et al., 2016), and a wide variety of other physical diseases and subjective health complaints (e.g., coronary heart disease; Deary, Weiss, & Batty, 2010; Gale et al., 2016; Iacovino, Bogdan, & Oltmanns, 2016; Jokela, Pulkki-Raback, et al., 2014; Lahey, 2009; Mund & Neyer, *in press*). Among nearly 7,000 older adults (mean age = 68.4 years) followed as part of the Health and Retirement study, a one standard-deviation increase in dispositional negativity prospectively increased the odds of being diagnosed with a heart condition by 24%, lung disease by 29%, and hypertension by 37% during the four-year follow-up period (Weston, Hill, & Jackson, 2015). Increased morbidity partially reflects dispositional negativity's association with unhealthy behaviors, such as chronic tobacco and substance use (Gale et al., 2016; Hakulinen, Hintsanen, et al., 2015; Hengartner, Kawohl, et al., 2016; Kotov et al., 2010; Leventhal et al., 2012; Malouff et al., 2007; Soldz & Vaillant, 1999), and likely contributes to premature mortality among individuals with a more negative disposition (Chapman, Fiscella, Kawachi, & Duberstein, 2010; Jackson, Connolly, Garrison, Leveille, & Connolly, 2015; Terracciano, Lockenhoff, Zonderman, Ferrucci, & Costa, 2008; R. S. Wilson et al., 2005).

Dispositional negativity is also a key risk factor for anxiety disorders, depression, and substance abuse (Clauss & Blackford, 2012; Conway, Craske, Zinbarg, & Mineka, 2016; Grav, Stordal, Romild, & Hellzen, 2012; Hakulinen, Elovainio, et al., 2015; Hengartner, Kawohl, et al., 2016; Kendler & Gardner, 2014; Soldz & Vaillant, 1999; Watson & Naragon-Gainey, 2014; S. Wilson, Vaidyanathan, Miller, McGue, & Iacono, 2014)—psychiatric disorders that are highly prevalent, debilitating, and often challenging to treat (Bystritsky, 2006; Collins et al., 2011; DiLuca & Olesen, 2014; Griebel & Holmes, 2013; Insel, 2012; Kessler, Petukhova, Sampson, Zaslavsky, & Wittchen, 2012; Whiteford et al., 2013). The magnitude of these associations is substantial: a meta-analysis incorporating 175 studies reported that the mean Cohen's d across mood, anxiety, and substance use disorders was 1.65, ranging from $d \cong 2$ for dysthymia

and anxiety disorders to $d = .77$ for alcohol use disorder (Kotov et al., 2010). Elevated dispositional negativity is among the strongest prospective predictors of future internalizing disorders (see also D. A. Clark, Durbin, Hicks, Iacono, & McGue, *in press*; $k = 46$ studies; mean Cohen's $d = .63$; Ormel et al., 2013). For example, data from the Zurich Cohort Study ($n = 591$) indicates that a one standard-deviation increase in dispositional negativity at the time of the baseline assessment in 1988 increased the odds of developing a major depressive episode by 41% and an anxiety disorder by 32% during the twenty-year follow-up period (Hengartner, Ajdacic-Gross, Wyss, Angst, & Rossler, 2016). These relations are evident after eliminating overlapping item content (Uliaszek et al., 2009). They also appear to be strengthened by exposure to stress (Kopala-Sibley et al., *in press*; Kopala-Sibley et al., 2016; Vinkers et al., 2014), suggesting that high levels of dispositional negativity represent a diathesis (Monroe & Simons, 1991) for the internalizing spectrum of disorders (i.e., anxiety and depression). Among patients with a history of internalizing disorders, higher levels of dispositional negativity are associated with a greater number of co-morbid diagnoses (Hengartner, Kawohl, et al., 2016) and a more pessimistic prognosis (Berlanga, Heinze, Torres, Apiquian, & Cabellero, 1999; Duggan, Lee, & Murray, 1990; Faravelli, Ambonetti, Pallanti, & Pazzagli, 1986; Hirschfeld, Klerman, Andreasen, Clayton, & Keller, 1986; Kendler, Neale, Kessler, & Heath, 1993; Ormel, Oldehinkel, & Vollebergh, 2004; Quilty et al., 2008; Scott, Williams, Brittlebank, & Ferrier, 1995; Weissman, Prusoff, & Klerman, 1978). For example, Steunenberg and colleagues found that individuals with above-median levels of dispositional negativity were 2.8-times more likely to relapse or experience a new depressive episode across a six-year follow-up period (Steunenberg, Beekman, Deeg, & Kerkhof, 2010). Among parents, higher levels of dispositional negativity are also associated with elevated clinician and teacher ratings of internalizing symptoms in their offspring (Ellenbogen & Hodgins, 2004). Determining the biological and psychological mechanisms underlying this intergenerational transmission of psychopathology remains an important challenge for future research.

Given this panoply of adverse, often co-morbid outcomes, dispositional negativity imposes a tremendous burden on healthcare providers and the global economy (Goodwin, Hoven, Lyons, & Stein, 2002; ten Have, Oldehinkel, Vollebergh, & Ormel, 2005). A recent Dutch study estimated that each individual in the upper-quartile of the dispositional negativity distribution is associated with \$6,362 in excess costs each year, largely due to increased use of health services and loss of productivity (in 2015 dollars; Cuijpers et al., 2010). In a population the size of the United States, this would translate to \$388 billion annually or nearly one-third of federal discretionary spending.

Trait-State Links Inferred from Self-Report and Behavior

Despite its profound significance for health and wealth, the processes linking individual differences in dispositional negativity (trait) to heightened negative affect (state) have only recently started to come into focus. As shown schematically in **Figure 1**, self-report data reported over the past several decades suggest that three inter-related processes explain most of the heightened negative affect characteristic of individuals with a more negative disposition:

1. **Increased stressor reactivity:** Dispositionally negative individuals report elevated negative affect in response to a range of stressors, including negative life events, daily hassles, interpersonal conflicts, and aversive laboratory challenges.
2. **Increased negative affect in the absence of clear stressors:** Dispositionally negative individuals frequently report exaggerated apprehension and distress in relaxed and familiar settings, when potential stressors are remote, diffuse, or altogether absent. This pervasive, context-independent negative affect has been described as a ‘tonic’ or ‘endogenous’ effect of

temperament, given the absence of a clear external source of distress (Gross, Sutton, & Ketelaar, 1998; Watson & Clark, 1984).

3. **Increased stressor exposure and generation:** Individuals with a more negative disposition tend to act in ways that increase the likelihood of experiencing hassles, conflict, and rejection, particularly during times of heightened stress. Increased exposure to stressors, in turn, promotes more frequent, intense, or persistent negative affect.

Increased stressor reactivity

Individuals with a more negative disposition are emotionally volatile and tend to over-react to novelty, threat, and other stressors. In one particularly compelling example, Hengartner and colleagues assessed reactions to an ‘active shooter’ incident that recently occurred on the campus of a Swiss university (Hengartner, van der Linden, Bohleber, & von Wyl, 2016). During the incident, an alarm sounded continuously while more than 100 heavily armed police officers secured the site. Three hours later, the ‘all clear’ signal was given, and students and staff were allowed to leave their shelters. On-line surveys revealed that individuals with a more negative disposition retrospectively (6-26 days later) reported experiencing elevated levels of fear, worry, and terror during the incident.

Experience-sampling studies show that individuals with high levels of dispositional negativity also tend to report elevated levels of negative affect in response to more mundane hassles and interpersonal conflicts in the home, school, and workplace (Bolger & Schilling, 1991; Gable et al., 2000; Komulainen et al., 2014; Leger, Charles, Turiano, & Almeida, *in press*; Mroczek & Almeida, 2004; Suls & Martin, 2005; Tan et al., 2012; Zautra et al., 2005). Heightened reactivity to everyday stressors, in turn, predicts the onset of future internalizing symptoms and episodes, characterizes patients with acute anxiety disorders and

depression, and remits with pharmacological therapy (Farmer & Kashdan, 2015; Tan et al., 2012; van Winkel et al., 2015; Wichers et al., 2009).

In the laboratory, individuals with a more negative disposition report elevated distress in response to standardized aversive challenges (e.g., amputation film clips; Gross et al., 1998; Matthews et al., 2009), suggesting that heightened emotional reactivity is not an artifact of systematic response biases, mnemonic distortions, or differences in stressor exposure. These self-report data are consistent with evidence that dispositionally negative children, adults, and monkeys show exaggerated behavioral (e.g., avoidance, crying, inhibition), psychophysiological (e.g., startle, skin conductance), and neuroendocrine (e.g., cortisol) reactions to novelty and potential threat (Brooker et al., *in press*; Buss et al., 2003a; Hengartner, van der Linden, et al., 2016; Kagan, Snidman, Kahn, & Towsley, 2007; Norris, Larsen, & Cacioppo, 2007; Oler et al., 2016; Schmidt & Fox, 1998; Shackman et al., 2013; Vaidyanathan, Patrick, & Cuthbert, 2009).

Taken together, these observations indicate that dispositional negativity represents a diathesis that serves to enhance the likelihood, intensity, or duration of negative affect elicited by a range of common stressors. Heightened stressor reactivity also appears to causally contribute to the development and recurrence of pathological anxiety and depression.

Increased negative affect in the absence of clear stressors

Dispositionally negative individuals often report heightened negative affect in the absence of clear and imminent stressors (Watson & Clark, 1984). In controlled laboratory settings, dispositionally negative adolescents and adults report more intense or frequent negative thoughts and feelings at 'baseline,' while viewing emotionally-neutral control stimuli or simply relaxing (Craske et al., 2009; Glue, Wilson,

Coupland, Ball, & Nutt, 1995; Gross et al., 1998; Larsen & Ketelaar, 1989, 1991; Watson & Clark, 1984). Likewise, children with a more negative disposition show elevated heart rate at 'baseline' (Reznick et al., 1986). In their daily lives, dispositionally negative adults report elevated negative affect in comfortable, familiar settings, such as their home (Suls & Martin, 2005). In a seminal study, Bolger and Schilling (1991) leveraged 6 weeks of daily reports collected from more than 300 individuals to show that individuals with a more negative disposition report elevated distress in daily life. Next, they used statistical decomposition techniques (i.e., hierarchical linear modeling) to show that nearly 60% of this effect reflects 'tonic' differences in distress, in settings where their subjects did not report a clear concurrent source of stress, more than double the variance attributable to either stressor reactivity or stressor exposure.

Heightened negative affect in the absence of clear, exogenous stressors may reflect dispositionally negative individuals' tendency to experience spill-over of negative affect across sequential moments, contexts, or days (Houben et al., 2015; Judge, Simon, Hurst, & Kelley, 2014; Koval & Kuppens, 2012; Suls & Martin, 2005). Data from several large U.S. studies ($n > 1,000$) show that individuals with heightened levels of dispositional negativity tend to carry negative affect from work to home and vice versa (Horwitz, Luong, & Charles, 2008; Wayne et al., 2004). Spill-over and emotional inertia has also been observed in patients with internalizing disorders (Houben et al., 2015; Newman & Fisher, 2013; Peeters, Nicolson, Berkhof, Delespaul, & deVries, 2003). Among patients, spill-over and inertia predict the severity of symptoms (Brose, Schmiedek, Koval, & Kuppens, 2015; Houben et al., 2015; Koval, Kuppens, Allen, & Sheeber, 2012; Newman & Fisher, 2013), foreshadow future episodes (van de Leemput et al., 2014), improves with treatment (Newman & Fisher, 2013), and predict treatment response (Newman & Fisher, 2013). This may reflect maladaptive emotion regulation. Individuals with a more negative disposition are prone to worrying about the future and ruminating about the past (Grupe & Nitschke, 2013; Nolan, Roberts, & Gotlib, 1998; Wupperman & Neumann, 2006) and these maladaptive coping strategies tend

to promote pervasive negative affect in otherwise quiescent settings (Barlow et al., 2013; Nolen-Hoeksema, Wisco, & Lyubomirsky, 2008).

In short, several decades of self-report research demonstrates that persistently elevated, context-independent negative affect is a key feature of dispositional negativity. More recently acquired experience-sampling data suggest that spill-over and inertia of negative affect play a role in the development and recurrence of internalizing disorders.

Increased stressor exposure and generation

Like the anxiety disorders (American Psychiatric Association, 2013), dispositional negativity is thought to be associated with heightened avoidance of punishment and potential threat (Barlow et al., 2013; Cavanagh & Shackman, 2015; Gray & McNaughton, 2000; Grupe & Nitschke, 2013; Hengartner, van der Linden, et al., 2016). Yet there is also compelling evidence that dispositionally negative individuals are more frequently exposed to hassles, stressors, and more severe kinds of adversity. In fact, converging lines of prospective-longitudinal (Carter, Garber, Ciesla, & Cole, 2006; Fergusson & Horwood, 1987; Headey & Wearing, 1989; Kercher, Rapee, & Schniering, 2009; Lakdawalla & Hankin, 2008; Ludtke et al., 2011; Magnus, Diener, Fujita, & Pavot, 1993; Ormel & Wohlfarth, 1991; Specht, Egloff, & Schmukle, 2011; van Os, Park, & Jones, 2001; Vollrath, 2000; Wetter & Hankin, 2009; Whittington & Huppert, 1998), behavioral-genetic (Billig, Hershberger, Iacono, & McGue, 1996; Kandler, Bleidorn, Riemann, Angleitner, & Spinath, 2012; Middeldorp, Cath, Beem, Willemsen, & Boomsma, 2008; Power et al., 2013; Saudino, Pedersen, Lichtenstein, McClearn, & Plomin, 1997), and daily-diary data (Berry, Willingham, & Thayer, 2000; Bolger & Schilling, 1991; Bolger & Zuckerman, 1995; J. P. David, Green, Martin, & Suls, 1997; Gunthert, Cohen, & Armeli, 1999; Hankin, 2010; Hankin, Fraley, & Abela, 2005; Leger et al., *in press*; Marco & Suls, 1993; Suls, Green, & Hillis, 1998) acquired from more than 50,000 individuals in the U.S.

and abroad, some followed for as long as 16 years, demonstrate that adolescents and adults with high levels of dispositional negativity report more frequent personal difficulties and conflicts, particularly those of an interpersonal nature. Similar effects have been reported in children and adults with internalizing disorders (J. L. Allen & Rapee, 2009; Farmer & Kashdan, 2015; Hoehn-Saric, McLeod, Funderburk, & Kowalski, 2004; Kendler & Karkowski-Shuman, 1997; Kendler, Karkowski, & Prescott, 1999; Liu & Alloy, 2010; Liu, Kraines, Massing-Schaffer, & Alloy, 2014).

Increased stressor exposure has been observed using subjective ratings and more objective experimenter ratings of stressor intensity (Gleason, Powers, & Oltmanns, 2012; Iacovino et al., 2016; Jeronimus et al., 2014). For example, data from the Virginia Twin Registry ($n > 7,000$) indicates that dispositional negativity strongly predicts job loss, marital problems, and conflicts with family and co-workers, but is unrelated to random misfortunes (e.g., robbery; Kendler, Gardner, & Prescott, 2003). Analyses of nearly 1,000 datasets from the Dunedin longitudinal sample show that individuals with a more negative disposition at age 18 report progressively higher levels of conflict and abuse in romantic relationships across young adulthood (Robins et al., 2002). Using digital audio recorders, Mehl and colleagues showed that individuals with higher levels of dispositional negativity were more likely to argue in their daily lives (Mehl, Gosling, & Pennebaker, 2006). Leveraging survey data collected from 900 AmeriCorps volunteers pseudo-randomly assigned to 100 teams, Klein and colleagues (2014) demonstrated that levels of dispositional negativity at the time of employment strongly influenced the formation of spontaneous social networks 10 months later. Individuals with a more negative disposition tended to become central to what they termed 'adversarial networks' (i.e., are disliked by many teammates) and to show low centrality in both friendship and advice networks (Klein, Lim, Saltz, & Mayer, 2004). Among children, elevated levels of dispositional negativity are prospectively associated with peer rejection, social exclusion, victimization, and reduced friendship quality (Coplan, Arbeau, & Armer, 2007; Gazelle, 2008; Gazelle & Ladd, 2003; Ladd, Kochenderfer-Ladd, Eggum, Kochel, & McConnell, 2011; K. Rubin, Bowker, &

Gazelle, 2010; K. H. Rubin, Wojslawowicz, Rose-Krasnor, Booth-LaForce, & Burgess, 2006; Strauss, Frame, & Forehand, 1987). In short, there is compelling evidence that elevated levels of dispositional negativity are associated with increased exposure to a range of psycho-social stressors across the lifespan.

Other work suggests that dispositionally negative individuals play an instrumental role in generating stressors; that they tend to act in ways that increase the likelihood or chronicity of negative life events (e.g., divorce, financial difficulties) and daily hassles (e.g., interpersonal conflict, social rejection). In adulthood, their friends report more frequent conflict and heightened irritation (Berry et al., 2000), their romantic partners report reduced relationship security (Neyer & Voigt, 2004), their spouses report reduced marital and sexual satisfaction (Dyrenforth et al., 2010; Malouff et al., 2010; Solomon & Jackson, 2014; Watson, Hubbard, & Wiese, 2000; Watson & Humrichouse, 2006), and their offspring report more frequent parent-child conflict (Hutteman, Bleidorn, Kereste, et al., 2014). Likewise, the parents of children with a more negative disposition describe their relationship with their offspring as challenging and emotionally exhausting (Shamir-Essakow, Ungerer, Rapee, & Safier, 2004). Similar effects have been reported for the parents of children with anxiety disorders (Lebowitz, Scharfstein, & Jones, 2014; Lebowitz et al., 2013).

These kinds of informant reports are complemented by laboratory studies showing that randomly assigned social partners judge dispositionally negative adults to be moody, uncomfortable, and negative (Creed & Funder, 1998). This negativity begets negativity and random partners tend to respond with elevated levels of criticism, contempt, and hostility (Creed & Funder, 1998) and to judge the interaction more negatively (Heerey & Kring, 2007). Likewise, dispositionally negative children tend to evoke more negative reactions from unfamiliar peers (Stewart & Rubin, 1995; O. L. Walker, Degnan, Fox, & Henderson, *in press*).

Heightened interpersonal stress and social rejection may stem from dispositionally negative individuals' tendency to express lower levels of warmth and empathy; to be less responsive and disclosing to relationship partners; to overreact and escalate negative affect during conflicts (e.g., angry venting, hostile or aggressive confrontation); and to engage in toxic interpersonal behaviors (i.e., criticism, contempt, and sarcasm), particularly during periods of heightened stress (Ackerman & Corretti, 2015; L. A. Clark, Kochanska, & Ready, 2000; Connor-Smith & Flachsbart, 2007; de Haan, Dekovic, & Prinzie, 2012; Donnellan, Conger, & Bryant, 2004; Ellenbogen & Hodgins, 2004; Kendler & Karkowski-Shuman, 1997; Kochanska, Clark, & Goldman, 1997; McNulty, 2008; Neyer & Asendorpf, 2001; Prinzie et al., 2009; Romero-Canyas, Downey, Berenson, Ayduk, & J., 2010; Vater & Schröder-Abé, 2015; Wang, Repetti, & Campos, 2011)⁴. Likewise, individuals with anxiety disorders are prone to more frequent and intense conflict with their partners and spouses (Johnson, Cohen, Kasen, & Brook, 2004; Metz, Majdandzic, & Bogels, *in press*).

Studies of patients with social phobia in semi-structured 'getting acquainted' tasks suggests that these and other maladaptive expressive behaviors elicit negative affect in others which, in turn, promotes discord, alienation, and rejection (Alden & Taylor, 2004; Plasencia, Alden, & Taylor, 2011). Similar results have been found in more naturalistic observational and experience-sampling studies (Pasch, Bradbury, & Davila, 1997; Zaider, Heimberg, & Iida, 2010). Interventions targeting these maladaptive behaviors reduce conflict and rejection, indicating a causal role (Snyder & Halford, 2012; Taylor & Alden, 2011).

⁴ Drawing on a unique body of anthropological research conducted in subject's own homes, Wang and colleagues provide a vivid description of this negative interpersonal style: "Ed Anderson...scored high on...[dispositional negativity] and reported high job stress. He is married to Rhoda and together they have three young daughters...On Day 1 of videotaping, Ed returned home and was immediately very engaged with his family members...However, he appeared somewhat irritated during these interactions with Rhoda and his daughters, and often sighed, rubbed his head in annoyance, and used a mildly sarcastic tone. At the start of dinner, Ed learned from an apologetic Rhoda that the fish she had made would not be ready for another 20 minutes. Although Ed said "OK, I'll eat it later," he made several references to the dinner that showed a critical undertone...Several minutes into the meal, Ed silently left the table and began loudly crunching on chips. Rhoda continued to be apologetic about the delay with the fish, but Ed told her "I'm not even really hungry"...Although Ed continued to inquire when the fish would be ready, he ultimately rejected the fish when it came out of the oven. After dinner, Ed continued to show irritation (e.g., raised eyebrows, annoyed looks, sighs, sarcastic tone)" (Wang et al., 2011, p. 451).

In sum, individuals with a more negative disposition and patients with internalizing disorders actively, if unintentionally, shape their environment in ways that generate stress. Increased exposure to hassles, conflict, and other, more severe psycho-social stressors (e.g., divorce), in turn, tends to promote more intense or pervasive negative affect.

The Psychophysiology and Neurobiology of Dispositional Negativity

The self-report and behavioral data that we have reviewed suggest that the link between dispositional negativity and heightened levels of momentary negative affect reflects a combination of increased stressor reactivity, tonic or ‘endogenous’ increases in negative affect, and increased stressor exposure (**Figure 1**). As described in more detail below, brain imaging, neuropsychological, and more mechanistic kinds of data gleaned from animal models strongly corroborate the link binding dispositional negativity to heightened stressor reactivity. But biological data raise the possibility that seemingly ‘tonic’ increases in negative affect may actually reflect increased reactivity⁵ to stressors that are mild, remote, uncertain, or diffuse (**Figure 2**). This is likely to be exacerbated by stress-induced sensitization of brain regions, such as the amygdala, that play a key role in assembling states of fear and anxiety⁶. While the neurobiological mechanisms underlying increased stressor generation and exposure remain largely opaque, the existing neurobiological record suggests that the tendency to behave in ways that promote

⁵ This is similar to what others have conceptualized as a reduced threshold for responding (Davidson, 1998; Reznick et al., 1986).

⁶ A growing number of researchers draw a sharp distinction between states of ‘fear’ and ‘anxiety’ (e.g., Barlow, 2000; D. C. Blanchard & Pearson, *in press*; M. Davis, Walker, Miles, & Grillon, 2010; LeDoux, 2015). Yet lay people, scholars in other areas, the American Psychiatric Association’s Diagnostic and Statistical Manual (American Psychiatric Association, 2013), and even domain experts—at least in unguarded moments—often use these terms interchangeably or inconsistently. As one psychiatrist noted almost 40 years ago, “The word ‘anxiety’ has become confused. It has so many meanings in so many languages, that...it has come to be a synonym for the generic term ‘fear’” (Gaylin, 1979, p. 18). Other commentators have emphasized the difficulty of drawing sharp operational boundaries between the terms (Perusini & Fanselow, 2015). To avoid misunderstanding, we recommend that researchers eschew these problematic redefinitions of everyday language (Shackman & Fox, *in press*).

interpersonal conflict and evoke social rejection may reflect variation in the function of circuits that underlie the appraisal of emotionally-salient social cues.

Increased reactivity to aversive challenges in the laboratory

Decades ago, Gordon Allport suggested that “traits are cortical [or] subcortical...dispositions having the capacity to gate or guide specific phasic reactions” (1966). To this day, most neurobiologically grounded models of dispositional negativity remain rooted in the idea that temperament and personality reflect differences in the magnitude or likelihood of reactions to punctate, trait-relevant challenges (e.g., conflict, criticism, punishment, danger; Eysenck, 1967; Goldsmith et al., 1987; Kagan, Reznick, & Snidman, 1988; Reiss, 1997; Spielberger, 1966; Zuckerman, 1976)⁷.

Consistent with this perspective, there is clear evidence that humans and monkeys with a more negative disposition show heightened reactions to threat-related cues in a number of brain regions, including the amygdala, anterior hippocampus, anterior insula, bed nucleus of the stria terminalis (BST), mid-cingulate cortex, orbitofrontal cortex (OFC), and periaqueductal gray (PAG) (Avery et al., 2016; Calder et al., 2011; Cavanagh & Shackman, 2015; A. S. Fox & Kalin, 2014; A. S. Fox, Oler, Shackman, et al., 2015; A. S. Fox, Oler, Tromp, et al., 2015; Shackman & Fox, *in press*; Shackman et al., 2011).

Here, we focus on the most intensively scrutinized of these regions, the amygdala. The amygdala is a heterogeneous collection of nuclei buried beneath the temporal lobe (Freese & Amaral, 2009; Yilmazer-Hanke, 2012). As shown in **Figure 3**, the amygdala is poised to assemble a broad spectrum of emotional reactions via projections to the downstream regions that directly mediate the behavioral (e.g., passive and active avoidance), peripheral physiological (e.g., cardiovascular and neuroendocrine activity, startle),

⁷ The revised Reinforcement Sensitivity Theory is one notable exception (Corr, 2008; Gray & McNaughton, 2000; Reuter, Cooper, Smillie, Markett, & Montag, 2015).

and cognitive (e.g., vigilance) components of momentary negative affect (M. Davis & Whalen, 2001; Freese & Amaral, 2009). Lesion, imaging, and electrophysiological evidence demonstrate that the amygdala can trigger shifts of attention to threat-relevant social cues (e.g., eyes) and that reentrant projections from the basolateral (BL) nucleus of the amygdala to the visual cortex and superior colliculus play a crucial role in prioritizing the processing of threat-relevant cues (Shackman, Kaplan, et al., *in press*). The amygdala is also poised to promote non-specific states of vigilance via projections to ascending neurochemical systems (i.e., acetylcholine, dopamine, norepinephrine) in the basal forebrain and brainstem that can modulate the responsiveness of sensory cortex to incoming information (Arnsten, 2009, 2015; M. Davis & Whalen, 2001; Freese & Amaral, 2009).

Imaging studies show that dispositionally negative individuals show increased or prolonged activation in the dorsal amygdala in response to novelty or potential threat (Blackford, Avery, Shelton, & Zald, 2009; Calder et al., 2011; Fonzo et al., 2015; A. S. Fox & Kalin, 2014; Schuyler et al., 2012; Stein, Simmons, Feinstein, & Paulus, 2007) (**Figures 4a-b**). This is particularly evident following periods of acute stress (Everaerd, Klumpers, van Wingen, Tendolkar, & Fernandez, 2015). Amygdala reactivity also tends to habituate more slowly in adults and youth with a more negative disposition (Blackford, Allen, Cowan, & Avery, 2013; Blackford, Avery, Cowan, Shelton, & Zald, 2011; Hare et al., 2008).

Large-scale ($n = 238-592$) positron emission tomography (PET) studies in juvenile monkeys show that threat-related metabolic activity in the dorsal amygdala, in the region of the central (Ce) nucleus (**Figure 4c**), is stable over time and context (i.e., trait-like), heritable, and associated with heightened behavioral and neuroendocrine reactions to threat (A. S. Fox & Kalin, 2014; A. S. Fox, Oler, Shackman, et al., 2015; A. S. Fox et al., 2012; Oler et al., 2010; Shackman et al., 2013). For example, Fox and colleagues reported that metabolic activity in the Ce during prolonged exposure to an unfamiliar human intruder's profile (**Figure 2b**) showed an intra-class correlation (ICC) of 0.64 across three occasions over a 1.1 year span, similar to

the concurrent re-test stability of dispositional negativity in young monkeys (ICC = 0.72; A. S. Fox et al., 2012) and the 5-year stability of dispositional negativity in humans (partial $R = .60$; $n = 56,735$; Hakulinen, Elovainio, et al., 2015). Other work in nonhuman primates demonstrates that elevated amygdala activity is a core substrate for different presentations of dispositional negativity (**Figure 5**). Like humans, monkeys express dispositional negativity in different ways. Some individuals characteristically respond to potential threat with high levels of the stress-sensitive hormone cortisol and middling levels of behavioral inhibition, whereas others show the reverse profile. What all of these individuals share is heightened threat-related activity in the Ce (Shackman et al., 2013). Collectively, these observations show that individual differences in dispositional negativity partially reflect trait-like variation in the function of the amygdala.

The amygdala's contribution to dispositional negativity appears to be causal. In monkeys and rodents, selective lesions of the amygdala, particularly the Ce, markedly reduce the expression of fear and anxiety elicited by a broad spectrum of learned and innate threats (Calhoun & Tye, 2015; J. S. Choi & Kim, 2010; Izquierdo, Suda, & Murray, 2005; Janak & Tye, 2015; Kalin et al., *in press*; Kalin, Shelton, & Davidson, 2004; Mason, Capitanio, Machado, Mendoza, & Amaral, 2006; Tovote, Fadok, & Luthi, 2015). These experimental findings in animals are consistent with observations made in humans with circumscribed amygdala damage (Adolphs, *in press*; Feinstein, Adolphs, Damasio, & Tranel, 2011; Klumpers, Morgan, Terburg, Stein, & van Honk, *in press*). Patient SM, who has near-complete bilateral destruction of the amygdala, shows a profound lack of negative affect when exposed to frightening movies, haunted houses, tarantulas, snakes, and even real-world assaults (Feinstein et al., 2011)⁸. Importantly, she also reports abnormally low levels of dispositional negativity on standardized psychometric measures (Feinstein et

⁸ Over the past two decades, "She has been held up at knife point and at gun point, she was once physically accosted by a woman twice her size, she was nearly killed in an act of domestic violence, and on more than one occasion she has been explicitly threatened with death...What stands out most is that, in many of these situations, SM's life was in danger, yet her behavior lacked any sense of desperation or urgency." (Feinstein et al., 2011, p. 307).

al., 2011), consistent with informal clinician ratings of temperament (Tranel, Gullickson, Koch, & Adolphs, 2006).

Other work suggests that elevated amygdala reactivity contributes to the development of pathological anxiety and depression. Activity in the amygdala co-varies with changes in threat-elicited peripheral physiology (e.g., startle potentiation, skin conductance) and self-reported arousal (Cheng, Knight, Smith, & Helmstetter, 2006; Cheng, Richards, & Helmstetter, 2007; Knight, Nguyen, & Bandettini, 2005; Kragel & LaBar, 2015; LaBar, Gatenby, Gore, LeDoux, & Phelps, 1998; van Well, Visser, Scholte, & Kindt, 2012; Wood, Ver Hoef, & Knight, 2014). Amygdala reactivity is amplified by exposure to the same kinds of stressors and psychological pathogens that can precipitate acute mental illness, including combat and childhood maltreatment (Dannlowski et al., 2012; Seo, Tsou, Ansell, Potenza, & Sinha, 2014; Swartz, Williamson, & Hariri, 2015; van Wingen, Geuze, Vermetten, & Fernandez, 2011), and predicts the development of internalizing symptoms following exposure to stress or trauma (Admon et al., 2009; McLaughlin et al., 2014; Swartz, Knodt, Radtke, & Hariri, 2015). Amygdala reactivity is elevated in children and adults diagnosed with internalizing disorders (Etkin & Wager, 2007; Hamilton et al., 2012; Thomas et al., 2001) and is reduced by clinically effective treatments for anxiety and depression (Arce, Simmons, Lovero, Stein, & Paulus, 2008; Brown et al., 2015; Felmingham et al., 2007; Furmark et al., 2002; Harmer, Mackay, Reid, Cowen, & Goodwin, 2006; Paulus, Feinstein, Castillo, Simmons, & Stein, 2005; Phan et al., 2013; Sheline et al., 2001; Windischberger et al., 2010).

In sum, converging lines of epidemiological, physiological, and mechanistic evidence indicate that dispositionally negative individuals' exaggerated reactivity to threat and other aversive challenges partially reflects larger or longer-lasting responses in the dorsal amygdala. Like dispositional negativity, individual differences in amygdala reactivity are trait-like and elevated levels of reactivity predict the future development of internalizing symptoms among individuals exposed to stress.

Trait-like individual differences in reactivity are discernible at 'rest'

Although most neurobiological research has focused on reactivity to acute stressors and threat-related cues—aversive cues, faces, images, films and so on—individual differences in threat-reactivity can also be discerned in the brain's spontaneous or 'resting' activity. For example, monkeys with elevated metabolic activity in the amygdala at 'baseline'—in their home-cage with a familiar cage-mate—show increased freezing and elevated levels of cortisol when threat is encountered in other contexts (A. S. Fox, Shelton, Oakes, Davidson, & Kalin, 2008). Likewise, humans with higher levels of dispositional negativity show heightened activity in the amygdala at rest, as indexed by 18-fluorodeoxyglucose PET (FDG-PET) or perfusion functional magnetic resonance imaging (fMRI; Abercrombie et al., 1998; Canli et al., 2006; Kaczurkin et al., *in press*).

Monkeys, children, and adults with a more negative disposition also show greater electroencephalographic (EEG) activity over the right compared to the left prefrontal cortex (PFC) at 'rest' (**Figure 6a**) (Buss et al., 2003b; Davidson, Jackson, & Kalin, 2000; N. A. Fox, Henderson, Marshall, Nichols, & Ghera, 2005; Kalin, Larson, Shelton, & Davidson, 1998; Oler et al., 2016; Shackman, McMenamin, Maxwell, Greischar, & Davidson, 2009; Wacker et al., 2010). Like dispositional negativity and amygdala reactivity, individual differences in resting prefrontal EEG asymmetry first emerge early in life and are relatively stable over time, heritable, and predict the intensity of negative affect elicited by aversive laboratory challenges (Buss et al., 2003b; Davidson et al., 2000; N. A. Fox et al., 2005; Kalin et al., 1998; Smit, Posthuma, Boomsma, & De Geus, 2007; Tomarken, Davidson, Wheeler, & Kinney, 1992; Towers & Allen, 2009; Wheeler, Davidson, & Tomarken, 1993).

Like amygdala reactivity, individual differences in prefrontal EEG asymmetry also confer increased risk for the development of internalizing disorders. This asymmetric pattern of 'resting' activity prospectively predicts the first-onset of mood disorders (Nusslock et al., 2011), is exaggerated in patients with internalizing disorders (Thibodeau et al., 2006), and is normalized by anxiolytic drugs (Davidson, Kalin, & Shelton, 1993; Davidson, Kalin, & Shelton, 1992). Furthermore, neurofeedback interventions targeting this pattern of scalp electrical activity cause lasting reductions in stress reactivity (J. J. Allen, Harmon-Jones, & Cavender, 2001). Along with the pharmacological evidence, this suggests that the neural circuit or circuits responsible for generating this marker make a causal contribution to individual differences in dispositional negativity. Recent efforts to pinpoint the source of the scalp-recorded EEG asymmetry have highlighted the importance of the dorsolateral prefrontal cortex (dlPFC; Shackman et al., 2009) (**Figure 6b**), consistent with this region's role in regulating momentary negative affect (Buhle et al., 2014).

Other work by our group suggests that the dlPFC and Ce form a coherent, evolutionarily-conserved functional circuit. Reduced functional connectivity between the two regions is associated with pathological anxiety in children and heightened dispositional negativity in monkeys (Birn et al., 2014) (**Figure 6c**). Selective lesions of the amygdala, including the Ce, are associated with chronically reduced metabolism in the dlPFC (Machado, Snyder, Cherry, Lavenex, & Amaral, 2008), reinforcing the idea that these regions represent an integrated functional circuit. A key challenge for future studies will be to use mechanistic techniques to clarify the functional architecture and causal contribution of this circuit to extreme anxiety.

In sum, individuals with a more negative disposition, who are prone to hyper-react to potential threat, show altered activity in the amygdala and dlPFC in the absence of explicit threat. These observations suggest that variation in the basal activity and functional connectivity of these regions represents a diathesis for heightened negative affect, given an appropriate trait-relevant challenge. More broadly, they

indicate that reactive features of dispositional negativity can be discerned in the spontaneous, on-going activity of the brain, even under sedation (Birn et al., 2014).

Altered ‘resting’ activity—States or traits, reactive or tonic differences?

At present, it remains unclear whether alterations in ‘resting’ activity reflect heightened reactivity to the experimental setting, which often entails a mixture of acute and diffuse threats; ‘tonic’ differences in negative affect (Gross et al., 1998; Watson & Clark, 1984); or more likely some combination of the two processes. It is clear that most neurophysiological assays are intrusive and can elicit substantial negative affect. FDG-PET requires the injection of a radiotracer. EEG and PET studies in monkeys entail unexpected separation from cage-mates and manual restraint. In humans, resting EEG procedures have been shown to increase negative affect (Blackhart, Kline, Donohue, LaRowe, & Joiner, 2002). MRI data collection requires subjects to lie motionless in a spatially-confined, often dark tube while being bombarded by loud noise for tens of minutes. Not surprisingly, MRI procedures have been shown to elicit feelings ranging from mild apprehension to severe panic, to increase cortisol levels, and to activate the sympathetic nervous system in children and adults (Eatough, Shirtcliff, Hanson, & Pollak, 2009; Lueken, Muehlhan, Evens, Wittchen, & Kirschbaum, 2012; Melendez & McCrank, 1993; Muehlhan et al., 2013; Törnqvist, Månsson, Larsson, & Hallström, 2006; Tyc, Fairclough, Fletcher, Leigh, & Mulhern, 1995). Furthermore, MRI-induced negative affect is amplified in individuals with a more negative disposition (Harris, Cumming, & Menzies, 2004; Harris, Robinson, & Menzies, 2001), even with prior acclimation in a mock scanner (Shechner et al., 2013). Psychometric analyses (i.e., latent state-trait or generalizability-theory models) indicate that resting-state prefrontal EEG asymmetry reflects the joint contribution of traits and states, in about equal measure (Coan, Allen, & McKnight, 2006; Hagemann, Hewig, Seifert, Naumann, & Bartussek, 2005; Hagemann, Naumann, Thayer, & Bartussek, 2002; Tomarken et al., 1992).

But as yet, the relative contribution of traits and states to PET and fMRI measures of brain function remains unknown, making this another important avenue for future research.

'Tonic' increases in self-reported negative affect may reflect heightened reactivity to uncertain or diffuse threat

Questionnaire data collected in the laboratory and in the field indicate that dispositionally negative individuals often report heightened negative affect in the absence of clear and immediate external stressors. Although this may reflect a 'tonic' or direct effect of temperament on mood (Watson & Clark, 1984), a wealth of psychophysiological and behavioral evidence suggests that persistent negative affect may reflect increased reactivity to stressors that are mild, uncertain, remote, or diffuse (D. M. Clark, 2001; M. Davis et al., 2010; Grupe & Nitschke, 2013).

For example, children, adolescents, and adults with a more negative disposition show exaggerated psychophysiological responses (e.g., startle, skin conductance) and report heightened negative affect during periods of explicit safety (i.e., CS-, inter-cue interval, CS+ paired with a safety cue) embedded within instructed and associative fear learning paradigms—that is, during the interstitial periods before and after the randomized presentation of genuine threat (CS+) (Baas, van Ooijen, Goudriaan, & Kenemans, 2008; Barker, Reeb-Sutherland, & Fox, 2014, *in press*; Chan & Lovibond, 1996; Craske et al., 2009; Gazendam et al., 2015; Gazendam, Kamphuis, & Kindt, 2013; Grillon, 2002; Grillon & Ameli, 2001; Haaker et al., 2015; Jovanovic et al., 2014; Reeb-Sutherland et al., 2009; Schmidt & Fox, 1998). Conceptually similar effects have been found in monkeys (Shackman, Fox, et al., *in press*; Shiba et al., 2014) and in patients with anxiety disorders (Duits et al., 2015). In fact, a comprehensive recent meta-analysis incorporating data from more than 2,000 individuals showed that patients consistently respond

more strongly than controls to safety cues (CS-: Cohen's $d = .30, p < .001$), whereas the two groups do not consistently differ in their response to acute threat (CS+: Cohen's $d = .07, p = .41$) (Duits et al., 2015).

Research using more naturalistic challenges in children also highlights the importance of contextually inappropriate negative affect. For example, dispositionally negative children show elevated psychophysiological defensive responses to neutral faces (Waters, Neumann, Henry, Craske, & Ornitz, 2008). Relative to typical children, two-year-olds with an extremely negative disposition show only modestly elevated negative affect in response to high-threat challenges (e.g., approaching robotic spider: Typical Children: ~32% time; Negative Children: ~45% time), whereas group differences are dramatically larger during interspersed low-threat challenges (e.g., puppet show: Typical Children: ~28%; Negative Children: ~65% time; Buss, 2011). Moreover, contextually inappropriate negative affect in the laboratory at age 2 predicts elevated parent- and teacher-reports of anxiety in preschool and kindergarten and heightened wariness around unfamiliar peers at age 5 (Buss, 2011; Buss et al., 2013).

Among adults, indiscriminate or contextually inappropriate negative affect has also been associated with increased avoidance of the aversive learning context. Grillon (2002), for example, showed that subjects who were unable to correctly report the shock-cue contingency following the training phase of a simple differential fear conditioning paradigm (CS+ cue paired with noxious electric shock compared to an unpaired CS- cue) were nearly four times more likely to fail to return for a follow-up session ($n = 125$, risk ratio = 3.80, $p < .005$). Furthermore, well-established anxiolytic compounds, like ethyl alcohol and benzodiazepines, have been shown to dampen persistent negative affect elicited by uncertain threat in a dose-dependent manner, while sparing phasic reactions to cues associated with clear and imminent danger (Bradford, Shapiro, & Curtin, 2013; Glue et al., 1995; Grillon et al., 2006; Hefner & Curtin, 2012; Hefner, Jaber, Grant, & Curtin, 2009; Hefner, Moberg, Hachiya, & Curtin, 2013; Moberg & Curtin, 2009).

This body of psychophysiological and behavioral research motivates the hypothesis that seemingly endogenous increases in negative affect, as described in the self-report literature, may reflect heightened sensitivity to weak, distal, or uncertain stressors, rather than a fixed or ‘tonic’ consequence of temperament. At a more granular level, this may reflect difficulties discriminating threat from safety, overgeneralization of threat to perceptually similar safety cues (i.e., a broader tuning of threat-detection mechanisms and higher tolerance of false alarms; cf. Nettle & Bateson, 2012; Pollak & Kistler, 2002), or problems using or learning to use safety-related information to regulate momentary negative affect, leading to mood spillover or affective inertia (Davidson, Fox, & Kalin, 2007; Davidson et al., 2000; Grupe & Nitschke, 2013; Kheirbek, Klemenhagen, Sahay, & Hen, 2012; Lissek, 2012).

Mechanistic work in rodents (Calhoun & Tye, 2015; M. Davis et al., 2010; Gungor & Paré, *in press*; Tovote et al., 2015) suggests that defensive responses to diffuse, uncertain threats (**Figure 2a**) are organized by the central extended amygdala, an anatomical concept encompassing the Ce and the lateral division of BST (**magenta regions in Figure 3**) (Alheid & Heimer, 1988; A. S. Fox, Oler, Tromp, et al., 2015; Shackman & Fox, *in press*; Yilmazer-Hanke, 2012). Like humans with high levels of dispositional negativity, some rodents show poor discrimination of cues associated with danger and safety (i.e., elevated fear and anxiety in response to CS-; often termed ‘over-generalization’; Lissek, 2012) and heightened defensive responses during sustained exposure to diffusely threatening contexts (e.g., the cage paired with cued fear learning or the elevated plus-maze) (Duvarci, Bauer, & Paré, 2009). Selective lesions of the BST reduce these maladaptive emotional responses, while sparing more adaptive, phasic responses to cues signaling imminent danger (CS+) (Duvarci et al., 2009). Anatomically, both the Ce and BST are poised to orchestrate key features of sustained negative affect—including alterations in arousal, behavioral inhibition, and neuroendocrine activation—via dense mono- and poly-synaptic (e.g., via the medial division of the Ce) projections to brainstem and subcortical effector regions (M. Davis et al., 2010; M. Davis & Whalen, 2001; A. S. Fox, Oler, Tromp, et al., 2015; Freese & Amaral, 2009).

Recent imaging work in humans and monkeys demonstrates that individuals with a more negative disposition show heightened activity in the BST during prolonged periods of diffuse or uncertain threat (**Figures 2c and 7a, b**) (A. S. Fox et al., 2008; Shackman, Fox, et al., *in press*; Somerville, Whalen, & Kelley, 2010), with parallel effects reported for patients with anxiety disorders (Munsterkotter et al., *in press*; Straube, Mentzel, & Miltner, 2007; Yassa, Hazlett, Stark, & Hoehn-Saric, 2012). Work in nonhuman primates shows that threat-related metabolic activity in the BST is heritable and genetically correlated with individual differences in dispositional negativity, suggesting that it contributes to the inter-generational transmission of this dispositional phenotype (A. S. Fox, Oler, Shackman, et al., 2015) (**Figure 7b**). In humans and monkeys, BST activity and functional connectivity co-vary with self-reported negative affect, freezing, skin conductance, cardiovascular activity, and cortisol elicited by uncertain or diffuse threat (Alvarez et al., 2015; Banihashemi, Sheu, Midei, & Gianaros, 2015; Jahn et al., 2010; Kalin, Shelton, Fox, Oakes, & Davidson, 2005; McMenamin, Langeslag, Sirbu, Padmala, & Pessoa, 2014; Somerville et al., 2013), consistent with a causal contribution to momentary negative affect. At present, the consequences of selective BST lesions have yet to be unexplored in humans or nonhuman primates.

BST activity is often described as a ‘sustained’ response to uncertain threat. Yet, mechanistic work in rodents suggests that BST engagement could begin quite early, between 4 and 60 s following the onset of threat-related cues (M. Davis et al., 2010). Consistent with this, a number of human imaging studies have found transient activation in the BST in response to punctate threats (**Figure 7a**), such as a 4-sec video clip of an approaching tarantula (J. M. Choi, Padmala, & Pessoa, 2012; Grupe, Oathes, & Nitschke, 2013; Klumpers et al., 2015; Mobbs et al., 2010). Likewise, a recent large-scale imaging study ($n = 168$) reported phasic activation of the BST in response to 4-s shock-predictive cues (Klumpers et al., 2015). In fact, imaging studies of fear and anxiety consistently reveal activation in the central extended amygdala—including the BST and the dorsal amygdala in the region of the Ce—across a range of populations,

paradigms, and time-scales (**Figure 7c**). This and other recent work in humans and rodents highlights the importance of these two closely related regions across a broad spectrum of aversive challenges (Shackman & Fox, *in press*)⁹.

Individual differences in the function of the central extended amygdala may reflect altered communication with the orbitofrontal cortex (OFC) (**cf. “Regulatory/Evaluative” inputs in Figure 3**). Large-scale imaging studies in monkeys ($n = 592$) reveal that threat-related metabolic activity in the posterior OFC/anterior insula is heritable and, like the BST, genetically correlated with individual differences in dispositional negativity (A. S. Fox, Oler, Shackman, et al., 2015). Aspiration lesions of the OFC markedly reduce passive avoidance of potential threat (i.e., freezing) (Kalin, Shelton, & Davidson, 2007; Rudebeck, Saunders, Prescott, Chau, & Murray, 2013) and this appears to be mediated by downstream changes in BST metabolism (A. S. Fox et al., 2010). Reduced BST activity has also been found in humans with OFC damage (Motzkin, Philippi, Oler, et al., 2015), suggesting that this circuit is conserved across primate species. Interestingly, viral vector manipulations that increase metabolic activity in the Ce—the other major component of the central extended amygdala—are associated with elevated metabolic activity in the OFC, increased functional connectivity between the Ce and OFC, and heightened signs of fear and anxiety during prolonged exposure to threat (Kalin et al., *in press*). Conversely, Ce lesions are associated with reduced metabolic activity in the OFC (Machado et al., 2008). In other words, perturbations targeting one region (e.g., OFC or Ce damage) propagate to the others (e.g., reduced BST or

⁹ Inspired by the seminal work of Michael Davis, David Walker, and their colleagues (M. Davis, 1998, 2006; M. Davis, Walker, & Lee, 1997; M. Davis et al., 2010; Grillon, 2008; D. L. Walker & Davis, 2008; D. L. Walker, Miles, & Davis, 2009; D. L. Walker, Toufexis, & Davis, 2003), it is widely thought that phasic and sustained responses to threat reflect dissociable circuits centered on the Ce and the BST, respectively. Indeed, a version of this influential hypothesis has been incorporated into the National Institute of Mental Health (NIMH) Research Domain Criteria (RDoC) as Acute Threat (‘Fear’) and Potential Threat (‘Anxiety’) (cf. <https://www.nimh.nih.gov/research-priorities/rdoc/constructs/potential-threat-anxiety.shtml>; <https://www.nimh.nih.gov/research-priorities/rdoc/negative-valence-systems-workshop-proceedings.shtml>; Kozak & Cuthbert, 2016). But as described in greater detail elsewhere (Calhoun & Tye, 2015; Gungor & Paré, *in press*; Janak & Tye, 2015; Shackman & Fox, *in press*; Tovote et al., 2015), more recent observations in rodents, monkeys, and humans make it clear that the central extended amygdala forms a tightly interconnected functional unit and that claims of strict functional segregation are no longer tenable.

OFC metabolism) and damage to either the Ce or OFC reduces, but does not abolish, defensive responses to threat.

Collectively, these imaging and mechanistic findings suggest that the extended amygdala and OFC form a functionally integrated circuit that plays a crucial role in detecting and organizing persistent responses to uncertain threat¹⁰. Much remains unknown about this circuit, including the necessity of the primate BST to persistent negative affect, the differential contributions of its three constituents (Ce, BST, OFC), the nature of their interactions with one another and other brain regions associated with dispositional negativity (e.g., dlPFC, PAG), and the relevance of this circuit to persistent, contextually inappropriate negative affect in the real world.

'Tonic' increases in self-reported negative affect may reflect stress-induced sensitization

Self-report data indicate that individuals with a more negative disposition tend to carry negative affect from stressful to less stressful contexts (e.g., work to home) and to behave in ways that promote interpersonal conflict during times of heightened stress (Suls & Martin, 2005; Wang et al., 2011)⁴. Recent work in humans suggests that the amygdala could contribute to the spill-over of negative mood via stress-induced sensitization, consistent with models derived from animal research (Rosen & Schulkin, 1998). In particular, there is evidence that acute stressors (e.g., threat-of-shock, aversive film clips) potentiate defensive reactions (i.e., startle) to threat-related facial expressions (Grillon & Charney, 2011), cause persistent increases in spontaneous amygdala activity (Cousijn et al., 2010), and potentiate amygdala reactivity to threat-related faces (Pichon, Miendlarzewska, Eryilmaz, & Vuilleumier, 2015; van

¹⁰ Other mechanisms, including overgeneralization of threat to perceptually-similar safety cues or contexts (Camp et al., 2012; Ciochi et al., 2010; Ghosh & Chattarji, 2015; Kheirbek et al., 2012; Laufer, Israeli, & Paz, *in press*; Lissek et al., 2014; Shaban et al., 2006) and deficient filtering of threat-related information from working memory (Stout, Shackman, Johnson, & Larson, 2014; Stout, Shackman, & Larson, 2013), are also likely to be important. In fact, given the central role of working memory in goal-directed cognition (e.g., attention, imagination, memory retrieval), emotion regulation (e.g., worry), and behavior, aberrant gating of threat-related information from working memory storage may represent a key source of sustained apprehension and negative affect in humans (Mobbs, Hagan, Dalgleish, Silston, & Prevost, 2015; Stout et al., 2013).

Marle, Hermans, Qin, & Fernandez, 2009). Acute stressors produce even longer-lasting changes (i.e., minutes to hours) in amygdala functional connectivity (Vaisvaser et al., 2013; van Marle, Hermans, Qin, & Fernandez, 2010). Furthermore, these neurobiological spill-over effects are exaggerated in individuals with a more negative disposition. In particular, a recent large-scale imaging study ($n = 120$) showed that dispositionally negative individuals exhibit potentiated activation to threat-related faces following acute stressor exposure (Everaerd et al., 2015).

Persistent amygdala sensitization could promote negative affect either directly, by potentiating emotional reactions to mild threat (Grillon & Charney, 2011), or indirectly, by increasing the likelihood that attention will be allocated to threat-related cues in the surrounding environment. Circuits centered on the amygdala play a central role in allocating attentional resources to threat and a rapidly growing body of longitudinal (Perez-Edgar, Bar-Haim, et al., 2010; Perez-Edgar et al., 2011; White et al., *in press*) and clinical-intervention studies (Linetzky et al., 2015; MacLeod & Clarke, 2015) indicate that attentional biases to threat causally contribute to the development and maintenance of extreme anxiety and distress (Shackman, Kaplan, et al., *in press*). For example, computer-based interventions targeting attentional biases to threat have been shown to reduce self-reported negative affect, behavioral signs of anxiety, and intrusive thoughts elicited by aversive challenges, such as public speaking tasks (Dennis & O'Toole, 2014; MacLeod & Mathews, 2012). Likewise, psychological manipulations that potentiate amygdala reactivity (e.g., exposure to loud, unpredictable trains of auditory stimuli) also enhance attentional biases to threat (Herry et al., 2007).

Collectively, these observations indicate that exposure to stress alters the activity and connectivity of the amygdala in ways that could promote contextually inappropriate negative affect in individuals with a negative disposition. A key challenge for the future will be to understand whether this reflects

sensitization of circuits involved in orchestrating states of negative affect or a downstream consequence of changes in circuits involved in detecting and attending to threat.

Increased stressor exposure and generation may reflect maladaptive information processing and choices

Self-report data indicate that individuals with a more negative disposition tend to behave in ways that increase the likelihood of hassles, social conflict, inter-personal rejection, and other stressors (e.g., unemployment, divorce)⁴. At present, the neurobiological mechanisms underlying these kinds of recursive Temperament—Environment—Affect relations have received little attention and remain poorly understood. Nonetheless, it is known that the amygdala and other brain regions involved in dispositional negativity can influence information processing and bias decision-making in ways that could promote stress. Imaging studies provide evidence that amygdala activation is associated with reduced trust and is sensitive to potential threat and betrayal during simple economic bargaining games (Baumgartner, Heinrichs, Vonlanthen, Fischbacher, & Fehr, 2008; Bhatt, Lohrenz, Camerer, & Montague, 2012; Gaspic et al., 2011; Li, Xiao, Houser, & Montague, 2009). Conversely, amygdala damage is associated with heightened, even pathological levels of approach and trust (Adolphs, Tranel, & Damasio, 1998; Feinstein et al., 2011; Kennedy, Glascher, Tyszka, & Adolphs, 2009; van Honk, Eisenegger, Terburg, Stein, & Morgan, 2013). Patient studies suggest that the amygdala also plays a role in inhibiting approach in the face of potentially aversive outcomes (i.e., monetary loss aversion; De Martino, Camerer, & Adolphs, 2010). Imaging studies have linked the amygdala to reactive aggression and punishment (Coccaro, McCloskey, Fitzgerald, & Phan, 2007; Treadway et al., 2014), consistent with limited mechanistic work in rodents (Nelson & Trainor, 2007). These observations raise the possibility that stable individual differences in amygdala function contribute to dispositionally negative individuals' tendency to engage in behaviors—including reduced self-disclosure, heightened inhibition and

avoidance, escalation during conflict—that promote social discord and rejection. Via its role in pessimism (Sharot, Riccardi, Raio, & Phelps, 2007) and loss aversion, it is plausible that the amygdala contributes, albeit in a more distal way, to the development or maintenance of other stressors (e.g., lost opportunities for financial or social success, difficulties escaping or improving adverse situations and relationships, avoidance of medical professionals and other potential sources of ‘bad news’). While speculative, the development of more ecologically valid tasks (Jarcho et al., 2013; Somerville, Heatherton, & Kelley, 2006), in combination with longitudinal imaging studies, makes these hypotheses increasingly tractable.

An Integrative Perspective on the Processes Linking Dispositional Negativity to Momentary Emotional Experience and Behavior

The past two decades have witnessed the emergence of powerful tools for assaying emotion and brain function and a number of important insights into the pathways linking stable individual differences in temperament to momentary emotional experience, behavior, and peripheral physiology. These observations provide an integrative framework for understanding the role of dispositional negativity in the development of a broad spectrum of adverse outcomes.

First, there is clear evidence that dispositionally negative individuals respond more strongly to explicit stressors and aversive challenges in the laboratory and in their daily lives. Variation in threat-reactivity reflects stable individual differences in the sensitivity and functional connectivity of several brain regions, including the dorsal amygdala and dlPFC, that have been implicated in the etiology and maintenance of internalizing disorders and substance abuse (**Figures 3-6**) (Chase et al., 2011; Etkin & Wager, 2007; Hamilton et al., 2012; Kuhn & Gallinat, 2011; McLaughlin et al., 2014; Swartz, Knodt, et al., 2015; Tang et al., 2012; Thibodeau et al., 2006; R. A. Wise & Koob, 2014). These differences manifest as heightened activation in response to punctate challenges, but they are also evident in on-going brain

activity, in the absence of explicit tasks or mood manipulations (**Figure 6**). At present, it remains unclear whether temperament-related variation in resting-state activity and functional connectivity reflects increased reactivity to the measurement context, which is often novel or mildly aversive; tonic differences in neurophysiology; or more likely, some combination of these two pathways.

Second, although increased threat-reactivity is important, individuals with a more negative disposition often report elevated levels of negative affect in contexts where stressors are uncertain, psychologically diffuse, or temporally remote. Mechanistic work in rodents and imaging research in humans and monkeys suggests that persistent, contextually indiscriminate negative affect reflects alterations in a neural circuit encompassing the extended amygdala and OFC (**Figures 3 and 7**). Other work indicates that persistent changes in amygdala reactivity and functional connectivity following exposure to acute stress could contribute to the spill-over of negative mood across contexts.

Third, individuals with a more negative disposition tend to behave in ways that evoke or intensify interpersonal conflict and social rejection, increasing the likelihood that they will experience momentary negative affect. Although the neurobiological mechanisms underlying stressor generation have received scant attention, the existing evidence highlights the potential importance of an amygdala-centered circuit responsible for evaluating the trustworthiness and approachability of other individuals.

Of these three pathways, the tendency to experience sustained levels of heightened negative affect in response to diffuse, uncertain, or remote threat appears to be most central to daily function (Bolger & Schilling, 1991). The vast majority of negative affect experienced by dispositionally negative individuals in their daily lives appears to be indiscriminate and cannot be attributed to clear and immediate stressors. In the laboratory, heightened negative affect in explicitly or implicitly safe contexts is generally more discriminative of dispositional negativity and pathological anxiety than that elicited by overt threat

(M. Davis et al., 2010; Duits et al., 2015; Watson & Clark, 1984). This tendency is amplified by exposure to early adversity (e.g., abuse, neglect, loss of caregiver; Wolitzky-Taylor et al., 2014) and it predicts the first onset of anxiety disorders in adolescents (Craske et al., 2012) and the intensification of internalizing symptoms in children and adults (Barker, Reeb-Sutherland, Degnan, et al., *in press*; Lenaert et al., 2014).

This pervasive, context-insensitive emotional bias likely reinforces other components of the negative phenotype, such as elevated avoidance and hyper-vigilance, and may serve to promote the expression of deleterious behaviors that elicit social rejection (Barlow et al., 2013; Carter et al., 2006; Cramer et al., 2012a, 2012b; Grupe & Nitschke, 2013; Jeronimus et al., 2014; Nettle & Bateson, 2012). This reciprocal cycle of maladaptive social transactions may help to explain why dispositionally negative individuals experience lower levels of occupational, financial, and marital success and are more likely to develop internalizing disorders.

A Roadmap to Future Challenges

The data that we have reviewed provide important new insights into the psychological and neurobiological mechanisms that support individual differences in dispositional negativity and the processes that link this disposition to more transient states of negative affect. Yet, it is clear that our understanding remains far from complete. Throughout the review, we highlighted a number of specific conceptual challenges for future research in this area. Here, we outline some broader questions for the field and offer some methodological strategies for beginning to address them¹¹.

¹¹ For more general recommendations about best practices, see (Button et al., 2013a, 2013b; Chalmers et al., 2014; S. P. David et al., 2013; Howells, Sena, & Macleod, 2014; Ioannidis, Greenland, et al., 2014; Ioannidis, Munafò, Fusar-Poli, Nosek, & David, 2014; Lueken et al., 2016; Poldrack et al., 2016; Stelzer, Lohmann, Mueller, Buschmann, & Turner, 2014; Steward & Balice-Gordon, 2014).

How does the brain represent and respond to different kinds of threat?

A major challenge for future research is to understand how the central extended amygdala, OFC, and other key brain regions represent and respond to different kinds of threat (**Figure 2**). Threats differ along several key dimensions—including their certainty, imminence (i.e., physical distance or temporal latency), diffuseness (i.e., cues versus contexts), and duration (D.C. Blanchard, Griebel, & Blanchard, 2001; D. C. Blanchard & Pearson, *in press*; R. J. Blanchard, Blanchard, & Hori, 1989; M. Davis et al., 2010; Fanselow, 1989, 1994; Fanselow & Lester, 1988). There is compelling evidence that these dimensions are psychiatrically important (Craske et al., 2012; M. Davis et al., 2010; Duits et al., 2015; Shackman, Stockbridge, LeMay, & Fox, *in press*) and pharmacologically dissociable (D. C. Blanchard, Griebel, & Blanchard, 2003; Bradford et al., 2013; M. Davis et al., 2010; Kalin & Shelton, 1989). Yet, we know remarkably little about how the human brain represents and differentially responds to them. Although some important advances have been made (Mobbs et al., 2010; Somerville et al., 2013), progress has been slowed by the use of paradigms and assays that confound these dimensions (e.g., *if vs. when* threat will occur; *brief-cues vs. prolonged-contexts*).

Drawing strong inferences about the neural systems supporting phasic and sustained responses to different dimensions of threat requires the use of well-matched tasks, both in humans (Luck, 2005; Shackman et al., 2006) and in animals (Hammack, Todd, Kocho-Schellenberg, & Bouton, 2015). Conditions must be equated for motor requirements and perceptual characteristics, including paired reinforcers (e.g., shocks, aversive images). Investigators should be cautious when comparing neural activity across conditions that markedly differ in duration or number of trials (i.e., in the precision of the estimated activation), as in paradigms where long blocks of uncertain threat are compared to more punctate aversive challenges. Parametric manipulations of threat probability (*if* threat will occur), imminence (*when* or *where* it will occur), and duration (as in Bradford et al., 2013; Mobbs et al., 2010)

would be particularly useful. The use of dynamic parametric tasks (e.g., where threat imminence or probability is smoothly and continuously varied) would also afford powerful new opportunities for understanding the kinds of uncertainty most relevant to fear and anxiety (Bach & Dolan, 2012; de Berker et al., 2016) and for identifying circuits involved in triggering behavioral and physiological ‘phase transitions’ (e.g., from vigilance to behavioral inhibition to active defense; Mobbs et al., 2015; Mobbs & Kim, 2015). Putative double dissociations (e.g., regions sensitive to certain versus uncertain threat) need to be rigorously assessed by testing the appropriate Region \times Condition interaction (as in Somerville et al., 2010). Absent that, strong claims of anatomical dissociation are unwarranted. Likewise, concluding that a particular brain region is ‘not involved’ in a complex, multidimensional psychological function, like ‘fear,’ based on a null statistical test or a single assay is unwarranted.

Emotional disorders are defined and diagnosed on the basis of symptoms (i.e., feelings) and human imaging studies will be particularly valuable for understanding the neural mechanisms underlying the subjective experience of different kinds of threat-elicited negative affect (e.g., Chang, Gianaros, Manuck, Krishnan, & Wager, 2015; Satpute et al., 2013; Somerville et al., 2013), something that cannot be assessed in animal models (Anderson & Adolphs, 2014; LeDoux, 2015). To this end, it will be critical for human studies to verify the presence of target emotions separately for each task or condition (Shackman et al., 2006) and examine relations with on-going neural activity. For correlational techniques, such as fMRI and EEG, trial-by-trial relations between neural signals and emotional experience provide the strongest and most direct link between the brain and self-reported emotion (Atlas, Bolger, Lindquist, & Wager, 2010; Lim, Padmala, & Pessoa, 2009). Multi-voxel classifier approaches, in which machine learning techniques are used to identify patterns of activation predictive of subjective emotional states, are also likely to be fruitful (Chang et al., 2015; Krishnan et al., 2016; Wager et al., 2013; Woo et al., 2014).

Which brain circuits underlie individual differences in dispositional negativity?

It is widely believed that dispositional negativity, like other psychologically and psychiatrically relevant processes, reflects the coordinated activity of distributed brain circuits (Shackman, Fox, & Seminowicz, 2015). Here, we reviewed recent work highlighting the importance of five functional circuits: (a) Projections from the Ce and BST to the subcortical and brainstem regions that proximally mediate key somatomotor and neuroendocrine features of momentary negative affect (**Figure 3**), (b) projections from the Ce to neurotransmitter systems in the basal forebrain and brainstem that modulate sensory processing and promote vigilance (**Figure 3**), (c) projections from the BL to the visual cortex and superior colliculus that play a role in re-directing or amplifying the amount of attention allocated to threat-relevant cues, (d) a poly-synaptic circuit encompassing the Ce and dlPFC that is associated with differences in dispositional negativity in monkeys and anxiety disorders in children (**Figure 6c**), and (e) a circuit encompassing the central extended amygdala and OFC that co-varies with dispositional negativity and supports heightened reactivity to a broad spectrum of aversive challenges (**Figures 4, 5, and 7**), including contexts associated with diffuse or temporally remote threat. Naturally, other circuits are likely to prove important. More broadly, our understanding of the neural circuits underlying dispositional negativity and momentary negative affect remains in its infancy (Calhoun & Tye, 2015; Janak & Tye, 2015; Okon-Singer, Hendler, Pessoa, & Shackman, 2015; Shackman et al., 2015; Tovote et al., 2015). Overcoming this important barrier requires that we accelerate the transition from localization strategies to network-based approaches (Anticevic et al., 2013; Fornito, Zalesky, & Breakspear, 2015; McMenamin et al., 2014; Petersen & Sporns, 2015; Servaas et al., 2014; Turk-Browne, 2013).

It will also be important to determine the specificity of these circuits to different components of dispositional negativity (e.g., anxious distress vs. irritable distress)² and assess whether they are equally relevant to all disorders on the internalizing spectrum (e.g., generalized anxiety disorder vs. major depression). The existing evidentiary record suggests that many of the brain regions that we have

highlighted likely contribute to multiple components and disorders, a one-to-many mapping (Shackman et al., 2015). The amygdala, for example, has been linked to anxiety disorders (Etkin & Wager, 2007), depression (Hamilton et al., 2012), substance abuse (Chase et al., 2011; Kuhn & Gallinat, 2011; Tang et al., 2012), psychopathy (Koenigs, Baskin-Sommers, Zeier, & Newman, 2011), and autism (Nacewicz et al., 2006).

What mechanisms underlie individual differences in dispositional negativity?

Much of the data that we have reviewed comes from brain imaging studies. Aside from unresolved questions about the origins and significance of the measured signals (Logothetis, 2008), the most important limitation of these techniques is that they cannot address causation. A crucial challenge for future studies is to develop a mechanistic understanding of the neural circuits that underlie dispositional negativity and control the expression of negative affect. Addressing these fundamental questions mandates coordinated research efforts in humans and nonhuman animal models. This could be achieved by combining mechanistic approaches (e.g., viral vector, chemogenetic, or optogenetic techniques) in animals with the same whole-brain imaging strategies routinely used in humans, enabling the development of bidirectional translational models (Borsook, Becerra, & Hargreaves, 2006; Casey et al., 2013; Desai et al., 2011; Ferenczi et al., 2016; A. S. Fox et al., 2010). The enhanced precision afforded by optogenetic and chemogenetic techniques would also open the door to identifying the specific molecules, cells, and sub-regions of the central extended amygdala that mediate effects detected in imaging studies (cf. Ferenczi et al., 2016). Nonhuman primate models are likely to be particularly useful for modeling and understanding the neurobiology of dispositional negativity because monkeys and humans share similar genes and brains (Freese & Amaral, 2009; Gibbs et al., 2007; Preuss, 2007), which endow the two species with a common repertoire of complex socio-emotional responses to potential threat and enable the use of similar behavioral assays (A. S. Fox & Kalin, 2014; Kaiser & Feng, 2015; Oler et al., 2016).

In humans, imaging approaches can be applied to patients with circumscribed brain damage (Adolphs, 2016; Motzkin, Philippi, Oler, et al., 2015; Motzkin, Philippi, Wolf, Baskaya, & Koenigs, 2014, 2015; Spunt et al., 2015). Alternatively, fMRI or EEG can be combined with noninvasive perturbation techniques (Bestmann & Feredoes, 2013; Reinhart & Woodman, 2014), neurofeedback (deBettencourt, Cohen, Lee, Norman, & Turk-Browne, 2015; Greer, Trujillo, Glover, & Knutson, 2014; Stoeckel et al., 2014), pharmacological manipulations (Duff et al., 2015; Paulus et al., 2005), or cognitive-behavioral interventions (Britton et al., 2015; Schnyer et al., 2015). Prospective longitudinal imaging studies represent another fruitful approach to identifying candidate mechanisms, especially in relation to the development of internalizing disorders and other kinds of adverse outcomes (Admon, Milad, & Hendler, 2013; Burghy et al., 2012; Herringa et al., 2013; McLaughlin et al., 2014; Swartz, Williamson, et al., 2015). From a broader perspective, human studies will be crucial for identifying the features of animal models that are conserved across species and, hence, most relevant to developing improved interventions for human suffering (Birn et al., 2014). They also afford an opportunity to develop objective biomarkers of psychiatric disease or disease-risk (Borsook et al., 2006; Borsook, Hargreaves, Bountra, & Porreca, 2014; J. Davis et al., 2015; R. G. Wise & Preston, 2010) and for generating novel hypotheses that can be mechanistically assessed in animal models ('reverse translation;' Ferenczi et al., 2016; Janak & Tye, 2015).

What is the relevance of individual differences in brain function for negative affect in daily life?

Biological studies of dispositional negativity and negative affect in humans rely on a limited number of well-controlled, but highly artificial manipulations—fearful and angry faces, aversive sounds and images, electric shocks, and so on—collected under unnatural conditions. Most of these manipulations are only mildly threatening, when compared to the kinds of stressors regularly encountered in daily life or those

routinely used in animal models (LeDoux, 2015; Levenson, *in press*; Shackman et al., 2006), and are subject to self-selection biases, precluding the study of individuals with more extreme dispositions (Scherer, 1986). Consequently, the real-world significance of neural systems identified in the laboratory, including the five circuits that we have highlighted, remains unclear.

Given the limitations of ambulatory measures of brain activity—there is no ‘fMRI helmet’ as yet—addressing these fundamental questions requires integrating assays of brain function and behavior acquired in the scanner with measures of negative affect and motivated behavior assessed under naturalistic conditions in the laboratory (e.g., during semi-structured interactions; Creed & Funder, 1998; Laidlaw, Foulsham, Kuhn, & Kingstone, 2011; Perez-Edgar, McDermott, et al., 2010; Pfeiffer, Vogeley, & Schilbach, 2013) or in the field. Recent work combining fMRI with intensive experience-sampling techniques underscores the value of this approach for identifying the neural systems associated with naturalistic variation in mood and behavior, a central goal of psychology, psychiatry, and the behavioral neurosciences (Berkman & Falk, 2013; Forbes et al., 2009; Heller et al., *in press*; Lopez, Hofmann, Wagner, Kelley, & Heatherton, 2014; S. J. Wilson, Smyth, & MacLean, 2014). To our knowledge, this approach has yet to be applied to the study of dispositional negativity, although on-going work by our group has begun to do so.

The development of robust mobile eye trackers, the emergence of commercial software for automated analyses of facial expressions (Olderbak, Hildebrandt, Pinkpank, Sommer, & Wilhelm, 2014), and the widespread dissemination of smart-phone and other kinds of ‘wearable’ technology afford additional opportunities for objectively, efficiently, and unobtrusively quantifying social attention and context, negative affect, and motivated behavior *in situ* (Gosling & Mason, 2015; Onnela & Rauch, 2016; Sano et al., 2015; Wrzus & Mehl, 2015). Combining these measures with laboratory assays of brain function would open the door to discovering the neural systems underlying persistent, contextually inappropriate

negative affect and pathology-promoting behaviors (e.g., social withdrawal, avoidance, and hypervigilance) in the real world, close to clinically relevant end-points. This approach promises a depth of understanding that cannot be achieved using either animal models or isolated measures of brain function. Even in the absence of biological measures, these new tools promise important clues about the dynamics of negative affect in daily life (e.g., spill-over of mood across sequential contexts and assessments) and the social factors and coping behaviors that help govern the momentary expression of dispositional negativity.

Conclusions

Dispositional negativity has enormous consequences for global health, wealth, and happiness. Self-report data suggest that three key pathways—increased stressor reactivity, tonic increases in negative affect, and increased stressor exposure—explain most of the heightened negative affect characteristic of individuals with a more negative disposition. Of these three pathways, indiscriminate negative affect appears to be the most central to daily life and most relevant to the development and maintenance of anxiety disorders and depression. Mechanistic and imaging studies in humans, monkeys, and rodents suggest that sustained, indiscriminate negative affect reflects variation in the activity and connectivity of several key brain regions, including the central extended amygdala and OFC. These observations provide an integrative framework for understanding the cascading network of psychological and biological processes that bind emotional traits to emotional states and emotional disorders and, ultimately, for guiding the development of more effective prevention and treatment strategies.

FIGURES AND LEGENDS






Figure 1. Pathways linking dispositional negativity (trait) to increased momentary negative affect (state). Questionnaire and behavioral data suggest that three key pathways—increased stressor reactivity, increased tonic levels of negative affect, and more frequent stressors—explain most of the heightened negative affect characteristic of individuals with a negative disposition. Lines depict hypothesized fluctuations in momentary negative affect in individuals with high (red) and low (blue) levels of dispositional negativity, respectively. Acute stressors (e.g., daily hassles, social conflict, and negative life events) are indicated by black lightning bolts.

A. Rodents



D. Comparison of paradigms



B. Nonhuman primates



C. Humans



Figure 2. Different kinds of threat. A. Rodents (Green). In rats and mice, the open field test and the elevated plus maze (EPM) are commonly used to assess emotional responses to diffuse threat. In the open field, rodents are placed into a relatively large, brightly lit, and unfamiliar context. In the elevated plus maze, rodents are placed in a maze with two open arms and two arms enclosed by walls. Freezing and avoidance of the center of the open field or the open arms of the maze provide behavioral ‘read-outs’ of negative affect. Figure adapted with permission from (Tovote et al., 2015). **B. Nonhuman primates (Orange).** In monkeys, the Human ‘Intruder’ Paradigm (HIP) can be used to quantify naturalistic defensive behaviors, neuroendocrine activity, and brain metabolic activity associated with exposure to a

range of threats. In the 'Alone' (ALN) condition, the monkey is simply placed in the testing cage. This novel, diffusely threatening context elicits low levels of freezing and cortisol and a moderate frequency of alarm and separation calls. In the 'No Eye Contact' (NEC) condition (*A, left*), the intruder presents his or her profile while avoiding making eye contact. This elicits passive defenses, including freezing and vocal reductions, similar to procedures used for quantifying behavioral inhibition in children. Panel adapted with permission from (A. S. Fox & Kalin, 2014). **C. Humans (Purple).** In humans, a wide variety of paradigms have been used to probe responses to uncertain, diffuse, or remote threat. Often, these involve the unpredictable presentation of electric shocks or, as shown in the accompanying figure, aversive images. For example, Somerville and colleagues assessed neural activity associated with the temporally certain or uncertain presentation of aversive images. Figure adapted with permission from (Somerville et al., 2013). **D. Comparison of paradigms (Scatter plot).** Threats differ along several key dimensions, including certainty (*x*-axis), physical or temporal imminence, diffuseness (*y*-axis; specific cues vs. real or virtual contexts), and duration (dot size). Here we present some common paradigms used in rodents (green), nonhuman primates (orange), and humans (purple). Studies were chosen for illustrative purposes. We did not attempt a comprehensive review of the literature and, of necessity, the locations of particular paradigms along the two dimensions of the scatter plot are approximate and somewhat arbitrary. Interestingly, many paradigms confound multiple dimensions (e.g., *if* vs. *when* threat will occur) and, because of temporal constraints imposed by conventional fMRI techniques, human imaging studies have focused on the relatively brief (<2 min) anticipation of uncertain threat. **Studies [duration in sec.]—Rodents: Alone in brightly lit cage [900] (D. L. Walker & Davis, 1997); CS+, 100% probability [30] (Duvarci et al., 2009); CS-, 0% probability [30] (Duvarci et al., 2009); Elevated plus maze (EPM) [900] (Tye et al., 2011); Open field [1080] (Tye et al., 2011); Temporally unpredictable cued shock [mean=162] (Miles, Davis, & Walker, 2011); Unpredictable shock context [1200] (Luyten et al., 2012). **Monkeys: Alone (ALN) [1800] (A. S. Fox et al., 2008); CS+, 100% probability [4] (Winslow, Noble, & Davis, 2007); CS-, 0% probability [10] (Kalin, Shelton, Davidson, & Lynn, 1996); No eye contact (NEC) [1800] (A. S. Fox et al., 2008). **Humans: Alone in dark room [120] (Grillon, Pellowski, Merikangas, & Davis, 1997); Anticipating image with unpredictable valence [5] (Grupe et al., 2013); CS+, 100% probability [8] (Gazendam et al., 2013); CS+, 50% probability [3] (Buchel, Morris, Dolan, & Friston, 1998); CS-, 0% shock probability (conditioned safety cue) [8] (Gazendam et al., 2013); Tarantula (video clip of approach) [4] (Mobbs et al., 2010); Temporally unpredictable aversive image [115] (Somerville et al., 2013); Temporally unpredictable cued shock [mean=140] (Moberg & Curtin, 2009); Trier Social Stress Test (TSST) [780] (http://topics.sciencedirect.com/topics/page/Trier_social_stress_test); Virtual reality context paired with unpredictable shock [40] (Alvarez, Chen, Bodurka, Kaplan, & Grillon, 2011).******




Figure 3. Central extended amygdala circuitry. Simplified schematic of key inputs and outputs to the central extended amygdala (magenta) in humans and other primates. The central amygdala encompasses the central nucleus of the amygdala (Ce) and neighboring bed nucleus of the stria terminalis (BST). As shown by the translucent white arrow at the center of the figure, most sensory (yellow), contextual (blue), and regulatory (green) inputs to the central extended amygdala are indirect (i.e., poly-synaptic), and first pass through adjacent amygdala nuclei before arriving at the Ce. In primates, projections linking the Ce to the BST are predominantly unidirectional (Ce → BST). The Ce and BST are poised to orchestrate or trigger momentary negative affect via projections to downstream effector regions (orange). Portions of this figure were adapted with permission from the atlas of Mai and colleagues (Mai, Paxinos, & Voss, 2007). **Abbreviations:** Basolateral (BL), Basomedial (BM), Central (Ce), Lateral (La), and Medial (Me) nuclei of the amygdala; Bed nucleus of the stria terminalis (BST).




A. Adults**B. Adults with a childhood history****C. Young monkeys**

Figure 4. The dorsal amygdala is more reactive to acute threat-related cues in dispositionally negative individuals. **A. Adults with elevated dispositional negativity.** Meta-analysis of six published imaging studies reveals consistently elevated activation bilaterally in the vicinity of the dorsal amygdala (Calder et al., 2011). Significant relations with dispositional negativity (trait) are shown in blue; significant relations with momentary negative affect (state) are depicted in red; and the overlap is shown in purple. **B. Adults with a childhood history of elevated dispositional negativity.** Meta-analysis of seven published imaging studies reveals consistently elevated activation in the right dorsal amygdala (A. S. Fox, Oler, Tromp, et al., 2015). Six of eight amygdala peaks overlapped (yellow) in the dorsal amygdala; four of the peaks extended into the region shown in red. **C. Young monkeys.** Using high-resolution 18-fluorodeoxyglucose-positron emission tomography (FDG-PET) acquired from 238 young rhesus monkeys, Oler and colleagues (2010) demonstrated that threat-related activity in the right Ce (i.e., dorsal amygdala) predicts stable individual differences in dispositional negativity. Figure depicts regions identified by a voxelwise regression analysis (yellow; $p < .05$, whole-brain corrected). The peak voxel and corresponding 95% spatial confidence interval are depicted in white and magenta, respectively. Portions of this figure were adapted with permission from (Calder et al., 2011; A. S. Fox & Kalin, 2014; A. S. Fox, Oler, Tromp, et al., 2015).




Figure 5. Elevated amygdala activity is a shared substrate for different phenotypic presentations of dispositional negativity. Shackman and colleagues (2013) used a well-established monkey model of childhood dispositional negativity and high-resolution FDG-PET to demonstrate that individuals with different presentations of the negative phenotype show increased activity in the central (Ce) nucleus of the amygdala (orange ring). **Divergent phenotypic presentations:** To illustrate this, phenotypic profiles are plotted for groups (N = 80/group; Total N = 238) selected to be extreme on a particular dimension of the phenotype (Top tercile: solid lines; Bottom tercile: broken lines). The panels on the left illustrate how this procedure sorts individuals into groups with divergent presentations of dispositional negativity. **Convergent neural activity:** To illustrate the consistency of Ce activity across divergent presentations, mean neural activity for the extreme groups (\pm SEM) is shown on the right. Individuals with high levels of cortisol, freezing, or vocal reductions (and intermediate levels of the other two responses on average) evinced greater metabolic activity in the Ce compared with those with low levels ($ps < .05$). This figure was adapted with permission from (Shackman et al., 2013).






A. Resting-state prefrontal EEG**B. High-resolution EEG source model****C. Resting-state PFC-Ce functional connectivity assayed using fMRI**

Figure 6. Individuals with a more negative disposition show altered resting-state activity and functional connectivity in the right dorsolateral prefrontal cortex (PFC). **A. Resting-state prefrontal EEG.** Monkeys, children, and adults with a more negative disposition show greater resting-state activity on the scalp overlying the right compared to the left dorsolateral PFC. Figure depicts typical EEG scalp topography. **B. High-resolution EEG source model.** Shackman and colleagues (2009) used 128-channel EEG recordings and distributed source modeling techniques to provide evidence that this scalp-recorded asymmetry reflects increased activity in the right dorsolateral PFC (yellow-orange cluster). **C. Resting-state functional connectivity between the dorsolateral PFC and the Ce assayed using fMRI.** Birn and colleagues (2014) demonstrated that children with anxiety disorders (left) and young monkeys with elevated levels of dispositional negativity (right) both show reduced functional connectivity between the Ce (cyan region in the red rings) and right dorsolateral PFC (black arrows). Pediatric data were collected while patients were quietly resting. Nonhuman primate data were collected under sedation, eliminating potential individual differences in scanner-elicited apprehension. Portions of this figure were adapted with permission from (Birn et al., 2014; Nusslock et al., 2011; Shackman et al., 2009).

A. Adults

Somerville et al., 2010



Mobbs et al., 2010



B. Young monkeys**C. Automated meta-analysis**

Figure 7. Individuals with a more negative disposition show heightened activity in the bed nucleus of the stria terminalis (BST) during periods of diffuse or uncertain threat. Clusters in the vicinity of the BST are indicated by magenta arrows. **A. Adults.** Recent human fMRI studies reveal increased activation in the BST in response to uncertain threat (Mobbs et al., 2010; Somerville et al., 2010). **B. Young monkeys.** Using high-resolution FDG-PET acquired from 592 young rhesus macaques, Fox and colleagues demonstrated that activity in the right BST is heritable and mediates heritable individual differences in dispositional negativity (i.e., BST activity and dispositional negativity are 'genetically correlated;' A. S. Fox, Oler, Shackman, et al., 2015). Regions where activity predicted dispositional negativity are outlined in green. **C. Automated meta-analysis.** An automated Neurosynth (Yarkoni, Poldrack, Nichols, Van Essen, & Wager, 2011) meta-analysis of 312 brain imaging studies featuring the term 'anxiety' revealed several significant regions (red; $Z > 6.0$ and FDR $q < .05$, whole-brain corrected), including the BST. Portions of this figure were adapted with permission from (A. S. Fox, Oler, Shackman, et al., 2015; Mobbs et al., 2010; Somerville et al., 2010).

REFERENCES

- Abbott, R. A., Croudace, T. J., Ploubidis, G. B., Kuh, D., Richards, M., & Huppert, F. A. (2008). The relationship between early personality and midlife psychological well-being: evidence from a UK birth cohort study. *Social Psychiatry and Psychiatric Epidemiology*, *43*, 679-687.
- Abercrombie, H. C., Schaefer, S. M., Larson, C. L., Oakes, T. R., Lindgren, K. A., Holden, J. E., . . . Davidson, R. J. (1998). Metabolic rate in the right amygdala predicts negative affect in depressed patients. *Neuroreport*, *9*, 3301-3307.
- Ackerman, R. A., & Corretti, C. A. (2015). Pathological personality traits and intimacy processes within roommate relationships. *European Journal of Personality*, *29*, 152-172.
- Admon, R., Lubin, G., Stern, O., Rosenberg, K., Sela, L., Ben-Ami, H., & Hendler, T. (2009). Human vulnerability to stress depends on amygdala's predisposition and hippocampal plasticity. *Proceedings of the National Academy of Sciences of the United States of America*, *106*, 14120-14125.
- Admon, R., Milad, M. R., & Hendler, T. (2013). A causal model of post-traumatic stress disorder: disentangling predisposed from acquired neural abnormalities. *Trends Cogn Sci*, *17*, 337-347.
- Adolphs, R. (2016). Human lesion studies in the 21st century. *Neuron*, *90*, 1151-1153.
- Adolphs, R. (*in press*). Consequences of developmental bilateral amygdala lesions in humans. In D. G. Amaral & R. Adolphs (Eds.), *Living without an amygdala*. NY: Guilford Press.
- Adolphs, R., Tranel, D., & Damasio, A. R. (1998). The human amygdala in social judgment. *Nature*, *393*, 470-474.
- Alden, L. E., & Taylor, C. T. (2004). Interpersonal processes in social phobia. *Clinical Psychology Review*, *24*, 857-882.
- Aldinger, M., Stopsack, M., Ulrich, I., Appel, K., Reinelt, E., Wolff, S., . . . Barnow, S. (2014). Neuroticism developmental courses--implications for depression, anxiety and everyday emotional experience; a prospective study from adolescence to young adulthood. *BMC Psychiatry*, *14*.

- Alheid, G. F., & Heimer, L. (1988). New perspectives in basal forebrain organization of special relevance for neuropsychiatric disorders: the striatopallidal, amygdaloid, and corticopetal components of substantia innominata. *Neuroscience*, *27*, 1-39.
- Alisch, R. S., Chopra, P., Fox, A. S., Chen, K., White, A. T., Roseboom, P. H., . . . Kalin, N. H. (2014). Differentially methylated plasticity genes in the amygdala of young primates are linked to anxious temperament, an at risk phenotype for anxiety and depressive disorders. *Journal of Neuroscience*, *34*, 15548-15556.
- Allen, J. J., Harmon-Jones, E., & Cavender, J. H. (2001). Manipulation of frontal EEG asymmetry through biofeedback alters self-reported emotional responses and facial EMG. *Psychophysiology*, *38*, 685-693.
- Allen, J. L., & Rapee, R. M. (2009). Are reported differences in life events for anxious children and controls due to comorbid disorders? *Journal of Anxiety Disorders*, *23*, 511-518.
- Alvarez, R. P., Chen, G., Bodurka, J., Kaplan, R., & Grillon, C. (2011). Phasic and sustained fear in humans elicits distinct patterns of brain activity. *Neuroimage*, *55*, 389-400.
- Alvarez, R. P., Kirlic, N., Misaki, M., Bodurka, J., Rhudy, J. L., Paulus, M. P., & Drevets, W. C. (2015). Increased anterior insula activity in anxious individuals is linked to diminished perceived control. *Transl Psychiatry*, *5*, e591.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.).
- Anderson, D. J., & Adolphs, R. (2014). A framework for studying emotions across species. *Cell*, *157*, 187-200.
- Anticevic, A., Cole, M. W., Repovs, G., Savic, A., Driesen, N. R., Yang, G., . . . Krystal, J. H. (2013). Connectivity, pharmacology, and computation: toward a mechanistic understanding of neural system dysfunction in schizophrenia. *Front Psychiatry*, *4*, 169.

- Arce, E., Simmons, A. N., Lovero, K. L., Stein, M. B., & Paulus, M. P. (2008). Escitalopram effects on insula and amygdala BOLD activation during emotional processing. *Psychopharmacology (Berl)*, *196*(4), 661-672.
- Arnsten, A. F. (2009). Stress signalling pathways that impair prefrontal cortex structure and function. *Nature Reviews. Neuroscience*, *10*, 410-422.
- Arnsten, A. F. (2015). Stress weakens prefrontal networks: molecular insults to higher cognition. *Nature Neuroscience*, *18*, 1376-1385.
- Atlas, L. Y., Bolger, N., Lindquist, M. A., & Wager, T. D. (2010). Brain mediators of predictive cue effects on perceived pain. *Journal of Neuroscience*, *30*, 12964-12977.
- Avery, S. N., Clauss, J. A., & Blackford, J. U. (2016). The human BNST: Functional role in anxiety and addiction. *Neuropsychopharmacology*, *41*, 126-141.
- Baas, J. M., van Ooijen, L., Goudriaan, A., & Kenemans, J. L. (2008). Failure to condition to a cue is associated with sustained contextual fear. *Acta Psychologica*, *127*, 581-592.
- Bach, D. R., & Dolan, R. J. (2012). Knowing how much you don't know: a neural organization of uncertainty estimates. *Nature Reviews. Neuroscience*, *13*, 572-586.
- Banihashemi, L., Sheu, L. K., Midei, A. J., & Gianaros, P. J. (2015). Childhood physical abuse predicts stressor-evoked activity within central visceral control regions. *Soc Cogn Affect Neurosci*, *10*, 474-485.
- Barker, T. V., Reeb-Sutherland, B., Degnan, K. A., Walker, O. L., Chronis-Tuscano, A., Henderson, H. A., . . . Fox, N. A. (*in press*). Contextual startle responses moderate the relation between behavioral inhibition and anxiety in middle childhood. *Psychophysiology*.
- Barker, T. V., Reeb-Sutherland, B. C., & Fox, N. A. (2014). Individual differences in fear potentiated startle in behaviorally inhibited children. *Developmental Psychobiology*, *56*(1), 133-141.
- Barker, T. V., Reeb-Sutherland, B. C., & Fox, N. A. (*in press*). Individual differences in fear potentiated startle in behaviorally inhibited children. *Developmental Psychobiology*.

- Barlow, D. H. (2000). Unraveling the mysteries of anxiety and its disorders from the perspective of emotion theory. *American Psychologist, 55*, 1247-1263.
- Barlow, D. H., Ellard, K. K., Sauer-Zavala, S., Bullis, J. R., & Carl, J. R. (2014). The origins of neuroticism. *Perspectives on Psychological Science, 9*, 481-496.
- Barlow, D. H., Sauer-Zavala, S., Carl, J. R., Bullis, J. R., & Ellard, K. K. (2013). The nature, diagnosis, and treatment of neuroticism: Back to the future. *Clinical Psychological Science, 2*.
- Bastiaansen, J. A., Servaas, M. N., Marsman, J. B., Ormel, J., Nolte, I. M., Riese, H., & Aleman, A. (2014). Filling the gap: Relationship between the serotonin-transporter-linked polymorphic region and amygdala activation. *Psychol Sci, 25*, 2058-2066.
- Baumgartner, T., Heinrichs, M., Vonlanthen, A., Fischbacher, U., & Fehr, E. (2008). Oxytocin shapes the neural circuitry of trust and trust adaptation in humans. *Neuron, 58*, 639-650.
- Beck, J. (2015, February 24, 2015). Hard feelings: Science's struggle to define emotions, *The Atlantic*.
- Bennett, K., Manassis, K., Duda, S., Bagnell, A., Bernstein, G. A., Garland, E. J., . . . Wilansky, P. (2015). Preventing child and adolescent anxiety disorders: Overview of systematic reviews. *Depression and Anxiety, 32*, 909-918.
- Berkman, E. T., & Falk, E. B. (2013). Beyond brain mapping: Using neural measures to predict real-world outcomes. *Curr Dir Psychol Sci, 22*, 45-50.
- Berlanga, C., Heinze, G., Torres, M., Apiquian, R., & Cabellero, A. (1999). Personality and clinical predictors of recurrence in depression. *Psychiatric Services, 50*, 376-380.
- Berry, D. S., Willingham, J. K., & Thayer, C. A. (2000). Affect and personality as predictors of conflict and closeness in young adults' friendships. *Journal of Research in Personality, 34*, 84-107.
- Bestmann, S., & Feredoes, E. (2013). Combined neurostimulation and neuroimaging in cognitive neuroscience: past, present, and future. *Annals of the New York Academy of Sciences, 1296*, 11-30.

- Bhatt, M. A., Lohrenz, T., Camerer, C. F., & Montague, P. R. (2012). Distinct contributions of the amygdala and parahippocampal gyrus to suspicion in a repeated bargaining game. *Proceedings of the National Academy of Sciences of the United States of America*, *109*, 8728-8733.
- Billig, J. P., Hershberger, S. L., Iacono, W. G., & McGue, M. (1996). Life events and personality in late adolescence: Genetic and environmental relations. *Behavior Genetics*, *26*, 543-554.
- Birn, R. M., Shackman, A. J., Oler, J. A., Williams, L. E., McFarlin, D. R., Rogers, G. M., . . . Kalin, N. H. (2014). Evolutionarily conserved dysfunction of prefrontal-amygdalar connectivity in early-life anxiety. *Molecular Psychiatry*, *19*, 915-922.
- Blackford, J. U., Allen, A. H., Cowan, R. L., & Avery, S. N. (2013). Amygdala and hippocampus fail to habituate to faces in individuals with an inhibited temperament. *Soc Cogn Affect Neurosci*, *8*, 143-150.
- Blackford, J. U., Avery, S. N., Cowan, R. L., Shelton, R. C., & Zald, D. H. (2011). Sustained amygdala response to both novel and newly familiar faces characterizes inhibited temperament. *Soc Cogn Affect Neurosci*, *6*, 621-629.
- Blackford, J. U., Avery, S. N., Shelton, R. C., & Zald, D. H. (2009). Amygdala temporal dynamics: temperamental differences in the timing of amygdala response to familiar and novel faces. *BMC Neurosci*, *10*, 145.
- Blackhart, G. C., Kline, J. P., Donohue, K. F., LaRowe, S. D., & Joiner, T. E. (2002). Affective responses to EEG preparation and their link to resting anterior EEG asymmetry. *Personality and Individual Differences*, *32*, 167-174.
- Blanchard, D. C., Griebel, G., & Blanchard, R. J. (2001). Mouse defensive behaviors: pharmacological and behavioral assays for anxiety and panic. *Neuroscience and Biobehavioral Reviews*, *25*, 205-218.
- Blanchard, D. C., Griebel, G., & Blanchard, R. J. (2003). The mouse defense test battery: Pharmacological and behavioral assays for anxiety and panic. *European Journal of Pharmacology*, *463*, 97-116.

- Blanchard, D. C., & Pearson, B. L. (*in press*). Emotion as an evolutionary adaptive pattern: The roles of context and cognition. In A. S. Fox, R. C. Lapate, A. J. Shackman & R. J. Davidson (Eds.), *The nature of emotion. Fundamental questions* (2nd ed.). New York: Oxford University Press.
- Blanchard, R. J., Blanchard, D. C., & Hori, K. (1989). Ethoexperimental approaches to the study of defensive behavior. In R. J. Blanchard, P. F. Brain, D. C. Blanchard & S. Parmigiani (Eds.), *Ethoexperimental approaches to the study of behavior* (pp. 114–136). Dordrecht: Kluwer.
- Block, J. (1995). A contrarian view of the five-factor approach to personality description. *Psychological Bulletin*, *117*, 187-215.
- Block, J. (2010). The five-factor framing of personality and beyond: Some ruminations. *Psychological Inquiry*, *21*, 2-25.
- Bogdan, R., Pagliaccio, D., Baranger, D. A., & Hariri, A. R. (2016). Genetic moderation of stress effects on corticolimbic circuitry. *Neuropsychopharmacology*, *41*, 275-296.
- Boissy, A. (1995). Fear and fearfulness in animals. *Quarterly Review of Biology*, *70*, 165-191.
- Bolger, N., & Schilling, E. A. (1991). Personality and the problems of everyday life: the role of neuroticism in exposure and reactivity to daily stressors. *Journal of Personality*, *59*, 355-386.
- Bolger, N., & Zuckerman, A. (1995). A framework for studying personality in the stress process. *Journal of Personality and Social Psychology*, *69*, 890-902.
- Borsook, D., Becerra, L., & Hargreaves, R. (2006). A role for fMRI in optimizing CNS drug development. *Nature Reviews. Drug Discovery*, *5*, 411-424.
- Borsook, D., Hargreaves, R., Bountra, C., & Porreca, F. (2014). Lost but making progress—Where will new analgesic drugs come from? *Sci Transl Med*, *6*, 249sr243.
- Bradford, D. E., Shapiro, B. L., & Curtin, J. J. (2013). How bad could it be? Alcohol dampens stress responses to threat of uncertain intensity. *Psychol Sci*, *24*, 2541-2549.
- Britton, J. C., Suway, J. G., Clementi, M. A., Fox, N. A., Pine, D. S., & Bar-Haim, Y. (2015). Neural changes with attention bias modification for anxiety: a randomized trial. *Soc Cogn Affect Neurosci*, *10*, 913-920.

- Brooker, R. J., Buss, K. A., Lemery-Chalfant, K., Aksan, N., J., D. R., & Goldsmith, H. H. (*in press*). The development of stranger fear in infancy and toddlerhood: Normative development, individual differences, antecedents, and outcomes. *Developmental Science*.
- Brose, A., Schmiedek, F., Koval, P., & Kuppens, P. (2015). Emotional inertia contributes to depressive symptoms beyond perseverative thinking. *Cogn Emot*, *29*(3), 527-538.
- Brown, G. G., Ostrowitzki, S., Stein, M. B., von Kienlin, M., Liu, T. T., Simmons, A., . . . Paulus, M. (2015). Temporal profile of brain response to alprazolam in patients with generalized anxiety disorder. *Psychiatry Research*, *233*, 394-401.
- Buchel, C., Morris, J., Dolan, R. J., & Friston, K. J. (1998). Brain systems mediating aversive conditioning: an event-related fMRI study. *Neuron*, *20*, 947-957.
- Buckholtz, J. W., Callicott, J. H., Kolachana, B., Hariri, A. R., Goldberg, T. E., Genderson, M., . . . Meyer-Lindenberg, A. (2008). Genetic variation in MAOA modulates ventromedial prefrontal circuitry mediating individual differences in human personality. *Molecular Psychiatry*, *13*, 313-324.
- Buhle, J. T., Silvers, J. A., Wager, T. D., Lopez, R., Onyemekwu, C., Kober, H., . . . Ochsner, K. N. (2014). Cognitive reappraisal of emotion: a meta-analysis of human neuroimaging studies. *Cerebral Cortex*, *24*, 2981-2990.
- Burghy, C. A., Stodola, D. E., Ruttle, P. L., Molloy, E. K., Armstrong, J. M., Oler, J. A., . . . Birn, R. M. (2012). Developmental pathways to amygdala-prefrontal function and internalizing symptoms in adolescence. *Nature Neuroscience*, *15*, 1736-1741.
- Buss, K. A. (2011). Which fearful toddlers should we worry about? Context, fear regulation, and anxiety risk. *Developmental Psychology*, *47*, 804-819.
- Buss, K. A., Davis, E. L., Kiel, E. J., Brooker, R. J., Beekman, C., & Early, M. C. (2013). Dysregulated fear predicts social wariness and social anxiety symptoms during kindergarten. *Journal of Clinical Child and Adolescent Psychology*, *42*, 603-616.

- Buss, K. A., Schumacher, J. R., Dolski, I., Kalin, N. H., Goldsmith, H. H., & Davidson, R. J. (2003a). Right frontal brain activity, cortisol, and withdrawal behavior in 6-month-old infants. *Behavioral Neuroscience, 117*, 11-20.
- Buss, K. A., Schumacher, J. R. M., Dolski, I., Kalin, N. H., Goldsmith, H. H., & Davidson, R. J. (2003b). Right frontal brain activity, cortisol, and withdrawal behavior in 6-month-old infants. *Behavioral Neuroscience, 117*, 11-20.
- Button, K. S., Ioannidis, J. P., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S., & Munafò, M. R. (2013a). Confidence and precision increase with high statistical power. *Nature Reviews. Neuroscience, 14*, 585-586.
- Button, K. S., Ioannidis, J. P., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S., & Munafò, M. R. (2013b). Power failure: why small sample size undermines the reliability of neuroscience. *Nature Reviews. Neuroscience, 14*, 365-376.
- Bystritsky, A. (2006). Treatment-resistant anxiety disorders. *Molecular Psychiatry, 11*, 805-814.
- Calder, A. J., Ewbank, M. P., & Passamonti, L. (2011). Personality influences the neural responses to viewing facial expressions of emotion. *Philosophical Transactions of the Royal Society B: Biological Sciences, 366*, 1684-1701.
- Calhoun, G. G., & Tye, K. M. (2015). Resolving the neural circuits of anxiety. *Nature Neuroscience, 18*, 1394-1404.
- Camp, M. C., Macpherson, K. P., Lederle, L., Graybeal, C., Gaburro, S., Debrouse, L. M., . . . Holmes, A. (2012). Genetic strain differences in learned fear inhibition associated with variation in neuroendocrine, autonomic, and amygdala dendritic phenotypes. *Neuropsychopharmacology, 37*(6), 1534-1547.
- Canli, T., Qiu, M., Omura, K., Congdon, E., Haas, B. W., Amin, Z., . . . Lesch, K. P. (2006). Neural correlates of epigenesis. *Proceedings of the National Academy of Sciences of the United States of America, 103*(43), 16033-16038.

- Carter, J. S., Garber, J., Ciesla, J. A., & Cole, D. A. (2006). Modeling relations between hassles and internalizing and externalizing symptoms in adolescents: a four-year prospective study. *Journal of Abnormal Psychology, 115*, 428-442.
- Casey, B. J., Craddock, N., Cuthbert, B. N., Hyman, S. E., Lee, F. S., & Ressler, K. J. (2013). DSM-5 and RDoC: progress in psychiatry research? *Nature Reviews. Neuroscience, 14*, 810-814.
- Caspi, A., Roberts, B. W., & Shiner, R. L. (2005). Personality development: stability and change. *Annual Review of Psychology, 56*, 453-484.
- Cavanagh, J. F., & Shackman, A. J. (2015). Frontal midline theta reflects anxiety and cognitive control: Meta-analytic evidence. *Journal of Physiology, Paris, 109*, 3-15.
- Chalmers, I., Bracken, M. B., Djulbegovic, B., Garattini, S., Grant, J., Gulmezoglu, A. M., . . . Oliver, S. (2014). How to increase value and reduce waste when research priorities are set. *Lancet, 383*, 156-165.
- Chan, C. K., & Lovibond, P. F. (1996). Expectancy bias in trait anxiety. *Journal of Abnormal Psychology, 105*, 637-647.
- Chang, L. J., Gianaros, P. J., Manuck, S. B., Krishnan, A., & Wager, T. D. (2015). A sensitive and specific neural signature for picture-induced negative affect. *PLoS Biol, 13*, e1002180.
- Chapman, B. P., Fiscella, K., Kawachi, I., & Duberstein, P. R. (2010). Personality, socioeconomic status, and all-cause mortality in the United States. *American Journal of Epidemiology, 171*, 83-92.
- Chase, H. W., Eickhoff, S. B., Laird, A. R., & Hogarth, L. (2011). The neural basis of drug stimulus processing and craving: an activation likelihood estimation meta-analysis. *Biological Psychiatry, 70*, 785-793.
- Cheng, D. T., Knight, D. C., Smith, C. N., & Helmstetter, F. J. (2006). Human amygdala activity during the expression of fear responses. *Behavioral Neuroscience, 120*, 1187-1195.
- Cheng, D. T., Richards, J., & Helmstetter, F. J. (2007). Activity in the human amygdala corresponds to early, rather than late period autonomic responses to a signal for shock. *Learning & Memory, 14*, 485-490.

- Choi, J. M., Padmala, S., & Pessoa, L. (2012). Impact of state anxiety on the interaction between threat monitoring and cognition. *Neuroimage, 59*, 1912-1923.
- Choi, J. S., & Kim, J. J. (2010). Amygdala regulates risk of predation in rats foraging in a dynamic fear environment. *Proceedings of the National Academy of Sciences of the United States of America, 107*, 21773-21777.
- Christian, B. T., Fox, A. S., Oler, J. A., Vandehey, N. T., Murali, D., Rogers, J., . . . Kalin, N. H. (2009). Serotonin transporter binding and genotype in the nonhuman primate brain using [C-11]DASB PET. *Neuroimage, 47*, 1230-1236.
- Chronis-Tuscano, A., Rubin, K. H., O'Brien, K. A., Coplan, R. J., Thomas, S. R., Dougherty, L. R., . . . Wimsatt, M. (2015). Preliminary evaluation of a multimodal early intervention program for behaviorally inhibited preschoolers. *Journal of Consulting and Clinical Psychology, 83*, 534-540.
- Ciocchi, S., Herry, C., Grenier, F., Wolff, S. B., Letzkus, J. J., Vlachos, I., . . . Luthi, A. (2010). Encoding of conditioned fear in central amygdala inhibitory circuits. *Nature, 468*, 277-282.
- Clark, D. A., Durbin, C. E., Hicks, B. M., Iacono, W. G., & McGue, M. (*in press*). Personality in the age of lindustry: Structure, heritability, and correlates of personality in middle childhood from the perspective of parents, teachers, and children. *Journal of Research in Personality*.
- Clark, D. M. (2001). A cognitive perspective on social phobia. In W. R. Crozier & L. E. Alden (Eds.), *International handbook of social anxiety* (pp. 405-430). NY: Wiley.
- Clark, L. A., Kochanska, G., & Ready, R. (2000). Mothers' personality and its interaction with child temperament as predictors of parenting behavior. *Journal of Personality and Social Psychology, 79*, 274-285.
- Clark, L. A., & Watson, D. (2008). Temperament: An organizing paradigm for trait psychology. In O. P. John, R. W. Robins & L. A. Pervin (Eds.), *Handbook of personality: Theory and research* (3rd ed., pp. 265-286). NY: Guilford.

- Clauss, J. A., & Blackford, J. U. (2012). Behavioral inhibition and risk for developing social anxiety disorder: a meta-analytic study. *Journal of the American Academy of Child and Adolescent Psychiatry, 51*, 1066-1075
- Clifford, S., Lemery-Chalfant, K., & Goldsmith, H. H. (2015). The unique and shared genetic and environmental contributions to fear, anger, and sadness in childhood. *Child Development, 86*, 1538-1556.
- Coan, J. A., Allen, J. J. B., & McKnight, P. E. (2006). A capability model of individual differences in frontal EEG asymmetry. *Biological Psychology, 72*, 198-207.
- Coccaro, E. F., McCloskey, M. S., Fitzgerald, D. A., & Phan, K. L. (2007). Amygdala and orbitofrontal reactivity to social threat in individuals with impulsive aggression. *Biological Psychiatry, 62*, 168-178.
- Collins, P. Y., Patel, V., Joestl, S. S., March, D., Insel, T. R., Daar, A. S., . . . Stein, D. J. (2011). Grand challenges in global mental health. *Nature, 475*, 27-30.
- Connolly, J. J., Kavanagh, E. J., & Viswesvaran, C. (2007). The convergent validity between self and observer ratings of personality: A meta-analytic review. *International Journal of Selection and Assessment, 15*, 110-117.
- Connor-Smith, J. K., & Flachsbart, C. (2007). Relations between personality and coping: a meta-analysis. *Journal of Personality and Social Psychology, 93*, 1080-1107.
- Conway, C. C., Craske, M. G., Zinbarg, R. E., & Mineka, S. (2016). Pathological personality traits and naturalistic course of internalizing disorders among high-risk young adults. *Depression and Anxiety, 33*, 84-93.
- Coplan, R., Arbeau, K., & Armer, M. (2007). Don't fret, be supportive! Maternal characteristics linking child shyness to psychosocial and school adjustment in kindergarten. *Journal of Abnormal Child Psychology, 36*, 359-371.

- Corr, P. J. (Ed.). (2008). *The reinforcement sensitivity theory of personality*. NY: Cambridge University Press.
- Cousijn, H., Rijpkema, M., Qin, S., van Marle, H. J., Franke, B., Hermans, E. J., . . . Fernandez, G. (2010). Acute stress modulates genotype effects on amygdala processing in humans. *Proceedings of the National Academy of Sciences of the United States of America*, *107*, 9867-9872.
- Cramer, A. O. J., van der Sluis, S., Noordhof, A., Wichers, M., Geschwind, N., Aggen, S. H., . . . Borsboom, D. (2012a). Dimensions of normal personality as networks in search of equilibrium: You can't like parties if you don't like people. *European Journal of Personality*, *26*, 414-431.
- Cramer, A. O. J., van der Sluis, S., Noordhof, A., Wichers, M., Geschwind, N., Aggen, S. H., . . . Borsboom, D. (2012b). Measurable like temperature or mereological like flocking? On the nature of personality traits. *European Journal of Personality*, *26*, 451-459.
- Craske, M. G., Waters, A. M., Nazarian, M., Mineka, S., Zinbarg, R. E., Griffith, J. W., . . . Ornitz, E. M. (2009). Does neuroticism in adolescents moderate contextual and explicit threat cue modulation of the startle reflex? *Biological Psychiatry*, *65*, 220-226.
- Craske, M. G., Wolitzky-Taylor, K. B., Mineka, S., Zinbarg, R., Waters, A. M., Vrshek-Schallhorn, S., . . . Ornitz, E. (2012). Elevated responding to safe conditions as a specific risk factor for anxiety versus depressive disorders: evidence from a longitudinal investigation. *Journal of Abnormal Psychology*, *121*(2), 315-324.
- Creed, A. T., & Funder, D. C. (1998). Social anxiety: from the inside and outside. *Personality and Individual Differences*, *25*, 19-33.
- Cuijpers, P., Smit, F., Penninx, B. W., de Graaf, R., ten Have, M., & Beekman, A. T. (2010). Economic costs of neuroticism: a population-based study. *Archives of General Psychiatry*, *67*, 1086-1093.
- Damian, R. I., Su, R., Shanahan, M., Trautwein, U., & Roberts, B. W. (2015). Can personality traits and intelligence compensate for background disadvantage? Predicting status attainment in adulthood. *Journal of Personality and Social Psychology*, *109*, 473-489.

- Dannlowski, U., Stuhrmann, A., Beutelmann, V., Zwanzger, P., Lenzen, T., Grotegerd, D., . . . Kugel, H. (2012). Limbic scars: long-term consequences of childhood maltreatment revealed by functional and structural magnetic resonance imaging. *Biological Psychiatry, 71*, 286-293.
- David, J. P., Green, P. J., Martin, R., & Suls, J. (1997). Differential roles of neuroticism, extraversion, and event desirability for mood in daily life: an integrative model of top-down and bottom-up influences. *Journal of Personality and Social Psychology, 73*, 149-159.
- David, S. P., Ware, J. J., Chu, I. M., Loftus, P. D., Fusar-Poli, P., Radua, J., . . . Ioannidis, J. P. (2013). Potential reporting bias in fMRI studies of the brain. *PLoS ONE, 8*, e70104.
- Davidson, R. J. (1998). Affective style and affective disorders: Perspectives from affective neuroscience. *Cognition & Emotion, 12*, 307-330.
- Davidson, R. J., Fox, A. S., & Kalin, N. H. (2007). Neural bases of emotion regulation in nonhuman primates and humans. In J. J. Gross (Ed.), *Handbook of emotion regulation*. (pp. 47-68). NY: Guilford Press.
- Davidson, R. J., Jackson, D. C., & Kalin, N. H. (2000). Emotion, plasticity, context, and regulation: Perspectives from affective neuroscience. *Psychological Bulletin, 126*, 890-909.
- Davidson, R. J., Kalin, N. H., & Shelton, S. (1993). Lateralized response to diazepam predicts temperamental style in rhesus monkeys. *Behavioral Neuroscience, 107*, 1106-1110.
- Davidson, R. J., Kalin, N. H., & Shelton, S. E. (1992). Lateralized effects of diazepam on frontal brain electrical asymmetries in rhesus monkeys. *Biological Psychiatry, 32*, 438-451.
- Davis, J., Maes, M., Andreazza, A., McGrath, J. J., Tye, S. J., & Berk, M. (2015). Towards a classification of biomarkers of neuropsychiatric disease: from encompass to compass. *Molecular Psychiatry, 20*(2), 152-153.
- Davis, M. (1998). Are different parts of the extended amygdala involved in fear versus anxiety? *Biological Psychiatry, 44*, 1239-1247.
- Davis, M. (2006). Neural systems involved in fear and anxiety measured with fear-potentiated startle. *American Psychologist, 61*, 741-756.

- Davis, M., Walker, D. L., & Lee, Y. (1997). Roles of the amygdala and bed nucleus of the stria terminalis in fear and anxiety measured with the acoustic startle reflex. Possible relevance to PTSD. *Annals of the New York Academy of Sciences*, *821*, 305-331.
- Davis, M., Walker, D. L., Miles, L., & Grillon, C. (2010). Phasic vs sustained fear in rats and humans: Role of the extended amygdala in fear vs anxiety. *Neuropsychopharmacology*, *35*, 105-135.
- Davis, M., & Whalen, P. J. (2001). The amygdala: vigilance and emotion. *Molecular Psychiatry*, *6*, 13-34.
- de Berker, A. O., Rutledge, R. B., Mathys, C., Marshall, L., Cross, G. F., Dolan, R. J., & Bestmann, S. (2016). Computations of uncertainty mediate acute stress responses in humans. *Nat Commun*, *7*, 10996.
- De Bolle, M., De Fruyt, F., McCrae, R. R., Lockenhoff, C. E., Costa, P. T., Jr., Aguilar-Vafaie, M. E., . . . Terracciano, A. (2015). The emergence of sex differences in personality traits in early adolescence: A cross-sectional, cross-cultural study. *Journal of Personality and Social Psychology*, *108*, 171-185.
- de Haan, A. D., Dekovic, M., & Prinzie, P. (2012). Longitudinal impact of parental and adolescent personality on parenting. *Journal of Personality and Social Psychology*, *102*, 189-199.
- De Martino, B., Camerer, C. F., & Adolphs, R. (2010). Amygdala damage eliminates monetary loss aversion. *Proceedings of the National Academy of Sciences of the United States of America*, *107*, 3788-3792.
- Deary, I. J., Weiss, A., & Batty, G. D. (2010). Intelligence and personality as predictors of illness and death: How researchers in differential psychology and chronic disease epidemiology are collaborating to understand and address health inequalities. *Psychological Science in the Public Interest*, *11*, 53-79.
- deBettencourt, M. T., Cohen, J. D., Lee, R. F., Norman, K. A., & Turk-Browne, N. B. (2015). Closed-loop training of attention with real-time brain imaging. *Nature Neuroscience*, *18*, 470-475.
- Dennis, T. A., & O'Toole, L. (2014). Mental health on the go: Effects of a gamified attention bias modification mobile application in trait anxious adults. *Clin Psychol Sci*, *2*, 576-590.
- Desai, M., Kahn, I., Knoblich, U., Bernstein, J., Atallah, H., Yang, A., . . . Boyden, E. S. (2011). Mapping brain networks in awake mice using combined optical neural control and fMRI. *Journal of Neurophysiology*, *105*, 1393-1405.

- DiLuca, M., & Olesen, J. (2014). The cost of brain diseases: a burden or a challenge? *Neuron*, *82*, 1205-1208.
- Donnellan, M. B., Conger, R. D., & Bryant, C. M. (2004). The Big Five and enduring marriages. *Journal of Research in Personality*, *38*, 481-504.
- Duckworth, A. L., & Allred, K. M. (2012). Temperament in the classroom. In R. L. Shiner & M. Zentner (Eds.), *Handbook of temperament* (pp. 627-644). NY: Guilford.
- Duff, E. P., Vennart, W., Wise, R. G., Howard, M. A., Harris, R. E., Lee, M., . . . Smith, S. M. (2015). Learning to identify CNS drug action and efficacy using multistudy fMRI data. *Sci Transl Med*, *7*, 274ra216.
- Duggan, C. F., Lee, A. S., & Murray, R. M. (1990). Does personality predict long-term outcome in depression? *British Journal of Psychiatry*, *157*, 19-24.
- Duits, P., Cath, D. C., Lissek, S., Hox, J. J., Hamm, A. O., Engelhard, I. M., . . . Baas, J. M. (2015). Updated meta-analysis of classical fear conditioning in the anxiety disorders. *Depression and Anxiety*, *32*, 239-253.
- Duvarci, S., Bauer, E. P., & Paré, D. (2009). The bed nucleus of the stria terminalis mediates inter-individual variations in anxiety and fear. *Journal of Neuroscience*, *29*, 10357-10361.
- Dyrenforth, P. S., Kashy, D. A., Donnellan, M. B., & Lucas, R. E. (2010). Predicting relationship and life satisfaction from personality in nationally representative samples from three countries: The relative importance of actor, partner, and similarity effects. *Journal of Personality and Social Psychology*, *99*, 690-702.
- Eatough, E. M., Shirtcliff, E. A., Hanson, J. L., & Pollak, S. D. (2009). Hormonal reactivity to MRI scanning in adolescents. *Psychoneuroendocrinology*, *34*, 1242-1246.
- Ellenbogen, M. A., & Hodgins, S. (2004). The impact of high neuroticism in parents on children's psychosocial functioning in a population at high risk for major affective disorder: a family-environmental pathway of intergenerational risk. *Development and Psychopathology*, *16*, 113-136.

- Epstein, S. (1994). Trait theory as personality theory: Can a part be as great as the whole? *Psychological Inquiry*, 5, 120-122.
- Etkin, A., & Wager, T. D. (2007). Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *American Journal of Psychiatry*, 164, 1476-1488.
- Everaerd, D., Klumpers, F., van Wingen, G., Tendolkar, I., & Fernandez, G. (2015). Association between neuroticism and amygdala responsivity emerges under stressful conditions. *Neuroimage*, 112, 218-224.
- Eysenck, H. J. (1967). *The biological basis of personality*. Springfield, IL: Charles C. Thomas.
- Eysenck, H. J. (1983). Cicero and the state-trait theory of anxiety: Another case of delayed recognition. *American Psychologist*, 38, 114-115.
- Fanselow, M. S. (1989). The adaptive function of conditioned defensive behavior: An ecological approach to Pavlovian stimulus substitution theory. In R. J. Blanchard, P. F. Brain, D. C. Blanchard & S. Parmigiani (Eds.), *Ethoexperimental approaches to the study of behavior* (pp. 151-166). Boston: Kluwer.
- Fanselow, M. S. (1994). Neural organization of the defensive behavior system responsible for fear. *Psychonomic Bulletin & Review*, 1, 429-438.
- Fanselow, M. S., & Lester, L. S. (1988). A functional behavioristic approach to aversively motivated behavior: Predatory imminence as a determinant of the topography of defensive behavior. In R. C. Bolles & M. D. Beecher (Eds.), *Evolution and learning* (pp. 185-211). Hillsdale, NJ: Erlbaum.
- Faravelli, C., Ambonetti, A., Pallanti, S., & Pazzagli, A. (1986). Depressive relapses and incomplete recovery from index episode. *American Journal of Psychiatry*, 7, 888-891.
- Farmer, A. S., & Kashdan, T. B. (2015). Stress sensitivity and stress generation in social anxiety disorder: a temporal process approach. *Journal of Abnormal Psychology*, 124, 102-114.

- Feinstein, J. S., Adolphs, R., Damasio, A., & Tranel, D. (2011). The human amygdala and the induction and experience of fear. *Current Biology, 21*, 1-5.
- Felmington, K., Kemp, A., Williams, L., Das, P., Hughes, G., Peduto, A., & Bryant, R. (2007). Changes in anterior cingulate and amygdala after cognitive behavior therapy of posttraumatic stress disorder. *Psychol Sci, 18*, 127-129.
- Ferenczi, E. A., Zalocusky, K. A., Liston, C., Grosenick, L., Warden, M. R., Amatya, D., . . . Deisseroth, K. (2016). Prefrontal cortical regulation of brainwide circuit dynamics and reward-related behavior. *Science, 351*, aac9698.
- Ferguson, E., Heckman, J. J., & Corr, P. J. (2011). Personality and economics: Overview and proposed framework. *Personality and Individual Differences, 51*, 201-209.
- Fergusson, D. M., & Horwood, L. J. (1987). Vulnerability to life events exposure. *Psychological Medicine, 17*, 739-749.
- Fleeson, W. (2001). Toward a structure- and process-integrated view of personality: Traits as density distributions of states. *Journal of Personality & Social Psychology, 80*, 1011-1027.
- Fonzo, G. A., Ramsawh, H. J., Flagan, T. M., Sullivan, S. G., Letamendi, A., Simmons, A. N., . . . Stein, M. B. (2015). Common and disorder-specific neural responses to emotional faces in generalised anxiety, social anxiety and panic disorders. *British Journal of Psychiatry, 206*, 206-215.
- Forbes, E. E., Hariri, A. R., Martin, S. L., Silk, J. S., Moyles, D. L., Fisher, P. M., . . . Dahl, R. E. (2009). Altered striatal activation predicting real-world positive affect in adolescent major depressive disorder. *American Journal of Psychiatry, 166*, 64-73.
- Fornito, A., Zalesky, A., & Breakspear, M. (2015). The connectomics of brain disorders. *Nature Rev Neurosci, 16*, 159-172.
- Fox, A. S., & Kalin, N. H. (2014). A translational neuroscience approach to understanding the development of social anxiety disorder and its pathophysiology. *American Journal of Psychiatry, 171*, 1162-1173.

- Fox, A. S., Oler, J. A., Shackman, A. J., Shelton, S. E., Raveendran, M., McKay, D. R., . . . Kalin, N. H. (2015). Intergenerational neural mediators of early-life anxious temperament. *Proceedings of the National Acadademy of Sciences USA, 112*, 9118-9122.
- Fox, A. S., Oler, J. A., Shelton, S. E., Nanda, S. A., Davidson, R. J., Roseboom, P. H., & Kalin, N. H. (2012). Central amygdala nucleus (Ce) gene expression linked to increased trait-like Ce metabolism and anxious temperament in young primates. *Proceedings of the National Academy of Sciences of the United States of America, 109*, 18108-18113.
- Fox, A. S., Oler, J. A., Tromp, D. P., Fudge, J. L., & Kalin, N. H. (2015). Extending the amygdala in theories of threat processing. *Trends in Neurosciences, 38*, 319-329.
- Fox, A. S., Shelton, S. E., Oakes, T. R., Converse, A. K., Davidson, R. J., & Kalin, N. H. (2010). Orbitofrontal cortex lesions alter anxiety-related activity in the primate bed nucleus of stria terminalis. *Journal of Neuroscience, 30*, 7023-7027.
- Fox, A. S., Shelton, S. E., Oakes, T. R., Davidson, R. J., & Kalin, N. H. (2008). Trait-like brain activity during adolescence predicts anxious temperament in primates. *PLoS ONE, 3*, e2570.
- Fox, N. A., Henderson, H. A., Marshall, P. J., Nichols, K. E., & Ghera, M. M. (2005). Behavioral inhibition: linking biology and behavior within a developmental framework. *Annual Review of Psychology, 56*, 235-262.
- Fraley, R. C., & Roberts, B. W. (2005). Patterns of continuity: a dynamic model for conceptualizing the stability of individual differences in psychological constructs across the life course. *Psychological Review, 112*, 60-74.
- Freese, J. L., & Amaral, D. G. (2009). Neuroanatomy of the primate amygdala. In P. J. Whalen & E. A. Phelps (Eds.), *The human amygdala* (pp. 3-42). NY: Guilford.
- Funder, D. C. (1994). Explaining traits. *Psychological Inquiry, 5*, 125-127.

- Furmark, T., Tillfors, M., Marteinsdottir, I., Fischer, H., Pissiota, A., Langstrom, B., & Fredrikson, M. (2002). Common changes in cerebral blood flow in patients with social phobia treated with citalopram or cognitive-behavioral therapy. *Archives of General Psychiatry*, *59*, 425-433.
- Gable, S. L., Reis, H. T., & Elliot, A. J. (2000). Behavioral activation and inhibition in everyday life. *Journal of Personality and Social Psychology*, *78*, 1135-1149.
- Gale, C. R., Hagenaars, S. P., Davies, G., Hill, W. D., Liewald, D. C., Cullen, B., . . . Harris, S. E. (2016). Pleiotropy between neuroticism and physical and mental health: findings from 108 038 men and women in UK Biobank. *Transl Psychiatry*, *6*, e791.
- Gaylin, W. (1979). *Feelings: Our vital signs*. NY: Harper & Row.
- Gazelle, H. (2008). Behavioral profiles of anxious solitary children and heterogeneity in peer relations. *Developmental Psychology*, *44*, 1604-1624.
- Gazelle, H., & Ladd, G. W. (2003). Anxious solitude and peer exclusion: A diathesis-stress model of internalizing trajectories in childhood. *Child Development*, *74*, 257-278.
- Gazendam, F. J., Kamphuis, J. H., Eigenhuis, A., Huizenga, H. M. H., Soeter, M., Bos, M. G. N., . . . Kindt, M. (2015). Personality predicts individual variation in fear learning: A multilevel growth modeling approach. *Clinical Psychological Science*, *3*, 175-188.
- Gazendam, F. J., Kamphuis, J. H., & Kindt, M. (2013). Deficient safety learning characterizes high trait anxious individuals. *Biological Psychology*, *92*(2), 342-352.
- Gendron, M., & Barrett, L. F. (2009). Reconstructing the past: A century of ideas about emotion in psychology. *Emotion Review*, *1*, 316-339.
- Ghosh, S., & Chattarji, S. (2015). Neuronal encoding of the switch from specific to generalized fear. *Nature Neuroscience*, *18*(1), 112-120.
- Gibbs, R. A., Rogers, J., Katze, M. G., Bumgarner, R., Weinstock, G. M., Mardis, E. R., . . . Zwiig, A. S. (2007). Evolutionary and biomedical insights from the rhesus macaque genome. *Science*, *316*, 222-234.

- Gleason, M. E., Powers, A. D., & Oltmanns, T. F. (2012). The enduring impact of borderline personality pathology: risk for threatening life events in later middle-age. *Journal of Abnormal Psychology, 121*, 447-457.
- Glue, P., Wilson, S., Coupland, N., Ball, D., & Nutt, D. (1995). The relationship between benzodiazepine receptor sensitivity and neuroticism. *Journal of Anxiety Disorders, 9*, 33-45.
- Goldsmith, H. H., Buss, A. H., Plomin, R., Rothbart, M. K., Thomas, A., Chess, S., . . . McCall, R. B. (1987). Roundtable: what is temperament? Four approaches. *Child Development, 58*, 505-529.
- Goodwin, R. D., Hoven, C. W., Lyons, J. S., & Stein, M. B. (2002). Mental health service utilization in the United States. The role of personality factors. *Social Psychiatry and Psychiatric Epidemiology, 37*, 561-566.
- Gosling, S. D. (2008). Personality in non-human animals. *Social and Personality Psychology Compass, 2*, 985-1001.
- Gosling, S. D., & Mason, W. (2015). Internet research in psychology. *Annual Review of Psychology, 66*, 877-902.
- Gospic, K., Mohlin, E., Fransson, P., Petrovic, P., Johannesson, M., & Ingvar, M. (2011). Limbic justice--amygdala involvement in immediate rejection in the Ultimatum Game. *PLoS Biol, 9*, e1001054.
- Grav, S., Stordal, E., Romild, U. K., & Hellzen, O. (2012). The relationship among neuroticism, extraversion, and depression in the HUNT Study: in relation to age and gender. *Issues in Mental Health Nursing, 33*, 777-785.
- Gray, J. A., & McNaughton, N. (2000). *The neuropsychology of anxiety* (2nd ed.). NY: Oxford University Press.
- Greer, S. M., Trujillo, A. J., Glover, G. H., & Knutson, B. (2014). Control of nucleus accumbens activity with neurofeedback. *Neuroimage, 96*, 237-244.
- Griebel, G., & Holmes, A. (2013). 50 years of hurdles and hope in anxiolytic drug discovery. *Nature Reviews. Drug Discovery, 12*, 667-687.

- Grillon, C. (2002). Associative learning deficits increase symptoms of anxiety in humans. *Biological Psychiatry*, *51*, 851-858.
- Grillon, C. (2008). Models and mechanisms of anxiety: evidence from startle studies. *Psychopharmacology*, *199*, 421-437.
- Grillon, C., & Ameli, R. (2001). Conditioned inhibition of fear-potentiated startle and skin conductance in humans. *Psychophysiology*, *38*(5), 807-815.
- Grillon, C., Baas, J. M., Pine, D. S., Lissek, S., Lawley, M., Ellis, V., & Levine, J. (2006). The benzodiazepine alprazolam dissociates contextual fear from cued fear in humans as assessed by fear-potentiated startle. *Biological Psychiatry*, *60*(7), 760-766.
- Grillon, C., & Charney, D. R. (2011). In the face of fear: anxiety sensitizes defensive responses to fearful faces. *Psychophysiology*, *48*, 1745-1752.
- Grillon, C., Pellowski, M., Merikangas, K. R., & Davis, M. (1997). Darkness facilitates the acoustic startle reflex in humans. *Biological Psychiatry*, *42*, 453-460.
- Gross, J. J., Sutton, S. K., & Ketelaar, T. (1998). Relations between affect and personality: Support for the affect-level and affective reactivity views. *Personality and Social Psychology Bulletin*, *24*, 279-288.
- Grupe, D. W., & Nitschke, J. B. (2013). Uncertainty and anticipation in anxiety: an integrated neurobiological and psychological perspective. *Nature Reviews. Neuroscience*, *14*, 488-501.
- Grupe, D. W., Oathes, D. J., & Nitschke, J. B. (2013). Dissecting the anticipation of aversion reveals dissociable neural networks. *Cerebral Cortex*, *23*, 1874-1883.
- Gungor, N. Z., & Paré, D. (*in press*). Functional heterogeneity in the bed nucleus of the stria terminalis. *Journal of Neuroscience*.
- Gunthert, K. C., Cohen, L. H., & Armeli, S. (1999). The role of neuroticism in daily stress and coping. *Journal of Personality and Social Psychology*, *77*, 1087-1100.

- Haaker, J., Lonsdorf, T. B., Schumann, D., Menz, M., Brassens, S., Bunzeck, N., . . . Kalisch, R. (2015). Deficient inhibitory processing in trait anxiety: Evidence from context-dependent fear learning, extinction recall and renewal. *Biological Psychology, 111*, 65-72.
- Hagemann, D., Hewig, J., Seifert, J., Naumann, E., & Bartussek, D. (2005). The latent state-trait structure of resting EEG asymmetry: replication and extension. *Psychophysiology, 42*, 740-752.
- Hagemann, D., Naumann, E., Thayer, J. F., & Bartussek, D. (2002). Does resting electroencephalograph asymmetry reflect a trait? An application of latent state-trait theory. *Journal of Personality and Social Psychology, 82*, 619-641.
- Hakulinen, C., Elovainio, M., Pulkki-Raback, L., Virtanen, M., Kivimaki, M., & Jokela, M. (2015). Personality and depressive symptoms: Individual participant meta-analysis of 10 cohort studies. *Depression and Anxiety, 32*, 461-470.
- Hakulinen, C., Hintsanen, M., Munafo, M. R., Virtanen, M., Kivimaki, M., Batty, G. D., & Jokela, M. (2015). Personality and smoking: individual-participant meta-analysis of nine cohort studies. *Addiction, 110*, 1844-1852.
- Hamilton, J. P., Etkin, A., Furman, D. J., Lemus, M. G., Johnson, R. F., & Gotlib, I. H. (2012). Functional neuroimaging of major depressive disorder: a meta-analysis and new integration of base line activation and neural response data. *American Journal of Psychiatry, 169*, 693-703.
- Hammack, S. E., Todd, T. P., Kocho-Schellenberg, M., & Bouton, M. E. (2015). Role of the bed nucleus of the stria terminalis in the acquisition of contextual fear at long or short context-shock intervals. *Behavioral Neuroscience, 129*, 673-678.
- Hankin, B. L. (2010). Personality and depressive symptoms: Stress generation and cognitive vulnerabilities to depression in a prospective daily diary study. *J Soc Clin Psychol, 29*, 369-401.
- Hankin, B. L., Fraley, R. C., & Abela, J. R. (2005). Daily depression and cognitions about stress: evidence for a traitlike depressogenic cognitive style and the prediction of depressive symptoms in a prospective daily diary study. *Journal of Personality and Social Psychology, 88*, 673-685.

- Hare, T. A., Tottenham, N., Galvan, A., Voss, H. U., Glover, G. H., & Casey, B. J. (2008). Biological substrates of emotional reactivity and regulation in adolescence during an emotional go-nogo task. *Biological Psychiatry, 63*, 927-934.
- Harmer, C. J., Mackay, C. E., Reid, C. B., Cowen, P. J., & Goodwin, G. M. (2006). Antidepressant drug treatment modifies the neural processing of nonconscious threat cues. *Biological Psychiatry, 59*, 816-820.
- Harris, L. M., Cumming, S. R., & Menzies, R. G. (2004). Predicting anxiety in magnetic resonance imaging scans. *International Journal of Behavioral Medicine, 11*, 1-7.
- Harris, L. M., Robinson, J., & Menzies, R. G. (2001). Predictors of panic symptoms during magnetic resonance imaging scans. *International Journal of Behavioral Medicine, 8*, 80-87.
- Headey, B., & Wearing, A. (1989). Personality, life events, and subjective well-being: Toward a dynamic equilibrium model. *Journal of Personality and Social Psychology, 57*, 731-739.
- Heerey, E. A., & Kring, A. M. (2007). Interpersonal consequences of social anxiety. *Journal of Abnormal Psychology, 116*, 125-134.
- Hefner, K. R., & Curtin, J. J. (2012). Alcohol stress response dampening: selective reduction of anxiety in the face of uncertain threat. *J Psychopharmacol, 26*(2), 232-244.
- Hefner, K. R., Jaber, J. N., Grant, A. M., & Curtin, J. J. (2009). Alcohol intoxication: Selective reduction of anxiety in the face of uncertain threat. *Psychophysiology, 46*, S64.
- Hefner, K. R., Moberg, C. A., Hachiya, L. Y., & Curtin, J. J. (2013). Alcohol stress response dampening during imminent versus distal, uncertain threat. *Journal of Abnormal Psychology, 122*(3), 756-769.
- Heineck, G. (2011). Does it pay to be nice? Personality and earnings in the United Kingdom. *Industrial and Labor Relations Review, 64*, 1020-1038.
- Heller, A. S., Fox, A. S., Wing, E., Mayer, K., Vack, N. J., & Davidson, R. J. (*in press*). Affective neurodynamics predict prolonged real-world emotional responses. *Journal of Neuroscience, 35*, 10503-10509.

- Hengartner, M. P., Ajdacic-Gross, V., Wyss, C., Angst, J., & Rossler, W. (2016). Relationship between personality and psychopathology in a longitudinal community study: a test of the predisposition model. *Psychological Medicine, 46*, 1693-1705.
- Hengartner, M. P., Kawohl, W., Haker, H., Rossler, W., & Ajdacic-Gross, V. (2016). Big Five personality traits may inform public health policy and preventive medicine: Evidence from a cross-sectional and a prospective longitudinal epidemiologic study in a Swiss community. *Journal of Psychosomatic Research, 84*, 44-51.
- Hengartner, M. P., van der Linden, D., Bohleber, L., & von Wyl, A. (2016). Big Five Personality Traits and the General Factor of Personality as Moderators of Stress and Coping Reactions Following an Emergency Alarm on a Swiss University Campus. *Stress Health*.
- Herrington, R. J., Birn, R. M., Ruttle, P. L., Burghy, C. A., Stodola, D. E., Davidson, R. J., & Essex, M. J. (2013). Childhood maltreatment is associated with altered fear circuitry and increased internalizing symptoms by late adolescence. *Proceedings of the National Academy of Sciences of the United States of America*.
- Herry, C., Bach, D. R., Esposito, F., Di Salle, F., Perrig, W. J., Scheffler, K., . . . Seifritz, E. (2007). Processing of temporal unpredictability in human and animal amygdala. *Journal of Neuroscience, 27*, 5958-5966.
- Hintsanen, M., Puttonen, S., Smith, K., Tornroos, M., Jokela, M., Pulkki-Raback, L., . . . Keltikangas-Jarvinen, L. (2014). Five-factor personality traits and sleep: evidence from two population-based cohort studies. *Health Psychology, 33*, 1214-1223.
- Hirschfeld, R. M., Klerman, G. L., Andreasen, N. C., Clayton, P. J., & Keller, M. B. (1986). Psycho-social predictors of chronicity in depressed patients. *British Journal of Psychiatry, 148*, 648-654.
- Hoehn-Saric, R., McLeod, D. R., Funderburk, F., & Kowalski, P. (2004). Somatic symptoms and physiologic responses in generalized anxiety disorder and panic disorder: an ambulatory monitor study. *Archives of General Psychiatry, 61*(9), 913-921.

- Horwitz, B. N., Luong, G., & Charles, S. T. (2008). Neuroticism and extraversion share genetic and environmental effects with negative and positive mood spillover in a nationally representative sample. *Personality & Individual Differences, 45*, 636-642.
- Houben, M., Van Den Noortgate, W., & Kuppens, P. (2015). The relation between short-term emotion dynamics and psychological well-being: A meta-analysis. *Psychological Bulletin*.
- Howells, D. W., Sena, E. S., & Macleod, M. R. (2014). Bringing rigour to translational medicine. *Nature Reviews Neurology, 10*, 37-43.
- Hutteman, R., Bleidorn, W., Kereste, G., Brkovic, I., Butkovic, A., & Denissen, J. J. A. (2014). Reciprocal associations between parenting challenges and parents' personality development in young and middle adulthood. *European Journal of Personality, 28*, 168-179.
- Hutteman, R., Bleidorn, W., Kerestes, G., Brkovic, I., Butkovic, A., & Denissen, J. J. A. (2014). Reciprocal associations between parenting challenges and parents' personality development in young and middle adulthood. *European Journal of Personality, 28*, 168-179.
- Iacovino, J. M., Bogdan, R., & Oltmanns, T. F. (2016). Personality predicts health declines through stressful life events during late mid-life. *Journal of Personality, 84*, 536-546.
- Insel, T. R. (2012). Next-generation treatments for mental disorders. *Sci Transl Med, 4*, 155ps119.
- Ioannidis, J., Greenland, S., Hlatky, M., Khoury, M., Macleod, M., Moher, D., . . . Tibshirani, R. (2014). Increasing value and reducing waste in research design, conduct, and analysis. *Lancet, 383*, 166-175.
- Ioannidis, J., Munafo, M., Fusar-Poli, P., Nosek, B., & David, S. (2014). Publication and other reporting biases in cognitive sciences: detection, prevalence, and prevention. *Trends in Cognitive Sciences, 18*, 235-241.
- Izard, C. E. (2010). The many meanings/aspects of emotion: Definitions, functions, activation, and regulation. *Emotion Review, 2*, 363-370.

- Izquierdo, A., Suda, R. K., & Murray, E. A. (2005). Comparison of the effects of bilateral orbital prefrontal cortex lesions and amygdala lesions on emotional responses in rhesus monkeys. *Journal of Neuroscience, 25*(37), 8534-8542.
- Jackson, J. J., Connolly, J. J., Garrison, S. M., Leveille, M. M., & Connolly, S. L. (2015). Your friends know how long you will live: a 75-year study of peer-rated personality traits. *Psychol Sci, 26*, 335-340.
- Jacobs, N., van Os, J., Derom, C., Thiery, E., Delespaul, P., & Wichers, M. (2011). Neuroticism explained? From a non-informative vulnerability marker to informative person-context interactions in the realm of daily life. *British Journal of Clinical Psychology, 50*, 19-32.
- Jahn, A. L., Fox, A. S., Abercrombie, H. C., Shelton, S. E., Oakes, T. R., Davidson, R. J., & Kalin, N. H. (2010). Subgenual prefrontal cortex activity predicts individual differences in hypothalamic-pituitary-adrenal activity across different contexts. *Biological Psychiatry, 67*, 175-181.
- Janak, P. H., & Tye, K. M. (2015). From circuits to behaviour in the amygdala. *Nature, 517*, 284-292.
- Jarcho, J. M., Leibenluft, E., Walker, O. L., Fox, N. A., Pine, D. S., & Nelson, E. E. (2013). Neuroimaging studies of pediatric social anxiety: paradigms, pitfalls and a new direction for investigating the neural mechanisms. *Biol Mood Anxiety Disord, 3*, 14.
- Jeronimus, B. F., Riese, H., Sanderman, R., & Ormel, J. (2014). Mutual reinforcement between neuroticism and life experiences: a five-wave, 16-year study to test reciprocal causation. *Journal of Personality and Social Psychology, 107*, 751-764.
- Johnson, J. G., Cohen, P., Kasen, S., & Brook, J. S. (2004). Paternal psychiatric symptoms and maladaptive paternal behavior in the home during the child rearing years. *Journal of Child and Family Studies, 13*, 421-437.
- Jokela, M., Hakulinen, C., Singh-Manoux, A., & Kivimaki, M. (2014). Personality change associated with chronic diseases: pooled analysis of four prospective cohort studies. *Psychological Medicine, 44*, 2629-2640.

- Jokela, M., Kivimaki, M., Elovainio, M., & Keltikangas-Jarvinen, L. (2009). Personality and having children: a two-way relationship. *J Pers Soc Psychol*, *96*, 218-230.
- Jokela, M., Pulkki-Raback, L., Elovainio, M., & Kivimaki, M. (2014). Personality traits as risk factors for stroke and coronary heart disease mortality: pooled analysis of three cohort studies. *Journal of Behavioral Medicine*, *37*(5), 881-889.
- Jovanovic, T., Nylocks, K. M., Gamwell, K. L., Smith, A., Davis, T. A., Norrholm, S. D., & Bradley, B. (2014). Development of fear acquisition and extinction in children: effects of age and anxiety. *Neurobiology of Learning and Memory*, *113*, 135-142.
- Judge, T. A., Simon, L. S., Hurst, C., & Kelley, K. (2014). What I experienced yesterday is who I am today: Relationship of work motivations and behaviors to within-individual variation in the five-factor model of personality. *Journal of Applied Psychology*, *99*, 199-221.
- Kaczurkin, A. N., Moore, T. M., Ruparel, K., Ciric, R., Calkins, M. E., Shinohara, R. T., . . . Sattherthwaite, T. D. (*in press*). Elevated amygdala perfusion mediates developmental sex differences in trait anxiety. *Biological Psychiatry*.
- Kagan, J., Reznick, J. S., & Snidman, N. (1988). Biological bases of childhood shyness. *Science*, *240*, 167-171.
- Kagan, J., Snidman, N., Kahn, V., & Towsley, S. (2007). The preservation of two infant temperaments into adolescence. *Monographs of the Society for Research in Child Development*, *72*, 1-75.
- Kaiser, T., & Feng, G. (2015). Modeling psychiatric disorders for developing effective treatments. *Nature Medicine*, *21*, 979-988.
- Kalin, N. H., Fox, A. S., Kovner, R., Riedel, M. K., Fekete, E. M., Roseboom, P. H., . . . Oler, J. A. (*in press*). Overexpressing corticotropin-releasing hormone in the primate amygdala increases anxious temperament and alters its neural circuit. *Biological Psychiatry*.

- Kalin, N. H., Larson, C., Shelton, S. E., & Davidson, R. J. (1998). Asymmetric frontal brain activity, cortisol, and behavior associated with fearful temperament in rhesus monkeys. *Behavioral Neuroscience, 112*, 286-292.
- Kalin, N. H., & Shelton, S. E. (1989). Defensive behaviors in infant rhesus monkeys: environmental cues and neurochemical regulation. *Science, 243*, 1718-1721.
- Kalin, N. H., Shelton, S. E., & Davidson, R. J. (2004). The role of the central nucleus of the amygdala in mediating fear and anxiety in the primate. *Journal of Neuroscience, 24*, 5506-5515.
- Kalin, N. H., Shelton, S. E., & Davidson, R. J. (2007). Role of the primate orbitofrontal cortex in mediating anxious temperament. *Biological Psychiatry, 62*, 1134-1139.
- Kalin, N. H., Shelton, S. E., Davidson, R. J., & Lynn, D. E. (1996). A new method for aversive Pavlovian conditioning of heart rate in rhesus monkeys. *Physiology and Behavior, 60*, 1043-1046.
- Kalin, N. H., Shelton, S. E., Fox, A. S., Oakes, T. R., & Davidson, R. J. (2005). Brain regions associated with the expression and contextual regulation of anxiety in primates. *Biological Psychiatry, 58*, 796-804.
- Kandler, C., Bleidorn, W., Riemann, R., Angleitner, A., & Spinath, F. M. (2012). Life events as environmental states and genetic traits and the role of personality: A longitudinal twin study. *Behavior Genetics, 42*, 57-72.
- Karney, B. R., & Bradbury, T. N. (1995). The longitudinal course of marital quality and stability: a review of theory, method, and research. *Psychological Bulletin, 118*, 3-34.
- Kelly, E. L., & Conley, J. J. (1987). Personality and compatibility: A prospective analysis of marital stability and marital satisfaction. *Journal of Personality and Social Psychology, 52*, 27-40.
- Kendler, K. S., & Gardner, C. O. (2014). Sex differences in the pathways to major depression: a study of opposite-sex twin pairs. *American Journal of Psychiatry, 171*, 426-435.
- Kendler, K. S., Gardner, C. O., & Prescott, C. A. (2003). Personality and the experience of environmental adversity. *Psychological Medicine, 33*, 1193-1202.

- Kendler, K. S., & Karkowski-Shuman, L. (1997). Stressful life events and genetic liability to major depression: genetic control of exposure to the environment? *Psychological Medicine*, *27*, 539-547.
- Kendler, K. S., Karkowski, L. M., & Prescott, C. A. (1999). Causal relationship between stressful life events and the onset of major depression. *American Journal of Psychiatry*, *156*, 837-841.
- Kendler, K. S., Neale, M. C., Kessler, R. C., & Heath, A. C. (1993). A longitudinal twin study of personality and major depression in women. *Archives of General Psychiatry*, *50*, 853-862.
- Kennedy, D. P., Glascher, J., Tyszka, J. M., & Adolphs, R. (2009). Personal space regulation by the human amygdala. *Nature Neuroscience*, *12*, 1226-1227.
- Kercher, A. J., Rapee, R. M., & Schniering, C. A. (2009). Neuroticism, life events and negative thoughts in the development of depression in adolescent girls. *Journal of Abnormal Child Psychology*, *37*, 903-915.
- Kessler, R. C., Petukhova, M., Sampson, N. A., Zaslavsky, A. M., & Wittchen, H. U. (2012). Twelve-month and lifetime prevalence and lifetime morbid risk of anxiety and mood disorders in the United States. *Int J Methods Psychiatr Res*, *21*, 169-184.
- Kessler, R. C., Walters, E. E., & Forthofer, M. S. (1998). The social consequences of psychiatric disorders, III: Probability of marital stability. *American Journal of Psychiatry*, *155*, 1092-1096.
- Kheirbek, M. A., Klemenhagen, K. C., Sahay, A., & Hen, R. (2012). Neurogenesis and generalization: a new approach to stratify and treat anxiety disorders. *Nature Neuroscience*, *15*(12), 1613-1620.
- Klein, K. J., Lim, B.-C., Saltz, J. L., & Mayer, D. M. (2004). How do we get there? An examination of the antecedents of centrality in team networks. *Academy of Management Journal*, *47*, 952-963.
- Klumpers, F., Kroes, M. C., Heitland, I., Everaerd, D., Akkermans, S. E., Oosting, R. S., . . . Baas, J. M. (2015). Dorsomedial prefrontal cortex mediates the impact of serotonin transporter linked polymorphic region genotype on anticipatory threat reactions. *Biological Psychiatry*, *78*, 582-589.

- Klumpers, F., Morgan, B., Terburg, D., Stein, D. J., & van Honk, J. (*in press*). Impaired acquisition of classically conditioned fear-potentiated startle reflexes in humans with focal bilateral basolateral amygdala damage. *Social Cognitive and Affective Neuroscience*, nsu164.
- Knight, D. C., Nguyen, H. T., & Bandettini, P. A. (2005). The role of the human amygdala in the production of conditioned fear responses. *Neuroimage*, 26, 1193-1200.
- Kochanska, G., Clark, L. A., & Goldman, M. S. (1997). Implications of mothers' personality for their parenting and their young children's developmental outcomes. *Journal of Personality*, 65, 387-420.
- Koenigs, M., Baskin-Sommers, A., Zeier, J., & Newman, J. P. (2011). Investigating the neural correlates of psychopathy: a critical review. *Molecular Psychiatry*, 16, 792-799.
- Komulainen, E., Meskanen, K., Lipsanen, J., Lahti, J. M., Jylha, P., Melartin, T., . . . Ekelund, J. (2014). The effect of personality on daily life emotional processes. *PLoS ONE*, 9, e110907.
- Kopala-Sibley, D. C., Danzig, A. P., Kotov, R., Bromet, E. J., Carlson, G. A., Olino, T. M., . . . Klein, D. N. (*in press*). Negative emotionality and its facets moderate the effects of exposure to hurricane Sandy on children's postdisaster depression and anxiety symptoms. *Journal of Abnormal Psychology*.
- Kopala-Sibley, D. C., Kotov, R., Bromet, E. J., Carlson, G. A., Danzig, A. P., Black, S. R., & Klein, D. N. (2016). Personality diatheses and Hurricane Sandy: effects on post-disaster depression. *Psychological Medicine*, 46, 865-875.
- Kotov, R., Gamez, W., Schmidt, F., & Watson, D. (2010). Linking "big" personality traits to anxiety, depressive, and substance use disorders: a meta-analysis. *Psychological Bulletin*, 136, 768-821.
- Koval, P., & Kuppens, P. (2012). Changing emotion dynamics: individual differences in the effect of anticipatory social stress on emotional inertia. *Emotion*, 12(2), 256-267.
- Koval, P., Kuppens, P., Allen, N. B., & Sheeber, L. (2012). Getting stuck in depression: the roles of rumination and emotional inertia. *Cogn Emot*, 26(8), 1412-1427.
- Kozak, M. J., & Cuthbert, B. N. (2016). The NIMH research domain criteria initiative: Background, issues, and pragmatics. *Psychophysiology*, 53, 286-297.

- Kragel, P. A., & LaBar, K. S. (2015). Multivariate neural biomarkers of emotional states are categorically distinct. *Soc Cogn Affect Neurosci*, *10*, 1437-1448.
- Krishnan, A., Woo, C. W., Chang, L. J., Ruzic, L., Gu, X., Lopez-Sola, M., . . . Wager, T. D. (2016). Somatic and vicarious pain are represented by dissociable multivariate brain patterns. *Elife*, *5*.
- Kuhn, S., & Gallinat, J. (2011). Common biology of craving across legal and illegal drugs - a quantitative meta-analysis of cue-reactivity brain response. *European Journal of Neuroscience*, *33*, 1318-1326.
- Kurdek, L. A. (1993). Predicting marital dissolution: A 5-year prospective longitudinal study of newlywed couples. *Journal of Personality & Social Psychology*, *64*, 221-242.
- LaBar, K. S., Gatenby, J. C., Gore, J. C., LeDoux, J. E., & Phelps, E. A. (1998). Human amygdala activation during conditioned fear acquisition and extinction: a mixed-trial fMRI study. *Neuron*, *20*, 937-945.
- Laceulle, O. M., Nederhof, E., Karreman, A., Ormel, J., & Van Aken, M. A. G. (2011). Stressful events and temperament change during early and middle adolescence: The Trails study. *European Journal of Personality*, *26*, 276-284.
- Ladd, G. W., Kochenderfer-Ladd, B., Eggum, N. D., Kochel, K. P., & McConnell, E. M. (2011). Characterizing and comparing the friendships of anxious-solitary and unsociable preadolescents. *Child Development*, *82*, 1434-1453.
- Lahey, B. B. (2009). Public health significance of neuroticism. *American Psychologist*, *64*, 241-256.
- Laidlaw, K. E., Foulsham, T., Kuhn, G., & Kingstone, A. (2011). Potential social interactions are important to social attention. *Proceedings of the National Academy of Sciences of the United States of America*, *108*, 5548-5553.
- Lakdawalla, Z., & Hankin, B. L. (2008). Personality as a prospective vulnerability to dysphoric symptoms among college students: Proposed mechanisms. *J Psychopathol Behav Assess*, *30*, 121-131.
- Lake, R. I., Eaves, L. J., Maes, H. H., Heath, A. C., & Martin, N. G. (2000). Further evidence against the environmental transmission of individual differences in neuroticism from a collaborative study of 45,850 twins and relatives on two continents. *Behavior Genetics*, *30*, 223-233.

- Lapate, R. C., & Shackman, A. J. (*in press*). Afterword: What is an emotion? In A. S. Fox, R. C. Lapate, A. J. Shackman & R. J. Davidson (Eds.), *The nature of emotion: Fundamental questions* (2nd ed.). New York: Oxford University Press.
- Larsen, R. J., & Ketelaar, T. (1989). Extraversion, neuroticism and susceptibility to positive and negative mood induction procedures. *Personality & Individual Differences*, *10*, 1221-1228.
- Larsen, R. J., & Ketelaar, T. (1991). Personality and susceptibility to positive and negative emotional states. *Journal of Personality and Social Psychology*, *61*, 132-140.
- Laufer, O., Israeli, D., & Paz, R. (*in press*). Behavioral and neural mechanisms of overgeneralization in anxiety. *Current Biology*.
- Lebowitz, E. R., Scharfstein, L. A., & Jones, J. (2014). Comparing family accommodation in pediatric obsessive-compulsive disorder, anxiety disorders, and nonanxious children. *Depression and Anxiety*, *31*, 1018-1025.
- Lebowitz, E. R., Woolston, J., Bar-Haim, Y., Calvocoressi, L., Dauser, C., Warnick, E., . . . Leckman, J. F. (2013). Family accommodation in pediatric anxiety disorders. *Depression and Anxiety*, *30*, 47-54.
- LeDoux, J. E. (2015). *Anxious. Using the brain to understand and treat fear and anxiety*. NY: Viking.
- Leger, K. A., Charles, S. T., Turiano, N. A., & Almeida, D. M. (*in press*). Personality and stressor-related affect. *Journal of Personality & Social Psychology*.
- Lenaert, B., Boddez, Y., Griffith, J. W., Vervliet, B., Schruers, K., & Hermans, D. (2014). Aversive learning and generalization predict subclinical levels of anxiety: a six-month longitudinal study. *Journal of Anxiety Disorders*, *28*(8), 747-753.
- Levenson, R. W. (*in press*). What is the added value of studying the brain for understanding emotion? In A. S. Fox, R. C. Lapate, A. J. Shackman & R. J. Davidson (Eds.), *The nature of emotion. Fundamental questions* (2nd ed.). NY: Oxford University Press.

- Leventhal, A. M., Japuntich, S. J., Piper, M. E., Jorenby, D. E., Schlam, T. R., & Baker, T. B. (2012). Isolating the role of psychological dysfunction in smoking cessation: relations of personality and psychopathology to attaining cessation milestones. *Psychol Addict Behav, 26*, 838-849.
- Li, J., Xiao, E., Houser, D., & Montague, P. R. (2009). Neural responses to sanction threats in two-party economic exchange. *Proceedings of the National Academy of Sciences of the United States of America, 106*, 16835-16840.
- Lieberman, L., Liu, H., Huggins, A. A., Katz, A. C., Zvolensky, M. J., & Shankman, S. A. (*in press*). Comparing the validity of informant and self-reports of personality using laboratory indices of emotional responding as criterion variables. *Psychophysiology*.
- Lim, S. L., Padmala, S., & Pessoa, L. (2009). Segregating the significant from the mundane on a moment-to-moment basis via direct and indirect amygdala contributions. *Proceedings of the National Academy of Sciences of the United States of America, 106*, 16841-16846.
- Linetzky, M., Pergamin-Hight, L., Pine, D. S., & Bar-Haim, Y. (2015). Quantitative evaluation of the clinical efficacy of attention bias modification treatment for anxiety disorders. *Depression and Anxiety, 32*, 383-391.
- Lissek, S. (2012). Toward an account of clinical anxiety predicated on basic, neurally mapped mechanisms of Pavlovian fear-learning: the case for conditioned overgeneralization. *Depression and Anxiety, 29*, 257-263.
- Lissek, S., Bradford, D. E., Alvarez, R. P., Burton, P., Espensen-Sturges, T., Reynolds, R. C., & Grillon, C. (2014). Neural substrates of classically conditioned fear-generalization in humans: a parametric fMRI study. *Soc Cogn Affect Neurosci, 9*, 1134-1142.
- Liu, R. T., & Alloy, L. B. (2010). Stress generation in depression: A systematic review of the empirical literature and recommendations for future study. *Clinical Psychology Review, 30*, 582-593.
- Liu, R. T., Kraines, M. A., Massing-Schaffer, M., & Alloy, L. B. (2014). Rejection sensitivity and depression: mediation by stress generation. *Psychiatry, 77*(1), 86-97.

- Logothetis, N. K. (2008). What we can do and what we cannot do with fMRI. *Nature*, *453*, 869-878.
- Lopez, R. B., Hofmann, W., Wagner, D. D., Kelley, W. M., & Heatherton, T. F. (2014). Neural predictors of giving in to temptation in daily life. *Psychol Sci*, *25*(7), 1337-1344.
- Luck, S. J. (2005). Ten simple rules for designing ERP experiments. In T. C. Handy (Ed.), *Event-related potentials: A methods handbook* (pp. 17-32). Cambridge, MA: MIT Press.
- Ludtke, O., Roberts, B. W., Trautwein, U., & Nagy, G. (2011). A random walk down university avenue: life paths, life events, and personality trait change at the transition to university life. *Journal of Personality and Social Psychology*, *101*, 620-637.
- Lueken, U., Muehlhan, M., Evens, R., Wittchen, H. U., & Kirschbaum, C. (2012). Within and between session changes in subjective and neuroendocrine stress parameters during magnetic resonance imaging: A controlled scanner training study. *Psychoneuroendocrinology*, *37*, 1299-1308.
- Lueken, U., Zierhut, K. C., Hahn, T., Straube, B., Kircher, T., Reif, A., . . . Domschke, K. (2016). Neurobiological markers predicting treatment response in anxiety disorders: A systematic review and implications for clinical application. *Neuroscience and Biobehavioral Reviews*, *66*, 143-162.
- Luyten, L., Casteels, C., Vansteenwegen, D., van Kuyck, K., Koole, M., Van Laere, K., & Nuttin, B. (2012). Micro-positron emission tomography imaging of rat brain metabolism during expression of contextual conditioning. *Journal of Neuroscience*, *32*, 254-263.
- Machado, C. J., Snyder, A. Z., Cherry, S. R., Lavenex, P., & Amaral, D. G. (2008). Effects of neonatal amygdala or hippocampus lesions on resting brain metabolism in the macaque monkey: a microPET imaging study. *Neuroimage*, *39*, 832-846.
- MacLeod, C., & Clarke, P. J. F. (2015). The attentional bias modification approach to anxiety intervention. *Clinical Psychological Science*, *3*, 58-78.
- MacLeod, C., & Mathews, A. (2012). Cognitive bias modification approaches to anxiety. *Annu Rev Clin Psychol*, *8*, 189-217.

- Magnus, K., Diener, E., Fujita, F., & Pavot, W. (1993). Extraversion and neuroticism as predictors of objective life events: A longitudinal analysis. *Journal of Personality and Social Psychology, 65*, 1046-1053.
- Mai, J. K., Paxinos, G., & Voss, T. (2007). *Atlas of the human brain* (3rd ed.). San Diego, CA: Academic Press.
- Malouff, J. M., Thorsteinsson, E. B., Rooke, S. E., & Schutte, N. S. (2007). Alcohol involvement and the Five-Factor model of personality: a meta-analysis. *Journal of Drug Education, 37*, 277-294.
- Malouff, J. M., Thorsteinsson, E. B., Schutte, N. S., Bhullar, N., & Rooke, S. E. (2010). The five-factor model of personality and relationship satisfaction of intimate partners: A meta-analysis. *Journal of Research in Personality, 44*, 124-127.
- Marco, C. A., & Suls, J. (1993). Daily stress and the trajectory of mood: spillover, response assimilation, contrast, and chronic negative affectivity. *Journal of Personality and Social Psychology, 64*, 1053-1063.
- Markon, K. E., Krueger, R. F., & Watson, D. (2005). Delineating the structure of normal and abnormal personality: an integrative hierarchical approach. *Journal of Personality and Social Psychology, 88*, 139-157.
- Mason, W. A., Capitanio, J. P., Machado, C. J., Mendoza, S. P., & Amaral, D. G. (2006). Amygdalectomy and responsiveness to novelty in rhesus monkeys (*Macaca mulatta*): generality and individual consistency of effects. *Emotion, 6*, 73-81.
- Matthews, G., Deary, I. J., & Whiteman, M. C. (2009). Stable traits and transient states. In G. Matthews, I. J. Deary & M. C. Whiteman (Eds.), *Personality traits* (3rd ed., pp. 85-120). NY: Cambridge University Press.
- McCrae, R. R., & Costa, P. T. (1991). Adding liebe und arbeit: The full five-factor model and well-being. *Personality and Social Psychology Bulletin, 17*, 227-232.
- McCrae, R. R., & Costa, P. T., Jr. (1987). Validation of the five-factor model of personality across instruments and observers. *Journal of Personality and Social Psychology, 52*, 81-90.

- McLaughlin, K. A., Busso, D. S., Duys, A., Green, J. G., Alves, S., Way, M., & Sheridan, M. A. (2014). Amygdala response to negative stimuli predicts PTSD symptom onset following a terrorist attack. *Depression and Anxiety, 31*, 834-842.
- McMenamin, B. W., Langeslag, S. J., Sirbu, M., Padmala, S., & Pessoa, L. (2014). Network organization unfolds over time during periods of anxious anticipation. *Journal of Neuroscience, 34*, 11261-11273.
- McNulty, J. K. (2008). Neuroticism and interpersonal negativity: the independent contributions of perceptions and behaviors. *Personality and Social Psychology Bulletin, 34*, 1439-1450.
- Mehl, M. R., Gosling, S. D., & Pennebaker, J. W. (2006). Personality in its natural habitat: manifestations and implicit folk theories of personality in daily life. *Journal of Personality and Social Psychology, 90*, 862-877.
- Melendez, J. C., & McCrank, E. (1993). Anxiety-related reactions associated with magnetic resonance imaging examinations. *Journal of the American Medical Association, 270*, 745-747.
- Metz, M., Majdandzic, M., & Bogels, S. (in press). Concurrent and predictive associations between infants' and toddlers' fearful temperament, coparenting, and parental anxiety disorders. *Journal of Clinical Child and Adolescent Psychology*.
- Middeldorp, C. M., Cath, D. C., Beem, A. L., Willemsen, G., & Boomsma, D. I. (2008). Life events, anxious depression and personality: a prospective and genetic study. *Psychological Medicine, 38*(11), 1557-1565.
- Mihalopoulos, C., Vos, T., Rapee, R. M., Pirkis, J., Chatterton, M. L., Lee, Y. C., & Carter, R. (2015). The population cost-effectiveness of a parenting intervention designed to prevent anxiety disorders in children. *Journal of Child Psychology and Psychiatry and Allied Disciplines, 56*, 1026-1033.
- Miles, L., Davis, M., & Walker, D. (2011). Phasic and sustained fear are pharmacologically dissociable in rats. *Neuropsychopharmacology, 36*, 1563-1574.

- Mobbs, D., Hagan, C. C., Dalgleish, T., Silston, B., & Prevost, C. (2015). The ecology of human fear: survival optimization and the nervous system. *Front Neurosci*, *9*, 55.
- Mobbs, D., & Kim, J. J. (2015). Neuroethological studies of fear, anxiety, and risky decision-making in rodents and humans. *Current Opinion in Behavioral Sciences*, *5*, 8-15.
- Mobbs, D., Yu, R., Rowe, J. B., Eich, H., FeldmanHall, O., & Dalgleish, T. (2010). Neural activity associated with monitoring the oscillating threat value of a tarantula. *Proceedings of the National Academy of Sciences USA*, *107*, 20582-20586.
- Moberg, C. A., & Curtin, J. J. (2009). Alcohol selectively reduces anxiety but not fear: Startle response during unpredictable vs. predictable threat. *J Abnormal Psychol*, *118*, 335-347.
- Moffitt, T. E., Poulton, R., & Caspi, A. (2013). Lifelong impact of early self-control. *American Scientist*, *101*, 352-359.
- Monroe, S. M., & Simons, A. D. (1991). Diathesis-stress theories in the context of life stress research: implications for the depressive disorders. *Psychological Bulletin*, *110*(3), 406-425.
- Motzkin, J. C., Philippi, C. L., Oler, J. A., Kalin, N. H., Baskaya, M. K., & Koenigs, M. (2015). Ventromedial prefrontal cortex damage alters resting blood flow to the bed nucleus of stria terminalis. *Cortex*, *64*, 281-288.
- Motzkin, J. C., Philippi, C. L., Wolf, R. C., Baskaya, M. K., & Koenigs, M. (2014). Ventromedial prefrontal cortex lesions alter neural and physiological correlates of anticipation. *Journal of Neuroscience*, *34*(31), 10430-10437.
- Motzkin, J. C., Philippi, C. L., Wolf, R. C., Baskaya, M. K., & Koenigs, M. (2015). Ventromedial prefrontal cortex is critical for the regulation of amygdala activity in humans. *Biological Psychiatry*, *77*(3), 276-284.
- Mroczek, D. K., & Almeida, D. M. (2004). The effect of daily stress, personality, and age on daily negative affect. *Journal of Personality*, *72*, 355-378.

- Muehlhan, M., Lueken, U., Siegert, J., Wittchen, H. U., Smolka, M. N., & Kirschbaum, C. (2013). Enhanced sympathetic arousal in response to fMRI scanning correlates with task induced activations and deactivations. *PLoS ONE*, *8*, e72576.
- Mund, M., & Neyer, F. J. (*in press*). The winding paths of the lonesome cowboy: Evidence for mutual influences between personality, subjective health, and loneliness. *J Personality*.
- Munsterkotter, A. L., Notzon, S., Redlich, R., Grotegerd, D., Dohm, K., Arolt, V., . . . Dannlowski, U. (*in press*). Spider or no spider? Neural correlates of sustained and phasic fear in spider phobia. *Depression and Anxiety*.
- Nacewicz, B. M., Dalton, K. M., Johnstone, T., Long, M. T., McAuliff, E. M., Oakes, T. R., . . . Davidson, R. J. (2006). Amygdala Volume and Nonverbal Social Impairment in Adolescent and Adult Males With Autism. *Archives of General Psychiatry*, *63*, 1417-1428.
- Nelson, R. J., & Trainor, B. C. (2007). Neural mechanisms of aggression. *Nature Reviews. Neuroscience*, *8*, 536-546.
- Nettle, D., & Bateson, M. (2012). The evolutionary origins of mood and its disorders. *Current Biology*, *22*, R712-721.
- Newman, M. G., & Fisher, A. J. (2013). Mediated moderation in combined cognitive behavioral therapy versus component treatments for generalized anxiety disorder. *Journal of Consulting and Clinical Psychology*, *81*, 405-414.
- Neyer, F. J., & Asendorpf, J. B. (2001). Personality-relationship transaction in young adulthood. *Journal of Personality and Social Psychology*, *81*, 1190-1204.
- Neyer, F. J., & Voigt, D. (2004). Personality and social network effects on romantic relationships: A dyadic approach. *European Journal of Personality*, *18*, 279-299.
- Ng, T. W. H., Eby, L. T., Sorensen, K. L., & Feldman, D. C. (2005). Predictors of objective and subjective career success. A meta-analysis. *Personnel Psychology*, *58*, 367-408.

- Nolan, S. A., Roberts, J. E., & Gotlib, I. H. (1998). Neuroticism and ruminative response style as predictors of change in depressive symptomatology. *Cognitive Therapy and Research, 22*, 445-455.
- Nolen-Hoeksema, S., Wisco, B. E., & Lyubomirsky, S. (2008). Rethinking rumination. *Perspectives on Psychological Science, 3*, 400-424.
- Norris, C. J., Larsen, J. T., & Cacioppo, J. T. (2007). Neuroticism is associated with larger and more prolonged electrodermal responses to emotionally evocative pictures. *Psychophysiology, 44*, 823-826.
- Nusslock, R., Shackman, A. J., Harmon-Jones, E., Alloy, L. B., Coan, J. A., & Abramson, L. Y. (2011). Cognitive vulnerability and frontal brain asymmetry: common predictors of first prospective depressive episode. *Journal of Abnormal Psychology, 120*(2), 497-503.
- Okbay, A., Baselmans, B. M., De Neve, J. E., Turley, P., Nivard, M. G., Fontana, M. A., . . . Cesarini, D. (*in press*). Genetic variants associated with subjective well-being, depressive symptoms, and neuroticism identified through genome-wide analyses. *Nature Genetics*.
- Okon-Singer, H., Hendler, T., Pessoa, L., & Shackman, A. J. (2015). The neurobiology of emotion-cognition interactions: Fundamental questions and strategies for future research. *Frontiers in Human Neuroscience, 9*.
- Olderbak, S., Hildebrandt, A., Pinkpank, T., Sommer, W., & Wilhelm, O. (2014). Psychometric challenges and proposed solutions when scoring facial emotion expression codes. *Behav Res Methods, 46*, 992-1006.
- Oler, J. A., Fox, A. S., Shackman, A. J., & Kalin, N. H. (2016). The central nucleus of the amygdala is a critical substrate for individual differences in anxiety. In D. G. Amaral & R. Adolphs (Eds.), *Living without an amygdala*. NY: Guilford.
- Oler, J. A., Fox, A. S., Shelton, S. E., Christian, B. T., Murali, D., Oakes, T. R., . . . Kalin, N. H. (2009). Serotonin transporter availability in the amygdala and bed nucleus of the stria terminalis predicts anxious temperament and brain glucose metabolic activity. *Journal of Neuroscience, 29*, 9961-9966.

- Oler, J. A., Fox, A. S., Shelton, S. E., Rogers, J., Dyer, T. D., Davidson, R. J., . . . Kalin, N. H. (2010). Amygdalar and hippocampal substrates of anxious temperament differ in their heritability. *Nature*, *466*, 864-868.
- Onnela, J. P., & Rauch, S. L. (2016). Harnessing smartphone-based digital phenotyping to enhance behavioral and mental health. *Neuropsychopharmacology*, *41*, 1691-1696.
- Ormel, J., Jeronimus, B. F., Kotov, R., Riese, H., Bos, E. H., Hankin, B., . . . Oldehinkel, A. J. (2013). Neuroticism and common mental disorders: meaning and utility of a complex relationship. *Clinical Psychology Review*, *33*, 686-697.
- Ormel, J., Oldehinkel, A. J., & Vollebergh, W. (2004). Vulnerability before, during, and after a major depressive episode: a 3-wave population-based study. *Archives of General Psychiatry*, *61*, 990-996.
- Ormel, J., & Wohlfarth, T. (1991). How neuroticism, long-term difficulties, and life situation change influence psychological distress: A longitudinal model. *Journal of Personality and Social Psychology*, *60*, 744-755.
- Parker, P. D., Ludtke, O., Trautwein, U., & Roberts, B. W. (2012). Personality and relationship quality during the transition from high school to early adulthood. *Journal of Personality*, *80*, 1061-1089.
- Pasch, L. A., Bradbury, T. N., & Davila, J. (1997). Gender, negative affectivity, and observed social support behavior in marital interaction. *Personal Relationships*, *4*, 361-378.
- Paulus, M. P., Feinstein, J. S., Castillo, G., Simmons, A. N., & Stein, M. B. (2005). Dose-dependent decrease of activation in bilateral amygdala and insula by lorazepam during emotion processing. *Archives of General Psychiatry*, *62*, 282-288.
- Peeters, F., Nicolson, N. A., Berkhof, J., Delespaul, P., & deVries, M. (2003). Effects of daily events on mood states in major depressive disorder. *Journal of Abnormal Psychology*, *112*(2), 203-211.
- Perez-Edgar, K., Bar-Haim, Y., McDermott, J. M., Chronis-Tuscano, A., Pine, D. S., & Fox, N. A. (2010). Attention biases to threat and behavioral inhibition in early childhood shape adolescent social withdrawal. *Emotion*, *10*, 349-357.

- Perez-Edgar, K., McDermott, J. N., Korelitz, K., Degnan, K. A., Curby, T. W., Pine, D. S., & Fox, N. A. (2010). Patterns of sustained attention in infancy shape the developmental trajectory of social behavior from toddlerhood through adolescence. *Developmental Psychology, 46*, 1723-1730.
- Perez-Edgar, K., Reeb-Sutherland, B. C., McDermott, J. M., White, L. K., Henderson, H. A., Degnan, K. A., . . . Fox, N. A. (2011). Attention biases to threat link behavioral inhibition to social withdrawal over time in very young children. *Journal of Abnormal Child Psychology, 39*(6), 885-895.
- Perusini, J. N., & Fanselow, M. S. (2015). Neurobehavioral perspectives on the distinction between fear and anxiety. *Learning and Memory, 22*, 417-425.
- Petersen, S. E., & Sporns, O. (2015). Brain networks and cognitive architectures. *Neuron, 88*, 207-219.
- Pfeiffer, U. J., Vogeley, K., & Schilbach, L. (2013). From gaze cueing to dual eye-tracking: novel approaches to investigate the neural correlates of gaze in social interaction. *Neuroscience and Biobehavioral Reviews, 37*, 2516-2528.
- Phan, K. L., Coccaro, E. F., Angstadt, M., Kreger, K. J., Mayberg, H. S., Liberzon, I., & Stein, M. B. (2013). Corticolimbic brain reactivity to social signals of threat before and after sertraline treatment in generalized social phobia. *Biological Psychiatry, 73*(4), 329-336.
- Phillips, A. C., Batty, G. D., Weiss, A., Deary, I., Gale, C. R., Thomas, G. N., & Carroll, D. (2010). Neuroticism, cognitive ability, and the metabolic syndrome: The Vietnam Experience Study. *Journal of Psychosomatic Research, 69*(2), 193-201.
- Pichon, S., Miendlarzewska, E. A., Eryilmaz, H., & Vuilleumier, P. (2015). Cumulative activation during positive and negative events and state anxiety predicts subsequent inertia of amygdala reactivity. *Soc Cogn Affect Neurosci, 10*, 180-190.
- Pizzagalli, D. A. (2014). Depression, stress, and anhedonia: toward a synthesis and integrated model. *Annu Rev Clin Psychol, 10*, 393-423.
- Plasencia, M. L., Alden, L. E., & Taylor, C. T. (2011). Differential effects of safety behaviour subtypes in social anxiety disorder. *Behaviour Research and Therapy, 49*, 665-675.

- Polderman, T. J., Benyamin, B., de Leeuw, C. A., Sullivan, P. F., van Bochoven, A., Visscher, P. M., & Posthuma, D. (*in press*). Meta-analysis of the heritability of human traits based on fifty years of twin studies. *Nature Genetics*.
- Poldrack, R. A., Baker, C. I., Durnez, J., Gorgolewski, K. J., Matthews, P. M., Munafò, M., . . . Yarkoni, T. (2016). Scanning the horizon: Future challenges for neuroimaging research. *bioRxiv*.
- Pollak, S. D., & Kistler, D. J. (2002). Early experience is associated with the development of categorical representations for facial expressions of emotion. *Proceedings of the National Academy of Sciences of the United States of America*, *99*, 9072-9076.
- Power, R. A., & Pluess, M. (2015). Heritability estimates of the Big Five personality traits based on common genetic variants. *Transl Psychiatry*, *5*, e604.
- Power, R. A., Wingenbach, T., Cohen-Woods, S., Uher, R., Ng, M. Y., Butler, A. W., . . . McGuffin, P. (2013). Estimating the heritability of reporting stressful life events captured by common genetic variants. *Psychological Medicine*, *43*, 1965-1971.
- Pressman, S. D., Cohen, S., Miller, G. E., Barkin, A., Rabin, B. S., & Treanor, J. J. (2005). Loneliness, social network size, and immune response to influenza vaccination in college freshmen. *Health Psychology*, *24*, 297-306.
- Preuss, T. M. (2007). Primate brain evolution in phylogenetic context. In J. H. Kaas & T. M. Preuss (Eds.), *Evolution of Nervous Systems* (Vol. 4, pp. 3-34). NY: Elsevier.
- Prinz, P., Stams, G. J. J. M., Dekovic, M., Reijntjes, A. H. A., & Belsky, J. (2009). The relations between parents? Big Five personality factors and parenting: A meta-analytic review. *Journal of Personality and Social Psychology*, *97*, 351-362.
- Quilty, L. C., De Fruyt, F., Rolland, J. P., Kennedy, S. H., Rouillon, P. F., & Bagby, R. M. (2008). Dimensional personality traits and treatment outcome in patients with major depressive disorder. *Journal of Affective Disorders*, *108*, 241-250.

- Reeb-Sutherland, B. C., Helfinstein, S. M., Degnan, K. A., Perez-Edgar, K., Henderson, H. A., Lissek, S., . . . Fox, N. A. (2009). Startle response in behaviorally inhibited adolescents with a lifetime occurrence of anxiety disorders. *J Am Acad Child Adolesc Psychiatry, 48*, 610-617.
- Reinhart, R. M., & Woodman, G. F. (2014). Causal control of medial-frontal cortex governs electrophysiological and behavioral indices of performance monitoring and learning. *Journal of Neuroscience, 34*(12), 4214-4227.
- Reiss, S. (1997). Trait anxiety: it's not what you think it is. *Journal of Anxiety Disorders, 11*, 201-214.
- Reuter, M., Cooper, A. J., Smillie, L. D., Markett, S., & Montag, C. (2015). A new measure for the revised reinforcement sensitivity theory: psychometric criteria and genetic validation. *Front Syst Neurosci, 9*, 38.
- Reznick, J. S., Kagan, J., Snidman, N., Gersten, M., Baak, K., & Rosenberg, A. (1986). Inhibited and uninhibited children: A follow-up study. *Child Development, 57*, 660-680.
- Roberts, B. W., Caspi, A., & Moffitt, T. E. (2003). Work experiences and personality development in young adulthood. *Journal of Personality and Social Psychology, 84*, 582-593.
- Roberts, B. W., & DelVecchio, W. F. (2000). The rank-order consistency of personality traits from childhood to old age: a quantitative review of longitudinal studies. *Psychological Bulletin, 126*, 3-25.
- Roberts, B. W., Kuncel, N. R., Shiner, R., Caspi, A., & Goldberg, L. R. (2007). The power of personality. The comparative validity of personality traits, socioeconomic status, and cognitive ability for predicting important life outcomes. *Perspect Psychol Sci, 2*, 313-345.
- Roberts, B. W., & Mroczek, D. (2008). Personality trait change in adulthood. *Curr Dir Psychol Sci, 17*, 31-35.
- Roberts, B. W., Walton, K. E., & Viechtbauer, W. (2006). Patterns of mean-level change in personality traits across the life course: a meta-analysis of longitudinal studies. *Psychological Bulletin, 132*, 1-25.
- Robins, R. W., Caspi, A., & Moffitt, T. E. (2002). It's not just who you're with, it's who you are: Personality and relationship experiences across multiple relationships. *Journal of Personality, 70*, 925-964.

- Rogers, J., Raveendran, M., Fawcett, G. L., Fox, A. S., Shelton, S. E., Oler, J. A., . . . Kalin, N. H. (2013). CRHR1 genotypes, neural circuits and the diathesis for anxiety and depression. *Molecular Psychiatry*, *18*, 700-707.
- Romero-Canyas, R., Downey, G., Berenson, K., Ayduk, O., & J., K. (2010). Rejection sensitivity and the rejection-hostility link in romantic relationships. *Journal of Personality*, *78*, 119-148.
- Roseboom, P. H., Nanda, S. A., Fox, A. S., Oler, J. A., Shackman, A. J., Shelton, S. E., . . . Kalin, N. H. (2014). Neuropeptide Y receptor gene expression in the primate amygdala predicts anxious temperament and brain metabolism. *Biological Psychiatry*, *76*, 850-857.
- Rosen, J. B., & Schulkin, J. (1998). From normal fear to pathological anxiety. *Psychological Review*, *105*, 325-350.
- Rubin, K., Bowker, J., & Gazelle, H. (2010). Social withdrawal in childhood and adolescence: Peer relationships and social competence. In K. H. Rubin & R. J. Coplan (Eds.), *The development of shyness and social withdrawal* (pp. 131-154). NY: Guilford.
- Rubin, K. H., & Asendorpf, J. (1993). Social withdrawal, inhibition, and shyness in childhood: Conceptual and definitional issues. In K. H. Rubin & J. B. Asendorpf (Eds.), *Social withdrawal, inhibition and shyness in children* (pp. 3-17). Hillsdale, NJ: Erlbaum.
- Rubin, K. H., Wojslawowicz, J. C., Rose-Krasnor, L., Booth-LaForce, C., & Burgess, K. B. (2006). The best friendships of shy/withdrawn children: Prevalence, stability, and relationship quality. *Journal of Abnormal Child Psychology*, *34*, 143-157.
- Rudebeck, P. H., Saunders, R. C., Prescott, A. T., Chau, L. S., & Murray, E. A. (2013). Prefrontal mechanisms of behavioral flexibility, emotion regulation and value updating. *Nature Neuroscience*, *16*, 1140-1145.
- Sano, A., Phillips, A. J., Yu, A. Z., McHill, A. W., Taylor, S., Jaques, N., . . . Picard, R. W. (2015). *Recognizing academic performance, sleep quality, stress level, and mental health using personality traits*,

wearable sensors and mobile phones. Paper presented at the 12th International IEEE Conference on Wearable and Implantable Body Sensor Networks.

- Satpute, A. B., Wager, T. D., Cohen-Adad, J., Bianciardi, M., Choi, J. K., Buhle, J. T., . . . Barrett, L. F. (2013). Identification of discrete functional subregions of the human periaqueductal gray. *Proceedings of the National Academy of Sciences of the United States of America*, *110*, 17101-17106.
- Saudino, K. J., Pedersen, N. L., Lichtenstein, P., McClearn, G. E., & Plomin, R. (1997). Can personality explain genetic influences on life events? *Journal of Personality and Social Psychology*, *72*, 196-206.
- Scherer, K. R. (1986). Studying emotion empirically: Issues and a paradigm for research. In K. R. Scherer, H. G. Wallbott & A. B. Summerfield (Eds.), *Experiencing emotion: A cross-cultural study* (pp. 1-27). NY: Cambridge University Press.
- Schmidt, L. A., & Fox, N. A. (1998). Fear-potentiated startle responses in temperamentally different human infants. *Developmental Psychobiology*, *32*, 113-120.
- Schmitt, D. P., Realo, A., Voracek, M., & Allik, J. (2008). Why can't a man be more like a woman? Sex differences in Big Five personality traits across 55 cultures. *Journal of Personality and Social Psychology*, *94*, 168-182.
- Schnyer, D. M., Beevers, C. G., deBettencourt, M. T., Sherman, S. M., Cohen, J. D., Norman, K. A., & Turk-Browne, N. B. (2015). Neurocognitive therapeutics: from concept to application in the treatment of negative attention bias. *Biol Mood Anxiety Disord*, *5*, 1.
- Schuyler, B. S., Kral, T. R., Jacquart, J., Burghy, C. A., Weng, H. Y., Perlman, D. M., . . . Davidson, R. J. (2012). Temporal dynamics of emotional responding: amygdala recovery predicts emotional traits. *Soc Cogn Affect Neurosci*, *9*, 176-181.
- Scott, J., Williams, J. M., Brittlebank, A., & Ferrier, I. N. (1995). The relationship between premorbid neuroticism, cognitive dysfunction and persistence of depression: a 1-year follow-up. *Journal of Affective Disorders*, *33*, 167-172.

- Seo, D., Tsou, K. A., Ansell, E. B., Potenza, M. N., & Sinha, R. (2014). Cumulative adversity sensitizes neural response to acute stress: association with health symptoms. *Neuropsychopharmacology, 39*, 670-680.
- Servaas, M. N., Geerligs, L., Renken, R. J., Marsman, J. B., Ormel, J., Riese, H., & Aleman, A. (2014). Connectomics and neuroticism: An altered functional network organization. *Neuropsychopharmacology, 40*, 296-304.
- Shaban, H., Humeau, Y., Herry, C., Cassasus, G., Shigemoto, R., Ciochi, S., . . . Luthi, A. (2006). Generalization of amygdala LTP and conditioned fear in the absence of presynaptic inhibition. *Nature Neuroscience, 9*(8), 1028-1035.
- Shackman, A. J., & Fox, A. S. (*in press*). Contributions of the central extended amygdala to fear and anxiety. *Journal of Neuroscience*.
- Shackman, A. J., Fox, A. S., Oler, J. A., Shelton, S. E., Davidson, R. J., & Kalin, N. H. (2013). Neural mechanisms underlying heterogeneity in the presentation of anxious temperament. *Proceedings of the National Academy of Sciences of the United States of America, 110*, 6145-6150.
- Shackman, A. J., Fox, A. S., Oler, J. A., Shelton, S. E., Oakes, T. R., Davidson, R. J., & Kalin, N. H. (*in press*). Heightened extended amygdala metabolism following threat characterizes the early phenotypic risk to develop anxiety-related psychopathology. *Molecular Psychiatry*.
- Shackman, A. J., Fox, A. S., & Seminowicz, D. A. (2015). The cognitive-emotional brain: Opportunities and challenges for understanding neuropsychiatric disorders. *Behavioral and Brain Sciences, 38*, e86.
- Shackman, A. J., Kaplan, C. M., Stockbridge, M. D., Tillman, R. M., Tromp, D. P. M., Fox, A. S., & Gamer, M. (*in press*). The neurobiology of anxiety and attentional biases to threat: Implications for understanding anxiety disorders in adults and youth. *Journal of Experimental Psychopathology*.
- Shackman, A. J., McMenamin, B. W., Maxwell, J. S., Greischar, L. L., & Davidson, R. J. (2009). Right dorsolateral prefrontal cortical activity and behavioral inhibition. *Psychological Science, 20*, 1500-1506.

- Shackman, A. J., Salomons, T. V., Slagter, H. A., Fox, A. S., Winter, J. J., & Davidson, R. J. (2011). The integration of negative affect, pain and cognitive control in the cingulate cortex. *Nature Reviews Neuroscience, 12*, 154-167.
- Shackman, A. J., Sarinopoulos, I., Maxwell, J. S., Pizzagalli, D. A., Lavric, A., & Davidson, R. J. (2006). Anxiety selectively disrupts visuospatial working memory. *Emotion, 6*, 40-61.
- Shackman, A. J., Stockbridge, M. D., LeMay, E. P., & Fox, A. S. (*in press*). The psychological and neurobiological bases of dispositional negativity. In A. S. Fox, R. C. Lapate, A. J. Shackman & R. J. Davidson (Eds.), *The nature of emotion. Fundamental questions* (2nd ed.). NY: Oxford University Press.
- Shamir-Essakow, G., Ungerer, J., Rapee, R., & Safier, R. (2004). Caregiving representations of mothers of behaviorally inhibited and uninhibited preschool children. *Developmental Psychology, 40*, 899-910.
- Shanahan, M. J., Bauldry, S., Roberts, B. W., Macmillan, R., & Russo, R. (2014). Personality and the reproduction of social class. *Social Forces, 93*, 209-240.
- Sharot, T., Riccardi, A. M., Raio, C. M., & Phelps, E. A. (2007). Neural mechanisms mediating optimism bias. *Nature, 450*, 102-105.
- Shechner, T., Wakschlag, N., Britton, J. C., Jarcho, J., Ernst, M., & Pine, D. S. (2013). Empirical examination of the potential adverse psychological effects associated with pediatric fMRI scanning. *Journal of Child and Adolescent Psychopharmacology, 23*, 357-362.
- Sheline, Y. I., Barch, D. M., Donnelly, J. M., Ollinger, J. M., Snyder, A. Z., & Mintun, M. A. (2001). Increased amygdala response to masked emotional faces in depressed subjects resolves with antidepressant treatment: an fMRI study. *Biological Psychiatry, 50*(9), 651-658.
- Shiba, Y., Santangelo, A. M., Braesicke, K., Agustin-Pavon, C., Cockcroft, G., Haggard, M., & Roberts, A. C. (2014). Individual differences in behavioral and cardiovascular reactivity to emotive stimuli and their relationship to cognitive flexibility in a primate model of trait anxiety. *Front Behav Neurosci, 8*, 137.

- Shiner, R. L. (*in press-a*). Personality as lasting individual differences in emotions. In A. S. Fox, R. C. Lapate, A. J. Shackman & R. J. Davidson (Eds.), *The nature of emotion. Fundamental questions*. (2nd ed.). NY: Oxford University Press.
- Shiner, R. L. (*in press-b*). What develops in emotional development? In A. S. Fox, R. C. Lapate, A. J. Shackman & R. J. Davidson (Eds.), *The nature of emotion. Fundamental questions* (2nd ed.). NY: Oxford University Press.
- Shiner, R. L., Buss, K. A., McClowry, S., Putnam, S., Saudino, K., & Zentner, M. (2012). What is temperament now? Assessing progress in temperament research on the twenty-fifth anniversary of Goldsmith et al. (1987). *Child Development Perspectives*, 6, 436-444.
- Smit, D. J. A., Posthuma, D., Boomsma, D. I., & De Geus, E. J. (2007). The relation between frontal EEG asymmetry and the risk for anxiety and depression. *Biological Psychology*, 74(1), 26-33.
- Smith, D. J., Escott-Price, V., Davies, G., Bailey, M. E. S., Conde, L. C., Ward, J., . . . O'Donovan, M. (2015). Genome-wide analysis of over 106,000 individuals identifies 9 neuroticism-associated loci. *bioRxiv*.
- Snyder, D. K., & Halford, W. K. (2012). Evidence-based couple therapy: current status and future directions. *Journal of Family Therapy*, 34, 229-249.
- Soldz, S., & Vaillant, G. E. (1999). The big five personality traits and the life course: A 45-year longitudinal study. *Journal of Research in Personality*, 33, 208-232.
- Solomon, B. C., & Jackson, J. J. (2014). Why do personality traits predict divorce? Multiple pathways through satisfaction. *Journal of Personality and Social Psychology*, 106, 978-996.
- Somerville, L. H., Heatherton, T. F., & Kelley, W. M. (2006). Anterior cingulate cortex responds differentially to expectancy violation and social rejection. *Nature Neuroscience*, 9, 1007-1008.
- Somerville, L. H., Wagner, D. D., Wig, G. S., Moran, J. M., Whalen, P. J., & Kelley, W. M. (2013). Interactions between transient and sustained neural signals support the generation and regulation of anxious emotion. *Cerebral Cortex*, 23, 49-60.

- Somerville, L. H., Whalen, P. J., & Kelley, W. M. (2010). Human bed nucleus of the stria terminalis indexes hypervigilant threat monitoring. *Biological Psychiatry, 68*, 416-424.
- Soskin, D. P., Carl, J. R., Alpert, J., & Fava, M. (2012). Antidepressant effects on emotional temperament: toward a biobehavioral research paradigm for major depressive disorder. *CNS Neurosci Ther, 18*, 441-451.
- Soto, C. J., & John, O. P. (2014). Traits in transition: the structure of parent-reported personality traits from early childhood to early adulthood. *Journal of Personality, 82*, 182-199.
- Soto, C. J., & John, O. P. (*in press*). The next Big Five Inventory (BFI-2): Developing and assessing a hierarchical model with 15 Facets to enhance bandwidth, fidelity, and predictive power. *Journal of Personality and Social Psychology*.
- Soto, C. J., John, O. P., Gosling, S. D., & Potter, J. (2011). Age differences in personality traits from 10 to 65: Big Five domains and facets in a large cross-sectional sample. *Journal of Personality and Social Psychology, 100*, 330-348.
- Soto, C. J., & Luhmann, M. (2013). Who can buy happiness? Personality traits moderate the effects of stable income differences and income fluctuations on life satisfaction. *Social Psychological and Personality Science, 4*, 46-53.
- Specht, J., Egloff, B., & Schmukle, S. C. (2011). Stability and change of personality across the life course: the impact of age and major life events on mean-level and rank-order stability of the Big Five. *Journal of Personality and Social Psychology, 101*, 862-882.
- Spielberger, C. D. (1966). Theory and research on anxiety. In C. D. Spielberger (Ed.), *Anxiety and behavior* (pp. 3-22). NY: Academic Press.
- Spunt, R. P., Elison, J. T., Dufour, N., Hurlemann, R., Saxe, R., & Adolphs, R. (2015). Amygdala lesions do not compromise the cortical network for false-belief reasoning. *Proceedings of the National Academy of Sciences of the United States of America, 112*, 4827-4832.

- Staw, B. M., Bell, N. E., & Clausen, J. A. (1986). The dispositional approach to job attitudes: A lifetime longitudinal test. *Administrative Science Quarterly*, *31*, 56-77.
- Steel, P., Schmidt, J., & Shultz, J. (2008). Refining the relationship between personality and subjective well-being. *Psychological Bulletin*, *134*, 138-161.
- Stein, M. B., Simmons, A. N., Feinstein, J. S., & Paulus, M. P. (2007). Increased amygdala and insula activation during emotion processing in anxiety-prone subjects. *American Journal of Psychiatry*, *164*, 318-327.
- Stelzer, J., Lohmann, G., Mueller, K., Buschmann, T., & Turner, R. (2014). Deficient approaches to human neuroimaging. *Front Hum Neurosci*, *8*, 462.
- Steunenberg, B., Beekman, A. T., Deeg, D. J., & Kerkhof, A. J. (2010). Personality predicts recurrence of late-life depression. *Journal of Affective Disorders*, *123*, 164-172.
- Steward, O., & Balice-Gordon, R. (2014). Rigor or mortis: best practices for preclinical research in neuroscience. *Neuron*, *84*, 572-581.
- Stewart, S. L., & Rubin, K. H. (1995). The social problem-solving skills of anxious-withdrawn children. *Development and Psychopathology*, *7*, 323-336.
- Stoeckel, L. E., Garrison, K. A., Ghosh, S., Wightton, P., Hanlon, C. A., Gilman, J. M., . . . Evins, A. E. (2014). Optimizing real time fMRI neurofeedback for therapeutic discovery and development. *Neuroimage Clin*, *5*, 245-255.
- Stokes, J. J. (1985). The relation of social network and individual difference variables to loneliness. *Journal of Personality and Social Psychology*, *48*, 981-990.
- Stout, D. M., Shackman, A. J., Johnson, J. S., & Larson, C. L. (2014). Worry is associated with impaired gating of threat from working memory. *Emotion*, *15*, 6-11.
- Stout, D. M., Shackman, A. J., & Larson, C. L. (2013). Failure to filter: Anxious individuals show inefficient gating of threat from working memory. *Frontiers in Human Neuroscience*, *7*, doi: 10.3389/fnhum.2013.00058.

- Straube, T., Mentzel, H. J., & Miltner, W. H. R. (2007). Waiting for spiders: Brain activation during anticipatory anxiety in spider phobics. *Neuroimage*, *37*, 1427-1436.
- Strauss, C. C., Frame, C. L., & Forehand, R. (1987). Psychosocial impairment associated with anxiety in children. *J. Clin. Child Psychol*, *16*, 235-239.
- Suls, J., Green, P. J., & Hillis, S. (1998). Emotional reactivity to everyday problems, affective inertia, and neuroticism. *Personality and Social Psychology Bulletin*, *24*, 127-136.
- Suls, J., & Martin, R. (2005). The daily life of the garden-variety neurotic: Reactivity, stressor exposure, mood spillover, and maladaptive coping. *Journal of Personality*, *73*, 1485-1509.
- Sutin, A., Costa, R., R., M., & Eaton, W. (2009). Personality and career success: Concurrent and longitudinal relations. *European Journal of Personality*, *23*, 71-84.
- Swartz, J. R., Knodt, A. R., Radtke, S. R., & Hariri, A. R. (2015). A neural biomarker of psychological vulnerability to future life stress. *Neuron*, *85*(3), 505-511.
- Swartz, J. R., Williamson, D. E., & Hariri, A. R. (2015). Developmental change in amygdala reactivity during adolescence: effects of family history of depression and stressful life events. *American Journal of Psychiatry*, *172*(3), 276-283.
- Tan, P. Z., Forbes, E. E., Dahl, R. E., Ryan, N. D., Siegle, G. J., Ladouceur, C. D., & Silk, J. S. (2012). Emotional reactivity and regulation in anxious and nonanxious youth: a cell-phone ecological momentary assessment study. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *53*, 197-206.
- Tang, D. W., Fellows, L. K., Small, D. M., & Dagher, A. (2012). Food and drug cues activate similar brain regions: a meta-analysis of functional MRI studies. *Physiology and Behavior*, *106*, 317-324.
- Taylor, C. T., & Alden, L. E. (2011). To see ourselves as others see us: an experimental integration of the intra and interpersonal consequences of self-protection in social anxiety disorder. *Journal of Abnormal Psychology*, *120*(1), 129-141.
- ten Have, M., Oldehinkel, A., Vollebergh, W., & Ormel, J. (2005). Does neuroticism explain variations in care service use for mental health problems in the general population? Results from the

- Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Social Psychiatry and Psychiatric Epidemiology*, *40*, 425-431.
- Terracciano, A., Lockenhoff, C. E., Zonderman, A. B., Ferrucci, L., & Costa, P. T., Jr. (2008). Personality predictors of longevity: activity, emotional stability, and conscientiousness. *Psychosomatic Medicine*, *70*, 621-627.
- Thibodeau, R., Jorgensen, R. S., & Kim, S. (2006). Depression, anxiety, and resting frontal EEG asymmetry: a meta-analytic review. *Journal of Abnormal Psychology*, *115*, 715-729.
- Thomas, K. M., Drevets, W. C., Dahl, R. E., Ryan, N. D., Birmaher, B., Eccard, C. H., . . . Casey, B. J. (2001). Amygdala response to fearful faces in anxious and depressed children. *Archives of General Psychiatry*, *58*, 1057-1063.
- Tomarken, A. J., Davidson, R. J., Wheeler, R. E., & Kinney, L. (1992). Psychometric properties of resting anterior EEG asymmetry: Temporal stability and internal consistency. *Psychophysiology*, *29*, 576-592.
- Törnqvist, E., Månsson, Å., Larsson, E. M., & Hallström, I. (2006). It's like being in another world—patients' lived experience of magnetic resonance imaging. *Journal of Clinical Nursing*, *15*, 954-961.
- Tovote, P., Fadok, J. P., & Luthi, A. (2015). Neuronal circuits for fear and anxiety. *Nature Reviews Neuroscience*, *16*, 317-331.
- Towers, D. N., & Allen, J. J. B. (2009). A better estimate of the internal consistency reliability of frontal EEG asymmetry scores. *Psychophysiology*, *46*, 132-142.
- Tranel, D., Gullickson, G., Koch, M., & Adolphs, R. (2006). Altered experience of emotion following bilateral amygdala damage. *Cognitive Neuropsychiatry*, *11*, 219-232.
- Treadway, M. T., Buckholtz, J. W., Martin, J. W., Jan, K., Asplund, C. L., Ginther, M. R., . . . Marois, R. (2014). Corticolimbic gating of emotion-driven punishment. *Nature Neuroscience*, *17*, 1270-1275.
- Turk-Browne, N. B. (2013). Functional interactions as big data in the human brain. *Science*, *342*, 580-584.

- Turkheimer, E., Pettersson, E., & Horn, E. E. (2014). A phenotypic null hypothesis for the genetics of personality. *Annual Review of Psychology, 65*, 515-540.
- Turner, C. A., Clinton, S. M., Thompson, R. C., Watson, S. J., Jr., & Akil, H. (2011). Fibroblast growth factor-2 (FGF2) augmentation early in life alters hippocampal development and rescues the anxiety phenotype in vulnerable animals. *Proceedings of the National Academy of Sciences of the United States of America, 108*, 8021-8025.
- Tyc, V. L., Fairclough, D., Fletcher, B., Leigh, L., & Mulhern, R. K. (1995). Children's distress during magnetic resonance imaging procedures. *Children's Health Care, 24*, 5-19.
- Tye, K. M., Prakash, R., Kim, S. Y., Fenno, L. E., Grosenick, L., Zarabi, H., . . . Deisseroth, K. (2011). Amygdala circuitry mediating reversible and bidirectional control of anxiety. *Nature, 471*, 358-362.
- Uliaszek, A. A., Hauner, K. K., Zinbarg, R. E., Craske, M. G., Mineka, S., Griffith, J. W., & Rose, R. D. (2009). An examination of content overlap and disorder-specific predictions in the associations of neuroticism with anxiety and depression. *J Res Pers, 43*, 785-794.
- Uysal, S. D., & Pohlmeier, W. (2011). Unemployment duration and personality. *Journal of Economic Psychology, 32*, 980-992.
- Vaidyanathan, U., Patrick, C. J., & Cuthbert, B. N. (2009). Linking dimensional models of internalizing psychopathology to neurobiological systems: affect-modulated startle as an indicator of fear and distress disorders and affiliated traits. *Psychological Bulletin, 135*, 909-942.
- Vaisvaser, S., Lin, T., Admon, R., Podlipsky, I., Greenman, Y., Stern, N., . . . Hendler, T. (2013). Neural traces of stress: cortisol related sustained enhancement of amygdala-hippocampal functional connectivity. *Front Hum Neurosci, 7*, 313.
- van de Leemput, I. A., Wichers, M., Cramer, A. O., Borsboom, D., Tuerlinckx, F., Kuppens, P., . . . Scheffer, M. (2014). Critical slowing down as early warning for the onset and termination of depression. *Proceedings of the National Academy of Sciences of the United States of America, 111*(1), 87-92.

- van den Berg, S. M., de Moor, M. H., McGue, M., Pettersson, E., Terracciano, A., Verweij, K. J., . . . Boomsma, D. I. (2014). Harmonization of neuroticism and extraversion phenotypes across inventories and cohorts in the Genetics of Personality Consortium: an application of item response theory. *Behav Genet, 44*(4), 295-313.
- van Honk, J., Eisenegger, C., Terburg, D., Stein, D. J., & Morgan, B. (2013). Generous economic investments after basolateral amygdala damage. *Proceedings of the National Academy of Sciences of the United States of America, 110*, 2506-2510.
- van Marle, H. J., Hermans, E. J., Qin, S., & Fernandez, G. (2009). From specificity to sensitivity: how acute stress affects amygdala processing of biologically salient stimuli. *Biological Psychiatry, 66*, 649-655.
- van Marle, H. J., Hermans, E. J., Qin, S., & Fernandez, G. (2010). Enhanced resting-state connectivity of amygdala in the immediate aftermath of acute psychological stress. *Neuroimage, 53*, 348-354.
- van Os, J., Park, S. B., & Jones, P. B. (2001). Neuroticism, life events and mental health: evidence for person-environment correlation. *British Journal of Psychiatry. Supplement, 40*, s72-77.
- van Well, S., Visser, R. M., Scholte, H. S., & Kindt, M. (2012). Neural substrates of individual differences in human fear learning: evidence from concurrent fMRI, fear-potentiated startle, and US-expectancy data. *Cogn Affect Behav Neurosci, 12*, 499-512.
- van Wingen, G. A., Geuze, E., Vermetten, E., & Fernandez, G. (2011). Perceived threat predicts the neural sequelae of combat stress. *Molecular Psychiatry, 16*(6), 664-671.
- van Winkel, M., Nicolson, N. A., Wichers, M., Viechtbauer, W., Myin-Germeys, I., & Peeters, F. (2015). Daily life stress reactivity in remitted versus non-remitted depressed individuals. *Eur Psychiatry, 30*(4), 441-447.
- Vater, A., & Schröder-Abé, M. (2015). Explaining the link between personality and relationship satisfaction: Emotion regulation and interpersonal behavior in conflict discussions. *European Journal of Personality, 29*, 201-215.

- Viinikainen, J., Kokko, K., Pulkkinen, L., & Pehkonen, J. (2010). Personality and labour market income: Evidence from longitudinal data. *Labour, 24*, 201-220.
- Vinkers, C. H., Joels, M., Milaneschi, Y., Kahn, R. S., Penninx, B. W., & Boks, M. P. (2014). Stress exposure across the life span cumulatively increases depression risk and is moderated by neuroticism. *Depression and Anxiety, 31*, 737-745.
- Vollrath, M. (2000). Personality and hassles among university students: A three-year longitudinal study. *European Journal of Personality, 14*, 199-215.
- Vukasovic, T., & Bratko, D. (2015). Heritability of personality: A meta-analysis of behavior genetic studies. *Psychological Bulletin, 141*, 769-785.
- Wacker, J., Chavanon, M.-L., & Stemmler, G. (2010). Resting EEG signatures of agentic extraversion: New results and meta-analytic integration. *Journal of Research in Personality, 44*, 167-179.
- Wager, T. D., Atlas, L. Y., Lindquist, M. A., Roy, M., Woo, C. W., & Kross, E. (2013). An fMRI-based neurologic signature of physical pain. *New England Journal of Medicine, 368*(15), 1388-1397.
- Walker, D. L., & Davis, M. (1997). Double dissociation between the involvement of the bed nucleus of the stria terminalis and the central nucleus of the amygdala in startle increases produced by conditioned versus unconditioned fear. *Journal of Neuroscience, 17*, 9375-9383.
- Walker, D. L., & Davis, M. (2008). Role of the extended amygdala in short-duration versus sustained fear: a tribute to Dr. Lennart Heimer. *Brain Struct Funct, 213*, 29-42.
- Walker, D. L., Miles, L. A., & Davis, M. (2009). Selective participation of the bed nucleus of the stria terminalis and CRF in sustained anxiety-like versus phasic fear-like responses. *Progress in Neuro-Psychopharmacology and Biological Psychiatry, 33*, 1291-1308.
- Walker, D. L., Toufexis, D. J., & Davis, M. (2003). Role of the bed nucleus of the stria terminalis versus the amygdala in fear, stress, and anxiety. *European Journal of Pharmacology, 563*, 199-216.
- Walker, O. L., Degnan, K. A., Fox, N. A., & Henderson, H. A. (in press). Early social fear in relation to play with an unfamiliar peer: Actor and partner effects. *Developmental Psychology*.

- Wang, S. W., Repetti, R. L., & Campos, B. (2011). Job stress and family social behavior: the moderating role of neuroticism. *Journal of Occupational Health Psychology, 16*, 441-456.
- Waters, A. M., Neumann, D. L., Henry, J., Craske, M. G., & Ornitz, E. M. (2008). Baseline and affective startle modulation by angry and neutral faces in 4-8-year-old anxious and non-anxious children. *Biological Psychology, 78*, 10-19.
- Watson, D., & Clark, L. A. (1984). Negative affectivity: the disposition to experience aversive emotional states. *Psychological Bulletin, 96*, 465-490.
- Watson, D., & Clark, L. A. (1992). On traits and temperament: general and specific factors of emotional experience and their relation to the five-factor model. *Journal of Personality, 60*(2), 441-476.
- Watson, D., Hubbard, B., & Wiese, D. (2000). General traits of personality and affectivity as predictors of satisfaction in intimate relationships: Evidence from self- and partner-ratings. *Journal of Personality, 68*, 413-449.
- Watson, D., & Humrichouse, J. (2006). Personality development in emerging adulthood: Integrating evidence from self-ratings and spouse ratings. *Journal of Personality & Social Psychology, 91*, 959-974.
- Watson, D., & Naragon-Gainey, K. (2014). Personality, emotions, and the emotional disorders. *Clinical Psychological Science, 2*, 422-442.
- Wayne, J. H., Musisca, N., & Fleeson, W. (2004). Considering the role of personality in the work-family experience: Relationships of the big five to work-family conflict and facilitation. *Journal of Vocational Behavior, 64*, 108-130.
- Weissman, M. M., Prusoff, B. A., & Klerman, G. L. (1978). Personality and the prediction of long-term outcome of depression. *American Journal of Psychiatry, 135*, 797-800.
- Weston, S. J., Hill, P. L., & Jackson, J. J. (2015). Personality traits predict the onset of disease. *Social Psychological and Personality Science, 6*, 309-317.

- Wetter, E. K., & Hankin, B. L. (2009). Mediation pathways through which positive and negative emotionality contribute to anhedonic symptoms of depression: a prospective study of adolescents. *Journal of Abnormal Child Psychology, 37*, 507-520.
- Wheeler, R. E., Davidson, R. J., & Tomarken, A. J. (1993). Frontal brain asymmetry and emotional reactivity: A biological substrate of affective style. *Psychophysiology, 30*, 82-89.
- White, L. K., Degnan, K. A., Henderson, H. A., Pérez-Edgar, K. A., Walker, O. L., Shechner, T., . . . Fox, N. A. (in press). Developmental relations between behavioral inhibition, anxiety, and attention biases to threat and positive information. *Child Development*.
- Whiteford, H. A., Degenhardt, L., Rehm, J., Baxter, A. J., Ferrari, A. J., Erskine, H. E., . . . Vos, T. (2013). Global burden of disease attributable to mental and substance use disorders: findings from the Global Burden of Disease Study 2010. *Lancet, 382*, 1575-1586.
- Whittington, J. E., & Huppert, F. A. (1998). Neuroticism, psychiatric symptoms and life events. *Personality and Individual Differences, 24*, 97-107.
- Wichers, M. C., Barge-Schaapveld, D. Q., Nicolson, N. A., Peeters, F., de Vries, M., Mengelers, R., & van Os, J. (2009). Reduced stress-sensitivity or increased reward experience: the psychological mechanism of response to antidepressant medication. *Neuropsychopharmacology, 34*(4), 923-931.
- Widiger, T. A. (2009). Neuroticism. In M. R. Leary & R. H. Hoyle (Eds.), *Handbook of individual differences in social behavior* (pp. 129-146). NY: Guilford.
- Wilson, R. E., Harris, K. M., & Vazire, S. (2015). Personality and friendship satisfaction in daily life: Do everyday social interactions account for individual differences in friendship satisfaction. *European Journal of Personality, 29*, 173-186.
- Wilson, R. S., Krueger, K. R., Gu, L., Bienias, J. L., Mendes de Leon, C. F., & Evans, D. A. (2005). Neuroticism, extraversion, and mortality in a defined population of older persons. *Psychosomatic Medicine, 67*, 841-845.

- Wilson, S., Vaidyanathan, U., Miller, M. B., McGue, M., & Iacono, W. G. (2014). Premorbid risk factors for major depressive disorder: are they associated with early onset and recurrent course? *Development and Psychopathology, 26*(4 Pt 2), 1477-1493.
- Wilson, S. J., Smyth, J. M., & MacLean, R. R. (2014). Integrating ecological momentary assessment and functional brain imaging methods: new avenues for studying and treating tobacco dependence. *Nicotine Tob Res, 16 Suppl 2*, S102-110.
- Windischberger, C., Lanzenberger, R., Holik, A., Spindelegger, C., Stein, P., Moser, U., . . . Kasper, S. (2010). Area-specific modulation of neural activation comparing escitalopram and citalopram revealed by pharmacofMRI: a randomized cross-over study. *Neuroimage, 49*, 1161-1170.
- Winslow, J. T., Noble, P. L., & Davis, M. (2007). Modulation of fear-potentiated startle and vocalizations in juvenile rhesus monkeys by morphine, diazepam, and buspirone. *Biological Psychiatry, 61*, 389-395.
- Wise, R. A., & Koob, G. F. (2014). The development and maintenance of drug addiction. *Neuropsychopharmacology, 39*(2), 254-262.
- Wise, R. G., & Preston, C. (2010). What is the value of human FMRI in CNS drug development? *Drug Discovery Today, 15*, 973-980.
- Wolitzky-Taylor, K., Vrshek-Schallhorn, S., Waters, A. M., Mineka, S., Zinbarg, R., Ornitz, E., . . . Craske, M. G. (2014). Adversity in early and mid-adolescence is associated with elevated startle responses to safety cues in late adolescence. *Clin Psychol Sci, 2*, 202-213.
- Woo, C. W., Koban, L., Kross, E., Lindquist, M. A., Banich, M. T., Ruzic, L., . . . Wager, T. D. (2014). Separate neural representations for physical pain and social rejection. *Nat Commun, 5*, 5380.
- Wood, K. H., Ver Hoef, L. W., & Knight, D. C. (2014). The amygdala mediates the emotional modulation of threat-elicited skin conductance response. *Emotion, 14*, 693-700.
- Wrzus, C., & Mehl, M. R. (2015). Lab and/or field? Measuring personality processes and their social consequences. *European Journal of Personality, 29*, 250-271.

- Wupperman, P., & Neumann, C. S. (2006). Depressive symptoms as a function of sex-role, rumination, and neuroticism. *Personality and Individual Differences, 40*, 189-201.
- Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nat Methods, 8*, 665-670.
- Yassa, M. A., Hazlett, R. L., Stark, C. E., & Hoehn-Saric, R. (2012). Functional MRI of the amygdala and bed nucleus of the stria terminalis during conditions of uncertainty in generalized anxiety disorder. *Journal of Psychiatric Research, 46*, 1045-1052.
- Yilmazer-Hanke, D. M. (2012). Amygdala. In J. K. Mai & G. Paxinos (Eds.), *The human nervous system* (pp. 759-834). San Diego: Academic Press.
- Zaider, T. I., Heimberg, R. G., & Iida, M. (2010). Anxiety disorders and intimate relationships: a study of daily processes in couples. *Journal of Abnormal Psychology, 119*, 163-173.
- Zautra, A. J., Affleck, G. G., Tennen, H., Reich, J. W., & Davis, M. C. (2005). Dynamic approaches to emotions and stress in everyday life: Bolger and Zuckerman reloaded with positive as well as negative affects. *Journal of Personality, 73*, 1511-1538.
- Zuckerman, M. (1976). General and situation-specific traits and states: new approaches to assessment of anxiety and other constructs. In M. Zuckerman & C. D. Spielberger (Eds.), *Emotions and anxiety: new concepts, methods and applications* (pp. 133-174). Hillsdale, NJ: Lawrence Erlbaum.