

UCSF

UC San Francisco Previously Published Works

Title

Large-Scale Phenotype-Based Antiepileptic Drug Screening in a Zebrafish Model of Dravet Syndrome

Permalink

<https://escholarship.org/uc/item/3r77r0zg>

Journal

eNeuro, 2(4)

ISSN

2373-2822

Authors

Dinday, Matthew T
Baraban, Scott C

Publication Date

2015-07-01

DOI

10.1523/eneuro.0068-15.2015

Peer reviewed

Disorders of the Nervous System

Large-Scale Phenotype-Based Antiepileptic Drug Screening in a Zebrafish Model of Dravet Syndrome^{1,2,3}

Matthew T. Dinday,¹ and Scott C. Baraban^{1,2}DOI:<http://dx.doi.org/10.1523/ENEURO.0068-15.2015>

¹Department of Neurological Surgery, Epilepsy Research Laboratory, University of California San Francisco, San Francisco, California 94143, ²Eli and Edythe Broad Center of Regeneration Medicine and Stem Cell Research, University of California San Francisco, San Francisco, California 94143

Abstract

Mutations in a voltage-gated sodium channel (*SCN1A*) result in Dravet Syndrome (DS), a catastrophic childhood epilepsy. Zebrafish with a mutation in *scn1Lab* recapitulate salient phenotypes associated with DS, including seizures, early fatality, and resistance to antiepileptic drugs. To discover new drug candidates for the treatment of DS, we screened a chemical library of ~1000 compounds and identified 4 compounds that rescued the behavioral seizure component, including 1 compound (dimethadione) that suppressed associated electrographic seizure activity. Fenfluramine, but not huperzine A, also showed antiepileptic activity in our zebrafish assays. The effectiveness of compounds that block neuronal calcium current (dimethadione) or enhance serotonin signaling (fenfluramine) in our zebrafish model suggests that these may be important therapeutic targets in patients with DS. Over 150 compounds resulting in fatality were also identified. We conclude that the combination of behavioral and electrophysiological assays provide a convenient, sensitive, and rapid basis for phenotype-based drug screening in zebrafish mimicking a genetic form of epilepsy.

Key words: antiepileptic; drug discovery; epilepsy; high throughput; pharmacology; zebrafish

Significance Statement

Dravet syndrome is a catastrophic childhood epilepsy that is resistant to available medications. Current animal models for this disease are not amenable to high-throughput drug screening. We used a zebrafish model for Dravet syndrome and screened >1000 compounds. We report the identification of compounds with the ability to suppress seizure behavior and electrographic seizure activity. This approach provides an example of precision medicine directed to pediatric epilepsy.

Introduction

Dravet syndrome (DS) is a devastating genetic epileptic encephalopathy that has been linked to more than >300

de novo mutations in a neuronal voltage-gated sodium channel (*SCN*). Children with DS are at a higher risk for sudden unexplained death in epilepsy and episodes of

Received June 18, 2015; accepted August 4, 2015; First published August 20, 2015.

¹The authors declare no competing financial interests.

²Author contributions: M.T.D. and S.C.B. designed research; M.T.D. and S.C.B. performed research; M.T.D. and S.C.B. analyzed data; S.C.B. wrote the paper.

³Funding was provided by National Institutes of Health-National Institute of Neurological Disorders and Stroke EUREKA Grant 5R01-NS-079214 and The Joseph & Vera Long Foundation to (S.C.B.).

Acknowledgments: We thank B. Grone and D. Lowenstein for comments on earlier versions of this manuscript.

Correspondence should be addressed to Scott C. Baraban, Department of Neurological Surgery, Epilepsy Research Laboratory, University of California, San Francisco, San Francisco, CA 94143. E-mail: scott.baraban@ucsf.edu.

DOI:<http://dx.doi.org/10.1523/ENEURO.0068-15.2015>

Copyright © 2015 Dinday and Baraban

This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use,

uncontrolled status epilepticus (Dravet et al., 2005; Ceulemans et al., 2012). Delayed language development, disruption of autonomic function, and motor and cognitive impairment are also associated with this disease. Seizure management includes treatment with benzodiazepines, valproate, and/or stiripentol (Caraballo et al., 2005; Chiron and Dulac, 2011). Some reduction in seizure activity has been reported with the use of bromides and topiramate, or a ketogenic diet (Lotte et al., 2012; Wilmshurst et al., 2014; Dressler et al., 2015). Despite these options, available antiepileptic drugs (AEDs) do not achieve adequate seizure control in most DS patients (Dravet et al., 2005; Chiron and Dulac, 2011; Dressler et al., 2015), making the identification of new drugs a critical unmet need. High-throughput screening offers a powerful tool to identify new drug candidates for these patients. However, commonly available screening approaches rely on *in vitro* cell-based assays (Masimirembwa et al., 2001; Snowden and Green, 2008; Ko and Gelb, 2014) and do not recapitulate the complicated neural networks that generate seizures *in vivo*. Given the need for new treatments for children with DS, and the growing number of genetic epileptic encephalopathies that are medically intractable (Leppert, 1990; Epi4K Consortium, 2012; Ottman and Risch, 2012), we developed an alternative phenotype-based *in vivo* drug-screening strategy. While cell-based *in vitro* screening assays can efficiently identify compounds that bind specific targets, whole-organism-based screens are more likely to reliably predict therapeutic outcomes as they maintain the complex neural circuitry involved in the underlying disease process. Whole-organism screens do not require well validated targets to identify compounds that yield a desirable phenotypic outcome, but can be prohibitively costly and time consuming in mammals. As a simple vertebrate with significant genetic similarity to human, zebrafish are now recognized as an ideal cost-effective alternative to achieve rapid *in vivo* phenotype-based screening (Ali et al., 2011).

Using *scn1a* mutant zebrafish larvae with a gene homologous to human and spontaneously occurring seizures (Baraban et al., 2013), we screened, in a blinded manner, a repurposed library of ~1000 compounds for drugs that inhibit unprovoked seizure events. We also screened two compounds (huperzine A and fenfluramine) that were discovered in rodent-based assays using acquired seizure protocols and that were recently suggested as potential treatments for DS (Boel and Casaer, 1996; Coleman et al., 2008; Ceulemans et al., 2012; Bialer et al., 2015). Only 20 compounds in the repurposed drug library reduced swim behavior to control levels. However, many of these compounds were toxic or were not confirmed on retesting, and only four compounds advanced to a second-stage *in vivo* electrophysiology assay. Of these compounds (cytarabine, dimethadione, theobromine, and norfloxacin) only dimethadione, a T-type calcium channel antagonist previously reported to have anticonvulsant activity (Lowson et al., 1990; Zhang et al., 1996), reduced

ictal-like electrographic discharges seen in *scn1Lab* mutant larvae. This two-stage phenotype-based screening approach, using a genetic DS model with >75% genomic similarity to human, is a sensitive, rapid means to successfully identify compounds with antiepileptic activity.

Materials and Methods

Zebrafish

Zebrafish were maintained in a light- and temperature-controlled aquaculture facility under a standard 14:10 h light/dark photoperiod. Adult zebrafish were housed in 1.5 L tanks at a density of 5–12 fish per tank and fed twice per day (dry flake and/or flake supplemented with live brine shrimp). Water quality was continuously monitored: temperature, 28–30° C; pH 7.4–8.0; conductivity, 690–710 mS/cm. Zebrafish embryos were maintained in round Petri dishes (catalog #FB0875712, Fisher Scientific) in “embryo medium” consisting of 0.03% Instant Ocean (Aquarium Systems, Inc.) and 000002% methylene blue in reverse osmosis-distilled water. Larval zebrafish clutches were bred from wild-type (WT; TL strain) or *scn1Lab* (*didy*^{s552}) heterozygous animals that had been backcrossed to TL wild-type for at least 10 generations. Homozygous mutants ($n = 6544$), which have widely dispersed melanosomes and appear visibly darker as early as 3 d postfertilization (dpf; Fig. 1b), or WT larvae ($n = 71$) were used in all experiments at 5 or 6 dpf. Embryos and larvae were raised in plastic petri dishes (90 mm diameter, 20 mm depth) and density was limited to ~60 per dish. Larvae between 3 and 7 dpf lack discernible sex chromosomes. The care and maintenance protocols comply with requirements outlined in the *Guide for the Care and Use of Animals* (eBrary Inc., 2011) and were approved by the Institutional Animal Care and Use Committee (protocol #AN108659-01D).

Seizure monitoring

Zebrafish larvae were placed individually into 1 well of a clear flat-bottomed 96-well microplate (catalog #260836, Fisher Scientific) containing embryo media. Microplates were placed inside an enclosed motion-tracking device and acclimated to the dark condition for 10–15 min at room temperature. Locomotion plots were obtained for one fish per well at a recording epoch of 10 min using a DanioVision system running EthoVision XT software (DanioVision, Noldus Information Technology); threshold detection settings to identify objects darker than the background were optimized for each experiment. Seizure scoring was performed using the following three-stage scale (Baraban et al., 2005): Stage 0, no or very little swim activity; Stage I, increased, brief bouts of swim activity; Stage II, rapid “whirlpool-like” circling swim behavior; and Stage III, paroxysmal whole-body clonus-like convulsions, and a brief loss of posture. WT fish are normally scored at Stage 0 or I. Plots were analyzed for distance traveled (in millimeters) and mean velocity (in millimeters per second). As reported previously (Winter et al., 2008; Baraban et al., 2013), velocity changes were a more sensitive assay of seizure behavior. For electrophysiology studies, zebrafish larvae were briefly paralyzed with

distribution and reproduction in any medium provided that the original work is properly attributed.




Figure 1. Locomotion assay to identify drugs that rescue the *scn1Lab* mutant epilepsy phenotype. **a**, Schematic of the phenotype-based screening process. Chemical libraries can be coded and aliquoted in small volumes ($75 \mu\text{L}$) into individual wells containing one mutant fish. The 96-well microplate is arranged so that six fish are tested per drug; with one row of six fish maintained as an internal control (red circles) on each plate. **b**, Representative images for WT and *scn1Lab* mutant zebrafish larvae at 5 dpf. Note the morphological similarity but darker pigmentation in mutant larvae. **c**, Box plot of mean velocity (in millimeters per second) for two consecutive recordings of mutant larvae in embryo media. Experiments were performed by first placing the mutant larvae in embryo media and obtaining a baseline locomotion response; embryo media was then replaced with new embryo media (to mimic the procedure used for test compounds), and a second locomotion response was obtained. The percentage change in velocity from baseline (recording 1) versus experimental model (recording 2) is shown. In the box plot, the bottom and top of the box represent the 25th percentile and the 75th percentile, respectively. The line across the box represents the median value, and the vertical lines encompass the entire range of values. This plot represents normal changes in tracking activity in the absence of a drug challenge. **d**, Plot of locomotor seizure behavior for *scn1Lab* mutants at 5 dpf for the 1012 compounds tested. Threshold for inhibition of seizure activity (positive hits) was set as a reduction in mean swim velocity of $\geq 44\%$; the threshold for a proconvulsant or hyperexcitable effect was set at an increase in the mean swim velocity of $\geq 44\%$ (green dashed lines).

α -bungarotoxin (1 mg/ml) and immobilized in 1.2% agarose; field recordings were obtained from forebrain structures. Epileptiform events were identified *post hoc* in Clampfit (Molecular Devices) and were defined as multipike or polyspike upward or downward membrane deflections greater than three times the baseline noise level and >500 ms in duration. During electrophysiology experiments zebrafish larvae were continuously monitored for the presence (or absence) of blood flow and heart beat by direct visualization on an Olympus BX51WI upright microscope equipped with a CCD camera and monitor.

Drugs

Compounds for drug screening were purchased from MicroSource Discovery Systems, Inc. (PHARMAKON 1600) and were provided as 10 mM DMSO solutions (Table 1). Test compounds for locomotion or electrophysiology studies were dissolved in embryo media and were tested at an initial concentration of $100 \mu\text{M}$, with a final DMSO concentration of $<2\%$. In all drug library screen studies, compounds were coded and experiments were performed by investigators who were blind to the nature of the compound. Baseline recordings of seizure behavior

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen.

ABACAVIR SULFATE
 ABAMECTIN (avermectin B1a shown)
 ACADESINE
 ACARBOSE
 ACEBUTOLOL HYDROCHLORIDE
 ACECLIDINE
 ACECLOFENAC
 ACENOCOUMAROL
 ACETAMINOPHEN
 ACETOHYDROXAMIC ACID
 ACETOPHENAZINE MALEATE
 ACETRIAZOIC ACID
 ACETYLCHOLINE CHLORIDE
 ACETYLCYSTEINE
 ACIPIMOX
 ACONITINE
 ACRIFLAVINIUM HYDROCHLORIDE
 ACRISORCIN
 ACTARIT
 ACYCLOVIR
 ADAPALENE
 ADELMIDROL
 ADENINE
 ADENOSINE
 ADENOSINE PHOSPHATE
 ADIPHENINE HYDROCHLORIDE
 AKLOMIDE
 ALAPROCLATE
 ALBENDAZOLE
 ALBUTEROL (+/-)
 ALENDRONATE SODIUM
 ALEXIDINE HYDROCHLORIDE
 ALLANTOIN
 ALLOPURINOL
 ALMOTRIPTAN
 alpha-TOCHOPHEROL
 alpha-TOCHOPHERYL ACETATE
 ALPRAZOLAM
 ALRESTATIN
 ALTHIAZIDE
 ALTRETAMINE
 ALVERINE CITRATE
 AMANTADINE HYDROCHLORIDE
 AMCINONIDE
 AMIFOSTINE
 AMIKACIN SULFATE
 AMILORIDE HYDROCHLORIDE
 AMINACRINE
 AMINOCAPROIC ACID
 AMINOGLUTETHIMIDE
 AMINOHIPPURIC ACID
 AMINOLEVULINIC ACID HYDROCHLORIDE
 AMINOSALICYLATE SODIUM
 AMITRIPTYLINE HYDROCHLORIDE
 AMLEXANOX
 AMLODIPINE BESYLATE
 AMODIAQUINE DIHYDROCHLORIDE
 AMOROLFINE HYDROCHLORIDE
 AMOXICILLIN
 AMPHOTERICIN B
 AMPICILLIN SODIUM

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

AMPROLIUM
 AMSACRINE
 ANASTROZOLE
 ANCITABINE HYDROCHLORIDE
 ANETHOLE
 ANIRACETAM
 ANISINDIONE
 ANTAZOLINE PHOSPHATE
 ANTHRALIN
 ANTIPYRINE
 APOMORPHINE HYDROCHLORIDE
 APRAMYCIN
 ARGININE HYDROCHLORIDE
 ARMODAFINIL
 ARTEMETHER
 ARTEMOTIL
 ARTESUNATE
 ASCORBIC ACID
 ASPIRIN
 ATENOLOL
 ATORVASTATIN CALCIUM
 ATOVAQUONE
 ATROPINE SULFATE
 AUROTHIOGLUCOSE
 AVOBENZONE
 AZACITIDINE
 AZASERINE
 AZATADINE MALEATE
 AZATHIOPRINE
 AZELAIC ACID
 AZITHROMYCIN
 AZLOCILLIN SODIUM
 AZTREONAM
 BACAMPICILLIN HYDROCHLORIDE
 BACITRACIN
 BACLOFEN
 BALSALAZIDE DISODIUM
 BECLOMETHASONE DIPROPIONATE
 BEKANAMYCIN SULFATE
 BEMOTRIZINOL
 BENAZEPRIL HYDROCHLORIDE
 BENDROFLUMETHIAZIDE
 BENORILATE
 BENSERAZIDE HYDROCHLORIDE
 BENZALKONIUM CHLORIDE
 BENZETHONIUM CHLORIDE
 BENZOCAINE
 BENZOIC ACID
 BENZONATATE
 BENZOYL PEROXIDE
 BENZTHIAZIDE
 BENZYL ALCOHOL
 BENZYL BENZOATE
 BEPRIDIL HYDROCHLORIDE
 BERGAPTEN
 beta-CAROTENE
 BETAHISTINE HYDROCHLORIDE
 BETAINES HYDROCHLORIDE
 BETAMETHASONE
 BETAMETHASONE 17,21-DIPROPIONATE
 BETAMETHASONE VALERATE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

BETAMIPRON
 beta-NAPHTHOL
 BETAZOLE HYDROCHLORIDE
 BETHANECHOL CHLORIDE
 BEZAFIBRATE
 BICALUTAMIDE
 BIOTIN
 BISACODYL
 BISOCTRIZOLE
 BISORCIC
 BITHIONATE SODIUM
 BLEOMYCIN (bleomycin B2 shown)
 BRETILIUM TOSYLATE
 BRINZOLAMIDE
 BROMHEXINE HYDROCHLORIDE
 BROMOCRIPTINE MESYLATE
 BROMPHENIRAMINE MALEATE
 BROXYQUINOLINE
 BUDESONIDE
 BUMETANIDE
 BUPIVACAINE HYDROCHLORIDE
 BUPROPION
 BUSULFAN
 BUTACAINE
 BUTAMBEN
 BUTOCONAZOLE
 CAFFEINE
 CAMPHOR (1R)
 CANDESARTAN
 CANDESARTAN CILEXTIL
 CANDICIDIN
 CANRENOIC ACID, POTASSIUM SALT
 CANRENONE
 CAPECITABINE
 CAPREOMYCIN SULFATE
 CAPSAICIN
 CAPTAMINE
 CAPTOPRIL
 CARBACHOL
 CARBENICILLIN DISODIUM
 CARBENOXOLONE SODIUM
 CARBETAPENTANE CITRATE
 CARBIDOPA
 CARBINOXAMINE MALEATE
 CARBOPLATIN
 CARISOPRODOL
 CARMUSTINE
 CARNITINE (dl) HYDROCHLORIDE
 CARPROFEN
 CARVEDILOL
 CEFACTOR
 CEFADROXIL
 CEFAMANDOLE NAFATE
 CEFAMANDOLE SODIUM
 CEFAZOLIN SODIUM
 CEFDINIR
 CEFEPIME HYDROCHLORIDE
 CEFMENOXIME HYDROCHLORIDE
 CEFMETAZOLE SODIUM
 CEFOPERAZONE
 CEFORANIDE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

CEFOTAXIME SODIUM
 CEFOTETAN
 CEFOXITIN SODIUM
 CEFPIRAMIDE
 CEFsulODIN SODIUM
 CEFTIBUTEN
 CEFTIOFUR HYDROCHLORIDE
 CEFTRIAXONE SODIUM TRIHYDRATE
 CEFUROXIME AXETIL
 CEFUROXIME SODIUM
 CELECOXIB
 CEPHALEXIN
 CEPHALOTHIN SODIUM
 CEPHAPIRIN SODIUM
 CEPHRADINE
 CETYLPYRIDINIUM CHLORIDE
 CHENODIOL
 CHLORAMBUCIL
 CHLORAMPHENICOL
 CHLORAMPHENICOL HEMISUCCINATE
 CHLORAMPHENICOL PALMITATE
 CHLORCYCLIZINE HYDROCHLORIDE
 CHLORHEXIDINE
 CHLOROCHRESOL
 CHLOROQUANIDE HYDROCHLORIDE
 CHLOROQUINE DIPHOSPHATE
 CHLOROTHIAZIDE
 CHLOROXYNE
 CHLOROXYLENOL
 CHLORPHENIRAMINE (S) MALEATE
 CHLORPROMAZINE
 CHLORPROPAMIDE
 CHLORPROTHIXENE HYDROCHLORIDE
 CHLORTETRACYCLINE HYDROCHLORIDE
 CHLORTHALIDONE
 CHLORZOXAZONE
 CHOLECALCIFEROL
 CHOLESTEROL
 CHOLINE CHLORIDE
 CICLOPIROX OLAMINE
 CILOSTAZOL
 CIMETIDINE
 CINCOPHEN
 CINNARAZINE
 CINOXACIN
 CINTRIAMIDE
 CIPROFIBRATE
 CIPROFLOXACIN
 CISPLATIN
 CITALOPRAM HYDROBROMIDE
 CITICOLINE
 CLARITHROMYCIN
 CLAVULANATE LITHIUM
 CLEMASTINE
 CLIDINIUM BROMIDE
 CLINAFOXACIN HYDROCHLORIDE
 CLINDAMYCIN HYDROCHLORIDE
 CLIOQUINOL
 CLOBETASOL PROPIONATE
 CLOFARABINE
 CLOFIBRATE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

CLOMIPHENE CITRATE
 CLONIDINE HYDROCHLORIDE
 CLOPIDOGREL SULFATE
 CLORSULON
 CLOSANTEL
 CLOTRIMAZOLE
 CLOXACILLIN SODIUM
 CLOXYQUIN
 CLOZAPINE
 COENZYME B12
 COLCHICINE
 COLFORSIN
 COLISTIMETHATE SODIUM
 CORTISONE ACETATE
 COTININE
 CRESOL
 CROMOLYN SODIUM
 CRYOFLURANE
 CYCLAMIC ACID
 CYCLIZINE
 CYCLOBENZAPRINE HYDROCHLORIDE
 CYCLOHEXIMIDE
 CYCLOPENTOLATE HYDROCHLORIDE
 CYCLOPHOSPHAMIDE HYDRATE
 CYCLOSERINE (D)
 CYCLOSPORINE
 CYCLOTHIAZIDE
 CYPERMETHRIN
 CYPROTERONE ACETATE
 CYSTEAMINE HYDROCHLORIDE
 CYTARABINE
 DACARBAZINE
 DACTINOMYCIN
 DANAZOL
 DANTHRON
 DANTROLENE SODIUM
 DAPSONE
 DAPTOMYCIN
 DASATINIB
 DAUNORUBICIN
 DECIMEMIDE
 DEFEROXAMINE MESYLATE
 DEFLAZACORT
 DEHYDROACETIC ACID
 DEHYDROCHOLIC ACID
 DEMECLOCYCLINE HYDROCHLORIDE
 DERACOXIB
 DESIPRAMINE HYDROCHLORIDE
 DESOXYCORTICOSTERONE ACETATE
 DESVENLAFAXINE SUCCINATE
 DEXAMETHASONE
 DEXAMETHASONE ACETATE
 DEXAMETHASONE SODIUM PHOSPHATE
 DEXIBUPROFEN
 DEXLANSOPRAZOLE
 DEXPROPRANOLOL HYDROCHLORIDE
 DEXTROMETHORPHAN HYDROBROMIDE
 DIAVERIDINE
 DIBENZOTHIOPHENE
 DIBUCAINE HYDROCHLORIDE
 DICHLOROPHENE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

DICHLORVOS
 DICLAZURIL
 DICLOFENAC SODIUM
 DICLOXACILLIN SODIUM
 DICUMAROL
 DICYCLOMINE HYDROCHLORIDE
 DIENESTROL
 DIETHYLCARBAMAZINE CITRATE
 DIETHYLSTILBESTROL
 DIFLOXACIN HYDROCHLORIDE
 DIFLUNISAL
 DIGITOXIN
 DIGOXIN
 DIHYDROERGOTAMINE MESYLATE
 DIHYDROSTREPTOMYCIN SULFATE
 DILAZEP DIHYDROCHLORIDE
 DIMENHYDRINATE
 DIMERCAPROL
 DIMETHADIONE
 DIOXYBENZONE
 DIPHENHYDRAMINE HYDROCHLORIDE
 DIPHENYLPYRALINE HYDROCHLORIDE
 DIPYRIDAMOLE
 DIPYRONE
 DIRITHROMYCIN
 DISOPYRAMIDE PHOSPHATE
 DISULFIRAM
 DOBUTAMINE HYDROCHLORIDE
 DOCETAXEL
 DONEPEZIL HYDROCHLORIDE
 DOPAMINE HYDROCHLORIDE
 DOXEPIN HYDROCHLORIDE
 DOXOFYLLINE
 DOXYCYCLINE HYDROCHLORIDE
 DOXYLAMINE SUCCINATE
 DROFENINE HYDROCHLORIDE
 DROPERIDOL
 DROSPIRENONE
 DYCLONINE HYDROCHLORIDE
 DYPHYLLINE
 ECAMSULE TRIETHANOLAMINE
 ECONAZOLE NITRATE
 EDETATE DISODIUM
 EDITOL
 EDOXUDINE
 EMETINE
 ENALAPRIL MALEATE
 ENALAPRILAT
 ENOXACIN
 ENROFLOXACIN
 ENTACAPONE
 EPHEDRINE (1R,2S) HYDROCHLORIDE
 EPINEPHRINE BITARTRATE
 EPRINOMECTIN
 ERGOCALCIFEROL
 ERGONOVINE MALEATE
 ERYTHROMYCIN
 ERYTHROMYCIN ESTOLATE
 ERYTHROMYCIN ETHYLSUCCINATE
 ESCITALOPRAM OXALATE
 ESOMEPRAZOLE POTASSIUM

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

ESTRADIOL
 ESTRADIOL BENZOATE
 ESTRADIOL CYPIONATE
 ESTRADIOL DIPROPIONATE
 ESTRADIOL VALERATE
 ESTRAMUSTINE
 ESTRIOL
 ESTRONE
 ESTROPIPATE
 ETHACRYNIC ACID
 ETHAMBUTOL HYDROCHLORIDE
 ETHAVERINE HYDROCHLORIDE
 ETHINYL ESTRADIOL
 ETHIONAMIDE
 ETHISTERONE
 ETHOPROPazine HYDROCHLORIDE
 ETHYL PARABEN
 ETODOLAC
 ETOPOSIDE
 EUCALYPTOL
 EUCATROPINE HYDROCHLORIDE
 EUGENOL
 EVANS BLUE
 EXEMESTANE
 EZETIMIBE
 FAMCICLOVIR
 FAMOTIDINE
 FAMPRIDINE
 FASUDIL HYDROCHLORIDE
 FEBUXOSTAT
 FENBENDAZOLE
 FENBUFEN
 FENDILINE HYDROCHLORIDE
 FENOFIBRATE
 FENOPROFEN
 FENOTEROL HYDROBROMIDE
 FENSPIRIDE HYDROCHLORIDE
 FEXOFENADINE HYDROCHLORIDE
 FIPEXIDE HYDROCHLORIDE
 FIROCOXIB
 FLOXURIDINE
 FLUCONAZOLE
 FLUDROCORTISONE ACETATE
 FLUFENAMIC ACID
 FLUIDAROL
 FLUMEQUINE
 FLUMETHASONE
 FLUMETHAZONE PIVALATE
 FLUNARIZINE HYDROCHLORIDE
 FLUNISOLIDE
 FLUNIXIN MEGLUMINE
 FLUOCINOLONE ACETONIDE
 FLUOCINONIDE
 FLUOROMETHOLONE
 FLUOROURACIL
 FLUOXETINE
 FLUPHENAZINE HYDROCHLORIDE
 FLURANDRENOLIDE
 FLURBIPROFEN
 FLUROFAMIDE
 FLUTAMIDE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

FLUVASTATIN
 FOLIC ACID
 FOSCARNET SODIUM
 FOSFOMYCIN CALCIUM
 FTAXILIDE
 FULVESTRANT
 FURAZOLIDONE
 FUROSEMIDE
 FUSIDIC ACID
 GABOXADOL HYDROCHLORIDE
 GADOTERIDOL
 GALANTHAMINE
 GALLAMINE TRIETHIODIDE
 GANCICLOVIR
 GATIFLOXACIN
 GEFITINIB
 GEMFIBROZIL
 GENTAMICIN SULFATE
 GENTIAN VIOLET
 GLIMEPIRIDE
 GLUCONOLACTONE
 GLUCOSAMINE HYDROCHLORIDE
 GLUTAMINE (D)
 GRAMICIDIN
 GRANISETRON HYDROCHLORIDE
 GRISEOFULVIN
 GUAIFENESIN
 GUANABENZ ACETATE
 GUANETHIDINE SULFATE
 HALAZONE
 HALCINONIDE
 HALOPERIDOL
 HEPTAMINOL HYDROCHLORIDE
 HETACILLIN POTASSIUM
 HEXACHLOROPHENE
 HEXYLRESORCINOL
 HISTAMINE DIHYDROCHLORIDE
 HOMATROPINE BROMIDE
 HOMATROPINE METHYLBROMIDE
 HOMOSALATE
 HYCANTHONE
 HYDRALAZINE HYDROCHLORIDE
 HYDRASTINE (1R, 9S)
 HYDROCHLOROTHIAZIDE
 HYDROCORTISONE
 HYDROCORTISONE ACETATE
 HYDROCORTISONE BUTYRATE
 HYDROCORTISONE HEMISUCCINATE
 HYDROCORTISONE PHOSPHATE TRIETHYLAMINE
 HYDROFLUMETHIAZIDE
 HYDROQUINONE
 HYDROXYAMPHETAMINE HYDROBROMIDE
 HYDROXYCHLOROQUINE SULFATE
 HYDROXYPROGESTERONE CAPROATE
 HYDROXYTOLUIC ACID
 HYDROXYUREA
 HYDROXYZINE PAMOATE
 HYOSCYAMINE
 IBANDRONATE SODIUM
 IBUPROFEN
 IDOXURDINE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

IDOXURIDINE
 IMIPRAMINE HYDROCHLORIDE
 IMIQUIMOD
 INAMRINONE
 INDAPAMIDE
 INDOMETHACIN
 INDOPROFEN
 INOSITOL
 IODIPAMIDE
 IODIXANOL
 IODOQUINOL
 IOHEXOL
 IOPANIC ACID
 IOTHALAMIC ACID
 IOVERSOL
 IOXILAN
 IPRATROPIUM BROMIDE
 IRBESARTAN
 ISONIAZID
 ISOPROPAMIDE IODIDE
 ISOPROTERENOL HYDROCHLORIDE
 ISOSORBIDE DINITRATE
 ISOSORBIDE MONONITRATE
 ISOTRETINON
 ISOXICAM
 ISOXSUPRINE HYDROCHLORIDE
 ITOPRIDE HYDROCHLORIDE
 IVERMECTIN
 KANAMYCIN A SULFATE
 KETOCONAZOLE
 KETOPROFEN
 KETOROLAC TROMETHAMINE
 KETOTIFEN FUMARATE
 LABETALOL HYDROCHLORIDE
 LACTULOSE
 LAMIVUDINE
 LANATOSIDE C
 LANSOPRAZOLE
 LEFLUNOMIDE
 LETROZOLE
 LEUCOVORIN CALCIUM
 LEVAMISOLE HYDROCHLORIDE
 LEVOCETIRIZINE DIHYDROCHLORIDE
 LEVOFLOXACIN
 LEVOMENTHOL
 LEVONORDEFIN
 LEVONORGESTREL
 LEVOSIMENDAN
 LEVOTHYROXINE
 LIDOCAINE HYDROCHLORIDE
 LINCOMYCIN HYDROCHLORIDE
 LINDANE
 LINEZOLID
 LIOTHYRONINE
 LIOTHYRONINE (L- isomer) SODIUM
 LISINAPRIL
 LITHIUM CITRATE
 LOBELINE HYDROCHLORIDE
 LOFEXIDINE HYDROCHLORIDE
 LOMEFLOXACIN HYDROCHLORIDE
 LOMERIZINE HYDROCHLORIDE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

LOMUSTINE
 LORATADINE
 LORNOXICAM
 LOSARTAN
 LOVASTATIN
 LUMIRACOXIB
 MAFENIDE HYDROCHLORIDE
 MALATHION
 MANGAFODIPIR TRISODIUM
 MANIDIPINE HYDROCHLORIDE
 MANNITOL
 MAPROTILINE HYDROCHLORIDE
 MEBENDAZOLE
 MEBEVERINE HYDROCHLORIDE
 MEBHYDROLIN NAPHTHALENESULFONATE
 MECAMYLAMINE HYDROCHLORIDE
 MECHLORETHAMINE
 MECLIZINE HYDROCHLORIDE
 MECLOCYCLINE SULFOSALICYLATE
 MECLOFENAMATE SODIUM
 MECLOFENOXATE HYDROCHLORIDE
 MEDROXYPROGESTERONE ACETATE
 MEDRYSONE
 MEFENAMIC ACID
 MEFEXAMIDE
 MEFLOQUINE
 MEGESTROL ACETATE
 MEGLUMINE
 MELOXICAM SODIUM
 MELPERONE HYDROCHLORIDE
 MELPHALAN
 MEMANTINE HYDROCHLORIDE
 MENADIONE
 MEPARTRICIN
 MEPENZOLATE BROMIDE
 MEPHENESIN
 MEPHENTERMINE SULFATE
 MEPIVACAINE HYDROCHLORIDE
 MEQUINOL
 MERBROMIN
 MERCAPTOPYRINE
 MEROPENEM
 MESNA
 MESO-ERYTHRITOL
 MESTRANOL
 METAPROTERENOL
 METARAMINOL BITARTRATE
 METAXALONE
 METHACHOLINE CHLORIDE
 METHACYCLINE HYDROCHLORIDE
 METHAPYRILENE HYDROCHLORIDE
 METHAZOLAMIDE
 METHENAMINE
 METHICILLIN SODIUM
 METHIMAZOLE
 METHOCARBAMOL
 METHOTREXATE(+/-)
 METHOXAMINE HYDROCHLORIDE
 METHOXSALEN
 METHSCOPOLAMINE BROMIDE
 METHYLCLOTHIAZIDE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

METHYLBENZETHONIUM CHLORIDE
 METHYLDOPA
 METHYLERGONOVINE MALEATE
 METHYLPREDNISOLONE
 METHYLPREDNISOLONE SODIUM SUCCINATE
 METHYLTHIOURACIL
 METOCLOPRAMIDE HYDROCHLORIDE
 METOPROLOL TARTRATE
 METRONIDAZOLE
 MEXILETINE HYDROCHLORIDE
 MICONAZOLE NITRATE
 MIDODRINE HYDROCHLORIDE
 MIGLITOL
 MILNACIPRAN HYDROCHLORIDE
 MINAPRINE HYDROCHLORIDE
 MINOCYCLINE HYDROCHLORIDE
 MINOXIDIL
 MITOMYCIN C
 MITOTANE
 MITOXANTRONE HYDROCHLORIDE
 MOLSIDOMINE
 MONENSIN SODIUM (monensin A is shown)
 MONOBENZONE
 MORANTEL CITRATE
 MOXALACTAM DISODIUM
 MOXIFLOXACIN HYDROCHLORIDE
 MYCOPHENOLATE MOFETIL
 MYCOPHENOLIC ACID
 NABUMETONE
 NADIDE
 NADOLOL
 NAFACILLIN SODIUM
 NAFRONYL OXALATE
 NALBUPHINE HYDROCHLORIDE
 NALIDIXIC ACID
 NALOXONE HYDROCHLORIDE
 NALTREXONE HYDROCHLORIDE
 NAPHAZOLINE HYDROCHLORIDE
 NAPROXEN(+)
 NAPROXOL
 NATEGLINIDE
 NEFAZODONE HYDROCHLORIDE
 NEFOPAM
 NELARABIN
 NEOMYCIN SULFATE
 NEOSTIGMINE BROMIDE
 NEVIRAPINE
 NIACIN
 NICARDIPINE HYDROCHLORIDE
 NICERGOLINE
 NICLOSAMIDE
 NICOTINYL ALCOHOL TARTRATE
 NIFEDIPINE
 NIFURSOL
 NILUTAMIDE
 NIMODIPINE
 NITAZOXANIDE
 NITRENDIPINE
 NITROFURANTOIN
 NITROFURAZONE
 NITROMIDE
 NOCODAZOLE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

NOMIFENSINE MALEATE
 NOREPINEPHRINE
 NORETHINDRONE
 NORETHINDRONE ACETATE
 NORETHYNODREL
 NORFLOXACIN
 NORGESTREL
 NORTRIPTYLINE
 NOSCAPINE HYDROCHLORIDE
 NOVOBIOCIN SODIUM
 NYLIDRIN HYDROCHLORIDE
 NYSTATIN
 OCTOPAMINE HYDROCHLORIDE
 OFLOXACIN
 OLMESARTAN
 OLMESARTAN MEDOXOMIL
 OLSALAZINE SODIUM
 OLSELTAMIVIR PHOSPHATE
 OMEGA-3-ACID ESTERS (EPA shown)
 ONDANSETRON
 ORLISTAT
 ORPHENADRINE CITRATE
 OUABAIN
 OXACILLIN SODIUM
 OXALIPLATIN
 OXCARBAZEPINE
 OXETHAZAINE
 OXIBENDAZOLE
 OXIDOPAMINE HYDROCHLORIDE
 OXOLINIC ACID
 OXYBENZONE
 OXYMETAZOLINE HYDROCHLORIDE
 OXYPHENBUTAZONE
 OXYPHENCYCLIMINE HYDROCHLORIDE
 OXYQUINOLINE HEMISULFATE
 OXYTETRACYCLINE
 PACLITAXEL
 PALIPERIDONE
 PAPAVERINE HYDROCHLORIDE
 PARACHLOROPHENOL
 PARAROSANILINE PAMOATE
 PARGYLINE HYDROCHLORIDE
 PAROMOMYCIN SULFATE
 PAROXETINE HYDROCHLORIDE
 PEMETREXED
 PENCICLOVIR
 PENICILLAMINE
 PENICILLIN G POTASSIUM
 PENICILLIN V POTASSIUM
 PENTOLINIUM TARTRATE
 PENTOXIFYLLINE
 PERGOLIDE MESYLATE
 PERHEXILINE MALEATE
 PERICIAZINE
 PERINDOPRIL ERBUMINE
 PERPHENAZINE
 PHENACEMIDE
 PHENAZOPYRIDINE HYDROCHLORIDE
 PHENELZINE SULFATE
 PHENINDIONE
 PHENIRAMINE MALEATE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

PHENOLPHTHALEIN
 PHENTOLAMINE HYDROCHLORIDE
 PHENYL AMINOSALICYLATE
 PHENYLBUTAZONE
 PHENYLEPHRINE HYDROCHLORIDE
 PHENYLMERCURIC ACETATE
 PHENYLPROPANOLAMINE HYDROCHLORIDE
 PHENYTOIN SODIUM
 PHTHALYLSULFATHIAZOLE
 PHYSOSTIGMINE SALICYLATE
 PILOCARPINE NITRATE
 PIMOZIDE
 PINDOLOL
 PIOGLITAZONE HYDROCHLORIDE
 PIPERACETAZINE
 PIPERACILLIN SODIUM
 PIPERAZINE
 PIPERIDOLATE HYDROCHLORIDE
 PIPERINE
 PIPOBROMAN
 PIRACETAM
 PIRENPERONE
 PIRENZEPINE HYDROCHLORIDE
 PIROCTONE OLAMINE
 PIROXICAM
 PITAVASTATIN CALCIUM
 PIZOTYLIN MALATE
 POLYMYXIN B SULFATE
 POTASSIUM p-AMINOBENZOATE
 PRAMIPEXOLE DIHYDROCHLORIDE
 PRAMOXINE HYDROCHLORIDE
 PRASUGREL
 PRAZQUANTEL
 PRAZOSIN HYDROCHLORIDE
 PREDNICARBATE
 PREDNISOLONE
 PREDNISOLONE ACETATE
 PREDNISON
 PRILOCAINE HYDROCHLORIDE
 PRIMAQUINE DIPHOSPHATE
 PRIMIDONE
 PROADIFEN HYDROCHLORIDE
 PROBENECID
 PROBUCOL
 PROCAINAMIDE HYDROCHLORIDE
 PROCAINE HYDROCHLORIDE
 PROCARBAZINE HYDROCHLORIDE
 PROCHLORPERAZINE EDISYLATE
 PROCYCLIDINE HYDROCHLORIDE
 PROGESTERONE
 PROGLUMIDE
 PROMAZINE HYDROCHLORIDE
 PROMETHAZINE HYDROCHLORIDE
 PRONETALOL HYDROCHLORIDE
 PROPANFENONE HYDROCHLORIDE
 PROPANTHELIN BROMIDE
 PROPIOLACTONE
 PROPOFOL
 PROPYLTHIOURACIL
 PSEUDOEPHEDRINE HYDROCHLORIDE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

PUROMYCIN HYDROCHLORIDE
 PYRANTEL PAMOATE
 PYRAZINAMIDE
 PYRETHRINS
 PYRIDOSTIGMINE BROMIDE
 PYRILAMINE MALEATE
 PYRIMETHAMINE
 PYRITHIONE ZINC
 PYRONARIDINE TETRAPHOSPHATE
 PYRVINIUM PAMOATE
 QUETIAPINE
 QUINACRINE HYDROCHLORIDE
 QUINAPRIL HYDROCHLORIDE
 QUINESTROL
 QUINETHAZONE
 QUINIDINE GLUCONATE
 QUININE SULFATE
 QUIPAZINE MALEATE
 RACEPHEDRINE HYDROCHLORIDE
 RACTOPAMINE HYDROCHLORIDE
 RAMIPRIL
 RAMOPLANIN [A2 shown; 2mm]
 RANITIDINE
 RASAGILINE
 RESERPINE
 RESORCINOL
 RESORCINOL MONOACETATE
 RETAPAMULIN
 RETINOL
 RETINYL PALMITATE
 RIBAVIRIN
 RIFAMPIN
 RITANSERIN
 RITODRINE HYDROCHLORIDE
 RITONAVIR
 RIZATRIPTAN BENZOATE
 ROFECOXIB
 RONIDAZOLE
 ROPINIROLE
 ROSIGLITAZONE
 ROSUVASTATIN CALCIUM
 ROXARSONE
 ROXATIDINE ACETATE HYDROCHLORIDE
 ROXITHROMYCIN
 RUFLOXACIN HYDROCHLORIDE
 SACCHARIN
 SALICIN
 SALICYL ALCOHOL
 SALICYLAMIDE
 SALICYLANILIDE
 SALINOMYCIN, SODIUM
 SALSALATE
 SANGUINARINE SULFATE
 SCOPOLAMINE HYDROBROMIDE
 SELAMECTIN
 SEMUSTINE
 SENNOSIDE A
 SERATRODAST
 SERTRALINE HYDROCHLORIDE
 SEVOFLURANE
 SIBUTRAMINE HYDROCHLORIDE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

SILDENAFIL CITRATE
SIMVASTATIN
SIROLIMUS
SISOMICIN SULFATE
SODIUM DEHYDROCHOLATE
SODIUM NITROPRUSSIDE
SODIUM OXYBATE
SODIUM PHENYLACETATE
SODIUM PHENYLBUTYRATE
SODIUM SALICYLATE
SPARFLOXACIN
SPARTEINE SULFATE
SPECTINOMYCIN HYDROCHLORIDE
SPIPERONE
SPIRAMYCIN
SPIRAPRIL HYDROCHLORIDE
SPIRONOLACTONE
STAVUDINE
STREPTOMYCIN SULFATE
STREPTOZOSIN
SUCCINYLSULFATHIAZOLE
SULBACTAM
SULCONAZOLE NITRATE
SULFABENZAMIDE
SULFACETAMIDE
SULFACHLORPYRIDAZINE
SULFADIAZINE
SULFADIMETHOXINE
SULFAMERAZINE
SULFAMETER
SULFAMETHAZINE
SULFAMETHIZOLE
SULFAMETHOXAZOLE
SULFAMETHOXYPYRIDAZINE
SULFAMONOMETHOXINE
SULFANILATE ZINC
SULFANITRAN
SULFAPYRIDINE
SULFAQUINOXALINE SODIUM
SULFASALAZINE
SULFATHIAZOLE
SULFINPYRAZONE
SULFISOXAZOLE
SULINDAC
SULMAZOLE
SULOCTIDIL
SULPIRIDE
SUPROFEN
SURAMIN
TACROLIMUS
TAMOXIFEN CITRATE
TANDUTINIB
TANNIC ACID
TAZOBACTAM
TEGASEROD MALEATE
TELMISARTAN
TEMEFOS
TEMOZOLAMIDE
TENIPOSIDE
TENOXICAM
TERBUTALINE HEMISULFATE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

TERCONAZOLE
TERFENADINE
TESTOSTERONE
TESTOSTERONE PROPIONATE
TETRACAINE HYDROCHLORIDE
TETRACYCLINE HYDROCHLORIDE
TETRAHYDROZOLINE HYDROCHLORIDE
TETROQUINONE
THALIDOMIDE
THEOBROMINE
THEOPHYLLINE
THIABENDAZOLE
THIAMPHENICOL
THIMEROSAL
THIOGUANINE
THIORIDAZINE HYDROCHLORIDE
THIOSTREPTON
THIOTEPA
THIOTHIXENE
THIRAM
THONZONIUM BROMIDE
THONZYLAMINE HYDROCHLORIDE
TIAPRIDE HYDROCHLORIDE
TIBOLONE
TIGECYCLINE
TILARGININE HYDROCHLORIDE
TILETAMINE HYDROCHLORIDE
TILMICOSIN
TIMOLOL MALEATE
TINIDAZOLE
TOBRAMYCIN
TODRALAZINE HYDROCHLORIDE
TOLAZAMIDE
TOLAZOLINE HYDROCHLORIDE
TOLBUTAMIDE
TOLMETIN SODIUM
TOLNAFTATE
TOLPERISONE HYDROCHLORIDE
TOSYLCHLORAMIDE SODIUM
TRANEXAMIC ACID
TRANLYCYPROMINE SULFATE
TRAZODONE HYDROCHLORIDE
TRETINOIN
TRIACETIN
TRIAMCINOLONE
TRIAMCINOLONE ACETONIDE
TRIAMCINOLONE DIACETATE
TRIAMTERENE
TRICHLORMETHIAZIDE
TRIFLUOPERAZINE HYDROCHLORIDE
TRIFLUPROMAZINE HYDROCHLORIDE
TRIFLURIDINE
TRIHXYPHENIDYL HYDROCHLORIDE
TRILOSTANE
TRIMEPAZINE TARTRATE
TRIMETHOBENZAMIDE HYDROCHLORIDE
TRIMETHOPRIM
TRIMETOZINE
TRIMIPRAMINE MALEATE
TRIOXSALEN
TRIPLENNAMINE CITRATE

(Continued)

Table 1. List of compounds from the PHARMAKON 1600 library used in this screen. (continued)

TRIPROLIDINE HYDROCHLORIDE
TRISODIUM ETHYLENEDIAMINE TETRACETATE
TROLEANDOMYCIN
TROPICAMIDE
TROPISETRON HYDROCHLORIDE
TRYPTOPHAN
TUAMINOHEPTANE SULFATE
TUBOCURARINE CHLORIDE
TYROTHRIN
URACIL
URAPIDIL HYDROCHLORIDE
UREA
URETHANE
URSODIOL
VALDECOSIB
VALGANCICLOVIR HYDROCHLORIDE
VALPROATE SODIUM
VALSARTAN
VANCOMYCIN HYDROCHLORIDE
VENLAFAXINE
VIDARABINE
VINBLASTINE SULFATE
VINORELBINE
VINPOCETINE
VIOMYCIN SULFATE
VORICONAZOLE
VORINOSTAT
WARFARIN
XYLAZINE
XYLOMETAZOLINE HYDROCHLORIDE
YOHIMBINE HYDROCHLORIDE
ZALCITABINE
ZAPRINAST
ZIDOVUDINE [AZT]
ZIPRASIDONE MESYLATE
ZOMEPIRAC SODIUM
ZOPICLONE

were obtained from mutants bathed in embryo media, as described above; a second locomotion plot was then obtained following a solution change to a test compound and an equilibration period of 15–30 min. Criteria for a positive hit designation were as follows: (1) a decrease in mean velocity of $\geq 44\%$ (e.g., a value based on the trial-to-trial variability measured in control tracking studies; Fig. 1c); and (2) a reduction to Stage 0 or Stage I seizure behavior in the locomotion plot for at least 50% of the test fish. Each test compound classified as a “positive hit” in the locomotion assay was confirmed, under direct visualization on a stereomicroscope, as the fish being alive based on movement in response to external stimulation and a visible heartbeat following a 60 min drug exposure. Toxicity (or mortality) was defined as no visible heartbeat or movement in response to external stimulation in at least 50% of the test fish. Hyperexcitability was defined as a compound causing a $\geq 44\%$ increase in swim velocity and/or Stage III seizure activity in at least 50% of the test fish. Hits identified in the primary locomotion screen were selected from the PHARMAKON 1600 library and re-screened using the method described above. Select com-

pound stocks that were successful in two primary locomotion assays, and were not classified as toxic in two independent clutches of zebrafish, were then purchased separately from Sigma-Aldrich for further testing. Drug concentrations between 0.5 and 1 mM were used for electrophysiology assays to account for more limited diffusion in agar-embedded larvae.

Data analysis

Data are presented as the mean and SEM, unless stated otherwise. Pairwise statistical significance was determined with a Student’s two-tailed unpaired *t* test, ANOVA, or Mann–Whitney rank sum test, as appropriate, unless stated otherwise. Results were considered significant at $p < 0.05$, unless otherwise indicated.

Results

A first-stage behavioral screen for antiepileptic activity

Locomotion tracking is a reliable and rapid strategy with which to monitor behavioral seizures in freely swimming larval zebrafish (Baraban et al., 2005, 2013; Winter et al., 2008). In these locomotion plots, high-velocity movements of ≥ 20 mm/s correspond to paroxysmal whole-body convulsions, referred to as Stage III, and are consistently observed in unprovoked *scn1Lab* mutant larvae but not in age-matched wild-type siblings. Using automated locomotion tracking, we performed a phenotype-based screen to identify compounds that significantly reduce mutant swim behavior to levels associated with Stage 0 or Stage I (e.g., activity equivalent to that seen in normal untreated WT zebrafish). In a 96-well format, we tracked mutant swim activity at baseline, and then again after addition of a test compound (100 μM); each compound was tested on six individual mutant larvae (Fig. 1a), and larvae were sorted based on pigmentation differences (Fig. 1b). Mutant swim activity between two consecutive recording epochs in embryo media is tracked on every plate as an internal control. A box plot showing the change in swim velocity in untreated mutants is shown in Figure 1c ($n = 112$) and defined as the control. Based on an SD of 21.8 for these data, we set the detection threshold as any compound that inhibits movement (measured as a change in mean velocity) by > 2 SDs (or $\geq 44\%$). This approach was previously validated using standard antiepileptic drugs in this model (Baraban et al., 2013). Next, we screened a repurposed library in which all compounds have reached the clinical evaluation stage (PHARMAKON 1600 Collection; <http://www.msdiscovery.com/pharma.html>). Among the 1012 compounds screened (Fig. 1d) only 20 (or 1.97%) were found to significantly inhibit spontaneous seizure behavior in *scn1Lab* mutants. All 20 compounds were subsequently retested in a separate clutch of *scn1Lab* mutants at a concentration of 100 μM (Fig. 2a, trial 2; $N = 6$ fish/compound). A total of 154 compounds were classified as “toxic” (Table 2); 55 compounds were classified as “hyperexcitable” (Table 3). Representative locomotion tracking raw data plots for gemfibrozil, a toxic nonsteroid




Figure 2. Positive hits identified in the locomotion assay. **a**, Heat map showing the results of individual zebrafish trials (1–6) for compounds tested at a concentration of 100 μM in the locomotion-tracking assay. Raw data values for individual fish are shown within the color-coded boxes for one sample trial. Mean velocity data are shown at right for “trial 1” and “trial 2”; six fish per trial. Note: only drugs highlighted in bold type were classified as positive nontoxic hits in two independent trials and moved on to further testing. **b**, Representative raw locomotion data plots for six individual *scn1Lab* mutant larvae at baseline (top) and following the addition of a compound resulting in fatality (bottom, gemfibrozil) or hyperactivity (bottom, mepivacaine). Movement is color coded, with low-velocity movements shown in yellow, and high velocity movements shown in red.

nuclear receptor ligand, and mepivacaine, a hyperexcitable proconvulsant anesthetic, are shown in Figure 2b.

A second-stage electrophysiology assay for antiepileptic activity

Extracellular recording electrodes are a reliable, reproducible, and sensitive approach to monitor electroencephalographic activity in agar-immobilized larval zebrafish (Baraban et al., 2005; Baraban, 2013). Field electrodes offer high a signal-to-noise ratio and can be placed, using direct visualization in transparent larvae, into specific CNS structures (i.e., telencephalon or optic tectum). Using a local field electrode, we can efficiently monitor the occurrence of electrographic seizure events in the same zebrafish that were previously tested in the locomotion assay. Based on a positive nontoxic result in two independent locomotion assays, four drugs moved

on to electrophysiology testing at concentrations between 500 μM and 1 mM (Fig. 3). Consistent with a “false-positive” classification, spontaneous epileptiform discharge activity was observed for three of these drugs: norfloxacin, theobromine, and cytarabine. Dimethadione, previously shown to inhibit spontaneous epileptiform discharges in thalamocortical slices at concentrations between 1 and 10 mM (Zhang et al., 1996), suppressed burst discharge activity in *scn1Lab* mutant larvae (Fig. 3a,b). To identify whether any of these four compounds exert non-specific effects on behavior, they were also tested on freely swimming WT zebrafish larvae (5 dpf) at a concentration of 500 μM . Comparing the total distance moved during a 10 min recording epoch before, and after, the application of a test compound failed to reveal any significant changes in locomotor activity (Fig. 3c).

Table 2: List of compounds exhibiting toxicity.

ABACAVIR SULFATE
 ACIPIMOX
 ADENOSINE PHOSPHATE
 ALAPROCLATE
 AMLEXANOX
 AMOROLFINE HYDROCHLORIDE
 ANTAZOLINE PHOSPHATE
 ARTEMETHER
 ASCORBIC ACID
 ATORVASTATIN CALCIUM
 AUROTHIOGLUCOSE
 AZELAIC ACID
 BENORILATE
 BENZONATATE
 BETAINE HYDROCHLORIDE
 BETAMIPRON
 BROMHEXINE HYDROCHLORIDE
 BUDESONIDE
 BUPIVACAINE HYDROCHLORIDE
 BUSULFAN
 BUTOCONAZOLE
 CAPSAICIN
 CARPROFEN
 CEFORANIDE
 CEFOTAXIME SODIUM
 CEFOXITIN SODIUM
 CEPHALEXIN
 CHLORAMBUCIL
 CHLORAMPHENICOL HEMISUCCINATE
 CHLOROGUANIDE HYDROCHLORIDE
 CHLORPHENIRAMINE (S) MALEATE
 CINCHOPHEN
 CINNARAZINE
 CINTRIAMIDE
 CIPROFLOXACIN
 CLIDINIUM BROMIDE
 CLOZAPINE
 COLISTIMETHATE SODIUM
 CRYOFLURANE
 CYCLOPHOSPHAMIDE HYDRATE
 CYCLOTHIAZIDE
 CYPERMETHRIN
 DAUNORUBICIN
 DECIMEMIDE
 DEXTROMETHORPHAN HYDROBROMIDE
 DICHLOROPHENE
 DIETHYLCARBAMAZINE CITRATE
 DIOXYBENZONE
 DIRITHROMYCIN
 DISOPYRAMIDE PHOSPHATE
 DISULFIRAM
 ECONAZOLE NITRATE
 EDETATE DISODIUM
 EMETINE
 ENALAPRILAT
 ERYTHROMYCIN
 ETHINYL ESTRADIOL
 ETHIONAMIDE
 ETHOPROPAZINE HYDROCHLORIDE
 ETHYL PARABEN
 EUGENOL
 FIPEXIDE HYDROCHLORIDE

(Continued)

Table 2: List of compounds exhibiting toxicity.

FLUMETHASONE
 FLUNISOLIDE
 FLUVASTATIN
 GEMFIBROZIL
 GENTAMICIN SULFATE
 GLUCONOLACTONE
 HALAZONE
 HALCINONIDE
 HETACILLIN POTASSIUM
 HEXACHLOROPHENE
 HOMATROPINE METHYLBROMIDE
 HYDRASTINE (1R, 9S)
 HYDROXYAMPHETAMINE HYDROBROMIDE
 HYDROXYCHLOROQUINE SULFATE
 IODIXANOL
 IOHEXOL
 IRBESARTAN
 LEVOSIMENDAN
 LISINOPRIL
 LOMERIZINE HYDROCHLORIDE
 MANGAFODIPIR TRISODIUM
 MECLOFENOXATE HYDROCHLORIDE
 MESTRANOL
 METHACHOLINE CHLORIDE
 METHYLERGONOVINE MALEATE
 METRONIDAZOLE
 MIGLITOL
 MONENSIN SODIUM (monensin A is shown)
 MONOBENZONE
 MOXALACTAM DISODIUM
 NADOLOL
 NALBUPHINE HYDROCHLORIDE
 NALTREXONE HYDROCHLORIDE
 NAPHAZOLINE HYDROCHLORIDE
 NAPROXEN(+)
 NEOMYCIN SULFATE
 NIFEDIPINE
 NITAZOXANIDE
 NITROMIDE
 NORETHINDRONE
 OLMESARTAN MEDOXOMIL
 OXYMETAZOLINE HYDROCHLORIDE
 PARACHLOROPHENOL
 PAROMOMYCIN SULFATE
 PERHEXILINE MALEATE
 PHENTOLAMINE HYDROCHLORIDE
 PHENYLBUTAZONE
 PHENYLMERCURIC ACETATE
 PHYSOSTIGMINE SALICYLATE
 PIMOZIDE
 PIPERACILLIN SODIUM
 PIPERAZINE
 PIRACETAM
 PIRENZEPINE HYDROCHLORIDE
 PIROCTONE OLAMINE
 PITAVASTATIN CALCIUM
 PRIMAQUINE DIPHOSPHATE
 PROBENECID
 PROCARBAZINE HYDROCHLORIDE
 PROGLUMIDE
 PROMETHAZINE HYDROCHLORIDE
 PUROMYCIN HYDROCHLORIDE

(Continued)

Table 2: List of compounds exhibiting toxicity.

QUININE SULFATE
RETINYL PALMITATE
RIFAMPIN
RITONAVIR
ROFECOXIB
RUFLOXACIN HYDROCHLORIDE
SACCHARIN
SALICIN
SENNOSIDE A
STAVUDINE
STREPTOMYCIN SULFATE
SULFADIAZINE
SULINDAC
SULOCTIDIL
TANNIC ACID
TELMISARTAN
TENOICAM
THEOPHYLLINE
TILETAMINE HYDROCHLORIDE
TILMICOSIN
TIMOLOL MALEATE
TOLBUTAMIDE
TOLNAFTATE
TRAZODONE HYDROCHLORIDE
TRETINOIN
TRIFLUPROMAZINE HYDROCHLORIDE
TROPISETRON HYDROCHLORIDE
VALDECOXIB
VORINOSTAT
ZALCITABINE

Assessment of huperzine A and fenfluramine for antiepileptic activity

Next, we tested two additional compounds that were not in our drug library, but have recently been described as potential antiepileptic treatments for DS. Huperzine A, a small-molecule alkaloid isolated from Chinese club moss with NMDA-type receptor blocking and anticholinesterase activity, has purported antiepileptic actions against NMDA- or soman-induced seizures (Tonduli et al., 2001; Coleman et al., 2008). In the locomotion assay, huperzine A failed to significantly alter *scn1Lab* seizure behavior at any concentration tested (Fig. 4a,b). In contrast, huperzine A was effective at 1 mM in the acute pentylenetetrazole (PTZ) assay (Fig. 4b). Fenfluramine is an amphetamine-like compound that has been reported to successfully reduce seizure occurrence in children with DS as a low-dose add-on therapy (Ceulemans et al., 2012). In the locomotion assay, fenfluramine significantly reduced mutant mean swim velocity at concentrations between 100 and 500 μM (Fig. 4c,d); 1 mM fenfluramine was toxic in the *scn1Lab* and PTZ assays (Fig. 4d). The fenfluramine-treated *scn1Lab* mutant exhibited a suppression of spontaneous electrographic seizure discharge to levels similar to controls at 500 μM , but only a partial reduction in electrographic activity at 250 μM (Fig. 4e).

Discussion

Zebrafish and humans share extensive genomic similarity. With regard to disease, 84% of genes known to be associated with disease states in humans have a zebrafish

Table 3: List of compounds exhibiting hyperexcitable or pro-convulsant activity.

ADENOSINE PHOSPHATE
ALBUTEROL (+/-)
ALEXIDINE HYDROCHLORIDE
AMANTADINE HYDROCHLORIDE
AMINOHIPPURIC ACID
AMINOLEVULINIC ACID HYDROCHLORIDE
AUROTHIOGLUCOSE
AZACITIDINE
BENZOYL PEROXIDE
BETAZOLE HYDROCHLORIDE
BROMHEXINE HYDROCHLORIDE
BUSULFAN
CEFSULODIN SODIUM
CEFUROXIME AXETIL
CHLOROQUANIDE HYDROCHLORIDE
CYSTEAMINE HYDROCHLORIDE
ECAMSULE TRIETHANOLAMINE
ECONAZOLE NITRATE
EDOXUDINE
ENROFLOXACIN
ESTRADIOL CYPIONATE
ETHINYL ESTRADIOL
ETHOPROPAZINE HYDROCHLORIDE
ETOPOSIDE
FASUDIL HYDROCHLORIDE
FEBUXOSTAT
FLUMETHASONE
FLUROMETHOLONE
FURAZOLIDONE
GANCICLOVIR
GLUCONOLACTONE
GRANISETRON HYDROCHLORIDE
HALAZONE
HEXACHLOROPHENE
IODIPAMIDE
LABELTALOL HYDROCHLORIDE
MEPIVACAINE HYDROCHLORIDE
MITOXANTRONE HYDROCHLORIDE
MORANTEL CITRATE
NOCODAZOLE
OFLOXACIN
PENTOLINIUM TARTRATE
PERINDOPRIL ERBUMINE
PIOGLITAZONE HYDROCHLORIDE
PRAMIPEXOLE DIHYDROCHLORIDE
PROGLUMIDE
RIFAMPIN
SERATRODAST
SERTRALINE HYDROCHLORIDE
SIBUTRAMINE HYDROCHLORIDE
SUCCINYL-SULFATHIAZOLE
TACROLIMUS
TETROQUINONE
TIMOLOL MALEATE
URACIL

homolog (Howe et al., 2013). This genetic similarity and the characteristic of zebrafish larvae to exhibit quantifiable seizure behaviors or electrographic seizure discharge that is fundamentally similar to that recorded in humans (Jirsa et al., 2014) make this an ideal system for drug discovery. Behavioral assays customized for auto-




Figure 3. Electrophysiology assay to identify drugs that rescue the *scn1Lab* mutant epilepsy phenotype. **a**, Representative field electrode recording epochs (5 min in duration) are shown for the “positive” compounds identified in the locomotion assay. All recordings were obtained with an electrode placed in the forebrain of agar-immobilized *scn1Lab* larvae that was previously tested in the locomotion assay. A suppression of epileptiform electrographic discharge activity was noted in mutants exposed to dimethadione. **b**, Bar plot showing the mean number of epileptiform events in a 10 min recording epoch for *scn1Lab* larvae exposed to cytarabine ($N = 6$), dimethadione ($N = 6$), theobromine ($N = 6$), and norfloxacin ($N = 6$). The mean \pm SEM is shown. The fish shown were tested in the locomotion assay first. **c**, Bar plot showing the total distance traveled before (black bars) and after (white bars) exposure to a test compound; 10 min recording epoch and six fish per drug. The mean \pm SEM is shown.

mated evaluation of locomotion (Winter et al., 2008; Creton, 2009; Baxendale et al., 2012; Baraban et al., 2013; Raftery et al., 2014) make moderate-to-high-throughput phenotype-based drug screening in zebrafish possible. Using this approach and a zebrafish *scn1* mutant (Baraban et al., 2013), we successfully identified antiepileptic compounds. Here we report results from screening ~1000 compounds from a repurposed drug library and present data that will be periodically updated on-line using this open-access publishing mechanism.

As a model system, the *scn1Lab* mutant zebrafish has many advantages. First, in contrast to transient and variable knockdown of gene expression using antisense morpholino oligonucleotides (Teng et al., 2010; Finckbeiner et al., 2011; Mahmood et al., 2013), *scn1Lab* mutants carry a stable and heritable amino acid substitution at position 1208 in the third domain of *SCN1A* that shares 76% homology with humans (Schoonheim et al., 2010; Baraban et al., 2013). Mutations in this channel are one of

the primary genetic causes underlying DS (Claes et al., 2003; Escayg and Goldin, 2010; De Jonghe, 2011; Saitoh et al., 2012). As zebrafish possess two *scn1* genes (Novak et al., 2006), homozygous mutants for *scn1Lab* are comparable to the haploinsufficient clinical condition, and there is no variability from larvae to larvae, or clutch to clutch, with respect to gene inactivation, as is commonly observed with morpholino injections (Kok et al., 2015). Although crosses of heterozygotes produce only one-quarter homozygous *scn1Lab* mutants per mating, there are virtually no limitations on maintaining a large colony of healthy, adult breeders for these types of large-scale screens. Second, it is possible to observe and monitor seizure-like behavior consisting of rapid movements and whole-body convulsions in freely swimming *scn1Lab* mutants as early as 4 dpf that persist for the life of the larvae (~12 dpf). These behaviors are comparable to those observed with exposure to a common convulsant agent (PTZ) and classified as Stage III (Baraban et al., 2005). In




Figure 4. Evaluation of putative antiepileptic drugs in *scn1Lab* mutants. **a**, Locomotion tracking plots for *scn1Lab* zebrafish at baseline and following huperzine A administration. Total movement is shown for a 10 min recording epoch. **b**, Plot showing the change in mean velocity for three different huperzine A concentrations (blue bars). Each bar is the mean change for six fish. The threshold for a positive hit is shown as a dashed line. WT fish exposed to PTZ and huperzine A are shown in red ($N = 7$). **c**, **d**, Same for fenfluramine. Note that 1 mM fenfluramine was toxic, as indicated. **e**, Representative field recordings from *scn1Lab* mutant larvae at 5 dpf. Electrographic activity is shown for a 5 min recording epoch (top traces); high-resolution traces are shown below, as indicated. Note that abnormal burst discharge activity persists in *scn1Lab* mutants exposed to 250 μM fenfluramine. The fish shown were tested in the locomotion assay first.

addition, clear evidence for epileptiform discharge generated in the CNS of immobilized *scn1Lab* mutant larvae has been obtained at ages between 4 and 8 dpf (Baraban et al., 2013). Both zebrafish measures of seizure activity are sensitive to inhibition by AEDs commonly prescribed to children with DS (e.g., valproate, benzodiazepines, and

stiripentol), but are resistant to many antiepileptic compounds (e.g., phenytoin, carbamazepine, ethosuximide, decimemide, tiletamine, primidone, phenacemide, and vigabatrin). Pharmacoresistance is defined as the inability to control seizure activity with at least two different AEDs (Berg, 2009), and, with demonstrated resistance to eight

AEDs, our model clearly fits this definition. This level of model validation has not been possible with morpholinos probably owing to the high degree of variability, or off-target effects, associated with this technique (Kok et al., 2015).

Our screening results highlight the stringency of our approach with a positive hit rate of only 1.97% on the first-stage locomotion assay, and successful identification of 1 compound (of 1012 compounds) with known antiepileptic activity (i.e., dimethadione, a T-type channel antagonist). In additional testing, we confirmed an antiepileptic action for fenfluramine (serotonin uptake inhibitor). Similar to ethosuximide, a reduction in regenerative burst discharges associated with neuronal T-type calcium currents could be the underlying mechanism for dimethadione in DS mutants; however, it is worth noting that T-type channel blockers ethosuximide and flunarizine were not similarly effective (Baraban et al. 2013; this article), and that dimethadione can cause arrhythmia owing to blockade of cardiac human ether-a-go-go-related gene potassium channels (Azarbayjani and Danielsson, 2002; Danielsson et al., 2007). Modulation of serotonin [5-hydroxytryptamine (5-HT)] signaling by blocking uptake or increasing release from neurons by acting as substrates for 5-HT transporter (sertraline) proteins (Fuller et al., 1988; Gobbi and Mennini, 1999; Baumann et al., 2000; Rothman et al., 2010) may be the mechanism of action for fenfluramine in patients with DS, though a detailed analysis of precisely how fenfluramine modulates excitability via this signaling pathway has not been performed. Nonetheless, both drugs probably exert a direct effect on network excitability (at neuronal or synaptic levels, respectively) to suppress electrographic discharge and the associated high-velocity seizure behavior seen in *scn1Lab* mutants, and may be potential targets for clinical use. In contrast, three other drugs identified in the primary locomotion assay were not effective in suppressing electrical events and were designated as false positives. This is not altogether surprising given that the xanthine alkaloid (theobromine), chemotherapeutic (cytarabine), and antibiotic (norfloxacin) mechanisms for these compounds would not be consistent with seizure inhibition. Moreover, the variability inherent in behavioral experiments performed on different zebrafish larvae, in different microplates, and on different days may contribute to these false-positive designations in locomotion assays, and is evident in the range of mean velocity values seen during tracking episodes from control studies (Fig. 1c) or in the failure of many of the initial 20 lead compounds to be confirmed on subsequent retesting (see Fig. 2a). This is a limitation of locomotion-based screening assays and is another reason why a secondary electrophysiology assay on the same zebrafish is a critical advantage of our approach.

An additional advantage of *in vivo* screening with zebrafish larvae is the simultaneous identification of compounds resulting in toxicity. Zebrafish-based anticonvulsant drug-screening assays based primarily on *in situ* hybridization detection of early gene expression at 2 dpf (Baxendale et al., 2012) do not routinely monitor sponta-

neous swim behavior, heart rate, or response to external stimuli. Lacking these real-time measures of toxicity, compounds observed to induce fatality in a freely swimming *scn1Lab*-based behavioral assay (e.g., gemfibrozil, suloctidil, pimozide, or dioxybenzone) were mistakenly classified as seizure-suppressing compounds in the PTZ-based c-Fos *in situ* hybridization assay. Indeed, 41% of the “anticonvulsant” compounds positively identified at 2 dpf in Baxendale et al. (2012) were toxic, proconvulsant, or simply not effective in *scn1Lab* mutant assays at 5–6 dpf. Similarly, it is critical to monitor blood flow and heart activity even in the agar-immobilized electrophysiology assay as compounds effective in suppressing electrical activity can also be toxic. These discrepancies highlight the potential problems associated with zebrafish drug-screening strategies that do not encompass multiple readouts and suggest the need for a note of caution when comparing screening results from different laboratory groups. While any lead compound identified in a zebrafish-based screening assay will, ultimately, need to be independently replicated and/or validated in additional mammalian model systems, the ability to rapidly identify such compounds, while simultaneously identifying potential negative side effects, makes genetically modified zebrafish a unique resource for drug discovery in an age of personalized medicine.

References

- Ali S, Champagne DL, Spaink HP, Richardson MK (2011) Zebrafish embryos and larvae: a new generation of disease models and drug screens. *Birth Defects Res C Embryo Today* 93:115–133. [CrossRef Medline](#)
- Azarbayjani F, Danielsson BR (2002) Embryonic arrhythmia by inhibition of HERG channels: a common hypoxia-related teratogenic mechanism for antiepileptic drugs? *Epilepsia* 43:457–468. [Medline](#)
- Baraban SC, Dinday MT, Hortopan GA (2013) Drug screening in *Scn1a* zebrafish mutant identifies clemizole as a potential Dravet syndrome treatment. *Nat Commun* 4:2410. [CrossRef Medline](#)
- Baraban SC, Taylor MR, Castro PA, Baier H (2005) Pentylentetrazole induced changes in zebrafish behavior, neural activity and c-fos expression. *Neuroscience* 131:759–768. [CrossRef Medline](#)
- Baumann MH, Ayestas MA, Dersch CM, Brockington A, Rice KC, Rothman RB (2000) Effects of phentermine and fenfluramine on extracellular dopamine and serotonin in rat nucleus accumbens: therapeutic implications. *Synapse* 36:102–113. [CrossRef Medline](#)
- Baxendale S, Holdsworth CJ, Meza Santoscoy PL, Harrison MR, Fox J, Parkin CA, Ingham PW, Cunliffe VT (2012) Identification of compounds with anti-convulsant properties in a zebrafish model of epileptic seizures. *Dis Model Mech* 5:773–784. [CrossRef Medline](#)
- Berg AT (2009) Identification of pharmacoresistant epilepsy. *Neuro Clin* 27:1003–1013. [CrossRef Medline](#)
- Bialer M, Johannessen SI, Levy RH, Perucca E, Tomson T, White HS (2015) Progress report on new antiepileptic drugs: a summary of the Twelfth Eilat Conference (EILAT XII). *Epilepsy Res* 111:85–141. [CrossRef Medline](#)
- Boel M, Casaer P (1996) Add-on therapy of fenfluramine in intractable self-induced epilepsy. *Neuropediatrics* 27:171–173. [CrossRef Medline](#)
- Caraballo RH, Cersosimo RO, Sakr D, Cresta A, Escobal N, Fejerman N (2005) Ketogenic diet in patients with Dravet syndrome. *Epilepsia* 46:1539–1544. [CrossRef Medline](#)
- Ceulemans B, Boel M, Leyssens K, Van Rossem C, Neels P, Jorens PG, Lagae L (2012) Successful use of fenfluramine as an add-on treatment for Dravet syndrome. *Epilepsia* 53:1131–1139. [CrossRef Medline](#)

- Chiron C, Dulac O (2011) The pharmacologic treatment of Dravet syndrome. *Epilepsia* 52 Suppl 2:72-75. [CrossRef Medline](#)
- Claes L, Ceulemans B, Audenaert D, Smets K, Löfgren A, Del-Favero J, Ala-Mello S, Basel-Vanagaite L, Plecko B, Raskin S, Thiry P, Wolf NI, Van Broeckhoven C, De Jonghe P (2003) De novo SCN1A mutations are a major cause of severe myoclonic epilepsy of infancy. *Hum Mutat* 21:615-621. [CrossRef Medline](#)
- Coleman BR, Ratcliffe RH, Oguntayo SA, Shi X, Doctor BP, Gordon RK, Nambiar MP (2008) [+-]Huperzine A treatment protects against N-methyl-D-aspartate-induced seizure/status epilepticus in rats. *Chem Biol Interact* 175:387-395. [CrossRef Medline](#)
- Creton R (2009) Automated analysis of behavior in zebrafish larvae. *Behav Brain Res* 203:127-136. [CrossRef Medline](#)
- Danielsson BR, Danielsson C, Nilsson MF (2007) Embryonic cardiac arrhythmia and generation of reactive oxygen species: common teratogenic mechanism for IKr blocking drugs. *Reprod Toxicol* 24:42-56. [CrossRef Medline](#)
- De Jonghe P (2011) Molecular genetics of Dravet syndrome. *Dev Med Child Neurol* 53 Suppl 2:7-10. [CrossRef Medline](#)
- Dravet C, Bureau M, Oguni H, Fukuyama Y, Cokar O (2005) Severe myoclonic epilepsy in infancy: Dravet syndrome. *Adv Neurol* 95: 71-102. [Medline](#)
- Dressler A, Trimmel-Schwahofner P, Reithofer E, Mühlebner A, Gröppel G, Reiter-Fink E, Benninger F, Grassl R, Feucht M (2015) Efficacy and tolerability of the ketogenic diet in Dravet syndrome - Comparison with various standard antiepileptic drug regimen. *Epilepsy Res* 109:81-89. [CrossRef Medline](#)
- ebry Inc. (2011) Guide for the care and use of laboratory animals. Washington, DC: National Academy Press.
- Epi4K Consortium (2012) Epi4K: gene discovery in 4,000 genomes. *Epilepsia* 53:1457-1467. [CrossRef](#)
- Escayg A, Goldin AL (2010) Sodium channel SCN1A and epilepsy: mutations and mechanisms. *Epilepsia* 51:1650-1658. [CrossRef Medline](#)
- Finckbeiner S, Ko PJ, Carrington B, Sood R, Gross K, Dolnick B, Sufirin J, Liu P (2011) Transient knockdown and overexpression reveal a developmental role for the zebrafish *enosf1b* gene. *Cell Biosci* 1:32. [CrossRef Medline](#)
- Fuller RW, Snoddy HD, Robertson DW (1988) Mechanisms of effects of d-fenfluramine on brain serotonin metabolism in rats: uptake inhibition versus release. *Pharmacol Biochem Behav* 30:715-721. [Medline](#)
- Gobbi M, Mennini T (1999) Release studies with rat brain cortical synaptosomes indicate that tramadol is a 5-hydroxytryptamine uptake blocker and not a 5-hydroxytryptamine releaser. *Eur J Pharmacol* 370:23-26. [Medline](#)
- Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, Collins JE, Humphray S, McLaren K, Matthews L, McLaren S, Sealy I, Caccamo M, Churche C, Scott C, Barrett JC, Koch R, Rauch GJ, White S, Chow W, et al (2013) The zebrafish reference genome sequence and its relationship to the human genome. *Nature* 496:498-503.
- Jirsa VK, Stacey WC, Quilichini PP, Ivanov AI, Bernard C (2014) On the nature of seizure dynamics. *Brain* 137:2210-2230. [CrossRef Medline](#)
- Ko HC, Gelb BD (2014) Concise review: drug discovery in the age of the induced pluripotent stem cell. *Stem Cells Transl Med* 3:500-509. [CrossRef Medline](#)
- Kok FO, Shin M, Ni CW, Gupta A, Grosse AS, van Impel A, Kirchmaier BC, Peterson-Maduro J, Kourkoulis G, Male I, DeSantis DF, Sheppard-Tindell S, Ebarasi L, Betsholtz C, Schulte-Merker S, Wolfe SA, Lawson ND (2015) Reverse genetic screening reveals poor correlation between morpholino-induced and mutant phenotypes in zebrafish. *Dev Cell* 32:97-108. [CrossRef Medline](#)
- Leppert MF (1990) Gene mapping and other tools for discovery. *Epilepsia* 31 [Suppl 3]:S11-S18. [Medline](#)
- Lotte J, Haberlandt E, Neubauer B, Staudt M, Kluger GJ (2012) Bromide in patients with SCN1A-mutations manifesting as Dravet syndrome. *Neuropediatrics* 43:17-21. [CrossRef Medline](#)
- Lowson S, Gent JP, Goodchild CS (1990) Anticonvulsant properties of propofol and thiopentone: comparison using two tests in laboratory mice. *Br J Anaesth* 64:59-63. [Medline](#)
- Mahmood F, Mozere M, Zdebik AA, Stanescu HC, Tobin J, Beales PL, Kleta R, Bockenbauer D, Russell C (2013) Generation and validation of a zebrafish model of EAST (epilepsy, ataxia, sensorineural deafness and tubulopathy) syndrome. *Dis Model Mech* 6:652-660. [CrossRef Medline](#)
- Masimirembwa CM, Thompson R, Andersson TB (2001) In vitro high throughput screening of compounds for favorable metabolic properties in drug discovery. *Comb Chem High Throughput Screen* 4:245-263. [Medline](#)
- Novak AE, Jost MC, Lu Y, Taylor AD, Zakon HH, Ribera AB (2006) Gene duplications and evolution of vertebrate voltage-gated sodium channels. *J Mol Evol* 63:208-221. [CrossRef Medline](#)
- Ottman R, Risch N (2012) Genetic epidemiology and gene discovery in epilepsy. In: Jasper's basic mechanisms of the epilepsies (Noebels JL, Avoli M, Rogawski M, Olsen R, Delgado-Escueta A, eds). New York: Oxford UP, pp. 651-658.
- Rafferty TD, Isaacs GM, Yozzo KL, Volz DC (2014) High-content screening assay for identification of chemicals impacting spontaneous activity in zebrafish embryos. *Environ Sci Technol* 48:804-810. [CrossRef Medline](#)
- Rothman RB, Baumann MH, Blough BE, Jacobson AE, Rice KC, Partilla JS (2010) Evidence for noncompetitive modulation of substrate-induced serotonin release. *Synapse* 64:862-869. [Cross-Ref Medline](#)
- Saitoh M, Shinohara M, Hoshino H, Kubota M, Amemiya K, Takahashi JL, Hwang SK, Hirose S, Mizuguchi M (2012) Mutations of the SCN1A gene in acute encephalopathy. *Epilepsia* 53:558-564. [CrossRef Medline](#)
- Schoonheim PJ, Arrenberg AB, Del Bene F, Baier H (2010) Optogenetic localization and genetic perturbation of saccade-generating neurons in zebrafish. *J Neurosci* 30:7111-7120. [CrossRef Medline](#)
- Snowden M, Green DV (2008) The impact of diversity-based, high-throughput screening on drug discovery: "chance favours the prepared mind". *Curr Opin Drug Discov Devel* 11:553-558. [Medline](#)
- Teng Y, Xie X, Walker S, Rempala G, Kozlowski DJ, Mumm JS, Cowell JK (2010) Knockdown of zebrafish *Lgi1a* results in abnormal development, brain defects and a seizure-like behavioral phenotype. *Hum Mol Genet* 19:4409-4420. [CrossRef Medline](#)
- Tonduli LS, Testylier G, Masqueliez C, Lallement G, Monmaur P (2001) Effects of Huperzine used as pre-treatment against soman-induced seizures. *Neurotoxicology* 22:29-37. [Medline](#)
- Wilmshurst JM, Berg AT, Lagae L, Newton CR, Cross JH (2014) The challenges and innovations for therapy in children with epilepsy. *Nat Rev Neurol* 10:249-260. [CrossRef Medline](#)
- Winter MJ, Redfern WS, Hayfield AJ, Owen SF, Valentin JP, Hutchinson TH (2008) Validation of a larval zebrafish locomotor assay for assessing the seizure liability of early-stage development drugs. *J Pharmacol Toxicol Methods* 57:176-187. [CrossRef Medline](#)
- Zhang YF, Gibbs JW 3rd, Coulter DA (1996) Anticonvulsant drug effects on spontaneous thalamocortical rhythms in vitro: ethosuximide, trimethadione, and dimethadione. *Epilepsy Res* 23:15-36. [CrossRef](#)