

# UC Davis

## UC Davis Previously Published Works

### Title

Maternal autoantibody profiles as biomarkers for ASD and ASD with co-occurring intellectual disability

### Permalink

<https://escholarship.org/uc/item/3rr3b669>

### Journal

Molecular Psychiatry, 27(9)

### ISSN

1359-4184

### Authors

Ramirez-Celis, Alexandra

Croen, Lisa A

Yoshida, Cathleen K

et al.

### Publication Date

2022-09-01

### DOI

10.1038/s41380-022-01633-4

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

## ARTICLE OPEN



# Maternal autoantibody profiles as biomarkers for ASD and ASD with co-occurring intellectual disability

Alexandra Ramirez-Celis<sup>1</sup>, Lisa A. Croen<sup>2</sup>, Cathleen K. Yoshida<sup>2</sup>, Stacey E. Alexeeff<sup>2</sup>, Joseph Schauer<sup>1</sup>, Robert H. Yolken<sup>3</sup>, Paul Ashwood<sup>4,5</sup> and Judy Van de Water<sup>1,4</sup>✉

© The Author(s) 2022

Maternal autoantibody-related ASD (MAR ASD) is a subtype of autism in which pathogenic maternal autoantibodies (IgG) cross the placenta, access the developing brain, and cause neurodevelopmental alterations and behaviors associated with autism in the exposed offspring. We previously reported maternal IgG response to eight proteins (CRMP1, CRMP2, GDA, LDHA, LDHB, NSE, STIP1, and YBOX) and that reactivity to nine specific combinations of these proteins (MAR ASD patterns) was predictive of ASD risk. The aim of the current study was to validate the previously identified MAR ASD patterns (CRMP1 + GDA, CRMP1 + CRMP2, NSE + STIP1, CRMP2 + STIP1, LDHA + YBOX, LDHB + YBOX, GDA + YBOX, STIP1 + YBOX, and CRMP1 + STIP1) and their accuracy in predicting ASD risk in a prospective cohort employing maternal samples collected prior to parturition. We used prenatal plasma from mothers of autistic children with or without co-occurring intellectual disability (ASD = 540), intellectual disability without autism (ID = 184) and general population controls (GP = 420) collected by the Early Markers for Autism (EMA) study. We found reactivity to one or more of the nine previously identified MAR ASD patterns in 10% of the ASD group compared with 4% of the ID group and 1% of the GP controls (ASD vs GP: Odds Ratio (OR) = 7.81, 95% Confidence Interval (CI) 3.32 to 22.43; ASD vs ID: OR = 2.77, 95% CI (1.19–7.47)) demonstrating that the MAR ASD patterns are strongly associated with the ASD group and could be used to assess ASD risk prior to symptom onset. The pattern most strongly associated with ASD was CRMP1 + CRMP2 and increased the odds for an ASD diagnosis 16-fold (3.32 to >999.99). In addition, we found that several of these specific MAR ASD patterns were strongly associated with ASD with intellectual disability (ASD + ID) and others associated with ASD without ID (ASD-no ID). Prenatal screening for these MAR patterns may lead to earlier identification of ASD and facilitate access to the appropriate early intervention services based on each child's needs.

*Molecular Psychiatry* (2022) 27:3760–3767; <https://doi.org/10.1038/s41380-022-01633-4>

## INTRODUCTION

Autism spectrum disorder (ASD) is a complex neurodevelopmental disorder characterized by challenges in social communication, restricted interests, and repetitive behaviors [1] that currently affects over 2% of 8-year-olds in the US (1 in 44) [2, 3]. Autism often co-occurs with other conditions such as psychiatric disorders [4, 5] attention deficit hyperactivity disorder (ADHD), intellectual disability (ID), sensory processing, immune dysregulation and gastrointestinal issues [6–8]. Recently, the CDC reported that one-third (34%) of autistic children have intellectual disability (ASD + ID) [3]. It is well recognized that individuals with ASD + ID present significant deficits and challenges with adaptive behavior and therefore have different behavior intervention requirements than those individuals with ASD without ID or ID only [9, 10]. The incidence of autism has increased over the past 50 years; however, we still lack autism-specific biomarkers that would facilitate an earlier diagnosis allowing the provision of directed services based on the ASD sub-phenotypes and associated conditions.

Over the past three decades, multiple studies have suggested an association between maternal immune dysregulation during pregnancy and neurodevelopmental disorders (NDD) in the offspring including autism [11–19]. In particular, gestational exposure to maternal autoantibodies that cross-react with specific fetal brain proteins has been shown to be associated with increased autism risk [20]. We previously reported, in a post-natal sample set from the Childhood Autism Risk- Genes and Environment (CHARGE) study [21] where maternal samples were collected 2–5 years after the birth of the study child, the presence of maternal autoantibodies that recognize eight proteins that are highly expressed in the fetal brain and play significant roles during neurodevelopment. These antigens include collapsin response mediator proteins 1 and 2 (CRMP1, CRMP2), guanine deaminase (GDA), lactate dehydrogenase A and B (LDHA, LDHB), neuron-specific enolase (NSE), stress-induced phosphoprotein-1 (STIP1) and Y-box binding protein 1 (YBOX) [22, 23]. We observed reactivity to single antigens in both the case and control groups; however, reactivity to combinations of two or more specific antigens was associated with autism and was present in up to 18%

<sup>1</sup>Department of Internal Medicine, Division of Rheumatology, Allergy, and Clinical Immunology, One Shields Avenue, University of California, Davis, CA 95616, USA. <sup>2</sup>Kaiser Permanente Division of Research, 2000 Broadway, Oakland, CA 94612, USA. <sup>3</sup>Department of Psychiatry and Behavioral Science, Johns Hopkins University School of Medicine, Baltimore, MD, USA. <sup>4</sup>UC Davis MIND Institute, 2825 50th St, Sacramento, CA 95817, USA. <sup>5</sup>Department of Medical Microbiology and Immunology, One Shields Avenue, University of California, Davis, CA 95616, USA. ✉email: javandewater@ucdavis.edu

Received: 25 April 2022 Revised: 5 May 2022 Accepted: 12 May 2022

Published online: 26 May 2022

of the ASD cases; therefore, we termed this subtype of autism “Maternal Autoantibody Related Autism (MAR ASD)” and the patterns that can predict risk as MAR ASD + patterns. The specific patterns contained CRMP1 + GDA, CRMP1 + CRMP2, NSE + STIP1, CRMP2 + STIP1, LDHA + YBOX, LDHB + YBOX, GDA + YBOX, and each of these patterns had 100% positive predictive value (PPV) in their ability to predict ASD. Further, patterns containing STIP1 + YBOX and CRMP1 + STIP1 had 92% and 90% PPV respectively and were present in less than 10% of the typically developing controls [23].

In the current study, we aimed to conduct an external validation of our previous findings by testing the recently validated MAR ASD + patterns and their predictive potential for autism risk using maternal plasma samples from the prospective Early Markers for Autism (EMA) study [24]. The samples studied herein were collected during mid-pregnancy, allowing us to directly evaluate the relationship between gestational exposure to maternal autoantibodies and child neurodevelopmental outcome. We assessed the association between each MAR ASD + pattern and an ASD diagnosis as well as the phenotypic subgroups of ASD as defined by presence or absence of intellectual disability (ID). To evaluate the specificity of the association of these autoantibodies to ASD, we also assessed associations with an outcome of intellectual disability (ID) in the absence of ASD.

## MATERIALS AND METHODS

### Study subjects

The Early Markers for Autism (EMA) study is a population-based, nested case-control study aimed to investigate genetic and immune susceptibility and environmental exposures that contribute to autism risk. The maternal samples were collected between March 2000 and July 2003 from women participating in the prenatal extended  $\alpha$ -fetoprotein screening program (XAFP) and included subjects from urban, suburban, and rural areas with multicultural backgrounds in Southern California. Children with ASD or intellectual disability without autism (ID) were ascertained from the California Department of Developmental Services (DDS) that provides services through Regional Centers (RC) to people with autism and other developmental disabilities. All study procedures were approved by the Institutional Review Board at Kaiser Permanente and the State Committee for the Protection of Human Subjects as previously described [24–26].

### Diagnostic verification

The EMA study population and diagnosis classification was recently described in detail by ref. [24]. Briefly, RC records for all children receiving services for ASD or ID were reviewed by expert clinicians and final case status was determined according to the DSM-IV-TR criteria. The ASD group was further categorized into two subgroups based on cognitive scores: Autism without intellectual disability (ASD no-ID) and ASD with intellectual disability (ASD + ID). ID determination was based on RC records with composite scores <70 on all standardized cognitive/functional tests and these designations were reviewed by our expert clinicians. General population (GP) controls were randomly selected from birth certificate files and frequency matched to the ASD group on age, sex, and county of residence at birth. GP controls were not receiving services from a RC and were not verified as typically developing. A summary of the study population can be found in Table 1.

### Specimen collection

Maternal blood was collected at mid pregnancy (15–20 weeks of gestation) in citrate dextrose. Plasma was separated, labeled, and stored at  $-80^{\circ}\text{C}$ . Prior to use, samples were thawed at room temperature (RT), vortexed, and centrifuged at 13,000 RPM for 10 min.

### Enzyme linked immunosorbent assay (ELISA)

Maternal antibody cross-reactivity against the eight antigens was determined by Enzyme-Linked Immunosorbent Assay (ELISA) using custom-made and commercially available proteins [23, 27]. The protein concentration and plasma dilutions were optimized for each antigen as previously described [23, 27]. In summary, microplates were coated with 100  $\mu\text{l}$  of antigen in carbonate coating buffer pH 9.6, incubated overnight

**Table 1.** Group classifications of study population.

| Group                            | Subjects, N |
|----------------------------------|-------------|
| Autism spectrum disorder (ASD)   | 540         |
| Intellectual disability (ID)     | 184         |
| General Population (GP)          | 420         |
| ASD with intellectual disability |             |
| Yes (ASD + ID)                   | 285         |
| No (ASD no-ID)                   | 219         |
| Unknown                          | 36          |

Group classifications included in the Early Markers for Autism (EMA) study.

at  $4^{\circ}\text{C}$ , washed four times with Phosphate Buffered Saline Tween-20 (PBST) 0.05%, and blocked with 2% Super Block (Thermo Scientific, Rockford, IL) for 1 h at RT. Then, 100  $\mu\text{l}$  of the diluted sample was added to each well, incubated for 1.5 h, washed 4 times in PBST 0.05% and incubated for 1 hour with 1:10,000 goat anti-human IgG-HRP IgG (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, MA), followed by four washes with (PBST) 0.05%. We only assessed IgG reactivity as it is the only isotype that can cross the placenta. Finally, 100  $\mu\text{l}$  of BD optEIA liquid substrate for ELISA (BD Biosciences, San Jose, CA) was added to each well, and the reaction was stopped after 4 min with 50  $\mu\text{l}$  of 2 N HCl. The absorbance was measured at 490–450 nm using an iMark Microplate Absorbance Reader (Biorad, Hercules, CA, USA).

### Statistical analysis

For the ELISA, a positive cutoff value for each antigen was determined after plate-plate normalization using an ROC curve, and the Youden’s index as previously described [23, 27]. Comparisons of sociodemographic characteristics between ASD, ID and GP groups were calculated using  $\chi^2$  test (statistical significance threshold set at  $p < 0.05$ ). To evaluate the association between reactivity to each of the MAR ASD + patterns and child outcome, we performed group comparisons using Fisher Exact test (ASD vs GP, ASD vs ID, ID vs GP, and ASD + ID vs ASD no-ID) and calculated the odds ratios (ORs) with 95% confidence intervals (95% CIs) using exact logistic regression, which is appropriate for small or zero cell sizes. Note that a zero-cell size for the number exposed in the reference group results in an unbounded upper confidence interval limit for the OR, displayed as >999.99. Additionally, we used  $\chi^2$  tests ( $p < 0.05$ ) to examine associations between sociodemographic characteristics and the MAR ASD + patterns.

## RESULTS

### Population sociodemographic characteristics

Mothers of autistic children were more likely to be older, non-Hispanic, and have higher education compared with mothers of the ID and GP groups (Table 2). Mothers of ID children were more likely to be younger, less educated, Mexican-born, and deliver prematurely compared with the ASD and GP groups (Table 2). Within the autism group, the mothers of children with ASD + ID were more likely to be multiparous, Hispanic, born in Mexico and have a lower level of education compared with mothers of children with ASD no-ID (Table 3).

### Autoantibody reactivity against fetal brain antigens

Maternal autoantibody reactivity to at least one antigen was observed in more than half of the study participants in each study group (60% of ASD, 54% of ID, and 57% of GP;  $\chi^2 p = 0.34$ ). Table 4 presents maternal IgG reactivity to the MAR ASD + patterns that predicted ASD risk in our previous discovery study [23] and were validated in this dataset. We found that 10% of the ASD group had significant IgG reactivity to any of the previously identified ASD-specific patterns compared with 4% of the ID group and 1% of GP controls (ASD vs GP: OR = 7.81, 95% CI 3.32–22.43,  $p < 0.001$ ; ASD vs ID: OR = 2.77, 95% CI 1.19–7.47,  $p = 0.01$ ). Although not a

**Table 2.** Descriptive characteristics of the EMA study population.

|                        | <b>ASD</b><br><b>N = 540</b><br><b>N (%)</b> | <b>ID</b><br><b>N = 184</b><br><b>N (%)</b> | <b>GP</b><br><b>N = 420</b><br><b>N (%)</b> | <b>ASD vs GP</b><br>$\chi^2$ <b>p value</b> | <b>ASD vs ID</b><br>$\chi^2$ <b>p value</b> | <b>ID vs GP</b><br>$\chi^2$ <b>p value</b> |
|------------------------|--|---|---|---|---|--|
| Maternal age           |  |   |   | 0.08  | <b>&lt;0.001</b>                            | <b>&lt;0.001</b>                           |
| <20                    | 17 (3.15%)                                   | 26 (14.13%)                                 | 21 (5.0%)                                   |   |   |  |
| 20–24                  | 80 (14.81%)                                  | 41 (22.28%)                                 | 70 (16.67%)                                 |   |   |  |
| 25–29                  | 146 (27.04%)                                 | 51 (27.72%)                                 | 129 (30.71%)                                |   |   |  |
| 30–34                  | 196 (36.3%)                                  | 43 (23.37%)                                 | 145 (34.52%)                                |   |   |  |
| >=35                   | 101 (18.7%)                                  | 23 (12.5%)                                  | 55 (13.1%)                                  |   |   |  |
| Maternal race          |  |   |   | 0.20  | <b>&lt;0.001</b>                            | <b>0.01</b>                                |
| White                  | 399 (73.89%)                                 | 160 (86.96%)                                | 334 (79.52%)                                |   |   |  |
| Asian                  | 109 (20.19%)                                 | 11 (5.98%)                                  | 65 (15.48%)                                 |   |   |  |
| Other                  | 27 (5.0%)                                    | 12 (6.52%)                                  | 19 (4.52%)                                  |   |   |  |
| Missing                | 5 (0.93%)                                    | 1 (0.54%)                                   | 2 (0.48%)                                   |   |   |  |
| Maternal ethnicity     |  |   |   | 0.14  | <b>&lt;0.001</b>                            | <b>&lt;0.001</b>                           |
| Hispanic               | 220 (40.74%)                                 | 128 (69.57%)                                | 198 (47.14%)                                |   |   |  |
| Not Hispanic           | 315 (58.33%)                                 | 55 (29.89%)                                 | 219 (52.14%)                                |   |   |  |
| Missing                | 5 (0.93%)                                    | 1 (0.54%)                                   | 3 (0.71%)                                   |   |   |  |
| Parity                 |  |   |   | 0.02  | <b>0.005</b>                                | 0.30                                       |
| Multiparous            | 294 (54.44%)                                 | 122 (66.3%)                                 | 260 (61.9%)                                 |   |   |  |
| Primiparous            | 246 (45.56%)                                 | 62 (33.7%)                                  | 160 (38.1%)                                 |   |   |  |
| Maternal birth country |  |   |   | <b>0.05</b>                                 | <b>&lt;0.001</b>                            | <b>&lt;0.001</b>                           |
| US                     | 267 (49.44%)                                 | 83 (45.11%)                                 | 203 (48.33%)                                |   |   |  |
| Mexico                 | 130 (24.07%)                                 | 84 (45.65%)                                 | 127 (30.24%)                                |   |   |  |
| Other                  | 143 (26.48%)                                 | 17 (9.24%)                                  | 90 (21.43%)                                 |   |   |  |
| Maternal education     |  |   |   | <b>0.01</b>                                 | <b>&lt;0.001</b>                            | <b>&lt;0.001</b>                           |
| <High School           | 98 (18.15%)                                  | 78 (42.39%)                                 | 102 (24.29%)                                |   |   |  |
| High School Grad       | 114 (21.11%)                                 | 47 (25.54%)                                 | 112 (26.67%)                                |   |   |  |
| Undergrad College      | 220 (40.74%)                                 | 46 (25.0%)                                  | 144 (34.29%)                                |   |   |  |
| Post-Grad College      | 100 (18.52%)                                 | 11 (5.98%)                                  | 59 (14.05%)                                 |   |   |  |
| Unknown                | 8 (1.48%)                                    | 2 (1.09%)                                   | 3 (0.71%)                                   |   |   |  |
| Child characteristics  |  |   |   |   |   |  |
| Child Sex              |  |   |   | 0.62  | <b>&lt;0.001</b>                            | <b>&lt;0.001</b>                           |
| Male                   | 442 (81.85%)                                 | 102 (55.43%)                                | 349 (83.1%)                                 |   |   |  |
| Female                 | 98 (18.15%)                                  | 82 (44.57%)                                 | 71 (16.9%)                                  |   |   |  |
| Birth type             |  |   |   | 0.33  | 0.93  | 0.39                                       |
| Singleton              | 526 (97.41%)                                 | 179 (97.28%)                                | 413 (98.33%)                                |   |   |  |
| Multiple               | 14 (2.59%)                                   | 5 (2.72%)                                   | 7 (1.67%)                                   |   |   |  |
| Birth weight           |  |   |   | 0.10  | <b>&lt;0.001</b>                            | <b>&lt;0.001</b>                           |
| <1500 g                | 5 (0.93%)                                    | 14 (7.61%)                                  | 6 (1.43%)                                   |   |   |  |
| 1500–2499 g            | 37 (6.85%)                                   | 25 (13.59%)                                 | 16 (3.81%)                                  |   |   |  |
| >=2500 g               | 498 (92.22%)                                 | 145 (78.8%)                                 | 398 (94.76%)                                |   |   |  |
| Gestational age        |  |   |   | 0.92  | <b>&lt;0.001</b>                            | <b>&lt;0.001</b>                           |
| <32 weeks              | 11 (2.04%)                                   | 16 (8.7%)                                   | 9 (2.14%)                                   |   |   |  |
| 33–36 weeks            | 41 (7.59%)                                   | 26 (14.13%)                                 | 29 (6.9%)                                   |   |   |  |
| >=37 weeks             | 488 (90.37%)                                 | 142 (77.17%)                                | 382 (90.95%)                                |   |   |  |

Demographic differences between groups was calculated by  $\chi^2$  test and *p* values > 0.05 were bolded and considered significant. ASD Autism Spectrum Disorders, ID intellectual disability, GP general population.

statistically significant difference, IgG reactivity to any of the MAR ASD + patterns was nearly 3 times as common among the ID group compared with the GP control group (ID vs GP: OR = 2.72, 95% CI 0.77–9.96, *p* = 0.07).

The pattern with the highest odds associated with ASD was CRMP1 + CRMP2, which was present in ~3% of the ASD group versus 0% of the GP and ID groups. Maternal autoantibody reactivity to both CRMP1 and CRMP2 increased the odds of an

**Table 3.** Descriptive characteristics of ASD phenotypic subgroups.

|                        | <b>ASD + ID</b> | <b>ASD no-ID</b> | <b>ASD-ID vs<br/>ASD no-ID</b>     |
|------------------------|-----------------|------------------|------------------------------------|
|                        | <b>N = 285</b>  | <b>N = 219</b>   |                                    |
|                        | <b>N (%)</b>    | <b>N (%)</b>     | <b><math>\chi^2</math> p value</b> |
| Maternal age           |                 |                  | 0.18                               |
| <20                    | 12 (4.21%)      | 5 (2.28%)        |                                    |
| 20–24                  | 50 (17.54%)     | 25 (11.42%)      |                                    |
| 25–29                  | 71 (24.91%)     | 67 (30.59%)      |                                    |
| 30–34                  | 98 (34.39%)     | 81 (36.99%)      |                                    |
| >=35                   | 54 (18.95%)     | 41 (18.72%)      |                                    |
| Maternal race          |                 |                  | 0.34                               |
| White                  | 203 (71.23%)    | 170 (77.63%)     |                                    |
| Asian                  | 62 (21.75%)     | 37 (16.89%)      |                                    |
| Other                  | 16 (5.61%)      | 11 (5.02%)       |                                    |
| Missing                | 4 (1.4%)        | 1 (0.46%)        |                                    |
| Maternal ethnicity     |                 |                  | <b>0.03</b>                        |
| Hispanic               | 129 (45.26%)    | 77 (35.16%)      |                                    |
| Not Hispanic           | 152 (53.33%)    | 141 (64.38%)     |                                    |
| Missing                | 4 (1.4%)        | 1 (0.46%)        |                                    |
| Parity                 |                 |                  | <b>0.01</b>                        |
| Multiparous            | 167 (58.6%)     | 102 (46.58%)     |                                    |
| Primiparous            | 118 (41.4%)     | 117 (53.42%)     |                                    |
| Maternal birth country |                 |                  | <b>0.03</b>                        |
| US                     | 126 (44.21%)    | 123 (56.16%)     |                                    |
| Mexico                 | 77 (27.02%)     | 45 (20.55%)      |                                    |
| Other                  | 82 (28.77%)     | 51 (23.29%)      |                                    |
| Maternal education     |                 |                  | <b>0.00</b>                        |
| <High School           | 61 (21.4%)      | 26 (11.87%)      |                                    |
| High School Grad       | 69 (24.21%)     | 42 (19.18%)      |                                    |
| Undergrad College      | 105 (36.84%)    | 101 (46.12%)     |                                    |
| Post-Grad College      | 44 (15.44%)     | 49 (22.37%)      |                                    |
| Unknown                | 6 (2.11%)       | 1 (0.46%)        |                                    |
| Child characteristics  |                 |                  |                                    |
| Child sex              |                 |                  | 0.74                               |
| Male                   | 231 (81.05%)    | 180 (82.19%)     |                                    |
| Female                 | 54 (18.95%)     | 39 (17.81%)      |                                    |
| Birth type             |                 |                  | 0.64                               |
| Singleton              | 279 (97.89%)    | 213 (97.26%)     |                                    |

**Table 3.** continued

|                 | <b>ASD + ID</b> | <b>ASD no-ID</b> | <b>ASD-ID vs<br/>ASD no-ID</b>     |
|-----------------|-----------------|------------------|------------------------------------|
|                 | <b>N = 285</b>  | <b>N = 219</b>   |                                    |
|                 | <b>N (%)</b>    | <b>N (%)</b>     | <b><math>\chi^2</math> p value</b> |
| Multiple        | 6 (2.11%)       | 6 (2.74%)        |                                    |
| Birth weight    |                 |                  | 0.19                               |
| <1500 g         | 5 (1.75%)       | 0 (0%)           |                                    |
| 1500–2499 g     | 22 (7.72%)      | 14 (6.39%)       |                                    |
| >=2500 g        | 258 (90.53%)    | 205 (93.61%)     |                                    |
| Gestational age |                 |                  | 0.23                               |
| <32 weeks       | 8 (2.81%)       | 3 (1.37%)        |                                    |
| 33–36 weeks     | 24 (8.42%)      | 12 (5.48%)       |                                    |
| >=37 weeks      | 253 (88.77%)    | 204 (93.15%)     |                                    |

Demographic differences between groups was calculated by  $\chi^2$  test and *p* values > 0.05 were bolded and considered significant.

*ASD + ID* Autism spectrum disorder with co-occurring intellectual disability, *ASD no-ID* autism spectrum disorder without co-occurring intellectual disability.

ASD diagnosis to nearly 16-fold relative to the GP controls (ASD vs GP: OR = 15.68, 95% CI 3.32–>999.99, *p* < 0.001) and over 6-fold relative to the ID group (ASD vs ID: *p* = 0.04, OR = 6.46, 95% CI 1.32–>999.99). Other patterns significantly associated with ASD risk included CRMP2 + STIP1, LDHA + YBOX, GDA + YBOX, and CRMP1 + STIP1 when compared to the GP group (Table 4). These patterns did not display reactivity differences between ID and GP, suggesting that reactivity to MAR ASD + patterns is highly correlated with an autism diagnosis.

Further, we evaluated the MAR ASD + patterns for the ASD phenotypic subgroups based on the co-occurrence with ID (Table 5). Reactivity to any of MAR ASD + patterns significantly increased the odds for both ASD + ID (ASD + ID vs GP: OR 8.7, 95% CI 3.52–25.82, *p* = < 0.001,) and ASD no-ID (ASD no-ID vs GP: OR 7.29, 95% CI 2.79–22.44, *p* = < 0.001) with similar OR for both groups. The pattern associated with the highest odds of both ASD + ID and ASD no-ID diagnosis was CRMP1 + CRMP2 (ASD + ID vs GP: OR 18.91, 95% CI 3.81 to >999.99, *p* = < 0.001; ASD no-ID vs GP: OR 13.11, 95% CI 2.37 to >999.99, *p* < 0.01). LDHA + YBOX pattern also showed similar magnitudes of increased odds of both ASD + ID and ASD no-ID (ASD + ID vs GP: OR 7.46, 95% CI 0.83–354.71, *p* = < 0.04 and ASD no-ID vs GP: OR 7.77, 95% CI 0.76–384.79, *p* < 0.05). Although there were no statistically significant differences between ASD + ID vs ASD no-ID with respect to the various antigen patterns, CRMP2 + STIP1 and CRMP1 + GDA trended higher in the ASD + ID group (Table 5). In addition, CRMP2 + STIP1 and CRMP1 + GDA were strongly associated with ASD + ID vs GP but did not reach statistical significance for ASD no-ID vs GP. The CRMP1 + STIP1 and GDA + YBOX patterns were associated with ASD no-ID vs GP but did not reach statistical significance for ASD no-ID vs GP (Table 5).

Lastly, we explored if having an ASD MAR + pattern was associated with any sociodemographic factors (Supplementary fig. 1). There were no statistically significant differences in sociodemographic profiles between women with reactivity to any of the ASD-specific antigen combination and women with no reactivity to any combination. While some patterns were present only in Hispanic women (NSE + STIP1) or women who delivered male offspring (LDHA + YBOX and GDA + YBOX), the study was underpowered for statistical analysis of individual patterns.

**Table 4.** Analysis of previously identified maternal antibody patterns (ASD MAR + patterns).

| Antibody Pattern | ASD<br>N = 540<br>N (%) | ID<br>N = 184<br>N (%) | GP<br>N = 420<br>N (%) | ASD vs GP<br>Fisher's Exact<br>p value | ASD vs GP<br>OR 95% CI            | ASD vs ID<br>Fisher's Exact<br>p value | ASD vs ID<br>OR 95% CI           | ID vs GP<br>Fisher's Exact<br>p value | ID vs GP<br>OR 95% CI  |
|------------------|-------------------------|------------------------|------------------------|--|-----------------------------------|--|----------------------------------|---------------------------------------|------------------------|
| ANY COMBO        | 55 (10.19)              | 7 (3.8)                | 6 (1.43)               | <0.001                                 | <b>7.81 (3.32 to 22.43)</b>       | <b>0.01</b>                            | <b>2.86 (1.27 to 7.6)</b>        | 0.07                                  | 2.72 (0.77 to 9.96)    |
| CRMP1 + CRMP2    | 14 (2.59)               | 0 (0)                  | 0 (0)                  | <0.001                                 | <b>15.68 (3.32 to &gt;999.99)</b> | <b>0.03</b>                            | <b>6.86 (1.45 to &gt;999.99)</b> | NA                                    | NA                     |
| CRMP2 + STIP1    | 9 (1.67)                | 1 (0.54)               | 0 (0)                  | <b>0.01</b>                            | <b>9.85 (1.99 to &gt;999.99)</b>  | 0.47                                   | 3.1 (0.42 to 136.68)             | 0.30                                  | 2.28 (0.12 to >999.99) |
| LDHA + YBOX      | 10 (1.85)               | 2 (1.09)               | 1 (0.24)               | <b>0.03</b>                            | <b>7.89 (1.12 to 343.89)</b>      | 0.74                                   | 1.72 (0.36 to 16.25)             | 0.22                                  | 4.59 (0.24 to 272.28)  |
| GDA + YBOX       | 6 (1.11)                | 2 (1.09)               | 0 (0)                  | <b>0.04</b>                            | <b>6.4 (1.21 to &gt;999.99)</b>   | 1.00                                   | 1.02 (0.18 to 10.45)             | 0.09                                  | 5.53 (0.66 to >999.99) |
| CRMP1 + STIP1    | 9 (1.67)                | 0 (0)                  | 1 (0.24)               | <b>0.05</b>                            | <b>7.09 (0.98 to 312.00)</b>      | 0.12                                   | 4.31 (0.87 to >999.99)           | 1.00                                  | 2.28 (0. to 43.37)     |
| NSE + STIP1      | 5 (0.92)                | 0 (0)                  | 0 (0)                  | 0.07                                   | 5.26 (0.95 to >999.99)            | 0.34                                   | 2.30 (0.42 to >999.99)           | NA                                    | NA                     |
| CRMP1 + GDA      | 8 (1.48)                | 1 (0.5)                | 1 (0.24)               | 0.09                                   | 6.29 (0.84 to 280.23)             | 0.46                                   | 2.75 (0.36 to 122.77)            | 0.52                                  | 2.29 (0.03 to 180.15)  |
| LDHB + YBOX      | 4 (0.74)                | 0 (0)                  | 0 (0)                  | 0.14                                   | 4.13 (0.7 to >999.99)             | 0.58                                   | 1.81 (0.31 to >999.99)           | NA                                    | NA                     |
| STIP1 + YBOX     | 4 (0.74)                | 2 (1.09)               | 3 (0.71)               | 1.00                                   | 1.04 (0.17 to 7.12)               | 0.65                                   | 0.68 (0.10 to 7.57)              | 0.64                                  | 1.53 (0.13 to 13.44)   |

ASD Autism Spectrum Disorders, ID intellectual disability, GP general population, CRMP1 and CRMP2 collapsin response mediator proteins 1 and 2, GDA guanine deaminase, NSE neuron specific enolase, LDHA-B lactate dehydrogenase A and B, STIP1 stress induced phosphoprotein 1, YBOX Y-box binding protein 1. Fisher's exact test (two-sided) was used to evaluate the association of the patterns with ASD diagnosis and *p* values > 0.05 were bolded and considered significant.

The odds ratios (ORs) with 95% confidence intervals (95% CIs) using exact logistic regression, which is appropriate for small or zero cell sizes. Note that a zero-cell size for the number exposed in the reference group results in an unbounded upper confidence interval limit for the OR, displayed as >999.99.

**Table 5.** Maternal IgG reactivity to MAR ASD + patterns by ASD phenotypic subgroups.

| Antibody Pattern | ASD + ID<br>N = 285<br>N (%) | ASD no-ID<br>N = 219<br>N (%) | GP<br>N = 420<br>N (%) | ASD + ID vs GP<br>Fisher's Exact<br>p value | ASD + ID vs GP<br>OR 95% CI       | ASD no-ID vs GP<br>Fisher's Exact<br>p value | ASD no-ID vs GP<br>OR 95% CI      | ASD + ID vs ASD no-ID<br>Fisher's Exact<br>p value | ASD + ID vs ASD no-ID<br>OR 95% CI |
|------------------|------------------------------|-------------------------------|------------------------|---|-----------------------------------|--|-----------------------------------|--|------------------------------------|
| ANY COMBO        | 32 (11.23)                   | 21 (9.59)                     | 6 (1.43)               | <0.001                                      | <b>8.7 (3.52 to 25.82)</b>        | < 0.01                                       | <b>7.29 (2.79 to 22.44)</b>       | 0.66   | 1.19 (0.64 to 2.25)                |
| CRMP1 + CRMP2    | 9 (3.16)                     | 5 (2.28)                      | 0 (0)                  | <0.001                                      | <b>18.91 (3.81 to &gt;999.99)</b> | < 0.01                                       | <b>13.11 (2.37 to &gt;999.99)</b> | 0.60   | 1.39 (0.41 to 5.38)                |
| CRMP2 + STIP1    | 8 (2.81)                     | 1 (0.46)                      | 0 (0)                  | <0.001                                      | <b>16.66 (3.3 to &gt;999.99)</b>  | 0.34   | 1.92 (0.1 to >999.99)             | 0.08   | 6.28 (0.83 to 280.63)              |
| CRMP1 + GDA      | 7 (2.46)                     | 1 (0.46)                      | 1 (0.24)               | <b>0.01</b>                                 | <b>10.52 (1.34 to 476.6)</b>      | 1.00   | 1.92 (0.02 to 151.23)             | 0.15   | 5.48 (0.7 to 248.51)               |
| LDHA + YBOX      | 5 (1.75)                     | 4 (1.83)                      | 1 (0.24)               | <b>0.04</b>                                 | <b>7.46 (0.83 to 354.71)</b>      | <b>0.05</b>                                  | <b>7.77 (0.76 to 384.79)</b>      | 1.00   | 0.96 (0.2 to 4.9)                  |
| NSE + STIP1      | 2 (0.7)                      | 2 (0.91)                      | 0 (0)                  | 0.16  | 3.57 (0.42 to >999.99)            | 0.12   | 4.65 (0.55 to >999.99)            | 1.00   | 0.77 (0.06 to 10.66)               |
| LDHB + YBOX      | 2 (0.7)                      | 2 (0.91)                      | 0 (0)                  | 0.16  | 3.57 (0.42 to >999.99)            | 0.12   | 4.65 (0.55 to >999.99)            | 1.00   | 0.77 (0.06 to 10.66)               |
| CRMP1 + STIP1    | 4 (1.4)                      | 5 (2.28)                      | 1 (0.24)               | 0.16  | 5.95 (0.58 to 294.42)             | <b>0.02</b>                                  | <b>9.76 (1.08 to 464.1)</b>       | 0.51   | 0.61 (0.12 to 2.87)                |
| GDA + YBOX       | 1 (0.35)                     | 4 (1.83)                      | 0 (0)                  | 0.40  | 1.47 (0.08 to >999.99)            | <b>0.01</b>                                  | <b>10.26 (1.73 to &gt;999.99)</b> | 0.17   | 0.19 (0.02 to 1.94)                |
| STIP1 + YBOX     | 1 (0.35)                     | 2 (0.91)                      | 3 (0.71)               | 0.65  | 0.49 (0.01 to 6.14)               | 1.00   | 1.28 (0.11 to 11.27)              | 0.58   | 0.38 (0.01 to 7.4)                 |

ASD + ID Autism spectrum disorder with co-occurring intellectual disability, ASD no-ID autism spectrum disorder without co-occurring intellectual disability, GP general population, CRMP1 and CRMP2 collapsin response mediator proteins 1 and 2, GDA guanine deaminase, NSE neuron specific enolase, LDHA-B lactate dehydrogenase A and B, STIP1 stress induced phosphoprotein 1, YBOX Y-box binding protein 1.

Fisher's exact test (two-sided) was used to evaluate the association of the patterns with ASD diagnosis and *p* values > 0.05 were bolded and considered significant.

The odds ratios (ORs) with 95% confidence intervals (95% CIs) using exact logistic regression, which is appropriate for small or zero cell sizes. Note that a zero-cell size for the number exposed in the reference group results in an unbounded upper confidence interval limit for the OR, displayed as >999.99.

## DISCUSSION

During pregnancy, the body goes through numerous adaptations and changes [28] including the establishment of maternal-fetal immune homeostasis, which provides protection against pathogens while allowing the allogenic embryo to implant and develop [29, 30].

Among the five primary immunoglobulin isotypes, IgG, IgM, IgA, IgD, and IgE, only maternal IgG can cross the uninfamed placenta in appreciable quantities and provide passive protection to the fetus beginning in gestational week 14 [31]. However, pathogenic autoantibodies can also cross the placenta to react with antigens



in the fetal compartment, and may contribute to neonatal diseases such as neonatal lupus, neonatal anemia, neonatal pemphigus and neonatal myasthenia gravis [32] as well as neurodevelopmental disorders such as ID [33–35], ADHD [36], and ASD [15, 22, 23, 37]. This project aimed to expand our earlier discovery results and evaluate the previously described MAR ASD + patterns [23] as potential early biomarkers for ASD using maternal blood samples collected mid-gestation [24], allowing us to assess the pathological significance of gestational exposure to these maternal autoantibodies to child neurodevelopmental outcome.

Over half of the maternal samples in each study group had IgG reactivity to one or more antigens, indicating that reactivity to an individual protein-target is not associated with child outcome. However, reactivity to any of the previously described MAR ASD + patterns, which occurred in 10% of the ASD group and 1% of the GP group, was highly predictive of ASD risk, consistent with our recent reports [22, 23]. Of interest, 4% of the ID group also reacted to one or more of the patterns but this was not significantly different than the 1% among the GP population, suggesting that MAR ASD + patterns were more strongly associated with an ASD diagnosis. Consistent with our prior study that analyzed maternal samples collected 2–5 years after the birth of the study child, in the present study, maternal IgG reactivity during pregnancy to CRMP1 + CRMP2, CRMP2 + STIP1, LDHA + YBOX, GDA + YBOX, and CRMP1 + STIP1 significantly increased the odds of an ASD diagnosis in the exposed child. However, the MAR ASD + patterns found most frequently among the ASD group differed between the two studies. The most abundant patterns in our previous study using the CHARGE study samples [23], were CRMP1 + GDA, followed by CRMP1 + CRMP2, and NSE + STIP1. In the present study (EMA), CRMP1 + GDA and NSE + STIP1 were less prevalent and only trended towards significantly increasing the odds for ASD diagnosis. The distribution discrepancies between the CHARGE and EMA studies could be due to demographic or geographical differences of the study populations (Northern CA vs Southern CA), the years during which the pregnancies occurred (2002–2012 vs 2000–2003), and/or the time period during which the maternal samples were collected (2–5 years post-delivery vs. second trimester) [21, 24].

In the current study, the MAR ASD + pattern associated with the highest odds of ASD was CRMP1 + CRMP2, increasing odds of ASD nearly 16-fold compared to the GP controls. This pattern was associated with the highest odds for both ASD + ID and ASD no-ID compared with GP controls. In contrast, the patterns CRMP2 + STIP1 and CRMP1 + GDA increased the odds for an ASD + ID diagnosis suggesting that maternal autoantibodies against these protein combinations could target shared pathways between ASD and ID, altering both behavior and cognition. Interestingly, GDA + YBOX and CRMP1 + STIP1 increased the odds of the ASD no-ID phenotype. Each of the target proteins are biologically relevant due to their key role in brain development as dendritic arborization, organization and maintenance of neural network [38–41], brain metabolism [39], cognition/memory formation [42], neuroprotection, and CNS homeostasis [43, 44]. Previous studies have shown that mutations or deficits in these proteins are associated with neurological pathology, such as ASD [45, 46], schizophrenia [47, 48], ADHD [49, 50], and intellectual disability [51, 52]. Therefore, we hypothesize that gestational exposure to maternal autoantibodies that cross-react with relevant brain proteins could have an additive effect in altering neurodevelopmental pathways and contribute to ASD etiology and pathogenesis. Thus the different MAR ASD + patterns presented here could serve as biomarkers not only for ASD but for specific ASD phenotypes.

Other studies have looked at maternal IgG reactivity to single proteins and their association with NDD, reviewed in [15, 16, 20]. For example, other clinical studies and animal models have reported that maternal antibodies that target CASPR2, a potential ASD-risk biomarker, alter brain anatomy, function, and are related to autism manifestations and learning issues in the exposed

offspring [20, 35, 37, 53–55]. Using gestational plasma collected from a subset of a large Danish study with over 100,000 participants, Coutinho and collaborators investigated maternal autoantibody against multiple brain proteins (including NMDA and CASPR2) and their association with child outcome. They reported a strong association between autoantibody reactivity to NMDA and CASPR2 and ID, but not ASD [35]. Thus, the utility of using maternal reactivity to NMDA and CASPR2 antibodies as biomarkers of ID and ASD would need clinical additional validation.

Although ASD can be diagnosed as early as 18 months of age, most children receive a diagnosis after the three years of age [56] thereby delaying early intervention services that would improve life outcomes. There is a high co-occurrence of ASD with intellectual disability (30–70%) [3, 57], and the ability to distinguish between ID, ASD no-ID and ASD + ID in the first years of life would enable clinicians to provide more targeted behavioral interventions [10]. Therefore, it is of clinical importance to develop biomarkers that can not only identify risk of ASD, but provide information regarding the ASD phenotype, allowing the clinical intervention strategy to be better tailored to the child's specific needs and strengths [10]. The MAR ASD + patterns presented herein provide information regarding potential candidates for use as biomarkers of ASD risk to be further validated in future studies.

Some limitations of the current study deserve mention. First, we could not verify that all children in the GP control group were typically developing. While the GP controls had never been clients of a Regional Center, it is possible that some may have an undiagnosed developmental disorder, which could account for the presence of some of the MAR ASD + patterns in 1% of the GP group. A second limitation was the small number of maternal samples with MAR ASD + patterns in the ASD + ID and ASD no-ID groups. This reduced our ability to reach statistical significance for some of the less common patterns. Future studies are underway to expand the study population and include information about environmental factors which might increase the level of autoantibodies such as infection, medications, cigarette smoking, and gestational exposure to wildfires and farmland pesticides.

One of the greatest strengths of the current study is that the samples were collected during mid-pregnancy, demonstrating the predicative value of maternal IgG reactivity against MAR ASD + patterns and child outcomes. In addition, the EMA study included children with ASD no-ID and ASD + ID as well as children with other developmental disorders without ASD, allowing us to identify maternal autoantibody patterns predictive of specific neurodevelopmental phenotypes. Finally, we had relevant sociodemographic information for this diverse sample set that allowed us to make interesting observations that should be confirmed in future studies. Forthcoming research will include clinical validation of the MAR ASD + patterns in larger cohorts from different geographical regions, and the creation of in vitro and in vivo models to study the biological pathways involved in MAR ASD while considering ASD in co-occurrence with ID. We aim to develop accurate biomarkers to provide clinicians with additional tools for an earlier diagnosis of ASD, and to better tailor intervention services based on the ASD phenotype and the child's individual strengths and specific challenges.

## REFERENCES

1. Association AP. *Diagnostic and statistical manual of mental disorders: DSM-5™, 5th ed.* American Psychiatric Publishing, Inc.: Arlington, VA, US, 2013, xlv, 947–xlv, 947pp.
2. Baio J, Wiggins L, Christensen DL, Maenner MJ, Daniels J, Warren Z, et al. Prevalence of Autism Spectrum Disorder Among Children Aged 8 Years - Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2014. *MMWR Surveill Summ.* 2018;67:1–23.
3. Maenner MJ, Shaw KA, Baio J, EdS, Washington A, Patrick M, et al. Prevalence of Autism Spectrum Disorder Among Children Aged 8 Years - Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2016. *MMWR Surveill Summ.* 2020;69:1–12.

4. Bai D, Yip BHK, Windham GC, Sourander A, Francis R, Yoffe R, et al. Association of genetic and environmental factors with Autism in a 5-country cohort. *JAMA Psychiatry*. 2019;76:1035–43.
5. Croen LA, Zerbo O, Qian Y, Massolo ML, Rich S, Sidney S, et al. The health status of adults on the autism spectrum. *Autism*. 2015;19:814–23.
6. Soke GN, Maenner MJ, Christensen D, Kurzius-Spencer M, Schieve LA. Prevalence of co-occurring medical and behavioral conditions/symptoms among 4- and 8-year-old children with autism spectrum disorder in selected areas of the United States in 2010. *J Autism Dev Disord*. 2018;48:2663–76.
7. Houghton R, Ong RC, Bolognani F. Psychiatric comorbidities and use of psychotropic medications in people with autism spectrum disorder in the United States. *Autism Res*. 2017;10:2037–47.
8. Muskens JB, Velders FP, Staal WG. Medical comorbidities in children and adolescents with autism spectrum disorders and attention deficit hyperactivity disorders: a systematic review. *Eur Child Adolesc Psychiatry*. 2017;26:1093–103.
9. Hyman SL, Levy SE, Myers SM. Identification, evaluation, and management of children with Autism spectrum disorder. *Pediatrics*. 2020;145:e20193447.
10. Matson JL, Shoemaker M. Intellectual disability and its relationship to autism spectrum disorders. *Res Dev Disabil*. 2009;30:1107–14.
11. Brimberg L, Mader S, Fujieda Y, Arinuma Y, Kowal C, Volpe BT, et al. Antibodies as Mediators of Brain Pathology. *Trends Immunol*. 2015;36:709–24.
12. Diamond B, Huerta PT, Mina-Osorio P, Kowal C, Volpe BT. Losing your nerves? Maybe it's the antibodies. *Nat Rev Immunol*. 2009;9:449–56.
13. Edmiston E, Ashwood P, Van, de Water J. Autoimmunity, autoantibodies, and Autism spectrum disorder. *Biol Psychiatry*. 2017;81:383–90.
14. Estes ML, McAllister AK. Immune mediators in the brain and peripheral tissues in autism spectrum disorder. *Nat Rev Neurosci*. 2015;16:469–86.
15. Gata-Garcia A, Diamond B. Maternal antibody and ASD: clinical data and animal models. *Front Immunol*. 2019;10:1129–1129.
16. Hughes HK, Mills KoE, Rose D, Ashwood P. Immune dysfunction and autoimmunity as pathological mechanisms in Autism spectrum disorders. *Front Cell Neurosci*. 2018;12:405.
17. Meltzer A, Van de Water J. The role of the immune system in Autism spectrum disorder. *Neuropsychopharmacol: Off Publ Am Coll Neuropsychopharmacol*. 2017;42:284–98.
18. Allswede DM, Yolken RH, Buka SL, Cannon TD. Cytokine concentrations throughout pregnancy and risk for psychosis in adult offspring: a longitudinal case-control study. *Lancet Psychiatry*. 2020;7:254–61.
19. Cheslack-Postava K, Cremers S, Bao Y, Shen L, Schaefer CA, Brown AS. Maternal serum cytokine levels and risk of bipolar disorder. *Brain Behav Immun*. 2017;63:108–14.
20. Marks K, Vincent A, Coutinho E. Maternal-autoantibody-related (MAR) Autism: identifying neuronal antigens and approaching prospects for intervention. *J Clin Med*. 2020;9:2564.
21. Hertz-Picciotto I, Croen LA, Hansen R, Jones CR, Water JVD, Pessah IN. The CHARGE study: an epidemiologic investigation of genetic and environmental factors contributing to Autism. *Environ Health Perspect*. 2006;114:1119–25.
22. Braunschweig D, Krakowiak P, Duncanson P, Boyce R, Hansen RL, Ashwood P, et al. Autism-specific maternal autoantibodies recognize critical proteins in developing brain. *Transl Psychiatry*. 2013;3:e277.
23. Ramirez-Celis A, Becker M, Nuño M, Schauer J, Aghaepour N, Van de Water J. Risk assessment analysis for maternal autoantibody-related autism (MAR-ASD): a subtype of Autism. *Mol Psychiatry*. 2021;26:1551–60.
24. Lyall K, Ames JL, Pearl M, Traglia M, Weiss LA, Windham GC, et al. A profile and review of findings from the Early Markers for Autism study: unique contributions from a population-based case-control study in California. *Mol Autism*. 2021;12:24.
25. Heuer LS, Croen LA, Jones KL, Yoshida CK, Hansen RL, Yolken R, et al. An exploratory examination of neonatal cytokines and chemokines as predictors of Autism risk: the early markers for Autism study. *Biol Psychiatry*. 2019;86:255–64.
26. Jones KL, Croen LA, Yoshida CK, Heuer L, Hansen R, Zerbo O, et al. Autism with intellectual disability is associated with increased levels of maternal cytokines and chemokines during gestation. *Mol Psychiatry*. 2017;22:273–9.
27. Ramirez A, Edmiston E, Schauer J, Vu T, Van, de Water J. Peptides of neuron specific enolase as potential ASD biomarkers: from discovery to epitope mapping. *Brain Behav Immun*. 2020;84:200–8.
28. Soma-Pillay P, Nelson-Piercy C, Tolppanen H, Mebazaa A. Physiological changes in pregnancy. *Cardiovasc J Afr*. 2016;27:89–94.
29. Ander SE, Diamond MS, Coyne CB. Immune responses at the maternal-fetal interface. *Sci Immunol*. 2019;4:eaat6114.
30. Abu-Raya B, Michalski C, Sadarangani M, Lavoie PM. Maternal immunological adaptation during normal pregnancy. *Front Immunol*. 2020;11:575197.
31. Palmeira P, Quinello C, Silveira-Lessa AL, Zago CA, Carneiro-Sampaio M. IgG Placental Transfer in Healthy and Pathological Pregnancies. *Clin Developmental Immunol*. 2012;2012:985646.
32. Mozaffarian N, Shaw EA, Stevens AM. Chapter 10 - Maternally Mediated Neonatal Autoimmunity. In: Ohls RK, Maheshwari A (eds). *Hematology, Immunology and Infectious Disease: Neonatology Questions and Controversies (Second Edition)*. W.B. Saunders: Philadelphia, 2012, pp 129–70.
33. Ross G, Sammaritano L, Nass R, Lockshin M. Effects of mothers' autoimmune disease during pregnancy on learning disabilities and hand preference in their children. *Arch Pediatr Adolesc Med*. 2003;157:397–402.
34. Yousef Yengej FA, van Royen-Kerkhof A, Derksen RHWM, Fritsch-Stork RDE. The development of offspring from mothers with systemic lupus erythematosus. A systematic review. *Autoimmun Rev*. 2017;16:701–11.
35. Coutinho E, Jacobson L, Pedersen MG, Benros ME, Nørgaard-Pedersen B, Mortensen PB, et al. CASPR2 autoantibodies are raised during pregnancy in mothers of children with mental retardation and disorders of psychological development but not autism. *J Neurol Neurosurg Psychiatry*. 2017;88:718.
36. Ghassabian A, Bongers-Schokking JJ, de Rijke YB, van Mil N, Jaddoe VW, de Muinck Keizer-Schrama SM, et al. Maternal thyroid autoimmunity during pregnancy and the risk of attention deficit/hyperactivity problems in children: the Generation R Study. *Thyroid*. 2012;22:178–86.
37. Coutinho E, Menassa D, Jacobson L, West S, Domingos J, Moloney T, et al. Maternal CASPR2 antibodies and neurodevelopmental disorders in the offspring: epidemiological findings and an animal model. *Lancet*. 2017;389:518.
38. Akum BF, Chen M, Gunderson SI, Riefler GM, Scerri-Hansen MM, Firestein BL. Cypin regulates dendrite patterning in hippocampal neurons by promoting microtubule assembly. *Nat Neurosci*. 2004;7:145–52.
39. Di Liberto V, Mudò G, Garozzo R, Frinchi M, Fernandez-Dueñas V, Di Iorio P et al. The guanine-based purinergic system: the tale of an orphan neuromodulation. *Front Pharmacol*. 2016;7:158.
40. Hou S-T. The regulatory and enzymatic functions of CRMPs in neurite outgrowth, synaptic plasticity, and gene transcription. *Neurochemistry Int*. 2020;139:104795.
41. Quach TT, Honnorat J, Kolattukudy PE, Khanna R, Duchemin AM. CRMPs: critical molecules for neurite morphogenesis and neuropsychiatric diseases. *Mol Psychiatry*. 2015;20:1037–45.
42. Zuccarini M, Giuliani P, Frinchi M, Mudò G, Serio RM, Belluardo N, et al. Uncovering the signaling pathway behind extracellular guanine-induced activation of NO system: new perspectives in memory-related disorders. *Front Pharm*. 2018;9:110.
43. Lopes MH, Hajj GN, Muras AG, Mancini GL, Castro RM, Ribeiro KC, et al. Interaction of cellular prion and stress-inducible protein 1 promotes neurogenesis and neuroprotection by distinct signaling pathways. *J Neurosci*. 2005;25:11330–9.
44. Hashimoto-Torii K, Torii M, Fujimoto M, Nakai A, El Fatimy R, Mezger V, et al. Roles of heat shock factor 1 in neuronal response to fetal environmental risks and its relevance to brain disorders. *Neuron*. 2014;82:560–72.
45. De Rubeis S, He X, Goldberg AP, Poultney CS, Samocha K, Cicek AE, et al. Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature*. 2014;515:209–15.
46. Takata A, Miyake N, Tsurusaki Y, Fukai R, Miyatake S, Koshimizu E, et al. Integrative analyses of de novo mutations provide deeper biological insights into Autism spectrum disorder. *Cell Rep*. 2018;22:734–47.
47. Liu Y, Pham X, Zhang L, Chen PL, Burzynski G, McGaughey DM, et al. Functional variants in DPYSL2 sequence increase risk of schizophrenia and suggest a link to mTOR signaling. *G3 (Bethesda)*. 2014;5:61–72.
48. Lee H, Joo J, Nah SS, Kim JW, Kim HK, Kwon JT, et al. Changes in Dpysl2 expression are associated with prenatally stressed rat offspring and susceptibility to schizophrenia in humans. *Int J Mol Med*. 2015;35:1574–86.
49. Beraldo FH, Soares IN, Goncalves DF, Fan J, Thomas AA, Santos TG, et al. Stress-inducible phosphoprotein 1 has unique chaperone activity during development and regulates cellular response to ischemia via the prion protein. *Faseb J*. 2013;27:3594–607.
50. Beraldo FH, Thomas A, Kolisnyk B, Hirata PH, De Jaeger X, Martyn AC, et al. Hyperactivity and attention deficits in mice with decreased levels of stress-inducible phosphoprotein 1 (STIP1). *Dis Model Mech*. 2015;8:1457–66.
51. Jeanne M, Demory H, Moutal A, Vuillaume M-L, Blesson S, Thépault R-A, et al. Missense variants in DPYSL5 cause a neurodevelopmental disorder with corpus callosum agenesis and cerebellar abnormalities. *Am J Hum Genet*. 2021;108:951–61.
52. Zhang H, Kang E, Wang Y, Yang C, Yu H, Wang Q, et al. Brain-specific Crmp2 deletion leads to neuronal development deficits and behavioural impairments in mice. *Nat Commun*. 2016;7:11773.
53. Saint-Martin M, Joubert B, Pellier-Monnin V, Pascual O, Noraz N, Honnorat J. Contactin-associated protein-like 2, a protein of the neurexin family involved in several human diseases. *Eur J Neurosci*. 2018;48:1906–23.
54. Varea O, Martin-de-Saavedra MD, Kopeikina KJ, Schürmann B, Fleming HJ, Fawcett-Patel JM, et al. Synaptic abnormalities and cytoplasmic glutamate receptor aggregates in contactin associated protein-like 2/Caspr2 knockout neurons. *Proc Natl Acad Sci*. 2015;112:6176–81.
55. Coutinho E, Menassa DA, Jacobson L, West SJ, Domingos J, Moloney TC, et al. Persistent microglial activation and synaptic loss with behavioral abnormalities in



- mouse offspring exposed to CASPR2-antibodies in utero. *Acta Neuropathol.* 2017;134:567–83.
56. van 't Hof M, Tisseur C, van Berckeleer-Onnes I, van Nieuwenhuyzen A, Daniels AM, Deen M, et al. Age at autism spectrum disorder diagnosis: A systematic review and meta-analysis from 2012 to 2019. *Autism.* 2021;25:862–73.
57. La Malfa G, Lassi S, Bertelli M, Salvini R, Placidi GF. Autism and intellectual disability: a study of prevalence on a sample of the Italian population. *J Intellect Disabil Res.* 2004;48(Pt 3):262–7.

## ACKNOWLEDGEMENTS

We would like to thank the families that participated in the EMA study, and the staff at the Kaiser Permanente- Research. Special thanks to Lori Haapanen for experimental and technical support. This study was funded by the NIEHS Center for Children's Environmental Health and Environmental Protection Agency (EPA) grants (2P01ES011269-11, 83543201, respectively), the NIEHS-funded EMA study (R01ES016669), the NICHD funded IDDRC P50 (P50HD103526), and Consejo Nacional de Ciencia y Tecnologia (CONACYT- UC MEXUS) Doctoral Fellowships.

## AUTHOR CONTRIBUTIONS

LAC and JWV: conceptualization of research and experimental design. ARC, CKY, and SEA: Data analysis and experimental design. JS: Maternal sample testing, data generation. RHY and PA: Expertise input on data analysis and translation to results. ARC, LAC, and JWV created the first manuscript draft, which was edited and approved by all the authors.

## COMPETING INTERESTS

JVdeW has patents issued for this technology and has founded a UC Davis startup company to develop this technology for commercial use. The remaining authors have no conflicts of interest.

## ADDITIONAL INFORMATION

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41380-022-01633-4>.

**Correspondence** and requests for materials should be addressed to Judy Van de Water.

**Reprints and permission information** is available at <http://www.nature.com/reprints>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022