

# Lawrence Berkeley National Laboratory

## Recent Work

**Title**

AN ERYTHRON-DEPENDENT MODEL OF IRON KINETICS

**Permalink**

<https://escholarship.org/uc/item/3sd211mj>

**Author**

Nooney, Grove C.

**Publication Date**

1965-09-15

**University of California**  
**Ernest O. Lawrence**  
**Radiation Laboratory**

AN ERYTHRON-DEPENDENT MODEL OF IRON KINETICS

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

**Berkeley, California**

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to the Biophysical Journal

UCRL-16406 Rev.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory  
Berkeley, California

AEC Contract No. W-7405-eng-48

AN ERYTHRON-DEPENDENT MODEL OF IRON KINETICS

Grove C. Nooney

April 21, 1966

AN ERYTHRON-DEPENDENT MODEL OF IRON KINETICS\*

Grove C. Nooney

Lawrence Radiation Laboratory  
University of California  
Berkeley, California

April 21, 1966

ABSTRACT

The mathematical model here presented of steady-state iron kinetics in the human explicitly displays the influence of the maturative and proliferative behavior of erythrons (erythrocytes or their precursory cells). Based on a compartmented iron kinetics scheme, the model is in the form of a system of linear integrodifferential equations and describes the distribution of administered radioactive tracer as a function of time. Erythron behavior is characterized by three functions specifying iron uptake, proliferation, and survival. Simulative uses of the model are discussed and illustrated by an example.

## INTRODUCTION

This paper presents a mathematical formulation of steady-state iron kinetics of the human blood and blood-forming system. The formulation or model describes the distribution in the body of administered tracer iron ( $\text{Fe}^{59}$ ) so as to make evident the influence of maturation and proliferation of erythrocytes and their precursors, the erythropoietic cells. For both types of cells we use the collective term erythron.

A central role in iron metabolism and iron kinetics is played by these erythrons, but earlier mathematical models of iron kinetics ignore their detailed behavior (Huff and Judd, 1956; Pollycove and Mortimer, 1961; Sharney et al., 1963). A more recent paper (Garby et al., 1963) provides for very restricted consideration of the erythron. My model described here explicitly reflects the essential details of erythron maturative and proliferative behavior. The result is not only a description of iron kinetics that is more accurate than previous ones, but is a unified description as well, one which offers the opportunity for investigation of the behavior of erythron itself. We have presented a special case of the new model, chosen and modified for use with certain radioiron tracer procedures (Nooney, 1965, referred to in the following as IK), and we have indicated the use of simulative computations in connection with both the special case and the general model; the general model and its specializations are parts of a more extensive theory of age-dependent cellular processes, also applicable, for instance, to neutrophil kinetics (Nooney, 1964).

In the paper presented here, we establish the general iron-kinetics model as a system of integrodifferential equations based on a compart-

mented iron-kinetics system. This model is compatible with all observed erythropoietic and hemolytic mechanisms, as well as with most plausible conjectured mechanisms. The generality, accuracy of representation, and novelty of the model are provided by allowing the erythron to influence intercompartmental transfers explicitly and naturally.

We discuss application of the model and illustrate its use in an example.

### DEFINITIONS AND ASSUMPTIONS

This section departs but slightly from the corresponding parts of IK, where more detail is provided. In accordance with previous models (especially that of Pollycove and Mortimer, 1961), we assign iron in the blood system to six mutually exclusive compartments. For convenience in reference, we shall use the term erythrocyte for both circulating reticulocytes and erythrocytes, reserving the term erythropoietic cells for the erythropoietic cells of the marrow. The six compartments are (i) iron in circulating erythrocytes, (ii) iron in plasma, (iii) iron in non-circulating erythropoietic cells, (iv) bone-marrow iron, (v) iron in storage as ferritin, and (vi) iron in storage as hemosiderin. The bone-marrow iron, which constitutes the labile marrow pool introduced by Pollycove and Mortimer (1956), may include iron compounds on the exterior membranes of erythropoietic cells or the ferritin of certain observed reticular cells (Bessis and Breton-Gorius, 1957). The ferritin and hemosiderin storage may be considered as short- and long-term storage, or as labile and reserve storage, respectively, according to evidence presented by Pollycove and Mortimer (1961). The definition of these six compartments may be made unique by defining them maximally and in the order given. We

shall ignore other iron in the body, regarding the six compartments as being a closed system with regard to tracer, except for the introduction of tracer.

The following transfers between compartments are permitted. As we shall see, some of these transfers may be omitted, according to the demands of individual physiology or prejudice. Erythrons may enter the circulation with their iron. Upon the destruction of an erythrocyte or of an erythropoietic cell, its iron may go into ferritin storage, plasma, or bone marrow. Iron in bone marrow may go into plasma or erythropoietic cells; iron in plasma may go into ferritin storage, erythrocytes, or bone marrow. Iron in ferritin storage may go into plasma or hemosiderin storage; iron in hemosiderin storage may go only into ferritin storage. Our system has the form shown in Fig. 1, where the arrows denote directed transfers. The number below each compartment will be used to identify that compartment.

We shall assume the system to be in a steady state, by which is meant that the amount of iron in each compartment and the rate of each transfer are constant. Since we later introduce a probability distribution, a more precise and correct statement of the steady-state assumption is that the expected amount of iron in each compartment and the expected rate of each transfer remain constant. Note that this does not exclude certain disease states. The requirements for and the implications of the steady-state assumption have been discussed elsewhere (Sharney et al., 1963, and Sheppard, 1962).

Let us measure the age of an erythron from the time its earliest progenitor began iron accumulation, and let us assume for each steady state that the pertinent erythron behavior depends on age alone. Define



$P(s)$  to be the probability of survival to age  $s$  of an erythron and its descendents along a particular branch of their genealogical tree. Thus, if an erythron has divided only once before age  $s$ ,  $P(s)$  is the probability of survival to age  $s$  of each daughter. Define  $q(s)$  to be the number of erythrons of age  $s$  if no erythron death were to occur; define  $r(s)$  to be the rate of iron acquisition by an erythron of age  $s$ . We allow  $r$  to assume negative values to account for conceivable iron loss, but  $r$  is not to account for loss by mitosis, in which iron content is approximately halved. Thus,  $r$  describes the iron uptake of each erythron except at the instant of division. We set  $R(s) = q(s) r(s)$ . We further assume that erythrons pass into the circulation at the fixed age  $T$ .

Suppose administration of  $\text{Fe}^{59}$  into plasma is begun at time  $t = 0$ . We assume injection of isotope to occur instantaneously at  $t = 0$ , but our considerations are not restricted to this mode of isotope administration. Of course, the underlying steady state of the system is assumed undisturbed by the addition of isotope. Denote by  $I_j$  the constant amount of iron in compartment  $j$ . Denote by  $x_i(t)$  the amount of radioactive iron in compartment  $i$  at time  $t$ , by  $\dot{x}_i(t)$  the derivative of  $x_i(t)$  with respect to  $t$ , and by  $\dot{x}_{i,j}(t)$  that portion of  $\dot{x}_i(t)$  due to transfer from compartment  $i$  to compartment  $j$ . Some of these symbols will be used also to denote expected values of the corresponding quantities with respect to the probability function  $P$ . We assume six of the transfers to be of first order on the basis of chemical and tracer kinetics. That is, we assume the existence of constants  $k_{ij}$  such that

$$\dot{x}_{1,2}(t) = k_{12} x_1(t),$$

$$\dot{x}_{2,1}(t) = k_{21} x_2(t),$$

$$\dot{x}_{1,5}(t) = k_{15} x_1(t),$$

$$\dot{x}_{5,1}(t) = k_{51} x_5(t),$$

$$\dot{x}_{5,6}(t) = k_{56} x_5(t),$$

$$\dot{x}_{6,5}(t) = k_{65} x_6(t).$$

Note that these transfers are physiologically independent of erythron behavior.

A prime requirement for the validity of the assumed first-order relations is the homogeneity of tracer iron relative to total iron in each compartment appearing on the right sides of those relations. This requirement is probably satisfied for compartments 1 and 2, but its satisfaction may be doubtful for the storage compartments 5 and 6. Thus the foregoing first-order transfers from storage compartments must be regarded only as approximations to the true form of those transfers. More precise formulations of these transfers must take into account the specific mechanisms of storage and retrieval and will depend on the history as well as the amount of stored material. Since our present concern is with the influence of the details of erythron behavior on iron kinetics, we shall postpone consideration of storage details.

Some further comments are necessary relative to the ferritin and hemosiderin compartments and their transfers. We shall treat those compartments as labile and reserve storage, respectively, although such correspondence has not been rigorously established. It is natural then to require that reserve iron pass through the labile stage before further participation in the system. That hemosiderin is physiologically associated with ferritin and the iron of ferritin is more readily available for the synthesis of hemoglobin (Bell et al., 1961) supports such treatment. Our use of the ferritin-hemosiderin nomenclature is an attempt to specify the composition of the storage compartments, which otherwise are identified only functionally. The considerations that follow are not influenced by that nomenclature.

## DERIVATION OF THE EQUATIONS

The steady-state condition implies that there is transferred to each compartment a constant fraction of the iron made available through the death of erythrons. Thus, there are constants  $\alpha_i$  and  $\beta_i$ , all between zero and unity, such that

$$\begin{aligned}\dot{x}_{3,i}(t) &= \alpha_i [\dot{x}_{3,1}(t) + \dot{x}_{3,2}(t) + \dot{x}_{3,5}(t)], \\ \dot{x}_{4,i}(t) &= \beta_i [\dot{x}_{4,1}(t) + \dot{x}_{4,2}(t) + \dot{x}_{4,5}(t)], \quad \text{for } i=1, 2, 5.\end{aligned}$$

It follows, of course, that  $\alpha_1 + \alpha_2 + \alpha_5 = \beta_1 + \beta_2 + \beta_5 = 1$ .

At any time, the rate of transfer of iron from compartment 2 to erythropoietic cells in compartment 3 is

$$k_{23}(T) = \int_{s=0}^T P(s) R(s) ds.$$

Therefore

$$\dot{x}_{2,3}(t) = \frac{k_{23}(T)}{I_2} x_2(t).$$

Similarly,

$$\dot{x}_{1,4}(t) = \frac{k_{14}(T)}{I_1} x_1(t),$$

where

$$k_{14}(T) = \int_{s=T}^{\infty} P(s) R(s) ds.$$

An erythropoietic cell in compartment 3 reaching age  $u$  at time  $t$  was age  $s$  at time  $t - u + s$ . Therefore, if no death were to occur, the amount of radioactive iron in all such cells of age  $u$  at time  $t$  would be

$$C_2(u, t) = \frac{1}{I_2} \int_{s=0}^u R(s) x_2(t - u + s) ds.$$

Taking into account the probability of death, we write

$$\dot{x}_{3,4}(t) = P(T) C_2(T, t),$$

$$\dot{x}_{3,i}(t) = -a_i \int_{u=0}^T C_2(u, t) dP(u), \quad \text{for } i = 1, 2, 5.$$

An erythrocyte of age  $u$  at time  $t$  entered the circulation at time  $t - u + T$  and was age  $s$  at time  $t - u + s$ . Therefore, if no death of erythrocytes were to occur, the amount of radioactive iron in all erythrocytes of age  $u$  at time  $t$  would be  $C_2(T, t - u + T) + C_1(u, t)$ , where the term

$$C_1(u, t) = \frac{1}{I_1} \int_{s=T}^u R(s) x_1(t - u + s) ds$$

represents radioactive-iron acquisition in the circulation. Again, taking into account the probability of death, we write

$$\dot{x}_{4,i}(t) = -\beta_i \int_{u=T}^{\infty} [C_2(t, t - u + T) + C_1(u, t)] dP(u), \quad \text{for } i = 1, 2, 5.$$

We have now furnished expressions for all nonzero transfer components.

For initial injection of  $\text{Fe}^{59}$  into plasma, the sought system of integrodifferential equations is

$$\dot{x}_i(t) = \sum_j \dot{x}_{j,i}(t) - \sum_j \dot{x}_{i,j}(t), \quad \text{for } i, j = 1, 2, \dots, 6. \quad (1)$$

Due to the conservation of isotope, these equations contain only five independent  $x_i$ . For example, the first of these equations is

$$\begin{aligned}
\dot{x}_1(t) = & -(k_{12} + k_{15} + k_{14}(T)/I_1)x_1(t) + k_{21}x_2(t) + k_{51}x_5(t), \\
& - \frac{\alpha_1}{I_2} \int_{u=0}^T \int_{s=0}^u R(s) x_2(t-u+s) ds dP(u) \\
& - \frac{\beta_1}{I_2} \int_{u=T}^{\infty} \int_{s=0}^T R(s) x_2(t-u+s) ds dP(u) \\
& - \frac{\beta_1}{I_1} \int_{u=T}^{\infty} \int_{s=T}^u R(s) x_1(t-u+s) ds dP(u).
\end{aligned} \tag{2}$$

Since these equations are linear in the  $x_i$ , we may assume the normalized initial conditions  $x_1(0) = 1$ ,  $x_i(0) = 0$ , where  $i = 2, 3, \dots, 6$ .

If another mode of isotope introduction is used (e. g., infusion), with the rate of introduction given by  $f(t)$ , then we must add  $f(t)$  to the right-hand side of the equation for  $\dot{x}_1$  and change the corresponding initial condition to  $x_1(0) = 0$ .

#### DISCUSSION

Equations (1) uniquely define functions  $x_i$  if the rate parameters and erythron-behavior functions are given and physiologically acceptable (finite erythrocyte life span and piece wise continuous  $R$  and  $P$ , for instance). The proof may be made as a slight extension of the classical Picard-Lindelöf proof for ordinary differential equations; that proof shows that the solution of equations (1) may be obtained by successive approximations or by stepwise integration (Bieberbach, 1926). Thus, these equations may be used in the simulating of iron kinetics or, what is perhaps more important, in the testing of conjectures about erythron behavior. The conjectures must specify the erythron-behavior functions used in the model, but some parameters might be determined in the usual way, by fitting the model to experimental data. In a previous paper (IK) we have described a method for the calculation of the erythron-behavior functions

from such data. The present model verifies the calculation by allowing reproduction of the data if the calculated erythron behavior functions are correct.

Extensions of the model are immediately obvious. For instance, acquisition of iron from plasma by noncirculating erythrons could be accommodated by defining an extra iron-uptake function. The restriction to a fixed erythron-maturation time could be relaxed by introducing a stochastic maturation time; this would require replacement of all expressions involving  $T$  by their expected values, and would require also, as a referee has observed, the extension of our steady-state assumption to imply steady expected state of the system with respect to this additional stochastic variable. Such extensions complicate, but do not prohibit, calculation and simulation with the model.

For those who, with the author, are unsatisfied with storage participation in this model, an alternative approach is available. The function  $x_1$  has been measured for many subjects with various hematological disorders (e. g., Pollycove and Mortimer, 1956, 1961; Sharney, et al., 1963; Gevirtz et al., 1965). If  $x_1$  is taken as known from these measurements, then a review of equations (1) reveals that the explicit appearance of  $x_5$  and  $x_6$  may be suppressed and that attention may be confined to  $x_2$ ,  $x_3$ ,  $x_4$  and the known  $x_1$ . Such a procedure reduces the dimensions of the problem from 5 to 3.

#### EXAMPLE

Equations (1) have been implemented by a general and efficient computing machine program, SBLOOD, written by Thomas Mahan (1966) of the Lawrence Radiation Laboratory. The results given in the following

example were obtained with that program.

To illustrate application of the model, we have used equations (1) to calculate some effects of differences in severity and timing of intramedullary hemolysis. For this purpose we chose a system with  $k_{15} = 0$ ,  $\alpha_1 = 1$ , and for  $s > T$ ,  $R(s) = 0$  and  $P(s) = \text{constant}$ , so that storage plays no role in isotope distribution, all isotope released by intramedullary hemolysis is transferred to plasma, and erythrocytes neither acquire isotope nor release it by death in the time period considered. We set  $T = 5$  (days), a plausible maturation period (Harris, 1963), and specified  $k_{12} = 9.21$ ,  $k_{21} = 0.131$  ( $\text{days}^{-1}$ ), and  $I_2 = 84.9$  (mg). The last three values, apparently hematologically acceptable, are mean values for 13 normal subjects as obtained by Pollycove and Mortimer (1961). The mean daily delivery of 21.4 mg of iron to compartment 4 was taken from the same source. The iron uptake rate  $R$  was taken as linearly decreasing to zero at 5 days and was adjusted for each hemolytic situation to yield that same daily delivery. The four cases considered were then distinguished by the following hemolytic behavior: a 10 percent (case I) and a 50 percent (case II) hemolysis of erythropoietic cells midway in their maturation period, and a 10 percent (case III) and a 50 percent (case IV) hemolysis of erythropoietic cells at the end of their maturation period. More precisely,  $P(s)$  was specified as continuous and composed of three linear segments for each case:

$$\begin{array}{l}
 \text{Case I.} \quad P(s) = \begin{cases} 1 & s < 2.5 \\ 0.9 & s > 2.6 \end{cases} \\
 \text{Case II.} \quad P(s) = \begin{cases} 1 & s < 2.5 \\ 0.5 & s > 2.6 \end{cases} \\
 \text{Case III.} \quad P(s) = \begin{cases} 1 & s < 4.9 \\ 0.9 & s > 5 \end{cases} \\
 \text{Case IV.} \quad P(s) = \begin{cases} 1 & s < 4.9 \\ 0.5 & s > 5 \end{cases}
 \end{array}$$

It should be emphasized that none of the values used is known to be physiologically correct. Indeed, some were obtained with models that we have implied are faulty. We use these values only in an illustrative sense.

The calculated proportion in plasma of injected isotope as a function of time after injection is displayed for each case in Fig. 2 or Fig. 3. The most striking feature of the curves shown is the obvious departure from convexity (the appearance of bumps) in Fig. 3. Gevirtz and co-workers (1965) showed that such aberrations occur in the plasma curves of normal subjects; they attributed this to ineffective erythropoiesis and hemolysis. Our simple example suggests that if this phenomenon is due to the hemolysis of erythropoietic cells, then that hemolysis probably occurs late in the maturation period. A comparison of Fig. 3 with the data of Gevirtz further suggests that such hemolysis results in the death of 10 to 50 percent of the erythropoietic cells that would otherwise reach maturation.

The author acknowledges the aid of H. S. Winchell for providing hematological background.



REFERENCES

- \*Work done under the auspices of the U. S. Atomic Energy Commission.
- Bell, George H., Davidson, J. Norman, and Scarborough, Harold, 1961. Textbook of Physiology and Biochemistry, Baltimore, The Williams and Wilkins Co.
- Bessis, M. and Breton-Gorius, J., 1957. Étude au microscope électronique des granulations ferrugineuses des érythrocytes normaux et pathologiques. Anémies hémolytiques, Héminoglobinoopathies Saturnisme, Rev. Hémat. 12, 43.
- Bieberbach, Ludwig, 1926, Theorie der Differentialgleichungen, Berlin, Julius Springer.
- Garby, L., Schneider, W., Sundquist, O., and Vuille, J., 1963. A ferro-erythrokinetic model and its properties, Acta Physiol. Scand. 59, Suppl. 216.
- Gevirtz, N. R., Wasserman, L. R., Sharney, L., and Tandler, D., 1965. Studies of plasma Fe<sup>59</sup> disappearance—a manifestation of ineffective erythropoiesis and hemolysis. Blood 25, 976.
- Harris, J. W., 1963. The Red Cell. Cambridge, Harvard University Press.
- Huff, R. L., and Judd, O. J., 1956, Kinetics of iron metabolism, in Advances in Biological and Medical Physics (J. H. Lawrence and C. A. Tobias, editors), New York, Academic Press, vol. 4, 223.
- Mahan, T., 1966. SBLOOD, a CHIPPEWA FORTRAN program for CDC 6600 (unpublished material).
- Nooney, Grove C., 1964, Models of blood cell proliferation, Fifth IBM Scientific Computing Symposium, Yorktown Heights, N. Y.

Nooney, Grove C., 1965, Iron kinetics and erythron development,

Biophys. J. 5, 755.

Pollycove, M. and Mortimer, R., 1956. The quantitative determination

of individual iron kinetics and hemoglobin formation in human subjects. Clin. Res. Proc. 4, 51.

Pollycove, M., and Mortimer, R., 1961. The quantitative determination

of iron kinetics and hemoglobin synthesis in human subjects, J. Clin. Invest. 40, 753.

Sharney, L., Wasserman, L. R., Schwartz, L., and Tendler, D., 1963,

Multiple pool analysis as applied to erythrokinetics, Ann. N. Y. Acad. Sci. 108, 230.

Sheppard, C. W., 1962, Basic Principles of the Tracer Method, New


York, John Wiley & Sons.

FIGURE CAPTIONS


Fig. 1. Iron compartmentation and transfers.

Fig. 2. Proportion of isotope in plasma; hemolysis of 10% and 50% at erythron age 2.5 days.


Fig. 3. Proportion of isotope in plasma; hemolysis of 10% and 50% at erythron age 4.9 days.



MUB-8261



MUB-10530



MUB-10531

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

