

UC Davis

UC Davis Previously Published Works

Title

Association Between COVID-19 and Planned and Postponed Cancer Screenings Among American Indian Adults Residing in California and Oklahoma, March-December 2020.

Permalink

<https://escholarship.org/uc/item/3z06s7hm>

Authors

Dang, Julie HT

Chen, Sixia

Hall, Spencer

et al.

Publication Date

2024-06-04

DOI

10.1177/00333549241254226

Peer reviewed

Association Between COVID-19 and Planned and Postponed Cancer Screenings Among American Indian Adults Residing in California and Oklahoma, March–December 2020

Public Health Reports

1–10

© 2024, Association of Schools and
Programs of Public Health

All rights reserved.

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/00333549241254226

journals.sagepub.com/home/phr

Julie H.T. Dang, PhD¹ ; Sixia Chen, PhD²; Spencer Hall, MPH²;
Janis E. Campbell, PhD²; Moon S. Chen Jr, PhD³;
and Mark P. Doescher, MD⁴

Abstract

Objective: Little is known about how the COVID-19 pandemic affected cancer screenings among American Indian people residing in California and Oklahoma, 2 states with the largest American Indian populations. We assessed rates and factors associated with cancer screenings among American Indian adults during the pandemic.

Methods: From October 2020 through January 2021, we surveyed 767 American Indian adults residing in California and Oklahoma. We asked participants whether they had planned to obtain screenings for breast cancer, cervical cancer, and colorectal cancer (CRC) from March through December 2020 and whether screening was postponed because of COVID-19. We calculated adjusted odds ratios (AORs) for factors associated with reasons for planned and postponed cancer screening.

Results: Among 395 participants eligible for breast cancer screening, 234 (59.2%) planned to obtain the screening, 127 (54.3%) of whom postponed it. Among 517 participants eligible for cervical cancer screening, 357 (69.1%) planned to obtain the screening, 115 (32.2%) of whom postponed it. Among 454 participants eligible for CRC screening, 282 (62.1%) planned to obtain CRC screening, 80 of whom (28.4%) postponed it. In multivariate analyses, women who lived with a child (vs did not) had lower odds of planning to obtain a breast cancer screening (AOR=0.6; 95% CI, 0.3-1.0). Adherence to social distancing recommendations was associated with planning to have and postponement of cervical cancer screening (AOR=7.3; 95% CI, 0.9-58.9). Participants who received (vs did not receive) social or financial support had higher odds of planning to have CRC screening (AOR=2.0; 95% CI, 1.1-3.9).

Conclusion: The COVID-19 pandemic impeded completion of cancer screenings among American Indian adults. Interventions are needed to increase the intent to receive evidence-based cancer screenings among eligible American Indian adults.

Keywords

American Indian, COVID-19, cancer screening

¹ Department of Public Health Sciences, Division of Health Policy and Management, University of California, Davis School of Medicine, Sacramento, CA, USA

² Department of Biostatistics and Epidemiology, Hudson College of Public Health, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA

³ Department of Internal Medicine, Division of Hematology and Oncology, University of California, Davis School of Medicine, Sacramento, CA, USA

⁴ Stephenson Cancer Center, College of Medicine, Department of Family and Preventive Medicine, University of Oklahoma College of Health Science Center, Oklahoma City, OK, USA

Corresponding Author:

Julie H.T. Dang, PhD, University of California, Davis School of Medicine, Department of Public Health Sciences, Division of Health Policy and Management, 4501 X St, Ste 3003, Sacramento, CA 95817, USA.

Email: jtdang@ucdavis.edu

During the COVID-19 pandemic, federal agencies and professional medical societies recommended postponement of preventive care visits to prioritize health care resources.¹⁻³ In the United States, approximately 9.4 million screenings for breast, colorectal, and prostate cancer did not occur because of the COVID-19 pandemic.⁴ Timely cancer screening is essential to primary and secondary prevention of cancer.^{5,6}

Although rates of cancer screening and cancer diagnosis have improved since the initial phase of the COVID-19 pandemic, models predicted increased rates of advanced-stage cancer at the time of diagnosis and increased rates of morbidity and mortality because of decreased and delayed cancer screening in 2020.⁷⁻⁹ Delayed cancer screenings can exacerbate disparities among American Indian and Alaska Native (AI/AN) people, who have significantly higher rates of cancer incidence and mortality compared with non-Hispanic White people.¹⁰⁻¹⁴ Depending on geographic region and source, AI/AN people are more likely than non-Hispanic White people to acquire and die from cancers for which screenings exist.¹⁰⁻¹⁶ Studies documented sharp declines in the volume of screening for breast cancer, cervical cancer, and colorectal cancer (CRC) among AI/AN people compared with non-Hispanic White people during the COVID-19 pandemic.¹⁷⁻¹⁹

In this study, we used survey data to examine rates of screening for breast cancer, cervical cancer, and CRC during the COVID-19 pandemic among AI/AN adults residing in California and Oklahoma, 2 states with the largest AI/AN populations.²⁰ We aimed to assess factors associated with plans among AI/AN adults to obtain screening for breast cancer, cervical cancer, and CRC during the COVID-19 pandemic and whether those plans were postponed because of the pandemic. We also analyzed rates of planned and postponed cancer screenings among survey participants. The term American Indian (AI) rather than AI/AN is used throughout the remainder of this article in reference to our sample population, because few Alaska Native people reside in California and Oklahoma.

Methods

Sample Selection and Data Collection

We used survey responses from a subset of survey questions developed by the National Cancer Institute–funded Impact on COVID-19 on the Cancer Continuum Consortium (IC-4) to assess the effects of COVID-19 on cancer prevention, control, and survivorship. The University of California, Davis (UCD) Comprehensive Cancer Center, and the University of Oklahoma Stephenson Cancer Center (OUSC) administered the survey to 1068 AI adults. From this sample, we selected 3 distinct cohorts of participants to assess planned and postponed cancer screenings that included mammograms for breast cancer screening (women aged 40–74 y), Papanicolaou (Pap) tests for cervical cancer screening (women aged 21–65 y), and colonoscopy and stool blood tests (men and women

aged 50–75 y) for CRC screening. Respondents answered whether they had planned to have a specific cancer screening test conducted from March through December 2020 (yes or no); if respondents answered yes to that question, the follow-up question was whether they or their doctor postponed the test because of COVID-19 (yes or no). UCD and OUSC administered the survey from October 2020 through January 2021.

Covariates

Although the complete survey consisted of 58 core questions, this study reported on the 30 questions related to sociodemographic characteristics, COVID-19–related attitudes and behaviors, and cancer screening behaviors. The survey questions included sociodemographic variables found to be predictors of cancer screenings among racially and ethnically diverse populations.²¹⁻²³ The survey questions on health behaviors related to COVID-19 were based on the Health Belief Framework.²⁴ The Health Belief Framework posits that individuals are more or less likely to engage in preventive behaviors based on their perceived susceptibility to and severity of the sickness or disease. The COVID-19 questions aimed to assess whether individuals engaged in preventive measures to reduce their risk of getting COVID-19.²⁵

Sociodemographic Variables

On the survey, respondents indicated their age group (18–39, 40–59, or ≥ 60 y), sex (female or male), Hispanic ethnicity (yes or no), educational attainment (high school diploma/General Educational Development or \geq some college), marital status (single/never married, married/living together, or separated/divorced/widowed), lived in a house with a child (eg, a person aged < 18 y; yes or no), annual household income ($< \$35\,000$ or $\geq \$35\,000$), covered by private or public health insurance or another kind of health plan (Indian Health Service; yes or no), had a preexisting condition (the survey provided a list of conditions and a fill-in-the-blank option), general health status (excellent or good, fair or poor), and employment status before COVID-19 (full-time/part-time employment or unemployed/other occupation). The other occupational categories included students, retired people, homemakers, and people with disabilities.

COVID-19 Variables

The survey presented a list of social distancing activities, and respondents answered whether they performed the following activities all or most of the time (yes or no): staying at home except for going to work, outdoors to exercise, to the grocery store, to the pharmacy, or to get medical care; not having anyone come into their home; staying 6 feet away from people when they leave the house; wearing a face mask when outdoors; and wearing a face mask when inside a place besides

their home. Respondents also answered questions on the perceived importance of social distancing recommendations (very important to somewhat important or a little important to not important) and whether they had ever been tested for COVID-19 (yes or no); had been in close physical contact with a person with a positive COVID-19 test in the past 30 days (yes or no); had attended any gatherings, rallies, demonstrations, or other social gatherings with >2 people outside their household (yes or no); and had received support (eg, emotional, materials, financial) during the pandemic from family or friends.

Statistical Analysis

We expressed descriptive statistics as counts and percentages. We calculated the proportion of survey respondents who were eligible (according to age and sex) for each cancer screening test. We then calculated the proportion of respondents who had planned to obtain the cancer screening and who had planned to obtain the cancer screening but reported that screening was postponed.

We examined binary associations between covariate variables and outcome variables by using the Pearson χ^2 test or the Fisher exact test when the expected cell counts of some cells were <5. We conducted multivariate analysis between outcome variables and covariate variables by using multivariate logistic regression. We used stepwise model selection to select the predictors. First, we conducted a binary analysis to identify the list of significant predictors in each model. Second, we examined collinearity among the list of significant predictors. For our analysis, we retained predictors among the group of mutually correlated predictors with a correlation coefficient >0.6. Finally, we used stepwise model selection to select the final model, with $P < .05$ considered significant. We also checked for confounding and interaction between the predictors in the model. For CRC screening, we combined stool-based screening tests and colonoscopies into 1 outcome variable because of the small sample size. We included covariate variables in the final model for each outcome variable. We determined the adjusted odds ratios (AORs) for associations between dependent (ie, outcome) variables and independent (ie, predictor) variables. We excluded respondents with missing outcome values from the bivariate and multivariate analyses. We used SAS version 9.4 (SAS Institute Inc) for statistical analysis.

The university institutional review boards (IRBs) at UCD (IRB no. 1639547-4) and OUSC (IRB no. 12190) approved the study; in Oklahoma, the Choctaw Nation (IRB no. 2020-003) and Cherokee Nation (IRB no. 338) IRBs also approved the study. All survey participants gave written or verbal consent.

Results

The overall sample included 767 AIs who were eligible for breast cancer, cervical cancer, and/or CRC screenings.

Among respondents, 395 women were eligible for mammography, 517 women were eligible for Pap testing, and 454 people were eligible for stool-based screening and/or colonoscopies (Table 1). Of 395 women eligible for mammograms, 234 (59.2%) responded that they were planning to obtain one; however, 127 of 234 (54.3%) postponed mammograms because of COVID-19. Of 517 women eligible for Pap testing, 357 (69.1%) had planned to obtain one; however, 115 of 357 (32.2%) postponed Pap tests because of COVID-19. Of 454 adults eligible for CRC screening, 282 (62.1%) responded they were planning to obtain one; however, 80 of 282 (28.4%) postponed CRC screening because of COVID-19.

In bivariate analysis, significant predictors associated with planning to obtain a mammogram included having a preexisting condition other than cancer, health insurance status, and residing with a minor. Significant predictors associated with planning to obtain a Pap test included attending gatherings with more than 2 people outside of your household and the number of social distancing recommendations followed. The only significant predictor associated with planning to obtain a CRC screening was health insurance status (Table 2).

Multivariate analyses showed that eligible AI women with (vs without) a preexisting condition other than cancer had higher odds of planning to obtain a mammogram (AOR=2.3; 95% CI, 1.1-4.8) (Table 3). Women who lived with a child (vs did not live with a child) had lower odds of planning to get a mammogram (AOR=0.6; 95% CI, 0.3-1.0). Women who had been tested for COVID-19 (vs not tested for COVID-19) had higher odds of postponing their mammogram (AOR=5.7; 95% CI, 1.6-20.3).

Among eligible AI women, characteristics independently associated with higher odds of planning to have a Pap test included having excellent or good (vs fair or poor) health (AOR=1.7; 95% CI, 1.0-2.9) and attending (vs not attending) a gathering with >2 individuals from outside their household (AOR=1.7; 95% CI, 1.1-2.8). Women who followed ≥ 3 (vs ≤ 2) social distancing guidelines had higher odds of postponing a Pap test (AOR=7.3; 95% CI, 0.9-58.9). Women who attended (vs did not attend) a gathering with >2 people from outside their household had lower odds of postponing their Pap test (AOR=0.4; 95% CI, 0.2-0.9).

Among eligible AI adults, those who received (vs did not receive) support during the COVID-19 pandemic had higher odds of planning to have a CRC screening (AOR=2.0; 95% CI, 1.1-3.9). Adults with (vs without) health insurance coverage had higher odds of postponing their CRC screening (AOR=7.7; 95% CI, 1.0-58.6).

Discussion

According to our survey results, more than half of AI participants eligible for a cancer screening had planned to get screened, but more than one-quarter of eligible AI

Table 1. Demographic characteristics and COVID-19 safety behaviors and attitudes of American Indian adults residing in California and Oklahoma who were eligible for breast, cervical, and/or colorectal cancer screening, March through December 2020^a

Characteristic	No. (%) of participants
Residence (n = 767)	
Oklahoma	453 (59.1)
California	314 (40.9)
Age group, y (n = 760)	
18-39	219 (28.8)
40-59	270 (35.5)
≥60	271 (35.7)
Sex (n = 765)	
Female	622 (81.3)
Male	143 (18.7)
Hispanic (n = 673)	
No	564 (83.8)
Yes	109 (16.2)
Educational attainment (n = 762)	
High school diploma/GED	240 (31.5)
≥Some college	522 (68.5)
Annual household income (n = 695)	
<\$35 000	305 (43.9)
≥\$35 000	390 (56.1)
Health insurance (n = 758)	
Yes	651 (85.9)
No	107 (14.1)
Health status (n = 761)	
Excellent or good	536 (70.4)
Fair or poor	225 (29.6)
Marital status (n = 760)	
Single, never been married	105 (13.8)
Married or living together	430 (56.6)
Separated/divorced/widowed	225 (29.6)
Employment prepandemic (n = 695)	
Full- or part-time employment	305 (43.9)
Unemployed or other ^b	390 (56.1)
Child residing in household (n = 767)	
Yes	463 (60.4)
No	304 (39.6)
No. of social distancing guidelines followed (n = 713)	
≤2	89 (12.5)
≥3	624 (87.5)
Importance of social distancing during COVID-19 (n = 747)	
Very to somewhat	657 (88.0)
Little to not	90 (12.0)
Tested for COVID-19 (n = 754)	
Yes	492 (65.3)
No	262 (34.7)
Close physical contact with person who tested positive for COVID-19 (n = 706)	
Yes	135 (19.1)

(Continued)

Table 1. (continued)

Characteristic	No. (%) of participants
No	571 (80.9)
Supported during the COVID-19 lockdown (n = 713)	
Yes	203 (28.5)
No	510 (71.5)
Attended gatherings (>2 people outside of household) (n = 767)	
Yes	327 (42.6)
No	440 (57.4)
Have a preexisting condition (other than cancer) (n = 767)	
Yes	620 (80.8)
No	147 (19.2)
Eligible for mammography and screening planned (n = 395)	
Yes	234 (59.2)
No	161 (40.8)
Mammography planned and postponed because of COVID-19 (n = 234)	
Yes	127 (54.3)
No	107 (45.7)
Eligible for Pap test and screening planned (n = 517)	
Yes	357 (69.1)
No	160 (30.9)
Pap test planned and postponed because of COVID-19 (n = 357)	
Yes	115 (32.2)
No	242 (67.8)
Eligible for colorectal cancer screening and screening planned (n = 454)	
Yes	282 (62.1)
No	172 (37.9)
Colorectal cancer screening planned and postponed because of COVID-19 (n = 282)	
Yes	80 (28.4)
No	202 (71.6)

Abbreviations: GED, General Educational Development; Pap, Papanicolaou.

^a Data source: subset of survey questions developed by the National Cancer Institute–funded Impact on COVID-19 on the Cancer Continuum Consortium; participants were surveyed from October 2020 through January 2021.^b Other included student, retired, homemaker, and person with disability.

participants reported that their screening was postponed and/or delayed because of COVID-19. The percentage of eligible AI participants who reported having a planned cancer screening in our study was higher than reported by Dennis et al.¹⁸ In that study, in 2020, 35.4% of AI/AN women had a mammogram, 40.1% of AI/AN women had a Pap test, and 13.8% of AI/AN adults had a CRC screening.¹⁸ Further investigation is needed to assess whether people who had planned to get a cancer screening received the screening.

Table 2. Bivariate analysis of demographic characteristics and COVID-19 safety behaviors and attitudes among American Indian adults residing in California and Oklahoma who had planned a cancer screening, March through December 2020^a

Characteristic	Mammography planned (women aged 40-74 y) (n=234)		Pap test planned (women aged 21-65 y) (n=357)		CRC screening planned (adults aged 50-75 y) (n=282)	
	No. (%)	P value ^b	No. (%)	P value ^b	No. (%)	P value ^b
Age group, y		.09		.15		.08
18-39	—		196 (54.9)		—	
40-59	124 (53.0)		120 (33.6)		102 (36.2)	
≥60	105 (44.9)		36 (10.1)		180 (63.8)	
Sex		—		—		.22
Female	234 (100.0)		357 (100.0)		180 (63.8)	
Male	—		—		105 (37.2)	
Hispanic		.98		.29		.36
No	155 (66.2)		75 (21.0)		190 (67.4)	
Yes	22 (9.4)		240 (67.2)		24 (8.5)	
Educational attainment		.84		.85		.90
High school diploma/GED	104 (44.4)		113 (31.7)		129 (45.7)	
≥Some college	130 (55.6)		243 (68.1)		157 (55.7)	
Annual household income		.50		.93		.23
<\$35 000	96 (41.0)		137 (38.4)		119 (42.2)	
≥\$35 000	106 (45.3)		188 (52.7)		121 (42.9)	
Health insurance		.03		.28		.02
Yes	210 (89.7)		287 (80.4)		264 (93.6)	
No	21 (9.0)		66 (18.5)		19 (6.7)	
Health status		.98		.02		.44
Excellent or good	151 (64.5)		264 (73.9)		196 (69.5)	
Fair or poor	82 (35.0)		91 (25.5)		89 (31.6)	
Marital status		.46		.31		.79
Single, never been married	29 (12.4)		70 (19.6)		27 (9.6)	
Married or living together	103 (44.0)		211 (59.1)		136 (48.2)	
Separated/divorced/widowed	100 (42.7)		74 (20.7)		119 (42.2)	
Employment prepandemic		.89		.20		.46
Full-time or part-time	97 (41.5)		239 (66.9)		94 (33.3)	
Unemployed or other ^c	132 (56.4)		112 (31.4)		185 (65.6)	
Child in household		.01		.15		.51
Yes	155 (66.2)		195 (54.6)		66 (23.4)	
No	79 (33.8)		162 (45.4)		221 (78.4)	
No. of social distancing guidelines followed		.24		.42		.72
≤2	10 (4.3)		43 (12.0)		9 (3.2)	
≥3	209 (89.3)		286 (80.1)		263 (93.3)	
Importance of social distancing		.15		.34		.67
Very to somewhat	216 (92.3)		293 (82.1)		268 (95.0)	
Little to not	11 (4.7)		56 (15.7)		6 (2.1)	
Tested for COVID-19		.07		.38		.56
Yes	145 (62.0)		242 (67.8)		165 (58.5)	
No	85 (36.3)		112 (31.4)		118 (41.8)	
Attended gatherings (>2 people)		.68		.009		.57
Yes	126 (53.8)		213 (59.7)		129 (45.7)	
No	108 (46.2)		144 (40.3)		158 (56.0)	
Preexisting condition		.008		.31		.23
Yes	195 (83.3)		259 (72.5)		252 (89.4)	
No	39 (16.7)		98 (27.5)		35 (12.4)	
Supported during COVID-19 lockdown		.19		.40		.28
Yes	77 (32.9)		105 (29.4)		73 (25.9)	
No	142 (60.7)		236 (66.1)		196 (69.5)	

Abbreviations: —, does not apply; CRC, colorectal cancer; GED, General Educational Development; Pap, Papanicolaou.

^a Data source: subset of survey questions developed by the National Cancer Institute–funded Impact on COVID-19 on the Cancer Continuum Consortium; participants were surveyed from October 2020 through January 2021. Percentages may not total to 100 because of missing values.

^b Significant at $P < .05$ using Pearson χ^2 test.

^c Other included student, retired, homemaker, and person with disability.

Table 3. Multivariate regression analysis of predictors of planned and delayed cancer screenings among American Indian adults residing in California and Oklahoma, March through December 2020^a

Factor	Adjusted odds ratio (95% CI) [P value] ^b					
	Mammogram planned (n=234)	Mammogram delayed (n=127)	Pap test planned (n=357)	Pap test delayed (n=115)	CRC screening planned (n=279)	CRC screening delayed (n=287)
Having a preexisting condition other than cancer (yes vs no)	2.3 (1.1-4.8) [.02] ^b	—	—	—	—	—
Child residing in household (yes vs no)	0.6 (0.3-1.0) [.04] ^b	—	—	—	—	—
Had a COVID-19 test (yes vs no)	—	5.7 (1.6-20.3) [.007] ^b	—	—	—	—
Health status (excellent or good vs fair or poor)	—	—	1.7 (1.0-2.9) [.06]	—	—	—
Has been in any group of >2 people outside the household (yes vs no)	—	—	1.7 (1.1-2.8) [.02] ^b	0.4 (0.2-0.9) [.03] ^b	—	—
No. of social distancing guidelines followed (≤2 vs ≥3)	—	—	—	7.3 (0.9-58.9) [.06]	—	—
Support during COVID-19 pandemic (yes vs no)	—	—	—	—	2.0 (1.1-3.9) [.03] ^b	—
Health insurance status (yes vs no)	—	—	—	—	—	7.7 (1.0-58.6) [.05] ^b

Abbreviations: CRC, colorectal cancer; Pap, Papanicolaou.

^a Data source: subset of survey questions developed by the National Cancer Institute–funded Impact on COVID-19 on the Cancer Continuum Consortium; participants were surveyed from October 2020 through January 2021.

^b Significant at $P < .05$ using multivariate logistic regression.

Rates of postponed and/or delayed cancer screenings because of COVID-19 among AI adults in our study were also higher than in a study at another IC-4 consortium site.²⁶ In that study, 24.5% of women had planned and delayed a mammogram (vs 54.3% of respondents in our study), 27.1% of women had planned and delayed a Pap test (vs 32.2% of respondents in our study), and 28.3% of participants had planned and delayed CRC screening (vs 36.2% of respondents in our study).²⁶ Our findings align with findings from the Centers for Disease Control and Prevention, which estimated that, by June 30, 2020, about 32% of US adults reported avoiding routine medical care because of COVID-19 concerns.²⁷

Unlike other studies that examined general delays in cancer screenings, our study focused only on AI people and included data on demographic characteristics and COVID-19 behaviors.^{19,28-30} The higher rates of postponed and/or delayed cancer screenings that we found in our study compared with what has been reported for the general population are alarming as, historically, AI communities have faced additional challenges and barriers to obtaining recommended cancer screenings (eg, transportation, geographic isolation, culture, mistrust of the health care system).³¹⁻³³ Health care organizations should consider employing patient navigators to help reduce barriers to cancer screening for AI

people. Patient navigation has proven to be an effective and successful model to increase cancer screenings among underresourced (eg, high poverty and low income) and medically underserved communities.^{34,35}

Of the 3 cancer screenings, our study found the highest rates of postponement among AI women eligible for mammography. The Kaiser Family Foundation reported that 49% of women reported postponing some type of medical care during the pandemic, with 23% stating that they skipped preventive screenings.³⁶ Not surprisingly, in our study, the odds of having a planned mammogram were 2.3 times greater among AI women who had a preexisting condition than among AI women without a preexisting condition. Women with preexisting conditions may be more attuned to their health care needs and more vigilant of their preventive care than women without preexisting conditions, because they may have to seek continuous primary care for their other health conditions. In addition, we surmised that women who had a COVID-19 test, compared with women who did not have a COVID-19 test, would be more likely to postpone their mammogram because these women may be more concerned about being infected.

AI women who reported living with a child had 40% lower odds of planning to have a mammogram than AI women who did not live with a child. Women who live with

children may have childcare responsibilities that make it more logistically challenging for them to have a planned mammography compared with women who do not live with children. The COVID-19 pandemic resulted in a greater effect on mothers than on men and women without children.^{37,38} Effects included high rates of unemployment and the additional need for childcare because schools and daycares were closed.³⁹⁻⁴¹ In addition to their already disproportionate caretaking roles in families, many mothers assumed additional responsibilities during the COVID-19 pandemic, so they may have postponed their mammograms. In our survey, we did not ask AI women if they were mothers; data are lacking on the role of motherhood responsibilities on cancer screening practices during the COVID-19 pandemic.

The second highest rate of cancer screening postponement in our study was among AI women eligible for Pap testing; 32.2% of AI women postponed their tests because of COVID-19. Our finding is higher than the 27.1% delay in Pap screening reported by Zhang et al.²⁶ Among women eligible for a Pap test, several COVID-19–related behavioral factors were associated with plans and postponements of tests. Women who attended gatherings outside their household had higher odds of planned Pap testing than women who did not have gatherings outside their household. This finding may be attributable to the positive effects of social support among women on cervical cancer screening.^{42,43} Although not a direct measure of social support, attending social gatherings can be an opportunity to obtain social support. Conversely, AI women who followed ≥ 3 social distancing guidelines were more likely to postpone their Pap testing than AI women who followed fewer social distancing guidelines. The role of social support networks needs to be further explored as we did not ask questions about the types of gatherings attended by AI women.

Among AI adults eligible for CRC screening, 28.4% postponed screening because of COVID-19. Zhang et al²⁶ reported that 11% of participants delayed stool blood tests and 36% delayed colonoscopies during the COVID-19 pandemic.²⁶ Our study showed that AI adults who received support during the pandemic had twice the odds of having a planned CRC screening than those who did not. This finding is aligned with another study that reported the positive effect of social support on mental health during the COVID-19 pandemic.⁴⁴ Our study is the first to report on the positive effects of social support on CRC screening during the pandemic. Although social support has shown a positive effect on cancer screening during nonpandemic times,^{45,46} we need to further assess the effects of perceived and received social support during the pandemic on cancer screenings. Understanding these measures can help public health professionals and clinicians develop and plan for future interventions in times of disruptions to normal health care services. Not surprisingly, the odds of having a CRC screening were 7.7 times greater for those with health insurance than for those without health insurance, a finding consistent with previous work on predictors of CRC screening use in the United States.^{47,48}

Limitations

Our study had some limitations. First, we used a convenience sampling frame that collected cross-sectional, self-reported data. Second, we may not have been able to establish an association between cancer screening postponement and/or delay and the covariates (eg, COVID-19–related health behaviors) because the recall window for the screening questions (March–December 2020) and when the survey was administered (October 2020–January 2021) were not the same. Our findings may reflect a high level of health consciousness that is positively correlated with both social distancing and postponing and/or delaying cancer screening. Third, we did not ask why a cancer screening was postponed, whether due to health care provider or participant delay or postponement of the appointment. Fourth, we did not assess the prevalence of COVID-19 in our sample; having COVID-19 may have contributed to high rates of postponed and/or delayed cancer screenings. During the recall time frame for cancer screenings, the 7-day positivity rate for COVID-19 ranged from 0% to 14.4% in California and from 0% to 22.8% in Oklahoma.⁴⁹ More research is needed to assess whether the prevalence of COVID-19 in the community was associated with postponing and/or delaying routine cancer screenings. Fifth, our missing data disproportionately affected the sample of Hispanic respondents; therefore, findings on this group merit caution.

Our study also had several strengths, including a large sample size of AI adults residing in California and Oklahoma and a high response rate. Response rates of $\geq 60\%$ have been considered as excellent according to previous literature.⁵⁰⁻⁵² In addition, our study is the only one that focused exclusively on AI adults during COVID-19.

Conclusion

The effects of COVID-19 on cancer outcomes among AI adults will continue to be felt for years. Our study found high rates of postponement of cancer screenings among AI adults residing in California and Oklahoma. These delays could lead to an increase in late-stage cancer diagnosis and mortality, further exacerbating the disproportionate incidence of cancer among AI people. Cancer screening interventions and public health campaigns are needed to ensure that prior efforts to increase cancer screening in AI communities have not been reversed. Increased surveillance and monitoring of cancer-related outcomes among AI people are needed to ensure that pandemic-related delays in cancer screening do not lead to worse cancer outcomes in the AI population.

Acknowledgments

The authors acknowledge our tribal and community partners in California (Northern Valley Indian Health, Feather River Tribal Health) and in Oklahoma (Choctaw Nation of Oklahoma, Cherokee Nation of Oklahoma) and the Southern Plains Tribal Health Board.

Author Contributions

Drs Julie H.T. Dang and Sixia Chen are joint first authors of this article. Drs Moon S. Chen Jr and Mark P. Doescher are joint senior authors of this article.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This project was supported by National Cancer Institute grants P30CA225520 and P30CA093373. Julie H.T. Dang was supported by National Institutes of Health (NIH) grants 1K01CA258956-01A1, 5K12HD051958-17, and P30CA93373. Spencer Hall was partially supported by NIH grant P30CA225520, Oklahoma Shared Clinical and Translational Resources grant U54GM104938, and an Institutional Development Award from the National Institute of General Medical Sciences. Moon S. Chen Jr was funded by National Cancer Institute grant P30CA093373.

Disclaimer

The content of this article is solely the authors' responsibility and does not necessarily represent the official views of NIH.

ORCID iD

Julie H.T. Dang, PhD  <https://orcid.org/0000-0002-0597-7457>

References

- Centers for Medicare and Medicaid Services. Non-emergent, elective medical services, and treatment recommendations. April 4, 2020. Accessed May 31, 2023. <https://www.cms.gov/files/document/cms-non-emergent-elective-medical-recommendations.pdf>
- American Society of Clinical Oncology. Cancer screening, diagnosis, staging and surveillance. 2024. Accessed May 31, 2023. <https://www.asco.org/asco-coronavirus-resources/care-individuals-cancer-during-covid-19/cancer-screening-diagnosis-staging>
- Dietz JR, Moran MS, Isakoff SJ, et al. Recommendations for prioritization, treatment, and triage of breast cancer patients during the COVID-19 pandemic. The COVID-19 Pandemic Breast Cancer Consortium. *Breast Cancer Res Treat.* 2020;181(3):487-497. doi:10.1007/s10549-020-05644-z
- Chen RC, Haynes K, Du S, Barron J, Katz AJ. Association of cancer screening deficit in the United States with the COVID-19 pandemic. *JAMA Oncol.* 2021;7(6):878-884. doi:10.1001/jamaoncol.2021.0884
- Inadomi JM, Sonnenberg A. The impact of colorectal cancer screening on life expectancy. *Gastrointest Endosc.* 2000;51(5):517-523. doi:10.1016/s0016-5107(00)70282-3
- Yang Z, Liu J, Wang Q. Diagnose earlier, live longer? The impact of cervical and breast cancer screening on life span. *PLoS One.* 2022;17(7):e0270347. doi:10.1371/journal.pone.0270347
- Dinmohamed AG, Cellamare M, Visser O, et al. The impact of the temporary suspension of national cancer screening programmes due to the COVID-19 epidemic on the diagnosis of breast and colorectal cancer in the Netherlands. *J Hematol Oncol.* 2020;13(1):147. doi:10.1186/s13045-020-00984-1
- Ricciardiello L, Ferrari C, Cameletti M, et al. Impact of SARS-CoV-2 pandemic on colorectal cancer screening delay: effect on stage shift and increased mortality. *Clin Gastroenterol Hepatol.* 2021;19(7):1410-1417.e9. doi:10.1016/j.cgh.2020.09.008
- Maringe C, Spicer J, Morris M, et al. The impact of the COVID-19 pandemic on cancer deaths due to delays in diagnosis in England, UK: a national, population-based, modelling study [published correction appears in *Lancet Oncol.* 2021;22(1):e5]. *Lancet Oncol.* 2020;21(8):1023-1034. doi:10.1016/S1470-2045(20)30388-0
- Roubidoux MA, Kaur JS, Rhoades DA. Health disparities in cancer among American Indians and Alaska Natives. *Acad Radiol.* 2022;29(7):1013-1021. doi:10.1016/j.acra.2021.10.011
- White MC, Espey DK, Swan J, Wiggins CL, Ehemam C, Kaur JS. Disparities in cancer mortality and incidence among American Indians and Alaska Natives in the United States. *Am J Public Health.* 2014;104(suppl 3):S377-S387. doi:10.2105/AJPH.2013.301673
- Ko NY, Hong S, Winn RA, Calip GS. Association of insurance status and racial disparities with the detection of early-stage breast cancer. *JAMA Oncol.* 2020;6(3):385-392. doi:10.1001/jamaoncol.2019.5672
- American Cancer Society. *Cancer Facts and Figure, Special Section: Cancer in the American Indian and Alaskan Native Population.* American Cancer Society; 2022.
- Kratzer TB, Jemal A, Miller KD, et al. Cancer statistics for American Indian and Alaska Native individuals, 2022: including increasing disparities in early onset colorectal cancer. *CA Cancer J Clin.* 2023;73(2):120-146. doi:10.3322/caac.21757
- United States Cancer Statistics Working Group. U.S. Cancer Statistics data visualizations tool, based on 2021 submission data (1999-2019): US Department of Health and Human Services, Centers for Disease Control and Prevention, and National Cancer Institute. November 2022. Accessed May 31, 2023. <https://www.cdc.gov/cancer/dataviz>
- Haverkamp D, Melkonian SC, Jim MA. Growing disparity in the incidence of colorectal cancer among non-Hispanic American Indian and Alaska Native populations—United States, 2013-2017. *Cancer Epidemiol Biomarkers Prev.* 2021;30(10):1799-1806. doi:10.1158/1055-9965.EPI-21-0343
- Barsouk A, Saginala K, Aluru JS, Rawla P, Barsouk A. US cancer screening recommendations: developments and the impact of COVID-19. *Med Sci (Basel).* 2022;10(1):16. doi:10.3390/medsci10010016
- Dennis LK, Hsu CH, Arrington AK. Reduction in standard cancer screening in 2020 throughout the U.S. *Cancers (Basel).* 2021;13(23):5918. doi:10.3390/cancers13235918
- Fedewa SA, Star J, Bandi P, et al. Changes in cancer screening in the US during the COVID-19 pandemic. *JAMA Netw Open.* 2022;5(6):e2215490. doi:10.1001/jamanetworkopen.2022.15490
- US Census Bureau. A look at the largest American Indian and Alaska Native tribes and villages in the nation, tribal areas and states. 2023. Accessed March 8, 2024. <https://www.census.gov/library/stories/2023/10/2020-census-dhc-a-aian-population.html>

21. Shokar NK, Carlson CA, Weller SC. Factors associated with racial/ethnic differences in colorectal cancer screening. *J Am Board Fam Med.* 2008;21(5):414-426. doi:10.3122/jabfm.2008.05.070266
22. Swan J, Breen N, Coates RJ, Rimer BK, Lee NC. Progress in cancer screening practices in the United States: results from the 2000 National Health Interview Survey. *Cancer.* 2003;97(6):1528-1540. doi:10.1002/cncr.11208
23. Hall IJ, Tangka FKL, Sabatino SA, Thompson TD, Graubard BI, Breen N. Patterns and trends in cancer screening in the United States. *Prev Chronic Dis.* 2018;15:E97. doi:10.5888/pcd15.170465
24. Rosenstock IM. The health belief model and preventive health behavior. *Health Educ Monogr.* 1974;2(4):354-386.
25. Scarinci IC, Pandya VN, Kim YI, et al. Factors associated with perceived susceptibility to COVID-19 among urban and rural adults in Alabama. *J Community Health.* 2021;46(5):932-941. doi:10.1007/s10900-021-00976-3
26. Zhang X, Elsaid MI, DeGraffinreid C, Champion VL, Paskett ED. Impact of COVID-19 on Behaviors Across the Cancer Control Continuum in Ohio Group. Impact of the COVID-19 pandemic on cancer screening delays. *J Clin Oncol.* 2023; 41(17):3194-3202. doi:10.1200/JCO.22.01704
27. Czeisler MÉ, Marynak K, Clarke KE, et al. Delay or avoidance of medical care because of COVID-19-related concerns—United States, June 2020. *Morb Mortal Wkly Rep.* 2020;69(36):1250-1257. doi:10.15585/mmwr.mm6936a4
28. Fedewa SA, Bandi P, Smith RA, Silvestri GA, Jemal A. Lung cancer screening rates during the COVID-19 pandemic. *Chest.* 2022;161(2):586-589. doi:10.1016/j.chest.2021.07.030
29. Star J, Bandi P, Siegel RL, et al. Cancer screening in the United States during the second year of the COVID-19 pandemic. *J Clin Oncol.* 2023;41(27):4352-4359. doi:10.1200/JCO.22.02170
30. Star J, Bandi P, Nargis N, et al. Updated review of major cancer risk factors and screening test use in the United States, with a focus on changes during the COVID-19 pandemic. *Cancer Epidemiol Biomarkers Prev.* 2023;32(7):879-888. doi:10.1158/1055-9965.EPI-23-0114
31. Towne SD, Smith ML, Ory MG. Geographic variations in access and utilization of cancer screening services: examining disparities among American Indian and Alaska Native elders. *Int J Health Geogr.* 2014;13:18. doi:10.1186/1476-072X-13-18
32. Frerichs L, Rhode J, Bell R, et al. Perspectives of American Indians in Eastern North Carolina on socio-cultural factors that influence colorectal cancer screening decisions. *J Health Care Poor Underserved.* 2018;29(2):723-742. doi:10.1353/hpu.2018.0055
33. Harris R, Van Dyke ER, Ton TG, Nass CA, Buchwald D. Assessing needs for cancer education and support in American Indian and Alaska Native communities in the northwestern United States. *Health Promot Pract.* 2016;17(6):891-898. doi:10.1177/1524839915611869
34. Krebs LU, Burhansstipanov L, Watanabe-Galloway S, Pingatore NL, Petereit DG, Isham D. Navigation as an intervention to eliminate disparities in American Indian communities. *Semin Oncol Nurs.* 2013;29(2):118-127. doi:10.1016/j.soncn.2013.02.007
35. Neal CD, Weaver DT, Raphael TJ, et al. Patient navigation to improve cancer screening in underserved populations: reported experiences, opportunities, and challenges. *J Am Coll Radiol.* 2018;15(11):1565-1572. doi:10.1016/j.jacr.2018.03.001
36. Long M, Frederiksen B, Ranji U, Diep K, Salganicoff A. *Many Women Use Preventive Services, But Gaps in Awareness of Insurance Coverage Requirements Persist: Findings From the 2022 KFF Women's Health Survey.* Kaiser Family Foundation. February 22, 2023. Accessed May 31, 2023. <https://www.kff.org/womens-health-policy/issue-brief/many-women-use-preventive-services-but-gaps-awareness-insurance-coverage-requirements-persist-findings-from-2022-kff-womens-health-survey>
37. Yavorsky JE, Qian Y, Sargent AC. The gendered pandemic: the implications of COVID-19 for work and family. *Sociol Compass.* 2021;15(6):e12881. doi:10.1111/soc4.12881
38. Yacona K, Hanna MW, Niyazi S, et al. Can COVID-19 worsen racial disparities in breast cancer screening and diagnosis? *J Clin Imaging Sci.* 2022;12:35. doi:10.25259/JCIS_26_2022
39. Couch KA, Fairlie RW, Xu H. The evolving impacts of the COVID-19 pandemic on gender inequality in the US labor market: the COVID motherhood penalty. *Econ Inquiry.* 2022;60(2):485-507. doi:10.1111/ecin.13054
40. Zamorro G, Prados MJ. Gender differences in couples' division of childcare, work and mental health during COVID-19. *Rev Econ Househ.* 2021;19(1):11-40. doi:10.1007/s11150-020-09534-7
41. Cummins MW, Brannon GE. Mothering in a pandemic: navigating care work, intensive motherhood, and COVID-19. *Gender Issues.* 2022;39(2):123-141. doi:10.1007/s12147-022-09295-w
42. Wasserman MR, Bender DE, Lee SY, Morrissey JP, Mouw T, Norton EC. Social support among Latina immigrant women: bridge persons as mediators of cervical cancer screening. *J Immigr Minor Health.* 2006;8(1):67-84. doi:10.1007/s10903-006-6343-0
43. Documet P, Bear TM, Flatt JD, Macia L, Trauth J, Ricci EM. The association of social support and education with breast and cervical cancer screening. *Health Educ Behav.* 2015;42(1):55-64. doi:10.1177/1090198114557124
44. Grey I, Arora T, Thomas J, Saneh A, Tohme P, Abi-Habib R. The role of perceived social support on depression and sleep during the COVID-19 pandemic. *Psychiatry Res.* 2020;293:113452. doi:10.1016/j.psychres.2020.113452
45. Honda K, Kagawa-Singer M. Cognitive mediators linking social support networks to colorectal cancer screening adherence. *J Behav Med.* 2006;29(5):449-460. doi:10.1007/s10865-006-9068-1
46. Smalls BL, Aroh A, McQuerry K, Adegboyega A, Schoenberg N, Hatcher J. Social support and breast cancer screening in rural Appalachia. *Psychooncology.* 2018;27(9):2281-2288. doi:10.1002/pon.4828
47. Seeff LC, Nadel MR, Klabunde CN, et al. Patterns and predictors of colorectal cancer test use in the adult U.S. population. *Cancer.* 2004;100(10):2093-2103. doi:10.1002/cncr.20276
48. Swan J, Breen N, Graubard BI, et al. Data and trends in cancer screening in the United States: results from the 2005 National Health Interview Survey. *Cancer.* 2010;116(20):4872-4881. doi:10.1002/cncr.25215

49. Johns Hopkins University of Medicine, Coronavirus Resource Center. Testing hub: daily state-by-state testing trends. September 21, 2022. Accessed December 26, 2023. <https://coronavirus.jhu.edu/testing/individual-states>
50. Booker QS, Austin JD, Balasubramanian BA. Survey strategies to increase participant response rates in primary care research studies. *Fam Pract*. 2021;38(5):699-702. doi:10.1093/fampra/cmab070
51. Meyer VM, Benjamens S, Moumni ME, Lange JF, Pol RA. Global overview of response rates in patient and health care professional surveys in surgery: a systematic review. *Ann Surg*. 2022;275(1):e75-e81. doi:10.1097/SLA.0000000000004078
52. Murdoch M, Simon AB, Polusny MA, et al. Impact of different privacy conditions and incentives on survey response rate, participant representativeness, and disclosure of sensitive information: a randomized controlled trial. *BMC Med Res Methodol*. 2014;14:90.