

Lawrence Berkeley National Laboratory

Recent Work

Title

ELECTRON PARAMAGNETIC RESONANCE SPECTRUM OF O₂(1 g) u ITS 170 HYPERFINE COUPLING AND ELECTRONIC g AND ROTATIONAL g-VALUES

Permalink

<https://escholarship.org/uc/item/3z83d50k>

Authors

Arrington, C. Anthony
Falick, Arnold M.
Myers, Rollie J.

Publication Date

1971-04-01

Submitted to Journal of
Chemical Physics

UCRL-20518
Preprint c.2

RECEIVED
LAWRENCE
RADIATION LABORATORY

DOCUMENTS SECTION

ELECTRON PARAMAGNETIC RESONANCE SPECTRUM OF
 $O_2(^1\Delta_g)$ -- ITS ^{17}O HYPERFINE COUPLING AND
ELECTRONIC AND ROTATIONAL g-VALUES

C. Anthony Arrington, Jr., Arnold M. Falick and Rollie J. Myers

April 1971

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

4
LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-20518
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

ELECTRON PARAMAGNETIC RESONANCE SPECTRUM OF $O_2(^1\Delta_g)$ --
 ITS ^{17}O HYPERFINE COUPLING AND ELECTRONIC AND ROTATIONAL g-VALUES

C. Anthony Arrington, Jr.,* Arnold M. Falick† and Rollie J. Myers

Inorganic Materials Research Division of the Lawrence Radiation Laboratory,
 and Department of Chemistry, University of California,
 Berkeley, California 94720

ABSTRACT

The ^{17}O hyperfine coupling was measured for $O_2(^1\Delta_g)$ in its $J = 2$ angular momentum state. The coupling constant $2g_I\mu_N\mu_B \langle \sum_i l_{zi}/r_i^3 \rangle$ was measured to be -424 ± 1 MHz. From this value we can follow Harvey's convention and determine $\langle 1/r_\ell^3 \rangle = 39.6 \times 10^{-24} \text{cm}^{-3}$. The previous work by Miller et al. on $O_2(^3\Sigma_g^-)$ can be used to determine $\langle 1/r_s^3 \rangle = 43.6 \times 10^{-24} \text{cm}^{-3}$. Even though these two values refer to different electronic states, they are in the same ratio that Harvey found for $O(^3P)$, but they are both appreciably larger than the values he found for $O(^3P)$.

The g_J values were determined for both the $J = 2$ and $J = 3$ angular momentum states. From these values we obtain $g_L = 0.999860$ and $g_r = -1.70 \times 10^{-4}$. The g_L value is very close to the recent determination of T. Miller, but the g_r value does not agree with Miller's work.

* present address: Department of Chemistry, Furman University,
 Greenville, South Carolina

† present address: Space Sciences Laboratory, University of
 California, Berkeley, California

Previous work¹ has shown that the metastable ${}^1\Delta_g$ excited electronic state of O_2 can be detected by electron paramagnetic resonance (EPR) spectroscopy in the gas phase. It can be generated by an electrical discharge through O_2 and by a variety of chemical techniques. It appears to have considerable importance in oxidation reactions including those involved in the production of photochemical smog. We have continued our EPR work on this species to include the ${}^{17}O$ hyperfine interaction and a determination of the electronic and rotational g-values.

Our results for the hyperfine interaction of $O_2({}^1\Delta_g)$ will be compared with the previous work on the atomic and molecular ground states of oxygen, $O_2({}^3\Sigma_g^-)$ and $O({}^3P)$. The ${}^{17}O$ hyperfine coupling constant is a measure of $\langle 1/r^3 \rangle$ for the unpaired electrons. Accurate theoretical estimates of $\langle 1/r^3 \rangle$ are still not available for either electronic state of O_2 .

The electronic and rotational g-values depend upon a variety of characteristics of the electronic wave function. Since they involve absolute field measurements, the experimental determinations of these quantities are difficult. After our work was in manuscript stage we received a copy of a $O_2({}^1\Delta_g)$ g-value measurement by Miller.² Comparison will be made with Miller's results which appear to be of high accuracy.

Experimental

The ${}^1\Delta_g$ state of O_2 was generated in a flowing gas stream by an electrical discharge. For the ${}^{17}O$ work we used a 2450 MHz 100 watt unit

and cavity³ while for the g-value work we used a 13 MHz radio frequency electrodeless discharge. In the ¹⁷O work the gas was recycled by means of a mercury diffusion pump of conventional design. We started with enriched oxygen⁴ consisting of 50% oxygen-18, 30% oxygen-16 and 20% oxygen-17. At the end of each run the enriched O₂ was condensed into an activated charcoal trap at 77°K. The enriched O₂ could be returned to the flow system by warming the trap to room temperature.

Typical pressures in the ¹⁷O work were 0.8 torr. At this pressure it was possible to get the maximum ¹Δ_g signal and still keep the mercury recycling pump in operation. A single charge of enriched O₂ was good for about 20 hrs of discharge and recycling time. The O₂ slowly reacted with the hot mercury in the pump and the percentage of oxygen-16 slowly increased in the recycled gas. For the g-value work we simply used dry tank O₂ in a fast flow system at a pressure of about 0.6 torr.

The ¹⁷O measurements were made with a standard 100 KHz field modulated X-band EPR spectrometer. The sample was pumped through a quartz lined Varian V4533 cylindrical cavity. Field measurements were made with a commercial NMR gaussmeter. The cavity frequency in the TE₀₁₁ mode was 9.082 GHz.

The g-value measurements were made with a specially constructed S-band spectrometer. A rectangular resonant cavity was made from a section of S-band wave guide in which the center part of the broad wall was replaced with 5 x 10⁻⁴ inch silver foil. The gas flowed through a quartz pill box, with side arms, which almost filled the center section of the TE₁₀₄ cavity. The two modulation coils were mounted on the outside of

1/8 inch thick plastic sheets which supported the silver foil walls. The final cavity frequency was about 3.13 GHz, and its Q was close to 10,000.

The microwave source was a HP 8616A signal generator which was locked to the cavity frequency by means of photon coupling⁵ and frequency modulation with phase detection at 10 KHz. The cavity was critically coupled to the line and the microwave bridge was constructed using a 4-port S-band wave guide circulator. The magnet had 12 inch dia. pole caps and a 2 5/8 inch gap. The modulation coils were quite close to the pole caps which gave severe eddy current loss at 100 KHz. For this reason, the modulation coils were driven in a non-resonant circuit by a 600 ohm output impedance power amplifier.

The magnetic field was measured with a proton NMR which was locked to the magnetic field. The circuits for this device were similar to those of Maki and Volpicelli.⁶ Our primary standard was a proton probe doped to the 0.1-0.2 M level with $\text{Cu}(\text{NO}_3)_2$. With this level of Cu^{2+} the proton resonance is shifted from its pure water value by less than 0.1 ppm. We used the 1963 N.B.S. value of 4257.59 Hz G^{-1} to convert the proton resonant frequencies to magnetic field values. In order to correct for the positional field shift between the sample and the primary standard probe we used a commercial Harvey Wells NMR instrument and its associated probe. This probe was placed in the same position in the magnet that the sample occupied and the difference between the two NMR systems was recorded over the full range of magnetic fields employed. We did find that the Harvey Wells probe and our primary standard NMR

did not give the same frequency in the same magnetic field. The Harvey Wells was found to be shifted to higher frequencies by +10 ppm. It is probable that the encapsulated Harvey Wells probe is doped with Mn^{2+} , Fe^{3+} or some other ion which gives a large contact shift.⁷ The +10 ppm is a rather large shift and it may be due, in part, to the encapsulation.

Since we wanted to know the centers of the EPR lines for the g-value work as accurately as possible, we averaged the data from several spectra and fit them by a least squares procedure to a Lorentz line shape. We utilized the magnetic tape data acquisition system and least squares programs previously described.⁸ Since both the NMR and EPR signals can have a time shift, an average of the line centers for both up-field and down-field sweeps was used. A difference of 0.5 G was typical between the up- and down-field sweeps.

The EPR Spectrum

The basic features of the EPR spectrum for $O_2(^1\Delta_g)$ can be described in terms of an electronic angular momentum ($\Lambda = 2$) which is quantized along the internuclear axis (Hund's case a). Together with the rotational angular momentum, they form the total angular momentum J which is quantized along the magnetic field axis (M_J). The electronic magnetic moment along the internuclear axis has the value of $2g_L\mu_B$ where g_L should be close to unity.

The first-order Zeeman energy of this system is

$$\epsilon(\text{first-order}) = g_L\mu_B HM_J \quad (1)$$

where

$$g_J = \frac{4g_L}{J(J+1)} \quad (2)$$

Using the selection rule for magnetic dipole radiation of $\Delta M_J = \pm 1$, one predicts four absorption signals all at $g_J \sim 2/3$ for the lowest total-angular-momentum state with $J = 2$. The observed $J = 2$ spectrum¹ in fact consists of these four absorptions, but they are split apart by a higher order Zeeman effect. This splitting can be readily explained by a mixing of the J levels which depends upon the ratio of $\mu_B H$ to the rotational constant B .

For the ^{17}O nucleus with $I = 5/2$, we must consider the new states characterized by the values of M_I . Since the $^1\Delta_g$ state has a first-order Zeeman effect, it also has a large component of magnetic field, produced by the orbital moment, which is at the nuclei and parallel to H . This field is much larger than H itself and it is the dominant interaction for the ^{17}O nuclei. There is also a small nuclear quadrupole interaction and small second-order terms involving the J mixing. The $J = 2$ energy level pattern for a molecule with one ^{17}O nucleus is shown in Figure 1.

The exact calculation of these energy levels is simplified by the fact that Eq. (1) is the major term in the Zeeman energy. One can either use the older formalism of Condon and Shortley⁹ as explained by Van Vleck and Frosch and Foley¹⁰ or the newer and more complete, formulation of Carrington, Levy and Miller (CLM).¹¹ A complete analysis of the ^{17}O

hyperfine interaction is possible from the spectrum of a single J state, and the work reported here was done on the $J = 2$ state at X-band (9 GHz).

A complete analysis of the g -value problem is only possible if work is done on two J states, because one must separate the rotational-electronic contributions to the magnetic moment from the purely electronic ones. We chose to operate at S-band (3 GHz) so that both the $J = 2$ and $J = 3$ spectra would be within the range of our magnet.

^{17}O Results

In our enriched O_2 one can see EPR lines due to $^{16}\text{O}_2$, $^{18}\text{O}_2$, $^{16}\text{O}^{18}\text{O}$, $^{16}\text{O}^{17}\text{O}$ and $^{17}\text{O}^{18}\text{O}$. The spectral region also includes some strong absorptions due to O_2 in its ground $^3\Sigma_g^-$ state. One could expect to observe 48 lines from the two ^{17}O species with the selection rule $\Delta M_I = 0$. We could only resolve 28 of these. The signal to noise ratio was only about 3:1 for the ^{17}O lines and in addition overlapping of some of the many possible lines from the several species present made complete resolution impossible. The line positions were measured to ± 0.5 G for both the $M_J = -1$ to 0 and the $M_J = 0$ to +1 transitions, and to ± 1 G for the other sets of transitions.

For a $^1\Delta$ electronic state there is only one important term in the magnetic hyperfine part of the electronic Hamiltonian. Following CIM it can be written for one interacting nucleus α as

$$\mathcal{H}_{\text{hf}}^e = \sum_i a_i \bar{l}_i \cdot \bar{I}_\alpha \quad (3)$$

where for a single electronic state η

$$\langle \eta | a_i \bar{l}_i | \eta \rangle = \langle \eta | 2g_{\alpha} \mu_N \mu_B r^{-3} \bar{l}_i | \eta \rangle \quad (4)$$

In their evaluation of the electronic-rotational matrix, elements given by Eq. (3), CLM use a JIM_F representation. In ordinary magnetic fields Eq. (1) is almost exact with the result that the orbital angular momentum is very nearly quantized along the axis of the field. As a result a JIM_I representation is more appropriate for the evaluation of the matrix elements that result from Eq. (3). For $J = 2$ the result is that

$$c_{hf}(\text{first-order}) = (\langle a \rangle / 3) M_J M_I + g_I \mu_N H M_I \quad (5)$$

where the constant $\langle a \rangle$ is defined by

$$\langle a \rangle = \sum_i \langle \eta | a_i \bar{l}_{z_i} | \eta \rangle \quad (6)$$

where z is the internuclear axis. Higher order terms arise from the fact that J and M_I are not exact quantum numbers. These terms can produce shifts of the order of 10 G in the absorption lines while Eq. (5) predicts most of the hyperfine splittings of several hundred gauss as shown in Table I. The final calculations were done by diagonalizing a 72×72 matrix with $\langle a \rangle$ as an adjustable parameter. No least squares fitting was attempted. The agreement between the calculated and observed field positions shown in Table I was obtained with $\langle a \rangle / h = -424$ MHz.

Table I Observed and Calculated Hyperfine Structure for J = 2

Isotopes	M_J	$h\nu/\mu_B H$		Isotopes	M_I	$h\nu/\mu_B H$	
		obs. ^a	cal. ^b			obs. ^a	cal. ^b
$M_J = -2$ to $M_J = -1$							
16 - 17	5/2	.65174	.65177	17 - 18	1/2	.65788	.65792
16 - 17	-1/2	.68272	.68265	17 - 18	-1/2	.66819	.66826
17 - 18	1/2	.67288	.67276	17 - 18	-3/2	.67871	.67875
17 - 18	-1/2	.68377	.68325	17 - 18	-5/2	.68939	.68940
$M_J = -1$ to $M_J = 0$				$M_J = 1$ to $M_J = 2$			
16 - 17 ^c	5/2	.64472	.64472	16 - 17	5/2	.63140	.63135
16 - 17 ^c	1/2	.66506	.66506	16 - 17	3/2	.64130	.64138
16 - 17 ^c	-1/2	.67547	.67546	16 - 17	1/2	.65140	.65155
16 - 17 ^c	-3/2	.68597	.68603	16 - 17	-1/2	.66172	.66186
$M_J = 0$ to $M_J = 1$				16 - 17	-3/2	.67227	.67231
16 - 17	3/2	.64798	.64801	16 - 17	-5/2	.68284	.68290
16 - 17	1/2	.65814	.65819	17 - 18	5/2	.63079	.63074
16 - 17	-1/2	.66849	.66854	17 - 18	3/2	.64068	.64075
16 - 17	-3/2	.67900	.67904	17 - 18	-1/2	.66106	.66118
				17 - 18	-3/2	.67150	.67161
				17 - 18	-5/2	.68218	.68218

(a) With $\nu(\text{obs})$ from 9.0822 to 9.0818 GHz

(b) With $\langle a \rangle/h = -424$ MHz

(c) The 16 - 17 and 17 - 18 were not resolved so assumed $H(16-17) = H(\text{obs}) + 1G$.

The quantity of interest to be obtained from the ^{17}O results is $\langle 1/r^3 \rangle$ for the orbitally unpaired electrons which form the $^1\Delta$ term. Since the experimental parameter $\langle a \rangle$ contains the orbital angular momentum operators l_{z_i} , we have to make some assumptions about the wave functions of the orbitally unpaired electrons. If we assume one-electron wave functions which are derived from π orbitals, then the l_{z_i} are separately quantized in Eq.(6). With this assumption we obtain

$$\begin{aligned} \langle 1/r^3 \rangle &= \langle a \rangle 2g_I^{\mu} g_N^{\mu} \\ &= 39.6 \times 10^{-24} \text{ cm}^{-3} \end{aligned} \quad (7)$$

This same result can be obtained from the final equations of CLM, but not without some difficulty. They assume that the radial and orbital parts of Eq.(6) can always be separated and they express their final matrix elements in this form. If one follows their technique then the first term in Eq.(5) would be written as $(2a/3) M_J M_I$. Their constant a is now some kind of weighted average which results when the radial and orbital parts of $\langle a \rangle$ are separated. For a two-electron problem $a = \langle a \rangle / 2$, but it is still not rigorous to make this separation. The recent paper¹² on ^{33}S interaction in $\text{SO}(^3\Sigma)$ uses a hyperfine formalism which is similar to ours.

The separation of orbital and radial factors in $\langle a \rangle$ has been carefully investigated for O-atom by Schaeffer et al.¹³ They added a "polarization wave function" to the usual restricted Hartree-Fock treatment in order to include the effects of core polarization. The core

polarization was found to be important for both the spin and orbital terms in the hyperfine interaction of the 3P ground state of O-atom. The result of their calculations for the orbital term in the hyperfine interaction for O-atom is shown in Table II. When they included the effects of core polarization they obtained excellent agreement with the experimental value for $\langle a \rangle$ obtained for $^{17}O(^3P)$ by Harvey.¹⁴ They found that the H-F value for $\langle 1/r^3 \rangle$ was almost 10% larger than that predicted from the experimental value of $\langle a \rangle$ and that this was accounted for by the non-separability of the orbital and radial factors in $\langle a \rangle$.

Harvey also found that different values for $\langle 1/r^3 \rangle$ were obtained if he used the spin and orbital hyperfine interactions. From the orbital hyperfine interaction he determined what he called $\langle 1/r^3_\ell \rangle$ and from the spin he obtained what he called $\langle 1/r^3_s \rangle$. He found that $\langle 1/r^3_s \rangle$ was about 11% larger than $\langle 1/r^3_\ell \rangle$. The calculations of Schaeffer et al. account for this difference by core polarization. The value for $\langle 1/r^3 \rangle$ that we obtained in Eq. 7 should be more properly termed $\langle 1/r^3_\ell \rangle$, and it should be at least 10% larger than the H-F value for $\langle 1/r^3 \rangle$ in $O_2(^1\Delta_g)$. Table II also shows a $\langle 1/r^3 \rangle$ value obtained for $O_2(^3\Sigma_g^-)$ by Miller et al.¹⁵ In Harvey's notation this is a $\langle 1/r^3_s \rangle$ value since it is deduced from an electron spin hyperfine interaction. Since the $O_2(^3\Sigma_g^-)$ value for $\langle 1/r^3_s \rangle$ is about 10% larger than our $\langle 1/r^3_\ell \rangle$ value for $O_2(^1\Delta_g)$, we can conclude that they follow the same pattern that Harvey found for O-atoms. Under these circumstances we are forced to conclude that the ^{17}O hyperfine interaction

Table II Theoretical and Experimental Oxygen Hyperfine Parameters

<u>System</u>	<u>Source</u>	$\left\langle \sum_i \ell_{zi} / r_i^3 \right\rangle$ in 10^{-24} cm^{-3}	$\langle 1/r^3 \rangle$
$O(^3P)$	Theory ^a	31.3	33.6
	Expt. ^b	31.0	31.0 ^c
$O_2(^3\Sigma)$	Expt. ^d	—	43.6 ^e
$O_2(^1\Delta)$	our work	39.6	39.6 ^c

(a) Ref. 13

(b) Ref. 14

(c) More properly designated $\langle 1/r_\ell^3 \rangle$, see Ref. 14

(d) Ref. 15

(e) More properly designated $\langle 1/r_s^3 \rangle$, see Ref. 14

for O_2 in both $^3\Sigma$ and $^1\Delta$ is consistent within the uncertainties of the method, with the same value of $\langle 1/r^3 \rangle$. It should be possible in the near future to do accurate H-F calculations for O_2 and we predict that the values for $\langle 1/r^3 \rangle$ for both the $^3\Sigma$ and $^1\Delta$ states should be very close in value.

It is possible from Table II to see that the $\langle 1/r^3 \rangle$ values for O_2 are quite obviously larger than they are in O-atom. The readily available explanation for this is that the bonding in O_2 produces a contraction of the 2p orbitals so that $\langle 1/r^3 \rangle$ is increased. Confirmation of this will also have to await accurate H-F calculations for O_2 .

The g-Value Results

At least two parameters are required to fit the set of transitions for each J value. In the absence of higher-order terms in H, the center of each pattern depends only upon g_J . The higher-order terms give a shift of the center and a splitting into $2J$ transitions. These higher-order terms depend upon the rotational constants characteristic of each J value. Since a complete assignment of the rotational levels of $O_2(^1\Delta_g)$ has been made,¹⁶ we found that it was quite satisfactory to assume the rotational constants and to only fit a g_J value for each set of transitions.

Our older work¹ on $J = 2$ at X-band indicated that the spectroscopic rotational constant¹⁶ $B_0 = 1.4178 \text{ cm}^{-1}$ did not fully account for the splittings. We have subsequently found that this small discrepancy was

due to an improper calibration of the magnetic field at the sample. The difference between the field at the sample and at the NMR probe is field dependent. When this is properly corrected for, we obtain the $J = 2$ results shown in Table III. The field positions shown in this table were also calculated using the 1963 N.B.S. value for the NMR frequency conversion.

We have supplemented these old X-band (9 GHz) results with new data taken at S-band (3 GHz). We also fit the centers of each EPR line with our least-squares fitting technique.⁸ The spectrum shown in Fig. 2 was fitted assuming four Lorentz lines of arbitrary width, intensity and position. One can see that the fit was very good and the intensities were close to the 2:3:3:2 pattern expected for $J = 2$. The centers of these lines could be fitted to a precision of about 0.01 G.

Table IV shows the S-band result for $J = 2$ and $J = 3$ and the values of g_J necessary to fit the data. Only two of the six possible $J = 3$ lines were accurately measured. These two had the best signal-to-noise ratio. One can see that the g_J values from the S-band data are consistent to within the uncertainties in the magnetic field. Since the $J = 2$ values obtained at S-band are apparently more accurate than those at X-band given in Table III, we take a weighted average of 0.666630 for g_J for $J = 2$. The $J = 3$ average is obviously 0.333400.

The contribution of the rotational g -value (g_r) to the g_J values can be easily calculated, and for the two J values the results are

$$0.666630 = \frac{2}{3} g_L - \frac{1}{3} g_r \quad (8)$$

$$0.333400 = \frac{1}{3} g_L - \frac{2}{3} g_r \quad (9)$$

Table III Our Corrected Values for J = 2 at X-band

<u>Transition</u> M_J	<u>Field</u> (G)	<u>$h\nu/\mu_B H$</u>	
		<u>obs.</u> ^a	<u>calc.</u> ^b
-2 → -1	10,090.3	0.65597	0.65597
-1 → 0	9,988.8	0.66264	0.66264
0 → 1	9,885.9	0.66954	0.66954
1 → 2	9,781.9	0.67667	0.67667

(a) $\mu_B/h = 13.9960 \times 10^5 \text{ G}^{-1}\text{sec}^{-1}$

(b) With the spectroscopic rotational constants of ref. 16 and

$$g_J = 0.66661$$

Table IV Experimental S-band Results and Resultant g_J Values

<u>Transition</u>	<u>J = 2^a</u>		<u>J = 3^b</u>	
	<u>H(G)</u>	<u>g_J</u>	<u>H(G)</u>	<u>g_J</u>
-2 → -1	3,377.29	0.666633	—————	—————
-1 → 0	3,365.46	0.666631	6,693.36	0.333399
0 → 1	3,353.66	0.666632	6,734.34	0.333401
1 → 2	3,341.80	0.666634	—————	—————

(a) $\nu = 3.134236 \times 10^9$ Hz

(b) $\nu = 3.134283 \times 10^9$ Hz

If these two equations are solved we obtain

$$g_L = 0.999860 \quad (10)$$

and

$$g_r = -1.70 \times 10^{-4} \quad (11)$$

or

$$\mu_B g_r / \mu_N = -0.313. \quad (12)$$

In Eq. (11) we express the rotational g-value in units of the Bohr magneton, while in Eq. (12) it is expressed in the more natural unit of the nuclear magneton.

Our value of g_L agrees very well with Miller's² value of 0.999866 ± 0.000010 . As Miller points out the small decrease below unity is due to the relativistic contributions. The work on atoms has been summarized by Hughes.¹⁷ These contributions can be considered as the result of a power series in the fine structure constant α . They come from the terms in the series of the order $\alpha^2 \mu_B H$. As a result, we expect that g_L should differ from unity by something like α^2 and so our experimental difference of -1.40×10^{-4} is of the expected order of magnitude.

For O-atom the most important relativistic contribution to the orbital g-value comes from the "relativistic increase of mass." This has a very simple form. It is first, necessarily diamagnetic in that it decreases the value of g_L . In order to calculate this term one only needs to know the average kinetic energy of the orbitally unpaired electrons. For O-atoms, this quantity has been evaluated¹⁸ and one can

show that this term alone should change g_L by -1.3×10^{-4} . This is very close to the total change in g_L and the other terms must either not be important, or more likely, they partly cancel.

Our value for g_r differs from Miller's value of $-1.234 \pm 0.025 \times 10^{-4}$. We are clearly outside of Miller's estimate of his experimental error. It is quite remarkable that we can agree so well for g_L and still not agree for g_r . One can see from Eqs.(8) and (9) that the value of g_L is primarily dependant upon the value of g_J for $J = 2$ and g_r comes mainly from a comparison of the g_J values for both $J = 2$ and $J = 3$. In fact, g_r depends upon how close the g_J value is for $J = 3$ to one-half of the value for $J = 2$.

In Eq.(9) if we assume that our value of g_L is correct but that Miller's value for g_r is correct, then g_J for $J = 3$ would have to be decreased by $.31 \times 10^{-4}$. This is a $10^{-2}\%$ error in g_J or a 0.6 G error in our magnetic field measurement. Since Miller's measurements were made at X-band it would require an even larger error in his field measurement. These errors are about the size of the shift between the NMR probe and the sample, but this shift was corrected for in both sets of experiments. One can conclude that one possible source for the difference between the two g_r values would be an error in the field measurements for the $J = 3$ lines.

The ratio of the two values of g_r is close to 4:3. Our value of g_r is defined in the common manner so that $\bar{\mu}_r = g_r \mu_B \bar{N}$, where \bar{N} is the rotational angular momentum. The same definition was used for $O_2(^3\Sigma_g^-)$ by Bowers, Kamper and Lustig¹⁹ who obtained $g_r = -1.2 \times 10^{-4}$.

This is identical to Miller's value for g_r in $O_2(^1\Delta_g)$. Miller points out that his value for g_r gives equal electronic contributions to g_r for both the $^1\Delta_g$ and $^3\Sigma_g^-$ electronic states. This is a remarkable result since the perturbation formulae for the electronic part of g_r depend upon different excited electronic states for the $^1\Delta_g$ and $^3\Sigma_g^-$ states. It is to be expected that both values for $\mu_B g_r / \mu_N$ should lie between zero and -1, but one would not expect them to be as close as Miller's result indicates.

At some future date it should be possible for us to make S-band measurements on the $J = 4$ lines and thus to have an over-determined set of equations to solve for g_L and g_r . The possibility of a numerical error in our calculations was eliminated when Miller²⁰ checked our data with his least-squares programs. He obtained very nearly the same values as we quote for g_r and g_L using his programs on our data. Under these circumstances it is particularly clear that further work should be done in order to determine the exact source of our discrepancy.

Acknowledgements

We thank James J. Chang for assistance with several of the measurements. This work was supported by a grant from the U. S. Atomic Energy Commission.

References

1. A. M. Falick, B. H. Mahan and R. J. Myers, J. Chem. Phys. 42, 1837 (1965).
2. T. A. Miller, private communication, J. Chem. Phys. 54, 330 (1971).
3. F. C. Fehsenfeld, K. M. Evenson and H. P. Broida, Rev. Sci. Inst. 36, 294 (1965). We used cavity 5.
4. Obtained from Yeda Research and Development Co. of the Weizmann Institute of Science, Rehovoth, Israel.
5. A. M. Falick and R. J. Myers, Rev. Sci. Inst. 40, 1349 (1969).
6. A. H. Maki and R. J. Volpicelli, Rev. Sci. Inst. 36, 325 (1965).
7. Z. Luz and R. G. Shulman, J. Chem. Phys. 43, 3750 (1965).
8. A. Bauder and R. J. Myers, J. Molec. Spectr. 27, 110 (1968).
9. E. U. Condon and G. H. Shortley, "The Theory of Atomic Spectra," Cambridge University Press, Cambridge (1951).
10. J. H. Van Vleck, Rev. Mod. Phys. 23, 213 (1951); R. A. Frosch and H. M. Foley, Phys. Rev. 88, 1337 (1952).
11. A. Carrington, D. H. Levy and T. A. Miller, "Advances in Chemical Physics," Vol. XVIII, p. 149, I. Prigogine and S. A. Rice, Eds., Interscience Publ. (1970), New York; see also T. A. Miller, Mol. Phys. 16, 105 (1969).
12. T. Amano, E. Hirota and Y. Morino, J. Phys. Soc. Jap. 22, 399 (1967); see also T. A. Miller, J. Chem. Phys. 54, 1658 (1971).
13. H. F. Schaeffer, R. A. Klemm and F. E. Harris, Phys. Rev. 176, 49 (1968).
14. J. S. M. Harvey, Proc. Roy. Soc. (London) A285, 581 (1965).


15. S. L. Miller, C. H. Townes and M. Kotani, Phys. Rev. 90, 542 (1952).
16. L. Herzberg and G. Herzberg, Astrophys. J. 105, 353 (1947).
17. V. W. Hughes in Recent Research in Molecular Beams, I. Esterman Ed., Academic Press, New York (1959), p. 65.
18. A. Abragam and J. H. Van Vleck, Phys. Rev. 91, 1448 (1953);
F. R. Innes and C. W. Ufford, Phys. Rev. 111, 194 (1958).
19. K. D. Bowers, R. A. Kamper and C. P. Lustig, Proc. Roy. Soc. 251A, 565 (1959).
20. T. A. Miller, private communication. We would like to thank Dr. Miller for reducing our data with his programs.

Figure Captions

Fig. 1. The magnetic splittings for $^{16}\text{O}^{17}\text{O}$ with $J = 2$. The splittings are shown accurate to second-order perturbations. For a Δ term $\Lambda = 2$. The arrow shows a typical EPR transition.


Fig. 2. A least squares fit to the $J = 2$ lines at 3GHz. The error curve below has a scale expansion of 1.661. Each line was assumed to have a Lorentz line shape but arbitrary width, intensity and position.

Energy levels for $O_2 (^1\Delta_g), J=2$



MUB13098

Fig. 1.



XBL 701-202

ERRORS * 1.661

Fig. 2.