

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### Title

CALIFORNIUM ISOTOPES FROM BOMBARDMENT OF URANIUM WITH CARBON IONS

### Permalink

<https://escholarship.org/uc/item/4117x8kc>

### Authors

Ghiorso, A.  
Thompson, S.G.  
Street, K. Jr.  
[et al.](#)

### Publication Date

2008-06-25

UNIVERSITY OF CALIFORNIA - BERKELEY

DECLASSIFIED

UCLL-895

TWO-WEEK LOAN COPY  
*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

RADIATION LABORATORY

UCLL-895  
C.D.

UNIVERSITY OF CALIFORNIA  
Radiation Laboratory

Contract No. W-7405-eng-48

**DECLASSIFIED**

Californium Isotopes from Bombardment of Uranium with Carbon Ions

A. Ghiorso, S. G. Thompson, K. Street, Jr., and G. T. Seaborg

September 6, 1950

~~CONFIDENTIAL~~  
**CAUTION**  
This document contains information affecting the  
National Defense of the United States.  
Its transmission or the disclosure of its contents in  
any manner to an unauthorized person is prohibited and  
may result in severe criminal penalties under applicable  
Federal Laws.

Berkeley, California

~~CONFIDENTIAL~~

Standard Distribution	Copy Nos.
Argonne National Laboratory	1-10
Atomic Energy Commission, Washington	11-12
Brookhaven National Laboratory	13-16
Carbide and Carbon Chemicals Division (K-25 Plant)	17-18
Carbide and Carbon Chemicals Division (Y-12 Plant)	19
General Electric Company, Richland	20-25
Hanford Operations Office	26
Iowa State College	27
Kellex Corporation	28
Knolls Atomic Power Laboratory	29-32
Los Alamos	33-35
Mound Laboratory	36-38
Naval Radiological Defense Laboratory	39
NEPA Project	40
New York Operations Office	41-42
Oak Ridge National Laboratory	43-48
Patent Branch, Washington	49
Technical Information Division, ORE	50-64
UCLA Medical Research Laboratory (Warren)	65
University of California Radiation Laboratory	66-68
University of Rochester	69-70
Total	70

Information Division  
Radiation Laboratory  
University of California  
Berkeley, California

CALIFORNIUM ISOTOPES FROM BOMBARDMENT OF URANIUM WITH CARBON IONS

A. Ghiorso, S. G. Thompson, K. Street, Jr., and G. T. Seaborg  
Radiation Laboratory and Department of Chemistry  
University of California, Berkeley, California

September 6, 1950

DECLASSIFIED

The recent production and identification<sup>1</sup> of isotopes of elements with atomic numbers up to six higher than the target element through bombardment with hexapositive 120-Mev carbon ions made it seem worthwhile to apply this technique to the transuranium region.

Accordingly, small pieces of natural uranium metal (about 0.5 mil thick and 2.5 cm by 0.6 cm area) were irradiated in the internal carbon ion beam in the Berkeley 60-inch cyclotron. Following the irradiations, the uranium was dissolved in dilute hydrochloric acid containing hydrogen peroxide and a transplutonium fraction was isolated through the use of lanthanum fluoride, and lanthanum hydroxide precipitation steps followed by the ion exchange adsorption column procedure in which concentrated hydrochloric acid is used to separate the tripositive actinide elements from the rare earth elements.<sup>2</sup>

The transplutonium fractions in hydrochloric acid were evaporated as weightless films on platinum plates which were placed in the ionization chamber of the 48 channel pulse analyzer apparatus in order to measure the yield and energies of any alpha-particles which might be present. In the best experiment at about one hour after the end of the 90-minute bombardment, some 50 disintegrations per minute of the distinctive 7.1-Mev alpha-particles<sup>3</sup> of Cf<sup>244</sup> were observed to be present and to decay with the 45-minute half-life. The Cf<sup>244</sup> was presumably formed by the reaction  $U^{238}(C^{12},6n)$ .

After the decay of the alpha-particles due to Cf<sup>244</sup>, about five disintegrations per minute of alpha-particles with 6.8-Mev energy was observed and

this alpha-radioactivity decayed with a half-life of about 35 hours. A consideration of the systematics of alpha-radioactivity<sup>4</sup> leads us to the view that this activity is due to the new isotope Cf<sup>246</sup> formed by the reaction U<sup>238</sup>(C<sup>12</sup>,4n). The measured half-life agrees with the expected alpha half-life for the observed energy for an even-even isotope of the element with atomic number 98.

It is not possible at this time to obtain a good estimate of the intensity of the carbon ion beam. Therefore, the cross sections for the reactions given above cannot be calculated. However, it is interesting to note that the indicated yields for the (C,6n) and (C,4n) reactions are comparable.

If the isotopic assignment of the new 35-hour activity to Cf<sup>246</sup> is correct, these new data give a better idea as to the slope of the alpha-energy vs. mass number line for californium, which in turn makes it possible to make better predictions of the radioactive properties of the nuclides in this region.

We wish to express our appreciation to Professor J. G. Hamilton, J. F. Miller, G. B. Rossi, T. M. Putnam, Jr., M. T. Webb, and the operating crew of the 60-inch cyclotron in the Crocker Laboratory for their help in the carbon ion bombardments. We would also like to thank Mr. E. K. Hulet for his help in the chemical separations. This work was performed under the auspices of the U. S. Atomic Energy Commission.

---

<sup>1</sup>Miller, Hamilton, Putnam, Haymond, and Rossi, Phys. Rev. (in press).

<sup>2</sup>K. Street, Jr. and G. T. Seaborg, J. Am. Chem. Soc. 72, 2790 (1950).

<sup>3</sup>Thompson, Street, Jr., Ghiorso, and Seaborg, Phys. Rev. 78, 298 (1950).

<sup>4</sup>Perlman, Ghiorso, and Seaborg, Phys. Rev. 77, 26 (1950).