

UC Irvine

UC Irvine Previously Published Works

Title

A Vantage from Space Can Detect Earlier Drought Onset: An Approach Using Relative Humidity

Permalink

<https://escholarship.org/uc/item/4178d0tv>

Journal

Scientific Reports, 5(1)

ISSN

2045-2322

Authors

Farahmand, Alireza
AghaKouchak, Amir
Teixeira, Joao

Publication Date

2015

DOI

10.1038/srep08553

Peer reviewed



OPEN

A Vantage from Space Can Detect Earlier Drought Onset: An Approach Using Relative Humidity

Alireza Farahmand¹, Amir AghaKouchak¹ & Joao Teixeira²¹University of California, Irvine, California 92697, USA, ²Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California 91109, USA.

Each year, droughts cause significant economic and agricultural losses across the world. The early warning and onset detection of drought is of particular importance for effective agriculture and water resource management. Previous studies show that the Standard Precipitation Index (SPI), a measure of precipitation deficit, detects drought onset earlier than other indicators. Here we show that satellite-based near surface air relative humidity data can further improve drought onset detection and early warning. This paper introduces the Standardized Relative Humidity Index (SRHI) based on the NASA Atmospheric Infrared Sounder (AIRS) observations. The results indicate that the SRHI typically detects the drought onset earlier than the SPI. While the AIRS mission was not originally designed for drought monitoring, we show that its relative humidity data offers a new and unique avenue for drought monitoring and early warning. We conclude that the early warning aspects of SRHI may have merit for integration into current drought monitoring systems.

Droughts can be described and assessed using different climatic variables such as precipitation, runoff and soil moisture¹. For example, a meteorological drought is often described as a deficit in precipitation, an agricultural drought is expressed as a deficit in soil moisture, whereas a hydrological drought typically refers to below average surface or sub-surface water². Given that droughts can be described relative to different variables, numerous drought indices have been developed based on one or more climatic variables^{3,4}. For example, the Standardized Precipitation Index (SPI^{5,6}) is widely used as an indicator of meteorological drought, while the Standardized Soil Moisture Index (SSI⁷) and soil moisture percentiles have been used for agricultural drought monitoring. A number of multivariate or multi-index indicators have also been developed such as the Joint Deficit Index⁸ and the Multivariate Standardized Drought Index⁹.

Drought monitoring indices show substantial variation in their ability to detect drought onset and termination^{10,11}. Generally, precipitation measures detect drought onset earlier than other variables such as soil moisture and runoff^{9,12} because those variables have a delayed response to precipitation deficits. Consequently, the SPI detects the drought onset earlier than the SSI and soil moisture percentiles, and is thus more suitable for drought onset detection¹². This study explores whether even earlier drought onset detection can be accomplished by factoring in the meteorological variables that influence precipitation. It is hypothesized that near surface air relative humidity (hereafter, relative humidity) can detect drought onset earlier than indications provided by precipitation signals. Relative humidity is an important climate variable defined as the ratio of air vapor pressure to the saturated vapor pressure. Precipitation and relative humidity are related to each other in the sense that precipitation is not expected at low relative humidity¹³.

Limitations in ground-based observations¹⁴ make satellite observations important for monitoring drought-related variables^{15–18}. These limitations include uneven distribution of ground-based observations, temporal inconsistencies and spatial inhomogeneity in the records, and lack of observations in remote regions¹⁴. The Evaporative Stress Index¹⁵, the Global Terrestrial Drought Severity Index³, and the Global Integrated Drought Monitoring and Prediction System (GIDMaPS¹⁹) all highlight the value of remote sensing observations for monitoring drought.

We show that drought onset can be detected by standardizing relative humidity data via the relative humidity from the Atmospheric Infrared Sounder (AIRS²⁰) satellite mission. Importantly, this detection can be earlier than that indicated by measures of precipitation and soil moisture. The mission's Version 6 data sets are obtained from two instruments: The Atmospheric Infrared Sounder (AIRS) and the Advanced Microwave Sounding Unit

SUBJECT AREAS:

HYDROLOGY

CLIMATE AND EARTH SYSTEM
MODELLING

Received

13 May 2014

Accepted

21 January 2015

Published

25 February 2015

Correspondence and requests for materials should be addressed to A.A. (amir.a@uci.edu)



(AMSU). AIRS is an infrared spectrometer and radiometer with 2378 spectral channels ranging 3.7–15 μm . AMSU is a 15-channel microwave radiometer covering 23 to 89 GHz^{20,21}. AIRS's monthly surface relative humidity (over equilibrium phase) is utilized for drought onset detection (Version 6, Level 3 data). The relative humidity data are available globally at a 1° spatial resolution (2002–present). AIRS products are available from ascending and descending tracks, which refer to the direction of movement of the sub-satellite point in the satellite track. We used the descending AIRS data, in which the direction of the movement is from Northern Hemisphere to Southern Hemisphere, with an equatorial crossing time of 1:30 AM local time²². To evaluate drought detection using relative humidity, the SPI and SSI data from GIDMaPS¹⁹ are used as additional indicators.

Typically, drought onset assessment is based on a certain drought threshold. In this study, the D0-Drought (Abnormally Dry²³) condition is used as the drought onset threshold, which corresponds to a drought event with an approximately 30% probability of occurrence. As an example, the global SPI, SSI and SRHI maps for August 2010 are presented in Figure 1a, Figure 1b, and Figure 1c respectively. As shown, all three indices captured the Russian drought. This event and its accompanying heat waves resulted in thousands of casualties and significant economic losses in Russia and eastern Europe²⁴. The Amazon drought was another major event in 2010, which led to substantial water level decreases in major Amazon tributaries²⁵. At the other extreme, August 2010 was abnormally wet in eastern Australia. These patterns of wet and dry conditions are reflected

on all three indices. Overall, Figure 1 illustrates that SRHI is consistent with SPI at wet and dry conditions, though there are discrepancies primarily around neutral condition (SPI and SRHI around 0).

To analyze drought early detection, we investigated time series of the SPI, SSI and SRHI over three major drought events: the 2010 Russian drought (Figure 2a), the 2010–2011 Texas-Mexico drought (Figure 2b), and the 2012 United States drought (Figure 2c). As we show, in the 2010 Russian drought, the SRHI indicates the onset nearly two months before both the SPI (Figure 2a - compare indices relative to the D0 threshold identified by the green horizontal line). Note that for a more severe drought condition (e.g., a lower threshold of -1), the SRHI detects the drought's onset even earlier.

The same drought indicators over one location in the Texas-Mexico Drought are displayed in Figure 2b. This series confirms that the SRHI identifies onset of this drought earlier than the other indicators. Finally, Figure 2c shows the SRHI, SPI and SSI over a specific location in an area affected by the 2012 United States drought. The 2012 drought was one of the most devastating events in the modern times and led to billions of U.S. dollars in economic damage. This event in particular affected crop development and early detection could have reduced agricultural losses²⁶. As shown in Figure 2c, the SRHI detects the onset of this drought 3 to 4 months earlier than SPI. Such early detection in the growing season could potentially reduce the effects of droughts on agriculture and society²⁶.

In the top three panels in Figure 2, the SRHI is the earliest drought detector, followed by SPI and SSI. The results are consistent with previous studies indicating that SPI detects the drought onset earlier




Figure 1 | Global Standardized Precipitation Index (SPI), Standardized Soil Moisture Index (SSI), and Standardized Relative Humidity Index (SRHI) for August 2010. This map was generated using MATLAB.




Figure 2 | Time series of 3-month SPI, SSI and SRHI for several locations in areas affected by the 2010 Russian drought, 2010–2011 Texas-Mexico drought, and 2012 United States drought.

than SSI. However, the results also show that remotely sensed relative humidity can be used for even earlier drought detection. While the SRHI does show the potential to advance drought early detection, in some cases it may not detect the drought onset earlier than the SPI (e.g., see Figure 2d where the SRHI detects the drought onset later than the SPI). Nonetheless, in all four cases shown in Figure 2, the

SRHI is consistent with the SPI and SSI on showing the drought signal.

To assess the potential capability of AIRS relative humidity data in drought detection, we statistically evaluated the global SRHI values against SPI during 2002–2013 period. Figure 3a presents the probability of drought detection. Figure 3b and Figure 3c shows the false




Figure 3 | Probability of drought detection (a), false drought ratio (b), and missed drought ratio (v) for the SRHI relative to SPI. This map was generated using MATLAB.

drought ratio and missed drought ratio respectively. Figure 3a shows the fraction of the reference data (i.e., negative SPI) identified correctly by the SRHI (perfect score = 1), whereas Figure 3b describes the fraction of drought events identified by SRHI, but not confirmed with the SPI (perfect score = 0)²⁷. Figure 3c displays the fraction of drought events identified by SPI, but missed in SRHI (perfect score = 0). Given that there are limited number of droughts in each pixel during 2002–2013, the global statistics is derived for each 10×10 pixels to ensure the statistics is reliable.

An important question is in cases where a drought was detected by both SRHI and SPI, what fraction of events is detected earlier by SRHI. To answer this question, the drought onset based on SRHI (DO_{SRHI}) is evaluated against that of SPI (DO_{SPI}). To avoid unreliable statistics, only drought events longer than three months have been considered. Figure 4a shows the probability of drought detection (i.e., fraction of detected drought) when $DO_{SRHI} \leq DO_{SPI}$. As shown, in most parts of the globe this fraction ranges between 0.5 to 0.8, with the global average being approximately 0.6 (i.e., 60% of all events). Figure 4b displays the mean lead time for each pixel based on SRHI relative to SPI. The figure indicates that the mean lead time ranges between 1 to 3 months with the global average being approximately 1.9 months. The results presented in Figures 3 and 4 do not show a strong regional/geographical pattern. This indicates that in most

parts of the world the SRHI, combined with other indicators, can potentially improve early drought detection.

The SRHI's main limitation is the relatively short length of record (2002–present). However, there are other data sets with similar length of record that provide valuable drought information (e.g., GRACE observations, and Evaporative Stress Index data). SRHI can provide valuable information on current conditions but it cannot be used to put an extreme event in historical perspective. In a recent study, a Bayesian algorithm is proposed for combining multiple precipitation data to create a long-term climate data record¹⁶. Similar algorithms could be used to extend AIRS relative humidity data by combining it with reanalysis data sets (e.g., Modern-Era Retrospective Analysis for Research and Applications²⁸). Efforts are underway to create a long-term and real-time relative humidity data set for drought monitoring and assessment. On the other hand, the current resolution (1°) of the relative humidity data only allows regional to continental scale drought assessment.

We believe that drought monitoring should be based on multiple sources of information. The proposed SRHI is not meant to replace the currently available indicators. Rather, it should be used alongside other drought indicators. This paper does not claim that SRHI alone is always sufficient for early drought detection. In fact, previous studies highlight the limitations of individual drought indicators^{11,29}.




Figure 4 | Probability of drought detection (i.e., fraction of detected drought) when Drought Onset (DO) based on SRHI is less or equal to that of SPI ($DO_{SRHI} \leq DO_{SPI}$) (a), mean lead time based on SRHI relative to SPI (months) (b). This map was generated using MATLAB.

Having an additional source of information based on relative humidity can improve our understanding of the drought onset and development. Precipitation, relative humidity, water vapor, temperature and vapor pressure deficit are closely related. For this reason, satellite-based temperature, water vapor and vapor pressure deficit information can also be explored for further improving drought onset detection. Moreover, several studies argue that statistical seasonal drought prediction is very sensitive to the initial meteorological and land-surface conditions^{30–32}. Early drought detection can potentially lead to improvements in statistical seasonal drought prediction by providing additional information on the initial meteorological conditions. This issue, however, requires more in-depth research in the future.

Drought early onset detection is fundamental to local and regional mitigation plans, especially in the agriculture and water resources sectors. A water manager may need drought information months in advance for water resource planning, while for a farmer even few weeks of lead time is very important. Early detection, even by few weeks/months, allows farmers and local agencies to take adaptive measures that include purchasing less fertilizer and increasing insurance coverage, especially before or early in the growing season. The results highlight that the AIRS near surface air relative humidity data can potentially be used for drought early warning if integrated into currently available systems such as the U.S. Drought Monitor²³ or GIDMaPS¹⁹.

Methods

Standardized drought indices are often derived by normalization after fitting a parametric distribution function to the data⁵. However, a single parametric distribution may not fit data from different climatic regions³³. In this study, the Standardized Relative Humidity Index (SRHI) is proposed using a non-parametric standardization approach.

First, the empirical probabilities of the AIRS relative humidity data are computed for each grid, using the empirical Gringorten plotting position³⁴:

$$p(RH_i) = \frac{i - 0.44}{n + 0.12} \quad (1)$$

Where i is the rank of relative humidity (RH) data from the smallest, and n is the sample size. In this study, an empirical approach is used to avoid any assumption on the underlying distribution function of relative humidity data across space³⁵. The empirical probabilities of relative humidity ($p(RH_i)$) is then standardized as:

$$SRHI = \Phi^{-1}(p(RH_i)) \quad (2)$$

Where Φ^{-1} is the inverse standard normal distribution function with the mean of zero and standard deviation of one. Here, the standardization is based on the following approximation^{36,37}:

$$SRHI = \begin{cases} -\left(t - \frac{C_0 + C_1 t + C_2 t^2}{1 + d_1 t + d_2 t^2 + d_3 t^3}\right) & \text{if } 0 < p(RH_i) \leq 0.5 \\ +\left(t - \frac{C_0 + C_1 t + C_2 t^2}{1 + d_1 t + d_2 t^2 + d_3 t^3}\right) & \text{if } 0.5 < p(RH_i) \leq 1 \end{cases} \quad (3)$$

where $c_0 = 2.515517$; $c_1 = 0.802583$; $c_2 = 0.010328$; $d_1 = 1.432788$; $d_2 = 0.189269$; $d_3 = 0.001308$; and

$$t = \begin{cases} \sqrt{\ln \frac{1}{p(RH_i)^2}} & \text{if } 0 < p(RH_i) \leq 0.5 \\ \sqrt{\ln \frac{1}{(1 - p(RH_i))^2}} & \text{if } 0.5 < p(RH_i) \leq 1 \end{cases} \quad (4)$$

A negative SRHI is an indication of below average (climatology) relative humidity, and is proposed as a measure of dryness. One attractive feature of SRHI is that, similar to SPI, it can be derived for different time-scales (e.g., 1-, 3-, 6-month SRHI). For consistency and cross-comparison, the three indicators (SRHI, SPI and SSI) are computed using the same non-parametric approach and for a 3-month time scale.

1. Dracup, J. A., Lee, K. S. & Paulson Jr, E. G. On the definition of droughts. *Water Resour Res* **16**, 297–302 (1980).
2. Wilhite, D. *Drought: A global assessment* (Routledge, 2000).



3. Mu, Q., Zhao, M., Kimball, J. S., McDowell, N. G. & Running, S. W. A remotely sensed global terrestrial drought severity index. *B Am Meteorol Soc* **94**, 83–98 (2013).
4. Wilhite, D. A. *Drought and water crises: science, technology, and management issues*, vol. 86 (CRC Press, New York, 2005).
5. McKee, T., Doesken, N. & Kleist, J. The relationship of drought frequency and duration to time scales. In *In Proceedings of the 8th Conference of Applied Climatology, Anaheim, California, United States, American Meteorological Society* (1993, January 17–22).
6. Hayes, M., Svoboda, M., Wall, N. & Widhalm, M. The Lincoln declaration on drought indices: Universal meteorological drought index recommended. *B Am Meteorol Soc* **92**, 485–488 (2011).
7. Hao, Z. & AghaKouchak, A. A nonparametric multivariate multi-index drought monitoring framework. *J Hydrometeorol* **15**, 89–101 (2014).
8. Kao, S. & Govindaraju, R. A copula-based joint deficit index for droughts. *J Hydrol* **380**, 121–134 (2010).
9. Hao, Z. & AghaKouchak, A. Multivariate standardized drought index: A parametric multi-index model. *Adv Water Resour* **57**, 12–18 (2013).
10. Keyantash, J. & Dracup, J. The quantification of drought: An evaluation of drought indices. *B Am Meteorol Soc* **83**, 1167–1180 (2002).
11. Quiring, S. M. Developing objective operational definitions for monitoring drought. *J Appl Meteor Climatol* **48**, 1217–1229 (2009).
12. Mo, K. C. Drought onset and recovery over the United States. *J Geophys Res-Atmos* **116**, D20106 (2011).
13. Manabe, S. & Wetherald, R. T. Thermal equilibrium of the atmosphere with a given distribution of relative humidity. *J Atmos Sci* **24**, 241–259 (1967).
14. Easterling, D. Global data sets for analysis of climate extremes. In *Extremes in a Changing Climate* (Springer, Dordrecht, 2013). doi: 10.1007/978-94-007-4479-0_12.
15. Anderson, M. C. *et al.* Evaluation of drought indices based on thermal remote sensing of evapotranspiration over the Continental United States. *J Climate* **24**, 2025–2044 (2011).
16. AghaKouchak, A. & Nakhjiri, N. A near real-time satellite-based global drought climate data record. *Environ Res Lett* **7**, 044037 (2012). doi:10.1088/1748-9326/7/4/044037.
17. Entekhabi, D. *et al.* The hydrosphere state (HYDROS) satellite mission: An earth system pathfinder for global mapping of soil moisture and land freeze/thaw. *IEEE Trans Geosci Remote Sensing* **42**, 2184–2195 (2004).
18. Nemani, R. *et al.* Monitoring and forecasting ecosystem dynamics using the terrestrial observation and prediction system (tops). *Remote Sens Environ* **113**, 1497–1509 (2009).
19. Hao, Z., AghaKouchak, A., Nakhjiri, N. & Farahmand, A. Global integrated drought monitoring and prediction system. *Sci Data* **1**, 140001 (2014). doi: 10.1038/sdata.2014.1.
20. Aumann, H. H. *et al.* AIRS/AMSU/HSB on the Aqua mission: Design, science objectives, data products, and processing systems. *IEEE Trans Geosci Remote Sensing* **41**, 253–264 (2003).
21. Goldberg, M. D. *et al.* AIRS near-real-time products and algorithms in support of operational numerical weather prediction. *IEEE Trans Geosci Remote Sensing* **41**, 379–389 (2003).
22. Tian, B., Manning, E., Fetzner, E., Olsen, E. & Wong, S. AIRS/AMSU/HSB version 6 level 3 product user guide. *Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA* (2013).
23. Svoboda, M. *et al.* The drought monitor. *B Am Meteorol Soc* **83**, 1181–1190 (2002).
24. Matsueda, M. Predictability of Euro-Russian blocking in summer of 2010. *Geophys Res Lett* **38** (2011).
25. Marengo, J. A., Tomasella, J., Alves, L. M., Soares, W. R. & Rodriguez, D. A. The drought of 2010 in the context of historical droughts in the Amazon region. *Geophys Res Lett* **38**, L12703 (2011).
26. Crutchfield, S. *US drought 2012: farm and food impacts. Tech. Rep., United States Department of Agriculture (USDA), Economic Research Service* (2012).
27. AghaKouchak, A. & Mehran. Extended contingency table: Performance metrics for satellite observations and climate model simulations. *Water Resour Res* **49**, 7144–7149 (2013). doi:10.1002/wrcr.20498.
28. Rienecker, M. M. *et al.* MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications. *J Climate* **24**, 3624–3648 (2011).
29. Steinemann, A. C. & Cavalcanti, L. F. Developing multiple indicators and triggers for drought plans. *J Water Resour Plan Manage* **132**, 164–174 (2006).
30. AghaKouchak, A. A baseline probabilistic drought forecasting framework using standardized soil moisture index: Application to the 2012 United States drought. *Hydrol Earth Syst Sc* **18**, 2485–2492 (2014).
31. AghaKouchak, A. A multivariate approach for persistence-based drought prediction: Application to the 2010–2011 East Africa drought. *J Hydrol* (2014) Doi: 10.1016/j.jhydrol.2014.09.063.
32. Lyon, B. *et al.* Baseline probabilities for the seasonal prediction of meteorological drought. *J Appl Meteor Climatol* **51**, 1222–1237 (2012).
33. Quiring, S. Monitoring drought: an evaluation of meteorological drought indices. *Geogr Compass* **3**, 64–88 (2009).
34. Gringorten, I. I. A plotting rule for extreme probability paper. *J Geophys Res-Atmos* **68**, 813–814 (1963).
35. Farahmand, A. & AghaKouchak, A. A Generalized Framework for Deriving Nonparametric Standardized Drought Indicators. *Adv Water Resour* **76**, 140–145 (2015). doi: 10.1016/j.advwatres.2014.11.012.
36. Edwards, D. C. *Characteristics of 20th century drought in the United States at multiple time scales. Tech. Rep., Department of Atmospheric Science, Colorado State University, Fort Collins* (1997).
37. Naresh Kumar, M., Murthy, C., Sesha Sai, M. & Roy, P. On the use of standardized precipitation index (SPI) for drought intensity assessment. *Meteorol Appl* **16**, 381–389 (2009).

Acknowledgments

This study is supported by the National Aeronautics and Space Administration (NASA) Award No. NNX15AC27G.

Author contributions

A.A. conceived the study. A.F. carried out the data analysis, and conducted the experiment. A.F., A.A. and J.T. wrote the paper.

Additional information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Farahmand, A., AghaKouchak, A. & Teixeira, J. A Vantage from Space Can Detect Earlier Drought Onset: An Approach Using Relative Humidity. *Sci. Rep.* **5**, 8553; DOI:10.1038/srep08553 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>