

Lawrence Berkeley National Laboratory

LBL Publications

Title

Grana-Like Structures of Synechococcus Cedorum

Permalink

<https://escholarship.org/uc/item/42s1w6hs>

Authors

Calvin, M

Lynch, V

Publication Date

1951-10-01

UCRL- 1502

UNCLASSIFIED

UNIVERSITY OF CALIFORNIA - BERKELEY

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

RADIATION LABORATORY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

UNCLASSIFIED

GRANA-LIKE STRUCTURES OF SYNECHOCOCCUS CEDORUM

M. Calvin and V. Lynch

October 23, 1951

Berkeley, California

GRANA-LIKE STRUCTURES OF SYNECHOCOCCUS CEDORUM

M. Calvin and V. Lynch

Radiation Laboratory and Department of Chemistry

University of California, Berkeley*

During photosynthetic studies in this laboratory, the desire for an organism possessing a simple internal structure prompted us to investigate the blue-green algae. These organisms have occasionally been reported to contain grana or chloroplasts, but the most widely accepted opinion is that all of their pigments are uniformly distributed throughout the cytoplasm.¹

A pure culture of a unicellular blue-green alga, Synechococcus cedorum, was grown in an inorganic medium and one-day old cells were used to investigate the pigment distribution.

After the cells were harvested by centrifuging and washed with water, they were broken by grinding with alumina.² The mixture was diluted with water and centrifuged at 2,000 g. for 10 minutes to remove unbroken cells, cell debris and alumina. A blue-green supernatant having a strong Tyndall effect and slight fluorescence was obtained. This supernatant was then centrifuged in a refrigerated Spinco ultracentrifuge for 30 minutes at 36,000 g. A clear blue supernatant above a very minute green sediment was obtained. The absorption spectra of the intact cells and of the several

(*) The work described in this paper was sponsored by the U. S. Atomic Energy Commission.

For publication in Nature.

fractions were determined from 3,500 Å to 7,500 Å with a Cary Recording Spectrophotometer. The curves of the aqueous solutions or suspensions are shown in Figure 1.

The suspension of the green sediment (Figure 1, Curve D) gives an absorption curve for chlorophyll and carotenoids and the clear supernatant (Figure 1, Curve C) shows the absorption spectrum for phycocyanin with no chlorophyll. Centrifugation at 36,000 g. for 20 minutes gives a complete separation of the two groups of pigments, chlorophyll and carotenoids associated with particles and phycocyanin in solution. Breaking of the cells by either ultrasonics or nitrous oxide treatment³ gave the same results as obtained by alumina grinding.

Electron micrographs of the whole cells of Synechococcus cedorum and of the green sediment (Figure 1, Curve D) obtained after breaking the cells by ultrasonics and ultracentrifuging are shown in Figure 2. The dark areas of the whole cells are believed to correspond to the particles obtained by ultracentrifuging. The polystyrene standards have a diameter of 2,600 Å and from this the diameter of the particles in the whole cells appeared to be approximately 2,200 Å. Therefore, these particles are similar in size to the grana of chloroplasts.

These results show that all of the chlorophyll and carotenoids of Synechococcus cedorum are associated with particles that will centrifuge out at 36,000 g. and that the phycocyanin is in solution. On the basis of size and chlorophyll content, these particles of Synechococcus cedorum might be considered analogous to the grana of other green plants. There is also experimental evidence to indicate that the pigments of the photosynthetic bacteria, Chlorobium⁴ and Rhodospirillum,⁵ are also aggregated in particles.

We wish to thank Dr. Marybelle Allen for the pure culture of Synechococcus cedorum and Dr. Robley Williams for taking the electron micrographs.

References

- (1) F. E. Fritsch, The Structure and Reproduction of the Algae, Vol. II, Cambridge University Press, London (1945), Pg. 781.
- (2) H. McIlwain, J. Gen. Microbiol., 2, 288 (1948).
- (3) D. Fraser, Nature, 167, 33 (1951).
- (4) A. K. Bicknell, Lloydia, 12, 183 (1949).
- (5) A. B. Pardee, H. K. Schachman, R. Y. Stanier, In Press.

Captions to Figures

Figure 1 -- Absorption spectra of Synechococcus cedorum

Curve A - Whole cells.

Curve B - Supernatant (2,000 g.) from alumina-treated cells.

Curve C - Supernatant (36,000 g.) after ultracentrifugation.

Curve D - Resuspended particles obtained from ultracentrifugation.

Figure 2 -- Electron micrographs of Synechococcus cedorum

(a) Whole cells (the polystyrene standard is shown in the lower right corner).

(b) Particles from broken cells sedimenting at 36,000 g. Corresponding to Curve D of Figure 1. (The completely opaque black circles (2) are the polystyrene standards-diameter 2,600 Å.)




Fig. 1.





Fig. 2 (a)



101