

UC Berkeley

UC Berkeley Previously Published Works

Title

Search for the W-exchange decays $B_0 \rightarrow D_s^{(*)} - D_s^{(*)+}$

Permalink

<https://escholarship.org/uc/item/4384w4mw>

Journal

Physical Review D, 72(11)

ISSN

2470-0010

Authors

Aubert, B
Barate, R
Boutigny, D
[et al.](#)

Publication Date

2005-12-01

DOI

10.1103/physrevd.72.111101

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Search for the W -exchange decays $B^0 \rightarrow D_s^{(*)-} D_s^{(*)+}$

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ M. Pappagallo,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ D. Best,⁶ A. B. Breon,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomoisky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ T. Cuhadar-Donszelmann,¹⁰ B. G. Fulsom,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ M. Saleem,¹¹ L. Teodorescu,¹¹ A. E. Blinov,¹² V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² A. N. Yushkov,¹² M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ S. Curry,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ P. Lund,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfiel,¹⁴ A. J. R. Weinstein,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ G. P. Dubois-Felsmann,¹⁹ A. Dvoretzki,¹⁹ D. G. Hitlin,¹⁹ J. S. Minamora,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ R. Andreassen,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. C. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ J. F. Hirschauer,²¹ A. Kreisel,²¹ U. Nauenberg,²¹ A. Olivas,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² Q. Zeng,²² D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ B. Spaan,²³ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,²⁵ S. Schrenk,²⁵ Ch. Thiebaut,²⁵ G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ L. Piemontese,²⁷ F. Anulli,²⁸ R. Baldini-Feroli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,^{28,*} M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² R. L. Flack,³² J. R. Gaillard,³² J. A. Nash,³² M. B. Nikolich,³² W. Panduro Vazquez,³² X. Chai,³³ M. J. Charles,³³ W. F. Mader,³³ U. Mallik,³³ V. Ziegler,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ J. I. Yi,³⁴ G. Schott,³⁵ N. Arnaud,³⁶ M. Davier,³⁶ X. Giroux,³⁶ G. Grosdidier,³⁶ A. Höcker,³⁶ F. Le Diberder,³⁶ V. Lepeltier,³⁶ A. M. Lutz,³⁶ A. Oyanguren,³⁶ T. C. Petersen,³⁶ S. Plaszczynski,³⁶ S. Rodier,³⁶ P. Roudeau,³⁶ M. H. Schune,³⁶ A. Stocchi,³⁶ G. Wormser,³⁶ C. H. Cheng,³⁷ D. J. Lange,³⁷ M. C. Simani,³⁷ D. M. Wright,³⁷ A. J. Bevan,³⁸ C. A. Chavez,³⁸ I. J. Forster,³⁸ J. R. Fry,³⁸ E. Gabathuler,³⁸ R. Gamet,³⁸ K. A. George,³⁸ D. E. Hutchcroft,³⁸ R. J. Parry,³⁸ D. J. Payne,³⁸ K. C. Schofield,³⁸ C. Touramanis,³⁸ C. M. Cormack,³⁹ F. Di Lodovico,³⁹ W. Menges,³⁹ R. Sacco,³⁹ C. L. Brown,⁴⁰ G. Cowan,⁴⁰ H. U. Flaecher,⁴⁰ M. G. Green,⁴⁰ D. A. Hopkins,⁴⁰ P. S. Jackson,⁴⁰ T. R. McMahon,⁴⁰ S. Ricciardi,⁴⁰ F. Salvatore,⁴⁰ D. N. Brown,⁴¹ C. L. Davis,⁴¹ J. Allison,⁴² N. R. Barlow,⁴² R. J. Barlow,⁴² C. L. Edgar,⁴² M. C. Hodgkinson,⁴² M. P. Kelly,⁴² G. D. Lafferty,⁴² M. T. Naisbit,⁴² J. C. Williams,⁴² C. Chen,⁴³ W. D. Hulsbergen,⁴³ A. Jawahery,⁴³ D. Kovalskyi,⁴³ C. K. Lae,⁴³ D. A. Roberts,⁴³ G. Simi,⁴³ G. Blaylock,⁴⁴ C. Dallapiccola,⁴⁴ S. S. Hertzbach,⁴⁴ R. Kofler,⁴⁴ X. Li,⁴⁴ T. B. Moore,⁴⁴ S. Saremi,⁴⁴ H. Staengle,⁴⁴ S. Y. Willocq,⁴⁴ R. Cowan,⁴⁵ K. Koeneke,⁴⁵ G. Sciolla,⁴⁵ S. J. Sekula,⁴⁵ M. Spitznagel,⁴⁵ F. Taylor,⁴⁵ R. K. Yamamoto,⁴⁵ H. Kim,⁴⁶ P. M. Patel,⁴⁶ S. H. Robertson,⁴⁶ A. Lazzaro,⁴⁷ V. Lombardo,⁴⁷ F. Palombo,⁴⁷ J. M. Bauer,⁴⁸ L. Cremaldi,⁴⁸ V. Eschenburg,⁴⁸ R. Godang,⁴⁸ R. Kroeger,⁴⁸ J. Reidy,⁴⁸ D. A. Sanders,⁴⁸ D. J. Summers,⁴⁸ H. W. Zhao,⁴⁸ S. Brunet,⁴⁹ D. Côté,⁴⁹ P. Taras,⁴⁹ F. B. Viaud,⁴⁹ H. Nicholson,⁵⁰ N. Cavallo,⁵¹ G. De Nardo,^{51,†} F. Fabozzi,⁵¹ C. Gatto,^{51,†} L. Lista,⁵¹ D. Monorchio,⁵¹ P. Paolucci,⁵¹ D. Piccolo,⁵¹ C. Sciacca,⁵¹ M. Baak,⁵² H. Bulten,⁵² G. Raven,⁵² H. L. Snoek,⁵² L. Wilden,⁵² C. P. Jessop,⁵³ J. M. LoSecco,⁵³ T. Allmendinger,⁵⁴ G. Benelli,⁵⁴ K. K. Gan,⁵⁴ K. Honscheid,⁵⁴ D. Hufnagel,⁵⁴ P. D. Jackson,⁵⁴ H. Kagan,⁵⁴ R. Kass,⁵⁴ T. Pulliam,⁵⁴ A. M. Rahimi,⁵⁴ R. Ter-Antonyan,⁵⁴ Q. K. Wong,⁵⁴ N. L. Blount,⁵⁵

J. Brau,⁵⁵ R. Frey,⁵⁵ O. Igonkina,⁵⁵ M. Lu,⁵⁵ C. T. Potter,⁵⁵ R. Rahmat,⁵⁵ N. B. Sinev,⁵⁵ D. Strom,⁵⁵ J. Strube,⁵⁵ E. Torrence,⁵⁵ F. Galeazzi,⁵⁶ M. Margoni,⁵⁶ M. Morandin,⁵⁶ M. Posocco,⁵⁶ M. Rotondo,⁵⁶ F. Simonetto,⁵⁶ R. Stroili,⁵⁶ C. Voci,⁵⁶ M. Benayoun,⁵⁷ H. Briand,⁵⁷ J. Chauveau,⁵⁷ P. David,⁵⁷ L. Del Buono,⁵⁷ Ch. de la Vaissière,⁵⁷ O. Hamon,⁵⁷ M. J. J. John,⁵⁷ Ph. Leruste,⁵⁷ J. Malclès,⁵⁷ J. Ocariz,⁵⁷ L. Roos,⁵⁷ G. Therin,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ M. Biasini,⁵⁹ R. Covarelli,⁵⁹ S. Pacetti,⁵⁹ M. Pioppi,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ F. Bucci,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ R. Cenci,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ M. Rama,⁶⁰ G. Rizzo,⁶⁰ J. Walsh,⁶⁰ M. Haire,⁶¹ D. Judd,⁶¹ D. E. Wagoner,⁶¹ J. Biesiada,⁶² N. Danielson,⁶² P. Elmer,⁶² Y. P. Lau,⁶² C. Lu,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² A. V. Telnov,⁶² F. Bellini,⁶³ G. Cavoto,⁶³ A. D'Orazio,⁶³ E. Di Marco,⁶³ R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ G. Piredda,⁶³ F. Polci,⁶³ F. Safai Tehrani,⁶³ C. Voena,⁶³ H. Schröder,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ N. De Groot,⁶⁵ B. Franek,⁶⁵ G. P. Gopal,⁶⁵ E. O. Olaiya,⁶⁵ F. F. Wilson,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ G. Graziani,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Legendre,⁶⁶ G. W. London,⁶⁶ B. Mayer,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ J. R. Wilson,⁶⁷ F. X. Yumiceva,⁶⁷ T. Abe,⁶⁸ M. T. Allen,⁶⁸ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ O. L. Buchmueller,⁶⁸ R. Claus,⁶⁸ J. P. Coleman,⁶⁸ M. R. Convery,⁶⁸ M. Cristinziani,⁶⁸ J. C. Dingfelder,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ S. Fan,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ C. Hast,⁶⁸ T. Hryn'ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ A. Snyder,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ K. Suzuki,⁶⁸ S. K. Swain,⁶⁸ J. M. Thompson,⁶⁸ J. Va'vra,⁶⁸ N. van Bakel,⁶⁸ M. Weaver,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ K. Yi,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ S. A. Majewski,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ M. Ahmed,⁷⁰ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ R. Bula,⁷⁰ J. A. Ernst,⁷⁰ M. A. Saeed,⁷⁰ F. R. Wappler,⁷⁰ S. B. Zain,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² J. L. Ritchie,⁷² A. Satpathy,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ M. Bomben,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ L. Vitale,⁷⁵ V. Azzolini,⁷⁶ F. Martinez-Vidal,⁷⁶ R. S. Panvini,⁷⁷ Sw. Banerjee,^{78,‡} B. Bhuyan,⁷⁸ C. M. Brown,⁷⁸ D. Fortin,⁷⁸ K. Hamano,⁷⁸ R. Kowalewski,⁷⁸ J. M. Roney,⁷⁸ R. J. Sobie,⁷⁸ J. J. Back,⁷⁹ P. F. Harrison,⁷⁹ T. E. Latham,⁷⁹ G. B. Mohanty,⁷⁹ H. R. Band,⁸⁰ X. Chen,⁸⁰ B. Cheng,⁸⁰ S. Dasu,⁸⁰ M. Datta,⁸⁰ A. M. Eichenbaum,⁸⁰ K. T. Flood,⁸⁰ M. T. Graham,⁸⁰ J. J. Hollar,⁸⁰ J. R. Johnson,⁸⁰ P. E. Kutter,⁸⁰ H. Li,⁸⁰ R. Liu,⁸⁰ B. Mellado,⁸⁰ A. Mihalýi,⁸⁰ A. K. Mohapatra,⁸⁰ Y. Pan,⁸⁰ M. Pierini,⁸⁰ R. Prepost,⁸⁰ P. Tan,⁸⁰ S. L. Wu,⁸⁰ Z. Yu,⁸⁰ and H. Neal⁸¹

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Dipartimento di Fisica and INFN, Università di Bari, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁵University of California at Riverside, Riverside, California 92521, USA

¹⁶University of California at San Diego, La Jolla, California 92093, USA

¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁹California Institute of Technology, Pasadena, California 91125, USA

²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA

²¹University of Colorado, Boulder, Colorado 80309, USA

- ²²Colorado State University, Fort Collins, Colorado 80523, USA
²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
²⁵Ecole Polytechnique, LLR, F-91128 Palaiseau, France
²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁷Dipartimento di Fisica and INFN, Università di Ferrara, I-44100 Ferrara, Italy
²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
³⁰Harvard University, Cambridge, Massachusetts 02138, USA
³¹Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
³²Imperial College London, London, SW7 2AZ, United Kingdom
³³University of Iowa, Iowa City, Iowa 52242, USA
³⁴Iowa State University, Ames, Iowa 50011-3160, USA
³⁵Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany
³⁶Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³⁷Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁸University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁹Queen Mary, University of London, E1 4NS, United Kingdom
⁴⁰University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
⁴¹University of Louisville, Louisville, Kentucky 40292, USA
⁴²University of Manchester, Manchester M13 9PL, United Kingdom
⁴³University of Maryland, College Park, Maryland 20742, USA
⁴⁴University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁵Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴⁶McGill University, Montréal, Québec, Canada H3A 2T8
⁴⁷Dipartimento di Fisica and INFN, Università di Milano, I-20133 Milano, Italy
⁴⁸University of Mississippi, University, Mississippi 38677, USA
⁴⁹Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7
⁵⁰Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁵¹Dipartimento di Scienze Fisiche and INFN, Università di Napoli Federico II, I-80126, Napoli, Italy
⁵²NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵³University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁴Ohio State University, Columbus, Ohio 43210, USA
⁵⁵University of Oregon, Eugene, Oregon 97403, USA
⁵⁶Dipartimento di Fisica and INFN, Università di Padova, I-35131 Padova, Italy
⁵⁷Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
⁵⁸University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁹Dipartimento di Fisica and INFN, Università di Perugia, I-06100 Perugia, Italy
⁶⁰Dipartimento di Fisica, Università di Pisa, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶¹Prairie View A&M University, Prairie View, Texas 77446, USA
⁶²Princeton University, Princeton, New Jersey 08544, USA
⁶³Dipartimento di Fisica and INFN, Università di Roma La Sapienza, I-00185 Roma, Italy
⁶⁴Universität Rostock, D-18051 Rostock, Germany
⁶⁵Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁶DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁷University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁸Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁹Stanford University, Stanford, California 94305-4060, USA
⁷⁰State University of New York, Albany, New York 12222, USA
⁷¹University of Tennessee, Knoxville, Tennessee 37996, USA
⁷²University of Texas at Austin, Austin, Texas 78712, USA
⁷³University of Texas at Dallas, Richardson, Texas 75083, USA
⁷⁴Dipartimento di Fisica Sperimentale and INFN, Università di Torino, I-10125 Torino, Italy
⁷⁵Dipartimento di Fisica and INFN, Università di Trieste, I-34127 Trieste, Italy
⁷⁶IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
⁷⁷Vanderbilt University, Nashville, Tennessee 37235, USA

* Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy.

† Also with Università della Basilicata, Potenza, Italy.

‡ Deceased.

⁷⁸University of Victoria, Victoria, British Columbia, Canada V8W 3P6

⁷⁹Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

⁸⁰University of Wisconsin, Madison, Wisconsin 53706, USA

⁸¹Yale University, New Haven, Connecticut 06511, USA

(Received 20 October 2005; published 19 December 2005)

We report a search for the decays $B^0 \rightarrow D_s^- D_s^+$, $B^0 \rightarrow D_s^{*-} D_s^+$, and $B^0 \rightarrow D_s^{*-} D_s^{*+}$ in a sample of $232 \times 10^6 \Upsilon(4S)$ decays to $B\bar{B}$ pairs collected with the *BABAR* detector at the PEP-II asymmetric-energy e^+e^- storage ring. We find no significant signal and set upper bounds for the branching fractions: $\mathcal{B}(B^0 \rightarrow D_s^- D_s^+) < 1.0 \times 10^{-4}$, $\mathcal{B}(B^0 \rightarrow D_s^{*-} D_s^+) < 1.3 \times 10^{-4}$, and $\mathcal{B}(B^0 \rightarrow D_s^{*-} D_s^{*+}) < 2.4 \times 10^{-4}$ at 90% confidence level.

DOI: 10.1103/PhysRevD.72.111101

PACS numbers: 13.25.Hw

In the standard model, $B^0 \rightarrow D_s^{(*)-} D_s^{(*)+}$ decays are dominated by the W -exchange mechanism $\bar{b}d \rightarrow c\bar{c}$ as shown in Fig. 1, while the corresponding loop diagram is highly suppressed. The decay rates of W -exchange or annihilation processes are usually argued to be negligibly small due to the suppression from helicity and/or form factors [1]; however this assumption has not been well tested experimentally.

Recently, it has been pointed out that it is difficult to calculate these decay amplitudes using the factorization approach, and a perturbative QCD (pQCD) [2] model has been used to predict the branching fraction for these decays. Prediction of branching fractions from an alternative model [3] gives an estimate of nonfactorizable contributions coming from chiral loops (CL) and tree level amplitudes generated by soft gluon emission forming a gluon condensate (GC) and it differs from pQCD approach by large amounts, as shown in Table I.

It has been estimated that a CP asymmetry of the order of 10% could arise between $B^0 \rightarrow D_s^- D_s^+$ and its charge conjugate [4]. A measurement of the decay rates of $B^0 \rightarrow D_s^{(*)-} D_s^{(*)+}$ relative to those of $B^0 \rightarrow D^{(*)-} D^{(*)+}$ will provide an estimate of the W -exchange contribution to the latter decay, a crucial piece of information for extracting the CKM angle γ from $B^0 \rightarrow D^{(*)-} D^{(*)+}$ and $B^0 \rightarrow D^{(*)-} D_s^{(*)+}$ decays [5].

Using 211 fb^{-1} of data taken on the $\Upsilon(4S)$ resonance with the *BABAR* detector at the PEP-II asymmetric B factory, we report a search for $B^0 \rightarrow D_s^- D_s^+$, $B^0 \rightarrow D_s^{*-} D_s^+$, and $B^0 \rightarrow D_s^{*-} D_s^{*+}$ decays [6]. We use the D_s^{*-} decays into $D_s^- \gamma$ and D_s^- decays into $\phi \pi^-$, $K_s^0 K^-$, and

$K^{*0} K^-$. The ϕ , K_s , and K^{*0} mesons are reconstructed in their decays to $K^+ K^-$, $\pi^+ \pi^-$, and $K^+ \pi^-$, respectively.

The *BABAR* detector is described in detail elsewhere [7]. Tracking of charged particles is provided by a five-layer silicon vertex tracker (SVT) and a 40-layer drift chamber (DCH). Discrimination between charged pions and kaons relies upon ionization energy loss (dE/dx) in the DCH and SVT, and upon Cherenkov photons detected in a ring-imaging detector (DIRC). An electromagnetic calorimeter (EMC), consisting of 6580 thallium-doped CsI crystals, is used to identify electrons and photons. These detector subsystems are mounted inside a 1.5-T solenoidal superconducting magnet. Finally, the instrumented flux return of the magnet allows us to discriminate muons from other particles. We use the GEANT4 Monte Carlo (MC) [8] program to simulate the response of the detector, taking into account the varying accelerator and detector conditions.

Charged tracks used in the reconstruction of ϕ , $K^{*0}(892)^0$, and D_s meson candidates must have a distance of closest approach to the interaction point of less than 1.5 cm in the transverse plane and less than 10 cm along the beam axis. All kaon candidates must pass particle identification (PID) criteria, based on a neural-network algorithm which uses measurements of dE/dx in the DCH and the SVT, Cherenkov angles and the number of Cherenkov photons in the DIRC. No PID requirement is applied to the pion candidates. A ϕ candidate is composed of two identified kaons of opposite charge that are consistent with originating from a common vertex. We accept ϕ candidates with invariant mass $1.000 < m_{K^+ K^-} < 1.039 \text{ GeV}$. K_s^0 candidates are composed of two oppositely-charged tracks coming from a common vertex



 FIG. 1. W -exchange decay diagram for $B^0 \rightarrow D_s^{(*)-} D_s^{(*)+}$.

TABLE I. Summary of theoretical predictions of the branching fractions.

B decays	Branching fraction ($\times 10^{-5}$)	
	pQCD [2]	CL-GC [3]
$B^0 \rightarrow D_s^- D_s^+$	$7.8^{+2.0}_{-1.6}$	25.0
$B^0 \rightarrow D_s^{*-} D_s^+$	$6.0^{+1.6}_{-1.1}$	33.0
$B^0 \rightarrow D_s^{*-} D_s^{*+}$	$8.5^{+2.0}_{-1.8}$	54.0

with an invariant mass $0.491 < m_{\pi^+\pi^-} < 0.505$ GeV. $K^{*}(892)^0$ candidates are reconstructed from two oppositely-charged tracks, where one track is identified as a kaon, with an invariant mass $0.842 < m_{K^-\pi^+} < 0.942$ GeV.

We reconstruct D_s^- mesons from decays to $\phi\pi^-$, $K_s^0 K^-$, and $K^{*}(892)^0 K^-$ using tracks coming from a common vertex with a χ^2 probability greater than 0.1%. The reconstructed mass of D_s^- candidates is required to be within 2.7 to 3.0 standard deviations of the nominal mass; a typical mass resolution of D_s is about 5.1 MeV. The selected D_s candidates are then kinematically fit with their masses constrained to the nominal value [9]. In the decays $D_s^- \rightarrow \phi\pi^-$ [$K^{*}(892)^0 K^-$], the ϕ [$K^{*}(892)^0$] mesons are polarized longitudinally. Therefore the cosine of the decay angle θ_H between the direction of the K^- from ϕ [π^- from $K^{*}(892)^0$] and the D_s^- direction in the ϕ [$K^{*}(892)^0$] rest frame is expected to follow $\cos^2\theta_H$ distribution. Background events from random combinations are expected to be uniformly distributed in $\cos\theta_H$. We place a decay mode-dependent requirement on the minimum value of $|\cos\theta_H|$, which varies from 0.3 to 0.5 and rejects 13% to 24% of the combinatorial background.

D_s^{*-} candidates are formed by combining D_s^- and γ candidates with a mass difference $\Delta M = M_{D_s^{*-}} - M_{D_s^-}$ in the range of $0.125 < \Delta M < 0.160$ GeV. The photon energy measured in the EMC is required to be more than 100 MeV.

B^0 meson candidates are reconstructed by combining either (i) two oppositely charged D_s candidates, (ii) one D_s^* candidate and an oppositely charged D_s candidate, or (iii) two oppositely charged D_s^* candidates. Finally, two quantities are used to discriminate between B^0 -meson signal and background: the beam-energy-substituted mass $m_{ES} = \sqrt{E_b^{*2} - (\mathbf{p}_B^*)^2}$ and the energy difference $\Delta E = E_B^* - E_b^*$, where E_b^* is the beam energy in the center of mass (CM) frame, and \mathbf{p}_B^* (E_b^*) is the CM momentum (energy) of the B^0 -meson candidate. For signal events m_{ES} peaks at the B^0 -meson mass with a typical resolution of 2.5 MeV, dominated by the uncertainty of the beam energy, and ΔE peaks near zero indicating that the B decay candidate has a total energy consistent with the beam energy in the CM frame. Depending on the particular B^0 decay mode, the measured resolution for ΔE is 6.5 – 13.3 MeV.

Multiple candidates are found in 3% to 5% of the selected events in the three different B^0 decay modes. The best candidate in each event is selected based on the smallest χ^2 combination, where

$$\chi^2 \equiv \sum \left| \frac{m_{D_s^\pm} - \overline{m}_{D_s^\pm}}{\sigma_{m_{D_s^\pm}}} \right|^2 + \sum \left| \frac{\Delta M - \overline{\Delta M}}{\sigma_{\Delta M}} \right|^2, \quad (1)$$

and the sum is over $D_s^{(*)+}$ and $D_s^{(*)-}$ candidates participating in a particular B^0 decay. The mean values ($\overline{m}_{D_s^\pm}$ and

$\overline{\Delta M}$) are the nominal values given in Ref. [9] and the errors ($\sigma_{m_{D_s^\pm}}$, $\sigma_{\Delta M}$) are measured in a data control sample of $B^0 \rightarrow D^- D_s^{(*)+}$ decays.

A small source of remaining background is $e^+e^- \rightarrow q\bar{q}$ production, which is suppressed based on event topology. We restrict the angle (θ_T) between the thrust axis [10] of the B^0 meson candidate and the thrust axis of the rest of the particles in the event. In the CM frame, $B\bar{B}$ pairs are produced approximately at rest and form a nearly uniform distribution in $|\cos\theta_T|$. In contrast, hadrons in $q\bar{q}$ events are produced back-to-back in two jets, which results in a $|\cos\theta_T|$ distribution peaked at 1. Based on the background level of each mode, we require the value of $|\cos\theta_T|$ to be less than a mode-dependent upper limit, which ranges from 0.83 and 0.9. We require $R_2 < 0.4$, where R_2 is the ratio of the second Fox-Wolfram moment to the zeroth moment [11], both determined using charged tracks and unmatched neutral showers in the event.

For different B^0 meson decays, a signal region is defined in a two dimensional scatter plane of m_{ES} and ΔE as shown in Table II. Optimization of the selection is performed separately for each of the three B^0 decays [12] by maximizing a figure of merit, $S^2/(S+B)$, where S is the number of signal events in the signal box as derived from the MC simulation and B is the number of background events estimated from simulations of generic B decays and $q\bar{q}$ continuum. We use the same selection criteria for different B^0 decay modes if the figure of merit differs by less than 10%.

After the aforementioned selection, four possible background sources are considered. First, the amount of combinatorial background in the signal region is estimated from the grand sideband region: $-0.25 < \Delta E < 0.25$ GeV and $5.20 < m_{ES} < 5.27$ GeV. The second source of backgrounds arises from B meson decays such as $B^0 \rightarrow D^{(*)-} D_s^{(*)+}$ and $B^- \rightarrow D_s^{(*)-} D^{(*)0}$. These background events have the same m_{ES} distribution as the signal, but their reconstructed energy is higher than the beam energy. Third, the cross-feed background that may arise among the six combinations of $D_s D_s$ modes and the three

TABLE II. The number of signal candidates (N_{cand}), total estimated background (N_{bkg}), efficiency from MC simulation times the branching fraction ($\epsilon_i \times \mathcal{B}$), and 90% C.L. upper limit for $B^0 \rightarrow D_s^- D_s^+$, $B^0 \rightarrow D_s^{*-} D_s^+$, and $B^0 \rightarrow D_s^{*-} D_s^{*+}$ decay modes.

	$B^0 \rightarrow D_s^- D_s^+$	$B^0 \rightarrow D_s^{*-} D_s^+$	$B^0 \rightarrow D_s^{*-} D_s^{*+}$
$\Delta E(\text{MeV})$	-18.0–18.0	-25.0–20.0	-46.0–30.0
$m_{ES}(\text{GeV})$	5.27–5.29	5.27–5.29	5.27–5.29
$\sum_i \epsilon_i \mathcal{B}$	3.51×10^{-4}	1.47×10^{-4}	0.85×10^{-4}
N_{cand}	6	4	3
N_{bkg}	3.3 ± 1.0	3.9 ± 1.2	2.3 ± 0.9
U.L.	$< 1.0 \times 10^{-4}$	$< 1.3 \times 10^{-4}$	$< 2.4 \times 10^{-4}$

B. AUBERT *et al.*

reconstructed B^0 decay modes was studied with a large sample of signal MC and the corresponding contributions were found to be small. Finally, rare B decays into the same final state particles, such as nonresonant $B^0 \rightarrow D_s^- K^0 K^+$, have the same m_{ES} and ΔE distributions as the signal. This source of background is estimated to be negligible.

Figure 2 shows the distributions of candidates for (i) $B^0 \rightarrow D_s^- D_s^+$, (ii) $B^0 \rightarrow D_s^{*-} D_s^+$, and (iii) $B^0 \rightarrow D_s^{*-} D_s^{*+}$ decays in the ΔE versus m_{ES} plane after all selection criteria have been applied. We find 6, 4, and 3 candidate events in the signal boxes that survived the selection criteria for the $B^0 \rightarrow D_s^- D_s^+$, $B^0 \rightarrow D_s^{*-} D_s^+$, and $B^0 \rightarrow D_s^{*-} D_s^{*+}$ decay processes, respectively. The combinatorial background in the signal box (N_{bkg}^{comb}) is estimated from the number of events in the grand sideband region of the data. We compute the average number of background (N_{bkg}^{avg}) within the region $E_2 < \Delta E < E_1$ GeV and $5.20 < m_{ES} < 5.27$ GeV from a fit to the ΔE distribution of the data events in the grand sideband [described well by a first order polynomial function $P(\Delta E)$] as $N_{bkg}^{avg} = N_{GSB} \times \int_{E_1}^{E_2} P(\Delta E) / \int_{-0.25}^{0.25} P(\Delta E)$, where E_1 and E_2 are the ΔE energy bounds of the signal box as shown in Table II and N_{GSB} is the total number of events in the grand sideband region. The m_{ES} projection of these background events is modeled with the threshold function [13],

$$\frac{dN}{dx} = x \sqrt{1 - x^2/E_b^{*2}} \exp[\xi(1 - x^2/E_b^{*2})], \quad (2)$$

characterized by the shape parameter ξ , the endpoint parameter E_b^* fixed at 5.289 GeV and $x = m_{ES}$. N_{bkg}^{comb} in the signal box is then estimated from N_{bkg}^{avg} scaled by a factor: $\int_{5.27}^{5.29} \frac{dN}{dx} / \int_{5.2}^{5.27} \frac{dN}{dx}$. We vary E_b^* by ± 2 MeV to include its effect in the systematic uncertainties in N_{bkg}^{comb} . The measured uncertainties due to the choice of threshold parameter ξ , endpoint parameter E_b^* , and parameter of the polynomial fit are combined in quadrature with the Poisson fluctuation of the number of events in the grand sideband to obtain the total error on N_{bkg}^{comb} . This procedure does not account for any potential backgrounds that are enhanced in the signal region. The simulation indicates that only a small component of the background from the $B^0 \rightarrow D_s^{(*)-} D^{(*)+}$ and $B^0 \rightarrow D_s^{(*)-} D^{(*)0}$ decay exhibits a peaking m_{ES} distribution. This component, N_{bkg}^{peak} , is extracted from a binned likelihood fit to the m_{ES} distribution of simulated events using a combination of the threshold function and a Gaussian. The ξ parameter in the threshold function is fixed to the value we obtained from the fit to the data grand sideband. The mean and width of the Gaussian component is fixed to the fit values obtained from $B^0 \rightarrow D_s^- D^+$ decays in the data. Uncertainties in N_{bkg}^{peak} arising from the $D_s^{(*)-} D^{(*)+}$ and $D_s^{(*)-} D^{(*)0}$ branching fractions [9] are added to its statistical error obtained from the fit. N_{bkg}^{comb}




FIG. 2. Distributions of events in the ΔE versus m_{ES} plane, for (i) $B^0 \rightarrow D_s^- D_s^+$, (ii) $B^0 \rightarrow D_s^{*-} D_s^+$, and (iii) $B^0 \rightarrow D_s^{*-} D_s^{*+}$ decays after all selection criteria are applied. The box in each plot is the signal region based on studies with MC simulation as described in the text.

and N_{bkg}^{peak} are added to obtain the total estimated background, N_{bkg} , as quoted in Table II.

We consider the following sources of systematic uncertainty for the signal efficiencies. The particle reconstruction and identification efficiencies are obtained from simulation and cross-checked and corrected using large

data control samples. This results in systematic uncertainties of (1) 0.8% per charged track; (2) 2.5% per reconstructed K_s^0 candidate; (3) 2.5% per identified charged kaon; and (4) 1.8% per reconstructed photon. The uncertainty on the number of $B\bar{B}$ events is estimated to be 1.1%. Depending on the B submodes, the error from the MC statistics is 2% to 4.5%. The systematic errors are dominated by the 13.3% relative uncertainty on $\mathcal{B}(D_s^- \rightarrow \phi \pi^-)$ [14], and 15.8% and 9.8% errors in $\mathcal{B}(D_s^- \rightarrow K_s^0 K^-)$ and $\mathcal{B}(D_s^- \rightarrow K^{*0} K^-)$ relative to $\mathcal{B}(D_s^- \rightarrow \phi \pi^-)$, respectively [9]. The uncertainty in modeling the simulation of the ΔE , $|\cos\theta_T|$, $|\cos\theta_H|$ distributions is evaluated using a ratio of the signal yield from $B^0 \rightarrow D^- D_s^{(*)+}$ data control sample and generic $B\bar{B}$ MC. Each selection requirement is varied and the resulting relative change in the ratio is assigned as the systematic error. The error due to vertexing is obtained by taking the difference in the ratio with and without the vertex requirement in the D_s candidate selection. A summary of the systematic uncertainties in signal efficiency is given in Table III. Using the measured signal efficiency ($\sum_i \epsilon_i \mathcal{B}_i$), 211 fb^{-1} on-resonance data corresponding to $N_{B\bar{B}} = (231.8 \pm 2.6) \times 10^6$, the background estimation along with the uncertainties and the observed candidate events in the signal region N_{cand} , we determine the 90% confidence level (C.L.) upper limit using the procedure given in [15]. The systematic uncertainties are included following the prescription in Ref. [16]. In all branching fraction calculations we assume equal production of $B^0\bar{B}^0$ and B^+B^- pairs at the $Y(4S)$.

The search for $B^0 \rightarrow D_s^- D_s^+$, $B^0 \rightarrow D_s^{*-} D_s^+$, and $B^0 \rightarrow D_s^{*-} D_s^{*+}$ decays yields the 90% C.L. upper limits (Table II):

$$\begin{aligned}\mathcal{B}(B^0 \rightarrow D_s^- D_s^+) &< 1.0 \times 10^{-4}, \\ \mathcal{B}(B^0 \rightarrow D_s^{*-} D_s^+) &< 1.3 \times 10^{-4}, \\ \mathcal{B}(B^0 \rightarrow D_s^{*-} D_s^{*+}) &< 2.4 \times 10^{-4}.\end{aligned}$$

In conclusion, we have performed a measurement of the decay rates for $B^0 \rightarrow D_s^- D_s^+$, $B^0 \rightarrow D_s^{*-} D_s^+$, and $B^0 \rightarrow D_s^{*-} D_s^{*+}$ processes with a sensitivity needed to test the standard model prediction [17]. Our upper limits disfavor the branching fraction predictions in Ref. [3] for all three B^0 decays and accommodate the predictions of the pQCD

TABLE III. Summary of systematic uncertainties for signal efficiencies.

Systematics	$D_s^- D_s^+$ (%)	$D_s^{*-} D_s^+$ (%)	$D_s^{*-} D_s^{*+}$ (%)
Tracking eff.	4.3	4.3	4.3
K_s eff.	2.7	2.7	2.7
Kaon PID	9.2	9.2	9.2
Photon eff.	...	1.8	3.6
B counting	1.1	1.1	1.1
MC statistics	2.0	3.5	4.5
$D_s^{(*)}$ b.f.	26.0	26.0	26.0
Selection	5.4	5.4	6.0
Total	28.7	28.8	29.3

calculation [2] for all three B^0 decay modes. The possible existence of a significant W -exchange component in $B^0 \rightarrow D^- D^+$ [18] decays is not confirmed in this analysis.

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the U.S. Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), Institute of High Energy Physics (China), the Commissariat à l'Énergie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Science and Technology of the Russian Federation, and the Particle Physics and Astronomy Research Council (United Kingdom). Individuals have received support from CONACyT (Mexico), the A.P. Sloan Foundation, the Research Corporation, and the Alexander von Humboldt Foundation.

-
- [1] H. Fritzsch and P. Minkowski, Phys. Lett. **90B**, 455 (1980); D. Fakirov and B. Stech, Nucl. Instrum. Methods Phys. Res., Sect. B **133**, 315 (1978).
[2] Ying Li *et al.* J. Phys. G **31**, 273 (2005).
[3] J. O. Eeg *et al.* Eur. Phys. J. C **42**, 29 (2005).
[4] B. Blok *et al.*, Phys. Rev. Lett. **78**, 3999 (1997).
[5] A. Datta and D. London, Phys. Lett. B **584**, 81 (2004).
[6] Inclusion of charge conjugate modes is implied throughout this paper.
[7] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
[8] S. Agostinelli *et al.* (GEANT4 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
[9] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B

B. AUBERT *et al.*

PHYSICAL REVIEW D **72**, 111101 (2005)

- 592**, 1 (2004).
- [10] E. Farhi, Phys. Rev. Lett. **39**, 1587 (1977).
 - [11] G. C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
 - [12] pQCD predictions for the branching fractions are assumed for the selection optimization.
 - [13] H. Albrecht *et al.* (ARGUS Collaboration), Z. Phys. C **48**, 543 (1990).
 - [14] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **71**, 091104(R) (2005).
 - [15] R. Barlow, Comput. Phys. Commun. **149**, 97 (2002).
 - [16] R. D. Cousins and V. L. Highland, Nucl. Instrum. Methods Phys. Res., Sect. A **320**, 331 (1992).
 - [17] While this paper was being written, the Belle Collaboration released an upper limit on the decay rate for $B^0 \rightarrow D_s^- D_s^+$ consistent with our result: K. Abe *et al.* hep-ex/0508040.
 - [18] G. Majumder *et al.* (Belle Collaboration), Phys. Rev. Lett. **95**, 041 803 (2005).