

# UC Davis

## UC Davis Previously Published Works

### Title

Catalyst-Controlled Regiodivergence in Rearrangements of Indole-Based Onium Ylides

### Permalink

<https://escholarship.org/uc/item/45j740zt>

### Journal

Journal of the American Chemical Society, 143(24)

### ISSN

0002-7863

### Authors

Nair, Vaishnavi N  
Kojasoy, Volga  
Laconsay, Croix J  
[et al.](#)

### Publication Date

2021-06-23

### DOI

10.1021/jacs.1c00283

Peer reviewed



# HHS Public Access

Author manuscript

*J Am Chem Soc.* Author manuscript; available in PMC 2022 June 23.

Published in final edited form as:

*J Am Chem Soc.* 2021 June 23; 143(24): 9016–9025. doi:10.1021/jacs.1c00283.

## Catalyst-Controlled Regiodivergence in Rearrangements of Indole-Based Onium Ylides

**Vaishnavi N. Nair,**

Department of Biochemistry, The University of Texas Southwestern Medical Center, Dallas, Texas 75390-9038, United States

**Volga Kojasoy,**

Department of Chemistry, University of California, Davis, Davis, California 95616, United States

**Croix J. Laconsay,**

Department of Chemistry, University of California, Davis, Davis, California 95616, United States

**Wang Yeuk Kong,**

Department of Chemistry, University of California, Davis, Davis, California 95616, United States

**Dean J. Tantillo,**

Department of Chemistry, University of California, Davis, Davis, California 95616, United States

**Uttam K. Tambar**

Department of Biochemistry, The University of Texas Southwestern Medical Center, Dallas, Texas 75390-9038, United States

### Abstract

We have developed catalyst-controlled regiodivergent rearrangements of onium-ylides derived from indole substrates. Oxonium ylides formed *in situ* from substituted indoles selectively undergo [2,3]- and [1,2]-rearrangements in the presence of a rhodium and a copper catalyst, respectively. The combined experimental and density functional theory (DFT) computational studies indicate divergent mechanistic pathways involving a metal-free ylide in the rhodium catalyzed reaction favoring [2,3]-rearrangement, and a metal-coordinated ion-pair in the copper catalyzed [1,2]-rearrangement that recombines in the solvent-cage. The application of our methodology was demonstrated in the first total synthesis of the indole alkaloid ( $\pm$ )-sorazolone B, which enabled the

---

**Corresponding Authors Uttam K. Tambar** – Department of Biochemistry, The University of Texas Southwestern Medical Center, Dallas, Texas 75390-9038, United States; [uttam.tambar@utsouthwestern.edu](mailto:uttam.tambar@utsouthwestern.edu), **Dean J. Tantillo** – Department of Chemistry, University of California, Davis, Davis, California 95616, United States; [djtantillo@ucdavis.edu](mailto:djtantillo@ucdavis.edu).

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

The authors declare no competing financial interest.

ASSOCIATED CONTENT

Accession Codes

CCDC 2057316-2057318 and 2068211 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.


Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.1c00283>.

Experimental details, characterization data, spectral data, and computational results (PDF)

stereochemical reassignment of the natural product. Further functional group transformations of the rearrangement products to generate valuable synthetic intermediates were also demonstrated.

## Graphical Abstract



## INTRODUCTION

Molecular rearrangements are arguably some of the most effective reactions for the generation of new carbon–carbon bonds in the synthesis of complex molecules.<sup>1</sup> In recent years, advances in catalytic onium ylide rearrangements have paved the way for catalyst control of rearrangements that are traditionally unselective.<sup>2,3</sup> In this context, catalytic generation of onium ylides from diazocarbonyl compounds has served as a versatile platform for selective rearrangements (Scheme 1A). Despite many reports of catalytic onium ylide rearrangements of *aliphatic* systems,<sup>4,5</sup> only a few examples of analogous *aromatic* rearrangements are known which are limited to sulfonium (X = SR) and ammonium (X = NR<sub>2</sub>) ylides.<sup>6</sup> The challenge of developing this class of rearrangements is partially due to the energetic penalty associated with disruption of aromaticity in the sigmatropic [2,3]-rearrangement (Scheme 1A).<sup>7</sup> A pioneering study on catalytic thia-Sommelet-Hauser rearrangement was reported by Wang and co-workers in 2008.<sup>6c</sup> In recent years, examples of controlled [1,2]- and [2,3]-rearrangements of ylides in aromatic systems have appeared in the literature.<sup>8</sup> Pan and co-workers reported rearrangements of sulfonium ylides where the selectivity for [1,2]- vs [2,3]-rearrangement is controlled by the solvent and substrate.<sup>8a</sup> Another report from Koenigs and co-workers shows a solvent controlled approach in rearrangements of sulfonium ylides formed from donor/acceptor carbenes.<sup>8b</sup> Alternatively, catalytic ylide-formation and aromatic [2,3]-rearrangements of oxonium systems (X = OR) are not known, presumably because of side reactions through nonylide pathways such as C–H insertion and cyclopropanation that compete with facile ylide formation.<sup>9</sup> Also, the examples for the [1,2]-rearrangement of oxonium ylides in the literature are generally limited to cyclic ylides.<sup>4b–k</sup> Catalytic methods for the selective formation of either [1,2]- or [2,3]-rearrangement products of aromatic systems from the same starting materials would provide a valuable new strategy for the synthesis of complex molecules.

Herein, we report the first catalyst-controlled regiodivergent aromatic rearrangements of indole-based oxonium ylides (Scheme 1B). With the proper choice of catalyst system, we can selectively generate the [1,2]- or [2,3]-rearrangement product. As our initial target for the rearrangements, we chose the indole-scaffold because of its prevalence in many natural products and medicinally valuable compounds.<sup>10</sup> In addition to exploring the scope of this reaction, we also performed DFT calculations to examine mechanisms and the origins of catalyst-controlled regiodivergence. Finally, to showcase the utility of our method, we demonstrated the conversion of rearrangement products into the indole alkaloid sorazolon B and several valuable building blocks for drug discovery.

## RESULTS AND DISCUSSION

### Development of Regiodivergent Rearrangements.

Our initial investigations began with the screening of various catalysts that are generally applied in carbene chemistry,<sup>11</sup> using 3-(methoxymethyl)-1-tosyl-1*H*-indole (**1a**) and benzyl diazoester (**2a**) as substrates (Table 1). While we did not observe any reactivity in the presence of palladium, silver, or gold catalysts (entries 1–3), CuOTf benzene (5 mol %) afforded a mixture of [1,2]- and [2,3]-rearrangement products **3a** and **4a** (82:18 rr), respectively, in low yield (entry 4). Moreover, we were pleased to note that indoline **4a** bearing an exomethylene moiety (presumably from the [2,3]-rearrangement) was formed with excellent diastereoselectivity (>20:1 dr). Moving forward with this initial result that gave **3a** as the major product, we began optimizing the [1,2]-rearrangement of the oxonium ylides by screening additional copper sources. Other copper catalysts such as CuCl, CuCl<sub>2</sub>, and CuOAc/NaBAR<sub>F</sub> gave similar or slightly improved yields, but relatively lower regioselectivities (entry 5–7). We were delighted to see an improved yield (38%) as well as regioselectivity (86:14 rr) with [Cu(MeCN)<sub>4</sub>]PF<sub>6</sub> as catalyst (entry 8). The use of Cu(hfacac)<sub>2</sub> to perform the rearrangement further enhanced the yield (52%) and regioselectivity (92:8 rr) (entry 9). An examination of the conversion of the starting materials under these conditions revealed that the yield was limited by the incomplete consumption of indole **1a**, whereas the diazoester **2a** was completely consumed to give the desired products along with minor amounts of dibenzyl fumarate and dibenzyl maleate as the side products resulting from homodimerization. Increasing the amount of benzyl diazoester **2a** to 2.4 equiv resulted in >95% conversion of **1a** to provide the [1,2]-rearrangement product **3a** in 78% yield and >95:5 rr (entry 10).

Alternatively, when Rh<sub>2</sub>(OAc)<sub>4</sub> (2 mol %) was used as the catalyst, we observed a switch in the regioselectivity that gave the [2,3]-rearrangement product **4a** as the major product (>95:5 rr) in 42% yield and >20:1 dr (entry 11). The screening of several other dirhodium carboxylate catalysts commonly used in metal-carbene transformations, such as Rh<sub>2</sub>(cap)<sub>4</sub>, Rh<sub>2</sub>(TFA)<sub>4</sub>, Rh<sub>2</sub>(TPA)<sub>4</sub>, and Rh<sub>2</sub>(oct)<sub>4</sub>,<sup>12</sup> failed to improve the yield for the reaction (entries 12–15). Similar to the copper-catalyzed [1,2]-rearrangement, an incomplete consumption of indole **1a** (55%) was identified as the reason for the moderate yields. However, increasing the amount of diazoester **2a** to 2.4 equiv diminished the yield of the [2,3]-rearrangement product **4a** to 26% (entry 16). We speculate that the reason for the lower yield with excess diazoester might be the propensity of the exomethylene group in **4a** to undergo

cyclopropanation with excess highly reactive rhodium-carbene (see SI for details). We also examined other indole substrates besides methyl ether **1a** for optimization studies. While ethyl ether underwent the [2,3]-rearrangement, other alkyl and arylethers such as isopropyl and phenyl ethers did not give any conversion under the [2,3]-rearrangement conditions (see SI for details). Other N-protecting groups such as Me, Boc, and Ac were also screened during optimization for [2,3]-rearrangement, but most of these substrates did not show significant product formation (see SI for details). Further screening of different solvents also did not provide improvement in the yield (entries 17–19). As a result, the conditions with Rh<sub>2</sub>(OAc)<sub>4</sub> (2 mol %) in CH<sub>2</sub>Cl<sub>2</sub> at 23 °C were identified as optimal for the catalytic ylide-formation/aromatic [2,3]-rearrangement (entry 11).

### Substrate Scope of Regiodivergent Rearrangements.

With the optimized reaction conditions for both the copper-catalyzed ylide-formation/[1,2]-rearrangement and rhodium-catalyzed ylide-formation/[2,3]-rearrangement in hand (entries 10 and 11, Table 1), we next explored reaction scope (Table 2).

Indole substrates with a broad range of substituents at various positions on the heteroaromatic ring (**3a–3p**) worked efficiently under the [1,2]-rearrangement conditions. Electronrich 5- and 6-substituted indole substrates provided high yields and excellent regioselectivities (**3b–3e**). Substitution at the 2-position generated the desired [1,2]-rearrangement product **3f**, albeit in slightly lower yield, but nonetheless gave excellent regioselectivity (>95:5 rr). Several electron-withdrawing substituents on the indole ring, including fluoro, bromo, trifluoromethyl, and ester groups (**3g–3m**), provided good yields and high regioselectivities. The reaction also progressed smoothly to generate dihalogenated product **3n** in 51% yield and >95:5 rr. In addition, other alkyl and aryl diazoesters were shown to be competent in generating [1,2]-rearrangement products in moderate to good yields (**3o** and **3p**).

Next, we explored the scope of the rhodium-catalyzed ylide-formation/[2,3]-rearrangement to provide various substituted indolines (**4**) that would be difficult to access in high selectivity by conventional methods.<sup>13</sup> Several electron-deficient indole rings with different substitution patterns underwent selective aromatic [2,3]-rearrangement to generate indolines (**4b–4f**) in moderate yields and with excellent regioselectivities. Electron-donating substituents on the indole ring led to comparatively lower isolated yields of the products (**4g** and **4h**); however, the [2,3]-rearrangement proceeded with excellent regioselectivity.<sup>14</sup> The rearrangement products were generally stable to rearomatization, presumably because of the electron-withdrawing tosyl protecting group similar to other known indolines.<sup>13b,c</sup> Notably, the [2,3]-rearrangement of all the substrates exhibited high diastereoselectivity (>20:1 dr). The relative stereochemistry of the major *anti*-diastereomer of product **4e** was confirmed by X-ray crystallography, and the relative stereochemistry of the major diastereomer of all other 2-substituted indolines was assigned by analogy.

### Mechanistic Studies.

To gain insight into the divergent mechanisms of the catalyst controlled ylide-formation/rearrangement reactions, a series of computational studies were performed using

density functional theory at the PWPB95-D3(BJ)/def2-QZVPP//IEFPCM(CH<sub>2</sub>Cl<sub>2</sub>)-B3LYP/6-31G(d), SDD level (see SI for details).<sup>15</sup> The robustness of our chosen level of theory was evaluated through a series of tests with other functionals and basis sets (see SI for details); while there is some variation in predicted relative energies, these variations do not affect our mechanistic conclusions. 3-(Methoxymethyl)-1-tosyl-1*H*-indole **1a** was selected as the model substrate.

For the rhodium-catalyzed ylide-formation/[2,3]-rearrangement, we first examined the structure of the metal-bound oxonium ylide (Figure 1). Formation of the carbon-bound ylide **5** is predicted to be exergonic by 9.2 kcal/mol, whereas formation of the oxygen-bound ylide **6** is endergonic by 7.0 kcal/mol.

Our proposed pathway for a concerted [2,3]-rearrangement process is summarized in Figure 1. Dissociation of Rh<sub>2</sub>(OAc)<sub>4</sub> prior to rearrangement generates free oxonium ylide **7**. Early dissociation of rhodium(II) complexes from ylides has been reported for other diazocarbonyl-mediated reactions.<sup>4a,16</sup> The most probable pathway to the product involves a metal-free [2,3]-rearrangement of oxonium ylide **7**, which leads to the observed product **4a** with the experimentally observed relative stereochemistry.<sup>17</sup> As expected, we were not able to find a transition state structure for the symmetry-forbidden metal-free [1,2]-rearrangement of oxonium ylide **7**.<sup>18</sup> The *relative* stereochemistry of the major diastereomer of products arising from the [2,3]-rearrangement of indole-based onium ylides such as **7** is consistent with a preference for an *exo* transition state structure (**8**).<sup>4d</sup> Relative free energies calculated for the *exo* and *endo* transition states predicted a lower energy barrier for *exo* transition state **8** leading to the observed diastereomer *anti*-**4a**.

For the copper-catalyzed ylide-formation/[1,2]-rearrangement, we propose a mechanism that involves a stepwise process (Figure 2A).<sup>4e,19</sup> The preferential formation of the [1,2]-rearrangement product **3a** over the [2,3]-rearrangement product **4a** is an argument against pathways involving the early dissociation of copper from the initially generated metal-coordinated ylide **11**, since metal-free [1,2] rearrangement is predicted to have an extremely high barrier compared to that of the [2,3]-rearrangement (see SI). On the basis of our computational results, we favor an ion-pair fragmentation/recombination pathway for the copper-catalyzed reactions.<sup>20</sup> Other possible pathways were explored but were not consistent with our experimental results (see SI for details). For example, formation of simple radical-pairs cannot be ruled out on the basis of our computational results, but our experimental data argues against it. Specifically, cyclopropane containing substrate **1q** reacted with diazoester **2a** to yield the [1,2]-rearrangement product **3q** with the radical probe intact (Figure 2B).<sup>21,22</sup>

To gain insight into the key carbon-carbon bond formation event in the copper-catalyzed reaction, we considered ion-pair complexes (**12a**, **12b**, **12c**, **12d**, and **12e**) that could lead to the [1,2]-rearrangement product **3a** or the [2,3]-rearrangement product diastereomers *syn*-**4a** and *anti*-**4a** with minimal reorganization (Figure 2A).<sup>23</sup> In principle, these ion-pairs would be in equilibrium with each other and could recombine to form copper-bound recombination products (e.g., **13a**, Figure 2A).<sup>24</sup> However, recombination in a solvent cage is expected to be faster than equilibration between ion-pairs.<sup>20g,25</sup> Although a solvent cage was not explicitly modeled in our calculations, the formation of ion-pairs in a solvent

cage is consistent with experimentally determined results. When substrates **1i** and **1r** were simultaneously subjected to the [1,2]-rearrangement conditions, we did not detect crossover products **3s** and **3a** (Figure 2C).

We were able to find a transition state structure (TS<sub>12a</sub>) converting the ion-pair **12a** to **13a**, the Cu(hfacac)<sub>2</sub>-bound experimentally observed product, with a 3.1 kcal/mol barrier. Subsequent dissociation of Cu catalyst yields **3a**. If **12a** was formed preferentially on the dissociation of the copper-coordinated oxonium ylide **11**, we propose that this ion-pair could rapidly recombine to the experimentally observed product (**12a** → **3a**) before equilibration with other ion-pairs. Comparisons of the various ion-pairs and metal-ylide intermediate **11** do indeed reveal greater conformational similarity between **11** and **12a** than either **12b**, **12c**, **12d**, or **12e** (Figure 2A; see SI for details). We also investigated the proposed stepwise ion-pair mechanism with other copper catalysts (Cu(acac)<sub>2</sub>, CuCl<sub>2</sub>, Cu(hfacac)<sup>+</sup>, and Cu(acac)<sup>+</sup>), and all qualitatively lead to similar results (see SI for details).

In summary, on the basis of our combined experimental and computational data, we favor a mechanism for the rhodium-promoted reaction where early catalyst dissociation occurs at the ylide stage, and products are formed via a metal-free [2,3]-sigmatropic rearrangement. For the copper-promoted reaction, we favor a mechanism where a copper-coordinated ion-pair is formed and rapidly recombines in a solvent cage to form the observed [1,2]-rearrangement product.

### Synthetic Applications of Regiodivergent Rearrangements.

The products generated through the [2,3]-rearrangement proved to be versatile substrates to access building blocks that are potentially useful for the synthesis of complex molecules (Scheme 2). For example, rearrangement product **4a** can undergo ozonolysis to yield indoxyl product **14**. In the presence of acid, the rearrangement product **4a** is rearomatized to furnish 2,3-disubstituted indole **15**. In the presence of an electrophilic source of bromine, it is selectively converted to 3-bromomethyl indole **16**.

To demonstrate the synthetic utility of the [1,2]-rearrangement products, we incorporated this transformation into the first total synthesis of the indole alkaloid sorazolone B, which enabled a stereochemical reassignment of the natural product's structure that was reported in the original isolation paper (Scheme 3).<sup>26</sup> To commence the total synthesis, 3-(methoxymethyl)-1-tosyl-1*H*-indole (**1a**) was coupled with diazoester **2a** under the [1,2]-rearrangement conditions to furnish benzylester **3a** in 82% yield and >95:5 rr. The efficiency of the reaction was maintained on a gram scale. A two-step procedure converted benzylester **3a** to the Weinreb amide **17**. Subsequent removal of the *N*-tosyl group provided *N*-H indole **18** in 93% yield. To access the relative configuration for the proposed structure of sorazolone B (**21**), we treated the Weinreb amide **18** first with ethynylmagnesium bromide followed by methylmagnesium bromide, which generated tertiary alcohol **19** in 19:1 dr and 57% yield over the two steps. The relative configuration of the major diastereomer, which was confirmed by X-ray crystallography, was consistent with a Cram chelation controlled addition of methylmagnesium bromide.<sup>27</sup> Alcohol **19** was then converted to diol **20**, which was subjected to gold catalyzed 6-endo cyclization.<sup>28</sup> Although the resulting

tricyclic skeleton of **21** was consistent with the proposed structure of sorazolon B, the NMR data of our synthetic sample did not match the corresponding data for the natural product.

We hypothesized that the relative configuration of the two stereogenic centers in sorazolon B may have been misassigned. To test this hypothesis, we switched the order of addition of Grignard reagents to the Weinreb amide **18**. An initial addition of methylmagnesium bromide followed by a Cram chelation controlled addition of ethynylmagnesium bromide yielded the tertiary alcohol **22** in 9:1 dr. The relative configuration of the major diastereomer was confirmed by X-ray crystallography. The treatment of methyl ether **22** with bromodimethylborane and 2-methyl-2-butene resulted in the formation of diol **23**. In the presence of Au(MeCN)SbF<sub>6</sub> and JohnPhos, diol **23** was converted to tricycle **24**, which had spectroscopic data that were identical with the data reported for sorazolon B in the original isolation paper.<sup>26</sup>

## CONCLUSION

We developed catalyst-controlled regiodivergent rearrangements of onium-ylides derived from indole methyl ethers and diazoesters. While a copper catalyst promotes a regioselective [1,2]-rearrangement, a rhodium catalyst facilitates a regioselective and diastereoselective [2,3]-rearrangement. We present experimental and computational studies that support divergent mechanistic pathways for the two rearrangement processes. We also describe the synthetic utility of the two rearrangements by demonstrating the functional group tolerance and scope of the reactions as well as the transformation of the rearrangement products to several indole-containing products. Finally, we applied the copper-catalyzed [1,2]-rearrangement in the first total synthesis of the indole alkaloid sorazolon B, which enabled the stereochemical reassignment of the natural product.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGMENTS

Financial support was provided to U.K.T by W. W. Caruth, Jr. Endowed Scholarship, Welch Foundation (I-1748), National Institutes of Health (R01GM102604), American Chemical Society Petroleum Research Fund (59177-ND1), Teva Pharmaceuticals Marc A. Goshko Memorial Grant (60011-TEV), and Sloan Research Fellowship. Financial support was provided to V.N.N by Sarah and Frank McKnight Fund Graduate Fellowship. Support from the National Science Foundation (CHE-1856416 and XSEDE via CHE-030089) to D.J.T. is gratefully acknowledged. We acknowledge Dr. Vincent Lynch (Manager of the X-ray Diffraction Lab at UT Austin) for the X-ray structural analysis and Dr. Hamid Baniyasi (Director of the UT Southwestern Metabolomics Core Facility) for the high resolution mass spectrometry. We also thank our diverse group of lab members for creating an environment that supports our scientific endeavors.

## REFERENCES

- (1) (a). Brückner R 2,3-Sigmatropic Rearrangements. In *Comprehensive Organic Synthesis*; Fleming I, Ed.; Pergamon: Oxford, 1991; pp 873–908. (b) Ilardi EA; Stivala CE; Zakarian A [3,3]-Sigmatropic rearrangements: recent applications in the total synthesis of natural products. *Chem. Soc. Rev.* 2009, 38, 3133–3148. [PubMed: 19847347] (c) Seashore-Ludlow B; Somfai P *Sigmatropic Rearrangements in Stereoselective Synthesis*; John Wiley & Sons: New York, 2013; pp 475–499. (d) Jones AC; May JA; Sarpong R; Stoltz BM *Toward a Symphony of Reactivity*:



Cascades Involving Catalysis and Sigmatropic Rearrangements. *Angew. Chem., Int. Ed.* 2014, 53, 2556–2591.(e)Molecular Rearrangements in Organic Synthesis; Rojas CM, Ed.; John Wiley & Sons: New York, 2016.

- (2). Reviews on catalytic onium ylide rearrangements: (a) Murphy GK; Stewart C; West FG Intramolecular generation and rearrangement of oxonium ylides: methodology studies and their application in synthesis. *Tetrahedron* 2013, 69, 2667–2686.(b)Hodgson DM; Pierard FYTM; Stupp PA Catalytic enantioselective rearrangements and cycloadditions involving ylides from diazo compounds. *Chem. Soc. Rev.* 2001, 30, 50–61.(c)West TH; Spoehrle SSM; Kasten K; Taylor JE; Smith AD Catalytic Stereoselective [2,3]-Rearrangement Reactions. *ACS Catal.* 2015, 5, 7446–7479.(d)Murphy GK; West FG In Molecular Rearrangements in Organic Synthesis; Rojas C, Ed.; John Wiley Sons: Hoboken, NJ, 2015; pp 497–538.(e)Sheng Z; Zhang ZK; Chu C; Zhang Y; Wang J Transition metal-catalyzed [2,3]-sigmatropic rearrangements of ylides: An update of the most recent advances. *Tetrahedron* 2017, 73, 4011–4022.
- (3) (a). Books and reviews on classical non-catalytic onium ylide rearrangements:Pine SH In *Organic Reactions*; Dauben WG, Ed.; Wiley: New York, 1970; Vol. 18 pp 403–464.(b)Nakai T; Mikami K [2,3]-Wittig sigmatropic rearrangements in organic synthesis. *Chem. Rev.* 1986, 86, 885–902. (c)Sweeney JB Sigmatropic rearrangements of ‘onium’ ylids. *Chem. Soc. Rev.* 2009, 38, 1027–1038. [PubMed: 19421580]
- (4). Examples of catalytic oxonium ylide rearrangements of aliphatic systems: (a) Doyle MP; Tamblin WH; Bagheri V Highly Effective Catalytic Methods for Ylide Generation from Diazo Compounds. Mechanism of the Rhodium- and Copper-Catalyzed Reactions with Allylic Compounds. *J. Org. Chem.* 1981, 46, 5094–5102.(b)Roskamp EJ; Johnson CR Generation and Rearrangements of Oxonium Ylides. *J. Am. Chem. Soc.* 1986, 108, 6062–6063. [PubMed: 22175385] (c)Pirrung MC; Werner JA Intramolecular Generation and [2,3]-Sigmatropic Rearrangement of Oxonium Ylides. *J. Am. Chem. Soc.* 1986, 108, 6060–6062. [PubMed: 22175384] (d)Doyle MP; Bagheri V; Harn NK Facile catalytic methods for intermolecular generation of allylic oxonium ylides and their stereoselective [2,3]-sigmatropic rearrangement. *Tetrahedron Lett.* 1988, 29, 5119–5122.(e)Eberlein TH; West FG; Tester RW The Stevens [1,2]-Shift of Oxonium Ylides: A Route to Substituted Tetrahydrofuranones. *J. Org. Chem.* 1992, 57, 3479–3482.(f)Clark JS Diastereoselective Synthesis of 2,5-Dialkyl Tetrahydrofuran-3-ones by a Copper-Catalysed Tandem Carbenoid Insertion and Ylide Rearrangement Reaction. *Tetrahedron Lett.* 1992, 33, 6193–6196.(g)Clark JS; Krowiak SA; Street LJ Synthesis of Cyclic Ethers from Copper Carbenoids by Formation and Rearrangement of Oxonium Ylides. *Tetrahedron Lett.* 1993, 34, 4385–4388.(h)West FG; Eberlein TH; Tester RW O-Bridged Medium Sized Rings *via* Bicyclic Oxonium Ylides. *J. Chem. Soc., Perkin Trans. 1* 1993, 2857–2859.(i)West FG; Naidu BN; Tester RW Profound Catalyst Effects in the Generation and Reactivity of Carbenoid-Derived Cyclic Ylides. *J. Org. Chem.* 1994, 59, 6892–6894.(j)Doyle MP; Ene DG; Forbes DC; Tedrow JS Highly Enantioselective Oxonium Ylide Formation and Stevens Rearrangement Catalyzed by Chiral Dirhodium (II) Carboxamidates. *Tetrahedron Lett.* 1997, 38, 4367–4370. (k)Li Z; Davies HML Enantioselective C-C Bond Formation by Rhodium-Catalyzed Tandem Ylide Formation/[2,3]-Sigmatropic Rearrangement between Donor/Acceptor Carbenoids and Allylic Alcohols. *J. Am. Chem. Soc.* 2010, 132, 396–401. [PubMed: 19994854] (l)Li Z; Parr BT; Davies HML Highly Stereoselective C-C Bond Formation by Rhodium-Catalyzed Tandem Ylide Formation/[2,3]-Sigmatropic Rearrangement between Donor/Acceptor Carbenoids and Chiral Allylic Alcohols. *J. Am. Chem. Soc.* 2012, 134, 10942–10946. [PubMed: 22694052] (m)Doyle MP; Forbes DC; Vasbinder MM; Peterson CS Enantiocontrol in the Generation and Diastereoselective Reactions of Catalytically Generated Oxonium and Iodonium Ylides. Metal-Stabilized Ylides as Reaction Intermediates. *J. Am. Chem. Soc.* 1998, 120, 7653–7654. (n)Li Z; Boyarskikh V; Hansen JH; Autschbach J; Musaev DG; Davies HML Scope and Mechanistic Analysis of the Enantioselective Synthesis of Allenes by Rhodium-Catalyzed Tandem Ylide Formation/[2,3]-Sigmatropic Rearrangement between Donor/Acceptor Carbenoids and Propargylic Alcohols. *J. Am. Chem. Soc.* 2012, 134, 15497–15504. [PubMed: 22924394]
- (5). Examples of catalytic ammonium, sulfonium, and iodonium ylide rearrangements of aliphatic systems: (I) *Ammonium (X = N) ylides*: (a) West FG; Glaeske KW; Naidu BN One-Step Synthesis of Tertiary  $\alpha$ -Amino Ketones And  $\alpha$ -Amino Esters from Amines and Diazocarbonyl Compounds. *Synthesis* 1993, 1993, 977–980.(b)West FG; Naidu BN New Route to Substituted Piperidines *via* the Stevens [1,2]-Shift of Ammonium Ylides. *J. Am. Chem. Soc.* 1993,

115, 1177–1178.(c)Clark JS; Hodgson PB Intramolecular Generation and Rearrangement of Ammonium Ylides from Copper Carbenoids: A General Method for the Synthesis of Cyclic Amines. *J. Chem. Soc., Chem. Commun.* 1994, 2701–2702.(d)West FG; Naidu BN Piperidines via Ammonium Ylide [1,2]-Shifts: A Concise, Enantioselective Route to (–)-Epilupinine From Proline Ester. *J. Am. Chem. Soc.* 1994, 116, 8420–8421.(e)Clark JS; Hodgson PB; Goldsmith MD; Street LJ Rearrangement of ammonium ylides produced by intramolecular reaction of catalytically generated metal carbenoids. Part 1. Synthesis of cyclic amines. *J. Chem. Soc., Perkin Trans. 1* 2001, 3312–3324.(f)Clark JS; Hodgson PB; Goldsmith MD; Blake AJ; Cooke PA; Street LJ Rearrangement of ammonium ylides produced by intramolecular reaction of catalytically generated metal carbenoids. Part 2. Stereoselective synthesis of bicyclic amines. *J. Chem. Soc., Perkin Trans. 1* 2001, 3325–3337.(g)Clark JS; Middleton MD Synthesis of Novel  $\alpha$ -Substituted and  $\alpha,\alpha$ -Disubstituted Amino Acids by Rearrangement of Ammonium Ylides Generated from Metal Carbenoids. *Org. Lett.* 2002, 4, 765–768. [PubMed: 11869122] (h)Heath P; Roberts E; Sweeney JB; Wessel HP; Workman JA Copper(II)-Catalyzed [2,3]-Sigmatropic Rearrangement of *N*-Methyltetrahydropyridinium Ylides. *J. Org. Chem.* 2003, 68, 4083–4086. [PubMed: 12737596] (i)Sançon J; Sweeney JB Probing the Effect of Allylic Substitution on Cyclic Ammonium Ylid Rearrangements. *Synlett* 2010, 2010, 664–666 Sulfonium (X = S) ylides.(j)Doyle MP; Griffin JH; Chinn MS; van Leusen D Rearrangements of Ylides Generated from Reactions of Diazo Compounds with Allyl Acetals and Thioketals by Catalytic Methods. Heteroatom Acceleration of the [2,3]-Sigmatropic Rearrangement. *J. Org. Chem.* 1984, 49, 1917–1925.(k)Aggarwal VK; Ferrara M; Hainz R; Spey SE [2,3]-Sigmatropic rearrangement of allylic sulfur ylides derived from trimethylsilyldiazomethane (TMSD). *Tetrahedron Lett.* 1999, 40, 8923–8927.(l)Carter DS; Van Vranken DL Metal-Catalyzed Ylide Formation and [2,3] Sigmatropic Rearrangement of Allyl Sulfides with Trimethylsilyldiazomethane. *Tetrahedron Lett.* 1999, 40, 1617–1620.(m)Hock KJ; Mertens L; Hommelsheim R; Spitzner R; Koenigs RM Enabling iron catalyzed Doyle-Kirmse rearrangement reactions with *in situ* generated diazo compounds. *Chem. Commun.* 2017, 53, 6577–6580 (III) Iodonium (X = I) ylides:(n)Xu B; Tambar UK Ligand-Controlled Regiodivergence in the Copper-Catalyzed [2,3]- and [1,2]-Rearrangements of Iodonium Ylides. *J. Am. Chem. Soc.* 2016, 138, 12073–12076. [PubMed: 27603216] (o)Xu B; Gartman JA; Tambar UK Copper-catalyzed [1,2]-rearrangements of allylic iodides and aryl  $\alpha$ -diazoacetates. *Tetrahedron* 2017, 73, 4150–4159. [PubMed: 28966401] (p)Xu B; Tambar UK Copper-Catalyzed Enantio-, Diastereo-, and Regioselective [2,3]-Rearrangements of Iodonium Ylides. *Angew. Chem., Int. Ed.* 2017, 56, 9868–9871.


- (6). Catalytic aromatic rearrangements with *sulfonium* (X = S) ylides: (a) Kennedy AR; Taday MH; Rainier JD The Use of Sulfur Ylides in the Synthesis of Substituted Indoles. *Org. Lett.* 2001, 3, 2407–2409. [PubMed: 11463328] (b)Novikov AV; Kennedy AR; Rainier JD Sulfur Ylide-Initiated Thio-Claisen Rearrangements. The Synthesis of Highly Substituted Indolines. *J. Org. Chem.* 2003, 68, 993–996. [PubMed: 12558426] (c)Liao M; Peng L; Wang J Rh(II)-Catalyzed Sommelet-Hauser Rearrangement. *Org. Lett.* 2008, 10, 693–696. [PubMed: 18237182] (d)Boyarskikh V; Nyong A; Rainier JD Highly Diastereoselective Sulfonium Ylide Rearrangements to Quaternary Substituted Indolines. *Angew. Chem., Int. Ed.* 2008, 47, 5374–5377.(e)Li Y; Shi Y; Huang Z; Wu X; Xu P; Wang J; Zhang Y Catalytic Thia-Sommelet-Hauser Rearrangement: Application to the Synthesis of Oxindoles. *Org. Lett.* 2011, 13, 1210–1213. [PubMed: 21306152] (f)Li S-S; Wang J Cu(I)/Chiral Bisoxazoline-Catalyzed Enantioselective Sommelet-Hauser Rearrangement of Sulfonium Ylides. *J. Org. Chem.* 2020, 85, 12343–12358 Catalytic aromatic rearrangements with *ammonium* (X = N) ylides: [PubMed: 32881498] (g)Pan C; Guo W; Gu Z Unusual biaryl torsional strain promotes reactivity in Cu-catalyzed Sommelet-Hauser rearrangement. *Chem. Sci.* 2018, 9, 5850–5854. [PubMed: 30079198]
- (7) (a). Hansen HJ; Schmid H Aromatic Sigmatropic Rearrangements. *Chem. Br.* 1969, 5, 111–116. (b)Murray AW Molecular rearrangements. *Org. React. Mech.* 2002, 487–615.(c)Snape TJ In Aromatic Rearrangements in Which the Migrating Group Migrates to the Aromatic Nucleus: An Overview; John Wiley & Sons, Inc.: 2015; pp 485–510.
- (8) (a). Xu X; Li C; Xiong M; Tao Z; Pan Y Hemin-catalyzed sulfonium ylide formation and subsequently reactant-controlled chemoselective rearrangements. *Chem. Commun.* 2017, 53, 6219–6222.(b)Yang Z; Guo Y; Koenigs RM Solvent-dependent, rhodium catalysed rearrangement reactions of sulfur ylides. *Chem. Commun.* 2019, 55, 8410–8413.

- (9). Wang J In *Comprehensive Organometallic Chemistry III*; Crabtree RH; Mingos DMP, Eds.; Elsevier: Oxford, U.K., 2007; Vol. 11, pp 151–178.
- (10) (a). Kochanowska-Karamyan AJ; Hamann MT Marine Indole Alkaloids: Potential New Drug Leads for the Control of Depression and Anxiety. *Chem. Rev.* 2010, 110, 4489–4497. [PubMed: 20380420] (b)Kaushik N; Kaushik N; Attri P; Kumar N; Kim C; Verma A; Choi E Biomedical Importance of Indoles. *Molecules* 2013, 18, 6620–6662. [PubMed: 23743888] (c)Vitaku E; Smith DT; Njardarson JT Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles Among U. S. FDA Approved Pharmaceuticals. *J. Med. Chem.* 2014, 57, 10257–10274. [PubMed: 25255204]
- (11) (a). Doyle MP Catalytic Methods for Metal Carbene Transformations. *Chem. Rev.* 1986, 86, 919–939.(b)Liu L; Zhang J Gold-catalyzed transformations of  $\alpha$ -diazocarbonyl compounds: selectivity and diversity. *Chem. Soc. Rev.* 2016, 45, 506–516. [PubMed: 26658761] (c)DeAngelis A; Panish R; Fox JM Rh-Catalyzed Intermolecular Reactions of  $\alpha$ -Alkyl- $\alpha$ -Diazo Carbonyl Compounds with Selectivity Over  $\beta$ -Hydride Migration. *Acc. Chem. Res.* 2016, 49, 115–127. [PubMed: 26689221] (d)Zhao X; Zhang Y; Wang J Recent developments in copper-catalyzed reactions of diazo compounds. *Chem. Commun.* 2012, 48, 10162–10173.
- (12) (a). Doyle MP Metal Carbene Reactions from Dirhodium(II) Catalysts. *Top. Organomet. Chem.* 2004, 13, 203–222.(b)Davies HML; Parr BT Rhodium Carbenes. *Wiley Ser. React. Intermed. Chem. Biol.* 2013, 7, 363–403.
- (13). Select examples for the synthesis of 3-methyleneindolines: (a) Burns B; Grigg R; Sridharan V; Worakun T Palladium catalysed tandem cyclisation-anion capture processes. Hydride ion capture by vinylpalladium species. *Tetrahedron Lett.* 1988, 29, 4325–4328.(b)Zenner JM; Larock RC Palladium-Catalyzed, Asymmetric Hetero- and Carboannulation of Allenes Using Functionally-Substituted Aryl and Vinylic Iodides. *J. Org. Chem.* 1999, 64, 7312–7322.(c)Font M; Cendón B; Seoane A; Mascareñas JL; Gulías M Rhodium(III)-Catalyzed Annulation of 2-Alkenyl Anilides with Alkynes Through C-H Activation: Direct Access to 2-Substituted Indolines. *Angew. Chem., Int. Ed.* 2018, 57, 8255–8259.(d)Kondoh A; Terada M Brønsted Base-Catalyzed Umpolung Intramolecular Cyclization of Alkynyl Imines. *Chem. - Eur. J.* 2018, 24, 3998–4001. [PubMed: 29341333] (e)Li X; Zhou B; Yang R-Z; Yang F-M; Liang R-X; Liu R-R; Jia Y-X Palladium-Catalyzed Enantioselective Intramolecular Dearomative Heck Reaction. *J. Am. Chem. Soc.* 2018, 140, 13945–13951. [PubMed: 30256629]
- (14). When subjected to the [2,3]-rearrangement conditions, indole **3f** proceeded to give a mixture of [2,3]:[1,2] rearrangement products. We presume that the 2-methyl substituent hinders the [2,3]-rearrangement that forms the fully substituted carbon center and increases the propensity for the ylide to dissociate and generate the competitive [1,2]-rearrangement product (see SI for details).
- (15) (a). Becke AD Density-Functional Thermochemistry. III. The Role of Exact Exchange. *J. Chem. Phys.* 1993, 98, 5648–5652.(b)Lee C; Yang W; Parr RG Development of the Colle-Salvetti Correlation-Energy Formula into a Functional of the Electron Density. *Phys. Rev. B: Condens. Matter Mater. Phys.* 1988, 37, 785–789.(c)Grimme S; Ehrlich S; Goerigk L Effect of the Damping Function in Dispersion Corrected Density Functional Theory. *J. Comput. Chem.* 2011, 32, 1456–1465. [PubMed: 21370243] (d)Hariharan PC; Pople JA The Influence of Polarization Functions on Molecular Orbital Hydrogenation Energies. *Theor. Chim. Acta* 1973, 28, 213–222. (e)Igel-Mann G; Stoll H; Preuss H Pseudopotentials for Main Group Elements (IIIa through VIIa). *Mol. Phys.* 1988, 65, 1321–1328.(f)Hay PJ; Wadt WR Ab Initio Effective Core Potentials for Molecular Calculations. Potentials for the Transition Metal Atoms Sc to Hg. *J. Chem. Phys.* 1985, 82, 270–283.(g)Goerigk L; Grimme S Efficient and Accurate Double-Hybrid-Meta-GGA Density Functionals—Evaluation with the Extended GMTKN30 Database for General Main Group Thermochemistry, Kinetics, and Noncovalent Interactions. *J. Chem. Theory Comput.* 2011, 7, 291–309. [PubMed: 26596152] (h)Weigend F; Ahlrichs R Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy. *Phys. Chem. Chem. Phys.* 2005, 7, 3297–3305. [PubMed: 16240044] (i)Weigend F Accurate Coulomb-fitting basis sets for H to Rn. *Phys. Chem. Chem. Phys.* 2006, 8, 1057–1065. [PubMed: 16633586] (j)Hellweg A; Hättig C; Höfener S; Klopper W Optimized Accurate Auxiliary Basis Sets for RI-MP2 and RI-CC2 Calculations for the Atoms Rb to Rn. *Theor. Chem. Acc.* 2007, 117, 587–597.(k)Chmela J; Harding ME Optimized Auxiliary Basis


Sets for Density Fitted Post-Hartree-Fock Calculations of Lanthanide Containing Molecules. *Mol. Phys.* 2018, 116, 1523–1538.

- (16) (a). Hock KJ; Koenigs RM Enantioselective [2,3]-Sigmatropic Rearrangements: Metal-Bound or Free Ylides as Reaction Intermediates? *Angew. Chem., Int. Ed.* 2017, 56, 13566–13568. (b) Zhang Z; Sheng Z; Yu W; Wu G; Zhang R; Chu W-D; Zhang Y; Wang J Catalytic asymmetric trifluoromethylation via enantioselective [2,3]-sigmatropic rearrangement of sulfonium ylides. *Nat. Chem.* 2017, 9, 970–976. [PubMed: 28937673] (c) Harrison JG; Gutierrez O; Jana N; Driver TG; Tantillo DJ Mechanism of Rh<sub>2</sub>(II)-Catalyzed Indole Formation: The Catalyst Does Not Control Product Selectivity. *J. Am. Chem. Soc.* 2016, 138, 487–490. [PubMed: 26718908] (d) Nickerson LA; Bergstrom BD; Gao M; Shiue Y-S; Laconsay CJ; Culberson MR; Knauss WA; Fettinger JC; Shaw JT; Tantillo DJ Enantioselective synthesis of isochromans and tetrahydroisoquinolines by C-H insertion of donor/donor carbenes. *Chem. Sci.* 2020, 11, 494–498. [PubMed: 32874491] (e) Laconsay CJ; Tantillo DJ Metal Bound or Free Ylides as Reaction Intermediates in Metal-Catalyzed [2,3]-Sigmatropic Rearrangements? It Depends. *ACS Catal.* 2021, 11, 829–839. (f) Li F; Pei C; Koenigs RM Rhodium-catalyzed cascade reactions of triazoles with organoselenium compounds - a combined experimental and mechanistic study. *Chem. Sci.* 2021, 12, 6362–6369. [PubMed: 34084435]
- (17). In addition, when two distinct substrates were simultaneously subjected to the [2,3]-rearrangement conditions, no crossover products were detected (see SI for details). These observations are consistent with a metal-free symmetry-allowed, concerted, and unimolecular process.
- (18). Other stepwise mechanisms were considered and cannot be definitively ruled out, including a fragmentation/recombination pathway (see SI).
- (19) (a). Ollis WD; Rey M; Sutherland IO; Closs GL The Mechanism of Stevens Rearrangement. *J. Chem. Soc., Chem. Commun.* 1975, 543–545. (b) Chantrapromma K; Ollis WD; Sutherland IO Radical-coupling Products Derived from Ammonium Ylides. Competing [1,2], [1,3], and [1,4] Anionic Rearrangements. *J. Chem. Soc., Chem. Commun.* 1978, 670–671. (c) Heard GL; Frankcombe KE; Yates BFA Theoretical Study of the Stevens Rearrangement of Methylammonium Methylide and Methylammonium Formylmethylide. *Aust. J. Chem.* 1993, 46, 1375–1388. (d) Liu Z; Jin X; Dang Y Mechanistic Studies of Copper(I)-Catalyzed Stereoselective [2,3]-Sigmatropic Rearrangements of Diazoesters with Allylic Iodides/ Sulfides. *ACS Catal.* 2021, 11, 691–702.
- (20) (a). Hogen-Esch TE; Smid J Studies of Contact and Solvent-Separated Ion Pairs of Carbanions. I. Effect of Temperature, and Solvent. *J. Am. Chem. Soc.* 1966, 88, 307–318. (b) Roy S; Baer MD; Mundy CJ; Schenter GK Marcus Theory of Ion-Pairing. *J. Chem. Theory Comput.* 2017, 13, 3470–3477. [PubMed: 28715638] (c) Macchioni A Ion Pairing in Transition-Metal Organometallic Chemistry. *Chem. Rev.* 2005, 105, 2039–2073. [PubMed: 15941209] (d) Brak K; Jacobsen EN Asymmetric Ion-Pairing Catalysis. *Angew. Chem., Int. Ed.* 2013, 52, 534–561. (e) Chen Y; Liu Y; Li Z; Dong S; Liu X; Feng X Tandem Insertion-[1,3]-Rearrangement: Highly Enantioselective Construction of  $\alpha$ -Aminoketones. *Angew. Chem.* 2020, 132, 8129–8133. (f) Yao L; Ishihara K Enantioselective [1,3] O-to-C Rearrangement: Dearomatization of Alkyl 2-Allyloxy/Benzyloxy-1/3-Naphthoates Catalyzed by a Chiral  $\pi$ -Cu(II) Complex. *Chem. Sci.* 2019, 10, 2259–2263. [PubMed: 30881650] (g) Tsuji Y; Richard JP Reactions of Ion-Pair Intermediates of Solvolysis. *Chem. Rec.* 2005, 5, 94–106. [PubMed: 15825158]
- (21). Bowry VW; Luszyk J; Ingold KU Calibration of a New Horology of Fast Radical “Clocks”. Ring-Opening Rates for Ring- and  $\alpha$ -Alkyl-Substituted Cyclopropylcarbinyl Radicals and for the Bicyclo[2.1.0]pent-2-yl Radical. *J. Am. Chem. Soc.* 1991, 113, 5687–5698.
- (22). We do note, however, that some ion-pairs appear to have diradical character (see SI).
- (23). Computed relative free energies of ion-pairs **12a-12e** with respect to the reactants were calculated using PWPB95-D3(BJ)/def2-QZVPP//IEFPCM(CH<sub>2</sub>Cl<sub>2</sub>)-B3LYP/6–31G(d), SDD, which is used to more accurately capture dispersion effects, but the accuracy of the computed energies is limited because explicit solvent was not included.
- (24). de Azambuja F; Yang MH; Feoktistova T; Selvaraju M; Brueckner AC; Grove MA; Koley S; Cheong PHY; Altman RA Connecting remote C-H bond functionalization and decarboxylative coupling using simple amines. *Nat. Chem.* 2020, 12, 489–496. [PubMed: 32152476]

- (25) (a). Fang Y; Powell JA; Li E; Wang Q; Perry Z; Kirchon A; Yang X; Xiao Z; Zhu C; Zhang L; Huang F; Zhou H-C Catalytic Reactions within the Cavity of Coordination Cages. *Chem. Soc. Rev.* 2019, 48, 4707–4730. [PubMed: 31339148] (b) Stelson AC; Hong CM; Groenenboom MC; Little CAE; Booth JC; Orloff ND; Bergman RG; Raymond KN; Schwarz KA; Toste FD; Long CJ Measuring ion-pairing and hydration in variable charge supramolecular cages with microwave microfluidics. *Commun. Chem.* 2019, 2, 54.
- (26). Karwehl S; Jansen R; Huch V; Stadler M Sorazolons, Carbazole Alkaloids from *Sorangium cellulosum* Strain Soce375. *J. Nat. Prod.* 2016, 79, 369–375. [PubMed: 26866461]
- (27) (a). Cram DJ; Elhafez FAA Studies in Stereochemistry. X. The Rule of ‘Steric Control of Asymmetric Induction’ in the Syntheses of Acyclic Systems. *J. Am. Chem. Soc.* 1952, 74, 5828–5835. (b) Reetz MT Chelation or Non-Chelation Control in Addition Reactions of Chiral  $\alpha$ - and  $\beta$ -Alkoxy Carbonyl Compounds [New Synthetic Methods (44)]. *Angew. Chem., Int. Ed. Engl.* 1984, 23, 556–569. (c) Mengel A; Reiser O Around and beyond Cram’s Rule. *Chem. Rev.* 1999, 99, 1191–1223. [PubMed: 11749444]
- (28). Ferrer C; Echavarren AM Gold-Catalyzed Intramolecular Reaction of Indoles with Alkynes: Facile Formation of Eight-Membered Rings and an Unexpected Allenylation. *Angew. Chem., Int. Ed.* 2006, 45, 1105–1109.




**Figure 1.** Computed (PWPB95-D3(BJ)/def2-QZVPP//IEFPCM(CH<sub>2</sub>Cl<sub>2</sub>)-B3LYP/6-31G(d), SDD) relative free energies (kcal/mol, italics) for minima and TSSs involved in the Rh-promoted reaction of **1a** and **2a**.




**Figure 2.** (A) Computed (PWPB95-D3(BJ)/def2-QZVPP//IEFPCM( $\text{CH}_2\text{Cl}_2$ )-B3LYP/6-31G(d),SDD) relative free energies (kcal/mol, italics) for minima and TSSs involved in the Cu-promoted reaction of **1a** and **2a**. A selection of ion-pairs **12a-12e** were generated by scanning the bonds that form en route to the products. The energies for ion-pairs **12a-12e** are based on optimized complexes. Geometries of ion-pairs **12a-12e**, their preceding zwitterion **11**, and the recombination product **13a** shown in ball-and-stick images are included to facilitate comparison of overall shapes. For clarity in visual comparison, the enolate part of the ion-pair is highlighted in green and the indolyl part (highlighted in gray) is positioned the same way for each structure above. The solvent cage (not modeled explicitly) is depicted in blue with dotted lines. (B) Reaction conditions for radical probe experiment: indole **1q** (0.16 mmol), benzyl diazoester **2a** (1.2 equiv, added using syringe pump as 0.2 M solution in  $\text{CH}_2\text{Cl}_2$  at a rate of 2 mL/h),  $\text{Cu}(\text{hfacac})_2$  (5 mol %),  $\text{CH}_2\text{Cl}_2$ , 23 °C. (C) Reaction conditions for crossover experiment: indole **1i** (0.08 mmol, 0.5 equiv), indole **1r** (0.08 mmol, 0.5 equiv) benzyl diazoester **2a** (1.2 equiv, added using syringe pump as 0.2 M solution in  $\text{CH}_2\text{Cl}_2$  at a rate of 2 mL/h),  $\text{Cu}(\text{hfacac})_2$  (5 mol %),  $\text{CH}_2\text{Cl}_2$ , 23 °C.

**A. Catalytic Generation of Oxonium Ylides from Diazocarbonyl Compounds**




**B. Catalyst-Controlled Regiodivergent Aromatic Rearrangements of Indole-Based Oxonium Ylides (This Research)**




**Scheme 1.**  
Catalyst Control of Regioselectivity in Oxonium Ylide Rearrangements






**Scheme 2.**  
Synthetic Derivatization of [2,3]-Rearrangement Product



**Scheme 3.**  
 Synthesis and Stereochemical Reassignment of (±)-Sorazolon B from [1,2]-Rearrangement Product

Table 1.

Optimization of [1,2]- and [2,3]-Rearrangements<sup>a</sup>

Entry	Catalyst	Solvent	Conversion <sup>b</sup> (%)	Yield <sup>c</sup> (%)	3a:4a
1	PdCl <sub>2</sub>	CH <sub>2</sub> Cl <sub>2</sub>	<5	<5	
2	AgOTf	CH <sub>2</sub> Cl <sub>2</sub>	<5	<5	
3	Ph <sub>3</sub> PAuCl	CH <sub>2</sub> Cl <sub>2</sub>	<5	<5	
4	Cu(OTf)·benzene	CH <sub>2</sub> Cl <sub>2</sub>	5	5	82:18
5	CuCl	CH <sub>2</sub> Cl <sub>2</sub>	5	5	71:29
6	CuCl <sub>2</sub>	CH <sub>2</sub> Cl <sub>2</sub>	8	7	75:25
7	CuOAc/NaBAR <sub>F</sub>	CH <sub>2</sub> Cl <sub>2</sub>	26	18	76:24
8	[Cu(MeCN) <sub>4</sub> ] PF <sub>6</sub>	CH <sub>2</sub> Cl <sub>2</sub>	42	38 <sup>d</sup>	86:14
9	Cu(hfacac) <sub>2</sub>	CH <sub>2</sub> Cl <sub>2</sub>	67	52 <sup>d</sup>	92:8
10 <sup>e</sup>	Cu(hfacac) <sub>2</sub>	CH <sub>2</sub> Cl <sub>2</sub>	>95	78 <sup>d</sup>	>95:5
11	Rh <sub>2</sub> (OAc) <sub>4</sub>	CH <sub>2</sub> Cl <sub>2</sub>	55	42 <sup>d</sup>	<5:95
12	Rh <sub>2</sub> (cap) <sub>4</sub>	CH <sub>2</sub> Cl <sub>2</sub>	<5	<5	
13	Rh <sub>2</sub> (TFA) <sub>4</sub>	CH <sub>2</sub> Cl <sub>2</sub>	30	<5	
14	Rh <sub>2</sub> (TPA) <sub>4</sub>	CH <sub>2</sub> Cl <sub>2</sub>	56	23	32:68
15	Rh <sub>2</sub> (oct) <sub>4</sub>	CH <sub>2</sub> Cl <sub>2</sub>	44	33	10:90
16 <sup>e</sup>	Rh <sub>2</sub> (OAc) <sub>4</sub>	CH <sub>2</sub> Cl <sub>2</sub>	92	26	<5:95
17	Rh <sub>2</sub> (OAc) <sub>4</sub>	DCE	41	36	<5:95
18	Rh <sub>2</sub> (OAc) <sub>4</sub>	CHCl <sub>3</sub>	51	33	<5:95
19	Rh <sub>2</sub> (OAc) <sub>4</sub>	toluene	20	16	<5:95

<sup>a</sup>Reaction conditions: indole **1a** (0.16 mmol), benzyl diazoester **2a** (1.2 equiv, added using syringe pump as 0.2 M solution in solvent at a rate of 2 mL/h), copper catalyst (5 mol %) or rhodium catalyst (2 mol %).

<sup>b</sup>Conversion of **1a**.


<sup>c</sup>NMR yield using 1,3,5-trimethoxybenzene as internal standard.

<sup>d</sup>Isolated yield.

<sup>e</sup>2.4 equiv of **2a** was used.

Table 2.

Products Generated by Regiodivergent [1,2]- and [2,3]-Rearrangements

<sup>a</sup> rsm = recovered starting material 1.