

UCSF

UC San Francisco Previously Published Works

Title

The Effect of Altered Toll-like Receptor 4 Signaling on Cancer Cachexia

Permalink

<https://escholarship.org/uc/item/45v0n8hv>

Journal

JAMA Otolaryngology - Head & Neck Surgery, 133(12)

ISSN

2168-6181

Authors

Cannon, Trinitia Y
Guttridge, Denis
Dahlman, Jason
[et al.](#)

Publication Date

2007-12-01

DOI

10.1001/archotol.133.12.1263

Peer reviewed

The Effect of Altered Toll-like Receptor 4 Signaling on Cancer Cachexia

Trinitia Y. Cannon, MD; Denis Guttridge, PhD; Jason Dahlman, BS; Jonathan R. George, BA, MS; Victor Lai, MD; Carol Shores, MD, PhD; Petra Bůžková, PhD; Marion E. Couch, MD, PhD

Objective: To determine whether mice unable to mount an intact inflammatory response because of a Toll-like receptor (TLR) pathway defect will develop less severe cancer cachexia.

Design: Prospective animal study.

Setting: Academic research center.

Subjects: Six- to eight-week-old, female C3H/HeJ mice (17-18 g) and age-, weight-, and sex-matched wild-type C3H/HeN mice, differing in that the HeJ mice have nonfunctional TLR4 due to a TLR4 double mutation (TLR4^{dl/dl}).

Intervention: The mice were inoculated with equal numbers of SCCF-VII cells and housed in individual cages.

Main Outcome Measures: Food intake, body weight, pretumor and posttumor body composition, circulating cytokines, and levels of a marker of muscle atrophy were analyzed.

Results: The wild-type HeN mice weighed less on average than the TLR4^{dl/dl} mice (2.6 g vs 4.9 g) ($P = .01$). They consumed more food, had smaller tumors, and had less lean body mass and fat mass than the TLR4^{dl/dl} mice. Interleukin 1 β level was significantly elevated in the tumor-bearing HeN mice (mean gain of 259 pg/mL) but not in the TLR4^{dl/dl} mice ($P = .03$). Both mouse strains had evidence of muscle atrophy.

Conclusions: In spite of increased food intake and smaller tumors, the wild-type HeN mice had more severe cachexia than the TLR4^{dl/dl} mice. The impaired ability to secrete proinflammatory cytokines such as interleukin 1 β may protect these animals from developing severe cancer cachexia. This animal model represents a novel system in which the host contributions to cachexia may be further studied.

Arch Otolaryngol Head Neck Surg. 2007;133(12):1263-1269

CANCER CACHEXIA IS A wasting syndrome that develops in the setting of advanced malignancy including squamous cell carcinoma of the head and neck. It results in reduced quality of life, decreased survival, and increased complications from treatment.¹ Cachexia will be the main cause of death in approximately 20% to 30% of all patients with cancer.² Clinically, cachexia manifests as unintentional weight loss, marked asthenia, sarcopenia, and anemia. Cachexia differs from starvation in that it is not reversible by increased nutritional intake, involves preferential loss of lean body mass rather than only fat mass, and is characterized by a chronic inflammatory state.³ Mounting evidence from animal models suggests a compelling link between activation of the inflammatory pathway and the development of cachexia.⁴ In addition to tumor-derived catabolic factors, humoral activa-

tors of the inflammatory cascade have been proposed as mediators of cancer cachexia.⁵ Interleukin 1 β (IL-1 β) and interleukin 6 (IL-6) are 2 proinflammatory cytokines that have been shown to stimulate the loss of lean body mass seen in experimental models of cancer cachexia.^{1,3} Interleukin 1 β is necessary for the induction of IL-6-mediated cachexia in the colon-26 (C26) model, in which it also appears to contribute to the initiation of cachexia at the level of the tumor.⁶ Intratumoral injection of IL-1 receptor antagonist considerably attenuates weight loss and loss of lean body mass and fat mass.⁶

Both sterile and infectious inflammation may be mediated by a common pathway involving similar proinflammatory cytokines.⁷ The Toll-like receptor (TLR) system is potentially one such pathway. Toll-like receptors are a family of innate cellular pathogen-recognition receptors that are present on many cell types including macrophages, dendritic cells, T and B cells,

Author Affiliations are listed at the end of this article.

tumor cells, and muscle cells.⁸ The classic ligand for TLR4 is lipopolysaccharide (LPS), commonly seen on gram-negative bacterium, but TLR4 is now known to recognize other endogenous ligands such as fibrinogen, heat-shock proteins, and polysaccharide fragments of heparin sulfate.⁹ Signaling through the TLR pathway not only results in activation of antigen-specific immunity but also leads to the production of the same cytokine mediators involved in infectious inflammation such as sepsis.¹⁰ Indeed, activation of the TLR4 receptor system by LPS elaborates the same proinflammatory cytokines that are known to mediate inflammation in cancer cachexia. One cytokine, IL-1 β , is capable of stimulating physiologic and muscle protein wasting in experimental models of cancer cachexia.⁶ Skeletal muscles, which express TLR2 and TLR4, respond to LPS by initiating an immune response that includes IL-1, IL-6, and tumor necrosis factor α (TNF- α).^{11,12} However, the ability of LPS to stimulate cytokine production, including in the serum and skeletal muscle, was markedly decreased in mice with a mutation in TLR4.¹¹ Increased levels of serum proinflammatory cytokines are associated with decreased muscle synthesis and loss of lean body mass in many wasting conditions.¹² Therefore, we reasoned that mice with defective TLR4 signaling pathway would have less proinflammatory cytokine production and muscle wasting if cancer cachexia was mediated by this innate immune system.

C3H/HeJ mice have a mutation in a single gene on chromosome 4, *lps*, which leaves the macrophages unable to respond to LPS and produce cytokines.^{13,14} Mice with this mutation in the TLR4 system (TLR4^{d/d} C3H/HeJ) experience impaired release of IL-1 β and do not develop the characteristic inflammation-induced wasting usually seen in response to LPS challenge.¹⁵⁻¹⁸ Therefore, this strain of mice may represent an important model for the study of host cytokine contribution to the pathogenesis of cancer cachexia. In addition, host contribution to cancer cachexia may be studied in this model because mice with an impaired TLR4 signaling pathway appear to have different cytokine profiles than the congenic HeN mice with intact TLR signaling. Our hypothesis was that TLR4^{d/d} mice, with a defective TLR4 cytoplasmic domain, will have less severe cancer cachexia. In the present study, we compared the host response to tumor challenge using the SCCF-VII tumor cell line in both HeN (wild-type) mice and TLR4^{d/d} C3H/HeJ mice.

METHODS

TUMOR CELL LINES

The SCCF cell line is a spontaneously arising squamous cell carcinoma that forms a measurable tumor after being injected into immunocompetent syngeneic mice. Cultured cells were propagated in tissue culture flasks at 37°C, with 5% carbon dioxide in RPMI (Roswell Park Memorial Institute) 1640 medium supplemented with 10% fetal bovine serum, 1% penicillin-streptomycin sulfate, and 1% sodium glutamate. The SCCF cells were grown in culture to 80% confluence. On the day of tumor injection, cells were harvested using a short trypsin treatment (0.25% trypsin with 1.0-mmol/L EDTA) and then washed twice and resuspended in phosphate-buffered saline.

ANIMAL MODEL

Six- to eight-week-old, female TLR4^{d/d} C3H/HeJ mice (17-18 g) were purchased from Jackson Laboratories, Bar Harbor, Maine. Age-, weight-, and sex-matched C3H/HeN mice were purchased from Charles River Laboratories, Inc (Wilmington, Massachusetts). The animals were housed individually in sterilized plastic cages in a temperature-controlled room with a 12-hour dark-light cycle and were fed a defined AIN-93M diet (Granville Milling, Creedmoor, North Carolina) and received tap water ad libitum.

On day 0, the mice in each group received bilateral subcutaneous injections of the SCCF cells (1×10^5 cells/100 μ L). Food intake, body weight, and tumor size were measured every 1 to 2 days. On day 23, all mice were humanely killed and serum was drawn. Gastrocnemius and quadriceps muscle, tumor, and epididymal fat were excised. Then, all tumors were excised and final tumor weight was measured. Serum and tissues were snap frozen in liquid nitrogen and stored at -80°C. Final body weight was calculated by subtracting tumor weight from total body weight (weight of mice with tumors intact). All animal procedures were approved by the Institutional Animal Care and Use Committee (IACUC) of the University of North Carolina at Chapel Hill. Mice were allowed to grow tumors for 23 days or when the tumor burden had reached the accepted limit set by the Department of Laboratory Medicine and the IACUC. The animals were then humanely killed.

ASSESSMENT OF FOOD INTAKE, BODY WEIGHT, AND TUMOR SIZE

The animals were individually housed for accurate measurement of food consumption. The amount of ingested food was measured in all cages every 2 days by taking the daily weights (grams) of food. Mean daily food intake and cumulative intake were determined for each group of mice. Body weight and the length (A) and width (B) of the tumors were measured every 2 to 4 days. Tumor growth was assessed using calipers in 2 dimensions. The mean orthogonal diameter of the tumor was determined using the following formula: mean diameter = (A + B)/2. The gross size of the tumors was determined after harvesting the tumors at the completion of the study.

BODY COMPOSITION ANALYSIS

Prior to tumor injection, all animals underwent body composition analysis using a Lunar PIXImus densitometer 2.00 (software version 1.42.006.010; Lunar Corp, Madison, Wisconsin). The densitometer uses dual-energy x-ray absorptiometry to provide precise measures of fat mass, bone-free lean body mass, and total bone mineral density.¹⁹ The analysis was repeated on day 23 in live animals, prior to killing. Pretumor and posttumor results were compared in both groups.

SERUM CYTOKINE AND CHEMOKINE ASSAYS

Cytokine and chemokine analysis was determined using Bio-plex Protein Array system (BioRad, Hercules, California), a novel multiplexed, particle-based, flow cytometric assay that uses anti-cytokine monoclonal antibodies linked to microspheres incorporating distinct proportions of 2 fluorescent dyes. Our assay was customized to detect and quantify 18 cytokines and chemokines, including IL-1 β , IL-1 α , IL-6, TNF- α , and interferon- γ , among others. For each cytokine, 8 standards ranged from 2 to 32 000 pg/mL, and the minimum detectable dose was lower than 10 pg/mL. Prior to tumor inoculation and at the completion of the experiment, approximately 200 μ L of blood




Figure 1. Effect of tumor growth on body weight. A, Mean \pm SE weight gained in wild-type C3H/HeN (HeN) and TLR4^{dd} C3H/HeJ (HeJ) mice measured 23 days after bilateral subcutaneous injections of 1×10^5 SCCF-VII cells/100 μ L. Weight gain = final body weight (after tumor removal) - initial body weight. The HeN mice gained significantly less weight (2.6 ± 0.3 g [minus tumor]) than the HeJ mice (4.9 ± 0.5 g [minus tumor]) ($P = .01$), and the weight gain occurred early in the experiment. Mice were individually housed to monitor food consumption. B, Comparison of the change in body weight between the HeN and HeJ mice ($n = 5$ for each group) during 23 days (no tumors). In this separate experiment, body weight was measured every 2 days.

was collected from each mouse. Serum was harvested after the clotting of blood at room temperature for 1 hour and kept frozen (-80°C) until analysis. Pretumor and posttumor implantation cytokine levels were compared.

UBIQUITIN LIGASE EXPRESSION IN MUSCLES

On postimplantation day 23, the mice were humanely killed, and hind leg quadriceps and gastrocnemius muscles were rapidly dissected and snap-frozen in liquid nitrogen. Gastrocnemius and quadriceps muscles from tumor-bearing C3H/HeJ and C3H/HeN mice were homogenized. To confirm the presence of muscle atrophy, isolated muscle tissue was analyzed for the E3 ubiquitin ligase muscle ring finger 1 (MuRF1) using Northern blot analysis as previously described.²⁰ Expression of MuRF1 was quantified by phosphoimaging and compared with the expression of glyceraldehyde-3-phosphate dehydrogenase (GAPDH), a housekeeping gene.

DATA ANALYSIS

All quantitative data are represented as mean \pm SE, with 95% confidence intervals (CIs) when appropriate. An unpaired *t* test allowing for unequal variance was used to compare group means. The reported *P* values, including those for cytokine analysis, are not corrected for multiple comparisons. Because of small sample sizes (5 mice per group), statistical testing results (*P* values) should be interpreted with care and analysis should be considered mainly as exploratory rather than inferential.

RESULTS

INCREASED FOOD INTAKE AND WEIGHT LOSS IN TUMOR-BEARING WILD-TYPE MICE

When equal numbers of SCCF cells were injected into the flanks of both strains of mice, the wild-type HeN mice

Table 1. Food Intake, Body Weight, and Body Composition Analyses^a

Variable	C3H/HeN Mice With Tumor (n = 5)	C3H/HeJ (TLR4 ^{dd}) Mice With Tumor (n = 5)
Total food intake, g	102.5 \pm 5.8	76 \pm 0.9 ^b
Body weight		
Initial, g	16.4 \pm 0.2	17.8 \pm 0.2
Final, g	19.0 \pm 0.2	22.7 \pm 0.6
Weight change, %	15.7 \pm 1.9	27.6 \pm 2.7 ^c
Body composition, %		
Change in lean body mass	3.9 \pm 0.6	8.4 \pm 0.8 ^d
Change in fat mass	0.5 \pm 0.1	1.2 \pm 0.2 ^e
Clinical cachexia ^f	++	+

Abbreviations: ++, significant clinical cachexia; +, less severe clinical signs.

^aData are given as mean \pm SE value.

^b $P = .002$.

^c $P = .01$.

^d $P = .004$.

^e $P = .02$.

^fClinical cachexia is defined as muscle wasting, weight loss, and lethargy.

weighed less than the TLR4^{dd} mice, even after tumor size was subtracted. The mean \pm SE weight gain in the HeN mice was 2.6 ± 0.3 g (minus tumor), while the TLR4^{dd} mice gained 4.9 ± 0.5 g (minus tumor) ($P = .006$) (Figure 1A). Beginning 1 week after tumor injection, the HeN mice stopped gaining weight. Despite the weight gain due to large tumors growing on the flanks, the wild-type mice started to lose weight beginning postinjection day 10 and had a significantly smaller change in body weight ($15.7\% \pm 1.9\%$) compared with the TLR4^{dd} mice ($27.6\% \pm 2.7\%$) ($P = .01$) (Figure 1B). The wild-type HeN mice without tumors usually have a mean \pm SE weight gain of $27\% \pm 7\%$, while the C3H/HeJ (TLR4^{dd}) mice without tumors have a weight gain of $18\% \pm 5\%$ over the time course of these experiments. This suggests that the TLR4^{dd} strain of mice is less useful as a model for cancer cachexia because they do not exhibit the weight loss typically seen by the tumor-bearing C3H/HeN mice (Table 1).

The differences in body weight could not be attributed to anorexia or differences in food intake between the groups. The wild-type HeN mice consumed more food at every time point and had a mean \pm SE cumulative food intake of 102.5 ± 5.8 g, while the TLR4^{dd} mice consumed 76.4 ± 0.9 g ($P = .002$) (Figure 2A). In a separate experiment, HeN mice with and without tumors had their daily food intake monitored to rule out anorexia as a cause of the weight loss. The tumor-bearing mice consumed as much food as the controls (Figure 2B).

LESS TUMOR GROWTH IN WILD-TYPE MICE

The SCCF cell line is highly aggressive, with the formation of palpable tumors in mice by day 7 when injected in the flanks. Because of its reproducible rapid growth, tumors become so bulky that mice usually need to be killed




Figure 2. Increased daily and cumulative food intake. Wild-type C3H/HeN (HeN) mice and TLR4^{del} C3H/HeJ (HeJ) mice were individually housed for accurate measurement of food intake. Food was weighed every 2 days following tumor challenge. A, The HeN mice consumed more food per day and had a significantly higher cumulative food intake (102.5 ± 5.8 g vs 76.4 ± 0.9 g) ($P = .002$). B, Control and tumor-bearing HeN mice consume the same amount of food (86.2 ± 0.9 g vs 88.1 ± 4.6 g).

by day 20 to 23. There was a significant difference in tumor size when measuring the mean tumor volume between mice with an intact TLR-4 pathway (C3H/HeN) and the TLR4^{del} strain (**Figure 3**). The wild-type HeN mice had significantly smaller tumors, with a final mean ± SE tumor volume of 1.35 ± 0.3 cm² compared with TLR4^{del} mice that had tumor volumes of 3.04 ± 0.7 cm² ($P = .05$).

LESS LEAN BODY AND FAT MASS IN WILD-TYPE MICE

To determine whether the weight loss experienced by the mice was due to cachexia, each group of mice underwent pretumor and posttumor body composition analysis to examine the effect of SCCF tumor growth on lean body and fat mass. Body composition analysis revealed that tumor growth resulted in less lean body and fat mass in the wild-type mice than in the TLR4^{del} mice (**Figure 4**). The wild-type HeN mice had a mean ± SE percentage change in lean body mass of 27.5% ± 4.7%, while the TLR4^{del} mice had a 56.5% ± 6.3% lean body mass change ($P = .004$). The percentage change in fat mass was also less in the HeN mice compared with the TLR4^{del} mice,




Figure 3. Smaller tumor volume in wild-type C3H/HeN (HeN) mice. Wild-type HeN and TLR4^{del} C3H/HeJ (HeJ) mice ($n = 5$ per group) were injected with equal numbers of SCCF-VII cells. Tumors were measured in 2 dimensions with microcalipers every 2 days, and final tumor volume after 21 days was compared. * $P = .05$.

17.8% ± 3.9% vs 34.6% ± 7.3% ($P = .02$). Figure 4 summarizes the results for the 2 groups of mice.

DIFFERENT CYTOKINE PROFILES IN THE 2 STRAINS OF MICE

There is evidence that activation of TLR4 system leads to the expression of some of the same proinflammatory cytokines involved in the production of cachexia. Cytokine production was therefore examined in the 2 congenic strains of mice bearing the same tumor type and was compared to see whether there were differences in the host's ability to mount an inflammatory response. Of the 18 serum cytokine and chemokines analyzed, IL-1 β was found to be significantly elevated in tumor-bearing HeN mice compared with the tumor-bearing TLR4^{del} mice ($P = .03$) (**Table 2** and **Figure 5**). Tumor necrosis factor α decreased in both strains of tumor-bearing mice (-242 vs -5053) ($P = .34$). Interferon- γ decreased in both strains as well (-1 vs -11) ($P = .56$).

MUSCLE WASTING IN BOTH STRAINS OF MICE

The E3 ubiquitin ligase MuRF1 targets proteins for degradation in the ubiquitin-proteasome system. Muscles undergoing protein degradation have elevated levels. When hind leg quadriceps muscles were analyzed for MuRF1, both strains of mice had evidence of muscle atrophy when compared with the control mice without tumors (**Figure 6**). The same regulation was seen with gastrocnemius muscles (data not shown).

COMMENT

The genetic predisposition of the host to mount an inflammatory response may determine the magnitude of cancer cachexia. Animal studies suggest that cytokine activation plays a central role in the genesis of fat and muscle catabolism seen in cachexia.⁴ In addition, tumor-

derived mediators such as proteolysis-inducing factor and lipid-mobilizing factor serve as a stimulus for proinflammatory cytokine production and contribute to the development cachexia. Because LPS activation of the TLR4 pathway results in the production of the same inflammatory cytokines seen in cachexia, we hypothesized that the ability to activate the TLR pathway would lead to production of inflammatory mediators that may exacerbate cancer cachexia. Skeletal muscles express TLR4 and are known to mount an inflammatory response to LPS, with dose-dependent increases in muscle messenger RNA (mRNA) levels of IL-1 β and other cytokines.^{11,12} Because cachexia is thought to be mediated by many of the same proinflammatory cytokines, it follows that TLR4^{del/del} mice would develop less severe cancer cachexia.

In the present study, we demonstrate that TLR4^{del/del} mice have evidence of muscle wasting but the cachexia was less severe than in their TLR4-intact counterparts. The wild-type mice had less fat and lean body mass, resulting in a significant difference in weight. This was also associated with a molecular marker of muscle atrophy, the up-regulation of the E3 ligase MuRF1, and elevated IL-1 β levels.

To design effective therapeutic strategies to combat cachexia, a better understanding of the role of cytokines, both host and tumor derived, is essential. As such, we believe that C3H/HeN and C3H/HeJ (TLR4^{del/del}) mice could serve as a model system to study altered host cytokine production. We tested the serum of wild-type and mutant mice for the presence of 18 cytokines and chemokines before and after tumor challenge. None of the TLR4^{del/del} mice exhibited a significant intact inflammatory response after tumor challenge, which is consistent with results seen in experimental systems investigating the response to LPS-induced sepsis.¹¹ The fact that these mice did not exhibit cachexia that was as severe supports the hypothesis that cytokine activation is important in the pathogenesis of cachexia.

To our knowledge, this model, HeN mice with SCCF tumor cells, is the first immunocompetent model of head and neck cancer cachexia. Previous work demonstrated that the SCCF tumor cells overexpress lipid-mobilizing factor, and this is associated with decreased fat mass in the tumor-bearing HeN mice.²¹ The addition of the congenic C3H/HeJ (TLR4^{del/del}) strain creates a novel system in which the host contributions to cachexia may be further studied. For instance, the role of TLR4 could be elucidated in this system, along with other downstream mediators of the TLR system, including nuclear factor- κ B.

Several mechanisms have been proposed for the pathogenesis of wasting seen in cachexia, including the inappropriate production and release of cytokines, with or without activation of nuclear factor- κ B. Nuclear factor- κ B is a transcription factor that is activated by TNF- α , which reportedly inhibits skeletal muscle differentiation by suppressing myosin-D messenger RNA and protein. Reduced myosin-D expression impairs the repair of skeletal muscles, reducing lean body mass.²² Toll-like receptor 4 also induces activation of signaling molecules such as nuclear factor- κ B.²³ Because TLR4 is expressed on skeletal muscle cells and activation of TLR4 results in the activation of the same pathways seen in cytokine-induced protein degradation,




Figure 4. Comparison of lean body mass and fat mass in mice with tumors. A, An example of the body composition analysis results obtained using the Lunar PIXImus Densitometer 2.00 (software version 1.42.006.010; Lunar Corp, Madison, Wisconsin). BMC indicates bone mineral content; BMD, bone mineral density; and ROI, region of interest. B, Comparison of the percentage change in lean body mass in wild-type C3H/HeN (HeN) and TLR4^{del/del} C3H/HeJ (HeJ) mice. C, Comparison of percentage change in fat mass between HeN and HeJ mice after tumor challenge. Mice in each group underwent pretumor and posttumor body composition analysis. Percentage change = [(final % - initial %) \times 100 - 100]. For both lean body mass and fat mass, the changes are significant ($P < .001$).

we hypothesized that mice with intact TLR4 pathways would have more tumor-induced muscle wasting. Following tumor challenge, wild-type HeN mice exhibited the classic body composition changes seen in cachexia. Previous work demonstrated that hind leg weights are lower in wild-type mice with tumors than in control mice.²¹ Body composition analysis confirmed that wild-type mice exhibited less lean body and fat mass compared with TLR4^{del/del} mice. This was manifested as a significant difference in weight despite increased food consumption.

Although we have provided preliminary evidence that TLR4 may be a key determinant of the severity of tumor-induced cachexia, more research is needed to determine the exact mechanism by which this pathway contributes to tumor-associated inflammation and wasting. It is possible that IL-1 β mediates muscle wasting through the inhibition of protein synthesis and not the proteasome pathway because both strains of mice had up-regulation of MuRF1, yet the TLR4^{del/del} mice had more lean body mass. In addition, the matched TLR4^{del/del} mice had significantly larger tumors, which merits further investigation. This may demonstrate that targeted therapies for cancer cachexia may not necessarily be sound anticancer therapies. Therefore,

Table 2. Comparison of Pretumor and Posttumor Serum Cytokine and Chemokine Analyses^a

Cytokine	C3H/HeN Mice, Mean (Posttumor – Pretumor Level)	C3H/HeJ Mice, Mean (Posttumor – Pretumor Level)	Mean (95% CI) Difference of C3H/HeN – C3H/HeJ Levels	P Value
IL-1 β	259	-100	359 (50 to 668)	.03
IFN- γ	-1	-11	10 (-30 to 51)	.56
IL-6	27	-112	139 (-198 to 477)	.33
TNF- α	-242	-5053	4811 (-7517 to 17138)	.34
IL-1 α	-76	-224	148 (-204 to 501)	.32
IL-2	7	-12	18 (-1 to 37)	.06
IL-4	0	-1	1 (-1 to 2)	.22
IL-5	-3	-4	1 (-4 to 7)	.61
IL-10	89	-3	92 (-194 to 379)	.41
GM-CSF	5	-3	8 (-16 to 30)	.42
IL-3	-2	-1	-1 (-8 to 5)	.60
IL-12 (p40)	44	193	-148 (-937 to 641)	.65
IL-12 (p70)	70	-83	153 (-171 to 478)	.29
IL-17	129	-160	289 (-253 to 832)	.24
G-CSF	11	147	-136 (-544 to 271)	.45
KC	98	230	-132 (-413 to 150)	.31
MIP-1 α	20	-15	35 (-4 to 74)	.07
RANTES	143	233	-90 (-294 to 114)	.32

Abbreviations: CI, confidence interval; GM-CSF, granulocyte macrophage colony-stimulating factor; G-CSF, granulocyte colony-stimulating factor; IFN- γ , interferon γ ; IL, interleukin; TNF- α , tumor necrosis factor α ; KC, mouse chemokine N51; MIP-1 α , macrophage inflammatory protein 1 α ; RANTES, regulated on activation, normal T expressed and secreted.

^aUnless otherwise indicated, results are expressed as picogram per milliliter concentrations of the differences between the pretumor and posttumor levels. Serum collected from wild-type C3H/HeN and TLR4^{d/d} C3H/HeJ mice was analyzed for 18 different cytokines using the Bioplex protein array system (Biorad, Hercules, California). Only IL-1 β was found to be significantly elevated following tumor challenge.




Figure 5. Comparison of interleukin 1 β (IL-1 β) levels in wild-type C3H/HeN (HeN) and TLR4^{d/d} C3H/HeJ (HeJ) mice. Both strains of mice had the same mean levels of circulating IL-1 β before the tumor cells were inoculated, and IL-1 β level was increased only in the wild-type mice after tumors were present ($P = .03$).

careful studies of TLR antagonists and antiinflammatory agents will have to be performed.

In conclusion, in spite of increased food intake and smaller tumors, the mice with an intact TLR4 signaling pathway experienced more severe cachexia than the mice with deficient signaling. This was evidenced by lower body weights, less lean body mass and fat mass, and clinical evidence of wasting compared with the age- and weight-matched TLR4^{d/d} mice. These results are more striking because wild-type mice without tumors tend to gain slightly more weight than the TLR4^{d/d} mice. Interleukin 1 β appears to be an important mediator of cachexia in this model. Since the mice are congeneric except for this




Figure 6. Markers of muscle atrophy, E3 ligase MuRF1 and GAPDH, in the mice. Up-regulation of MuRF1 is seen in wild-type C3H/HeN (HeN) and TLR4^{d/d} C3H/HeJ (HeJ) mice with tumors but not in control mice without tumors.

double mutation in TLR4 pathway signaling and are inoculated with the same number of identical tumor cells, the impaired ability to secrete proinflammatory cytokines such as IL-1 β may protect these animals from de-

veloping more severe cancer cachexia. This animal model represents a novel system in which host contributions to cachexia may be further studied.

Submitted for Publication: October 27, 2006; final revision received March 23, 2007; accepted April 18, 2007.

Author Affiliations: Department of Otolaryngology–Head & Neck Surgery and The Lineberger Comprehensive Cancer Center (Drs Cannon, Shores, and Couch) and the Doris Duke Research Fellowship Program, General Clinical Research Center (Mr George and Dr Lai), School of Medicine, and Department of Biostatistics, School of Public Health (Dr Bůžková), University of North Carolina, Chapel Hill; and the Division of Human Cancer Genetics, Department of Molecular Virology, Immunology, and Medical Genetics, the Integrated Biomedical Graduate Program, Department of Pathology, and The Arthur G. James Comprehensive Cancer Center, The Ohio State University, Columbus (Dr Guttridge and Mr Dahlman).

Correspondence: Marion E. Couch, MD, PhD, Department of Otolaryngology–Head & Neck Surgery, G0412 Neurosciences Hospital, University of North Carolina School of Medicine, CB #7070, Chapel Hill, NC 27599-7070 (marion_couch@med.unc.edu).

Author Contributions: Dr Couch had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. *Study concept and design:* Cannon, Guttridge, Shores, and Couch. *Acquisition of data:* Cannon, Guttridge, Dahlman, Lai, and Couch. *Analysis and interpretation of data:* Cannon, George, Bůžková, and Couch. *Drafting of the manuscript:* Cannon, Guttridge, George, Bůžková, and Couch. *Critical revision of the manuscript for important intellectual content:* Cannon, Dahlman, Lai, Shores, and Couch. *Statistical analysis:* Bůžková and Couch. *Obtained funding:* Couch. *Administrative, technical, and material support:* Guttridge, Dahlman, George, Lai, and Couch. *Study supervision:* Couch.

Financial Disclosure: None reported.

Funding/Support: This research was supported in part by grant RR00046 from the Doris Duke Clinical Research Fellowship program and grant NIH DK056350 from the National Institutes of Health.

Previous Presentation: This article was presented at The American Head & Neck Society 2007 Annual Meeting; April 29, 2007; San Diego, California.

REFERENCES

1. Theologides A. Cancer cachexia. *Cancer*. 1979;43(5)(suppl):2004-2012.
2. Inagaki J, Rodriguez V, Bodey GP. Proceedings: causes of death in cancer patients. *Cancer*. 1974;33(2):568-573.
3. Tisdale MJ. Cancer cachexia. *Langenbecks Arch Surg*. 2004;389(4):299-305.
4. Argilés JM, Lopez-Soriano FJ. The role of cytokines in cancer cachexia. *Med Res Rev*. 1999;19(3):223-248.
5. Tisdale MJ. Cachexia in cancer patients. *Nat Rev Cancer*. 2002;2(11):862-871.
6. Strassmann G, Masui Y, Chizzonite R, Fong M. Mechanisms of experimental cancer cachexia: local involvement of IL-1 in colon-26 tumor. *J Immunol*. 1993;150(6):2341-2345.
7. Hoebe K, Jiang Z, Tabeta K, et al. Genetic analysis of innate immunity. *Adv Immunol*. 2006;91:175-226.
8. Kopp EB, Medzhitov R. The Toll-receptor family and control of innate immunity. *Curr Opin Immunol*. 1999;11(1):13-18.
9. Huang B, Zhao J, Li H, et al. Toll-like receptors on tumor cells facilitate evasion of immune surveillance [published correction appears in *Cancer Res*. 2005;65(19):9108. Chen, Shu-Hsia (added)]. *Cancer Res*. 2005;65(12):5009-5014.
10. Takeda K, Akira S. Toll-like receptors in innate immunity. *Int Immunol*. 2005;17(1):1-14.
11. Frost RA, Nystrom GJ, Lang CH. Lipopolysaccharide regulates proinflammatory cytokine expression in mouse myoblasts and skeletal muscle. *Am J Physiol Regul Integr Comp Physiol*. 2002;283(3):R698-R709.
12. Lang CH, Silvis C, Deshpande N, Nystrom G, Frost RA. Endotoxin stimulates in vivo expression of inflammatory cytokines tumor necrosis factor alpha, interleukin-1beta, -6, and high-mobility-group protein-1 in skeletal muscle. *Shock*. 2003;19(6):538-546.
13. Watson J, Kelly K, Lergen M, Taylor BA. The genetic mapping of a defective LPS response gene in C3H/HeJ mice. *J Immunol*. 1978;120(2):422-424.
14. Watson J, Riblet R. Genetic control of responses to bacterial lipopolysaccharides in mice, I: evidence for a single gene that influences mitogenic and immunogenic responses to lipopolysaccharides. *J Exp Med*. 1974;140(5):1147-1161.
15. Poltorak A, He X, Smirnova I, et al. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science*. 1998;282(5396):2085-2088.
16. Qureshi ST, Lariviere L, Leveque G, et al. Endotoxin-tolerant mice have mutations in Toll-like receptor 4 (Tlr4). *J Exp Med*. 1999;189(4):615-625.
17. Takeuchi O, Hoshino K, Kawai T, et al. Differential roles of TLR2 and TLR4 in recognition of gram-negative and gram-positive bacterial cell wall components. *Immunity*. 1999;11(4):443-451.
18. Segreti J, Gheusi G, Dantzer R, Kelley KW, Johnson RW. Defect in interleukin-1beta secretion prevents sickness behavior in C3H/HeJ mice. *Physiol Behav*. 1997;61(6):873-878.
19. Iida-Klein A, Lu SS, Yokoyama K, Dempster DW, Nieves JW, Lindsay R. Precision, accuracy, and reproducibility of dual X-ray absorptiometry measurements in mice in vivo. *J Clin Densitom*. 2003;6(1):25-33.
20. Acharyya S, Ladner KJ, Nelsen LL, et al. Cancer cachexia is regulated by selective targeting of skeletal muscle gene products. *J Clin Invest*. 2004;114(3):370-378.
21. Cannon TY, Shores C, Yin X, et al. Immunocompetent murine model of cancer cachexia for head & neck squamous cell carcinoma [published online ahead of print July 26, 2007]. *Head Neck*. doi:10.1002/hed.20698.
22. Li YP, Reid MB. Effect of tumor necrosis factor-alpha on skeletal muscle metabolism. *Curr Opin Rheumatol*. 2001;13(6):483-487.
23. Takeda K, Kaisho T, Akira S. Toll-like receptors. *Annu Rev Immunol*. 2003;21:335-376.