

UC Irvine

UC Irvine Previously Published Works

Title

Sequence and function of the two P domain potassium channels: implications of an emerging superfamily

Permalink

<https://escholarship.org/uc/item/47f4n73b>

Journal

Journal of Molecular Medicine, 76(1)

ISSN

0946-2716

Authors

Goldstein, SAN
Wang, K-W
Ilan, Nitza
[et al.](#)

Publication Date

1998

DOI

10.1007/s001090050186

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

REVIEW

Steve A.N. Goldstein · Ke-Wei Wang · Nitza Ilan
Mark H. Pausch

Sequence and function of the two P domain potassium channels: implications of an emerging superfamily

Received: 28 January 1997 / Accepted: 16 June 1997



STEVE A. N. GOLDSTEIN received his M.D. and Ph.D. from Harvard University. He completed his residency and fellowship training in Pediatric Medicine and Cardiology at the Children's Hospital of Boston. After a postdoctoral fellowship in the Department of Biochemistry of Brandeis University, he joined the Yale University School of Medicine, Department of Pediatrics in New Haven, Conn. Steve is an Associate Professor of Pediatrics and Cellular and Molecular Physiology and Chief of Developmental Biology and Biophysics.



MARK H. PAUSCH received his B.S. in Biochemistry from the University of Illinois at Urbana-Champaign and Ph.D. from the Department of Biochemistry, Molecular Biology, and Cell Biology at Northwestern University. After a postdoctoral fellowship in the Department of Molecular and Cellular Biology of the University of California at Berkeley, he is presently a Senior Research Biologist at the American Cyanamid Company's Agricultural Research Center in Princeton, N.J.

S.A.N. Goldstein (✉) · K.-W. Wang · N. Ilan
Section of Developmental Biology and Biophysics,
Departments of Pediatrics and Cellular and Molecular Physiology,
Boyer Center for Molecular Medicine,
Yale University School of Medicine, 295 Congress Avenue,
New Haven, CT 06536, USA

M.H. Pausch
American Cyanamid Agricultural Research Center, Princeton,
NJ, USA

Abstract A new superfamily of K^+ channels has emerged in the past 2 years. Notable for possessing two pore-forming P domains in each subunit, members of the superfamily have been recognized through phylogeny from micro-organisms to humans. Four subfamilies of two P domain channels have been isolated thus far; among these are the first cloned examples of outward rectifier and open rectifier (or leak) K^+ channels. The two P domain K^+ channels offer a new perspective from which to glimpse the molecular basis for function and dysfunction of K^+ -selective ion channels.

Key words Potassium channel · Two P domains · *TOK1* · d-ORK1 · HOHO1 · h-TPKC1

Abbreviations NCBI National Center for Biotechnology Information

Introduction

Ion channels are proteins that reside in the plasma membranes of all cells and control their electrical activity [1]. In response to stimuli, such as neurotransmitters, mechanical stress, and voltage changes, channels open a water-filled pore across the membrane through which selected ions passively diffuse [2]. In this simple fashion ion channels mediate rapid signaling events that grant us sight, sensation, movement, and thought and execute slower, but essential, cellular house-keeping duties such as fluid and electrolyte homeostasis.

The first K^+ channel gene to be isolated, Shaker, was identified as the cause of a motion disorder in fruit flies [3–5]. Over 80 related K^+ channel genes have been cloned based on their sequence homology to Shaker, each encoding a protein with a single pore-forming P domain and six probable transmembrane segments (1P/6TM protein subunits) [6]. More recently, genes for K^+ channels have been isolated by expression cloning, a method that does not select for homology but identifies genes whose products show ion channel function [7–10],

or by computer searches for channel-like motifs in the expanding global database of nucleotide and protein sequences [11, 12]. It is by these later methods that members of the two P domain superfamily of K^+ channels were discovered.

Here we describe examples of each of the four recognized lineages of the new superfamily. *TOK1* was the founding member of the superfamily [11]. Isolated from *Saccharomyces cerevisiae*, it is an outwardly rectifying K^+ channel with two P domains and eight proposed transmembrane segments (2P/8TM). d-ORK1 was isolated from *Drosophila melanogaster* [10]; it is an open-rectifier (or leak) K^+ channel with two P domains and only four proposed transmembrane segments (2P/4TM). h-TPKC1, from human brain, shares a probable 2P/4TM structure with d-ORK1 but, like Tok1, is an outward rectifier [13]. Finally, HOHO1 was cloned from human brain and has a probable 2P/4TM structure (this report); we find it to be nonfunctional in oocytes while others report that it is a weak inward rectifier [12]. This growing superfamily of two P domain K^+ channels is examined below in the context of knowledge about K^+ channels with a single P domain.

Potassium channel function

Potassium channels are both efficient and highly selective; some catalyze flux of 100 million K^+ ions each second through a single channel complex while passing only one Na^+ ion in error every 10,000 events [14]. Because K^+ ions are the predominant monovalent cation in mammalian cells, and extracellular K^+ concentrations are low, opening K^+ channels favors an outpouring of these positively charged ions at the positive voltages characteristic of excitation. Outflow of K^+ ions shifts the voltage across the cell membrane toward the equilibrium reversal potential for K^+ ions (E_K), that is, the voltage where the tendency for K^+ ions to move outwardly (down their con-


centration gradient) is balanced by their tendency to move inward (down their electrical gradient). By this mechanism open K^+ channels mediate recovery after activity in excitable tissues such as nerves, skeletal muscles, and the heart and stabilize cellular potential near E_K [15]. Before their genes were cloned, individual K^+ channels were studied in native tissues and categorized by their functional attributes [16, 17]. Cloning of genes for single P domain K^+ channels has revealed examples of many but not all functional categories of K^+ channels; numerous voltage-gated, inward-rectifier, and ligand-gated K^+ channels are now known.

Voltage-gated K^+ channels

Voltage-activated K^+ channels undergo changes in protein conformation in response to changes in transmembrane potential to produce an open channel state that allows ion permeation (Fig. 1). Channels in this group show marked variation in opening and closing rates, and this underlies their wide diversity of roles in vivo. *Delayed rectifiers* activate with a delay after membrane depolarization and close only after cells return to resting potential. By opening relatively slowly, these channels do not inhibit the early explosive rising phase of the action potential but act in a delayed fashion to repolarize cells and prepare them for subsequent stimulation (Fig. 1b). On the other hand, *A currents* are voltage-gated channels that inactivate rapidly after opening, bringing channels to a nonconducting state unique from the closed state (Fig. 1a). These channels control the rate at which cells reach firing threshold and thus the interval between excitatory events. Both delayed rectifier and A-type voltage-gated K^+ channel subunits have a 1P/6TM topology (Fig. 1d). Indeed, these two phenotypes can be interconverted by site-directed mutations [18], heteromultimeric channel formation [19–21], and interaction with accessory subunits (see below) to alter the kinetics of inactivation.

Heteromeric subunit association is required for function of the slow delayed rectifier of human heart I_{K_S} [22–25] and perhaps for its rapid counterpart I_{K_r} as well [26]. These channels form through association of 1P/6TM protein subunits (KvLQT1 and HERG, respectively) and minK, a unique 130 residue protein with one transmembrane stretch that appears to contribute to for-

Fig. 1a–d Drawings to represent common attributes of voltage-gated K^+ channels. **a** Idealized A-type currents in whole oocytes in response to a family of voltage pulses from a holding voltage of -80 mV to command voltages of -40 to $+40$ mV in 10 -mV steps. **b** Idealized delayed rectifier currents subjected to the protocol in **a**. **c** Peak current/voltage relationships. **d** Probable 1P/6TM membrane topology. Scale bars, 1 μ A and 5 ms



mation of the channel pore [27]. Mutations in KvLQT1 or HERG that decrease channel function and slow cardiac repolarization have been found in some patients with genetically determined “long QT syndrome,” a dysrhythmia that prolongs action potential duration, predisposing to torsade de pointe and sudden death [28–31].

Inward rectifier K⁺ channels

Ion channels that preferentially pass K⁺ ions inward are called inward rectifier K⁺ channels. They show a steep voltage dependence, but unlike voltage-gated K⁺ channels which have a fixed response to voltage, inward rectifier channels shift the voltage at which they pass current with changes in external K⁺ ion concentration (Fig. 2c). This reflects the mechanism underlying opening and rectification in these channels – intracellular magnesium and polyamines tonically occlude the ion conduction pore and are expelled only when voltage or changing K⁺ ion concentration favors inward movement of K⁺ ions. Inward rectifiers stabilize the membrane near E_K and thus counter the hyperpolarizing effects of electrogenic sodium pump activity and the depolarizing effects of pacemaker currents [1]. These K⁺ channel subunits have a 1P/2TM topology (Fig. 2d). A unique inward rectifier from renal cortical tubular cells that exhibits marked sensitivity to pH, RACTK1, has a modified P domain and two transmembrane stretches [32, 33].

Inwardly rectifying currents can also be produced by some 1P/6TM voltage-gated channels with a particular type of gating behavior – rapid, voltage-dependent inactivation [34, 35]. Such channels have limited outflow of K⁺ ions at positive potentials because they rapidly enter an inactive state after opening in response to depolarization. When the membrane repolarizes, the channels recover from inactivation and reopen as they return to the closed state. During the reopened period large inward currents are seen as K⁺ ions move down their electro-

chemical gradient into the cell. These channels are present in cells of widely disparate origin including the guard cells of plants [36] and mammalian cardiac myocytes [35].

Ligand-gated K⁺ channels

Some K⁺ channels are classified by the molecules that regulate their function. Three examples include calcium-activated K⁺ channels that are opened by elevations in intracellular calcium or calcium and voltage in concert; these contain 1P/6TM subunits [37, 38]. ATP-sensitive K⁺ channels are blocked by ATP, activated by ADP, and appear to act as sensors for the metabolic state of the cell; these channels are heteromultimers of 1P/2TM subunits and regulatory sulfonylurea receptors, members of the ABC transporter family [39] and carry ligand binding sites on both subunit types [40]. Acetylcholine-sensitive K⁺ channels change their open probability in response to binding G_{βγ} liberated by muscarinic receptor stimulation; these channels are heteromultimers of 1P/2TM channel subunits [41].


Accessory subunits

Single P domain subunits function in macromolecular complexes containing one or more accessory subunits. Required for integrated channel function, accessory subunits regulate channel expression levels, modify gating kinetics, mediate responses to ligands, and in some cases are required for activity [42]. Thus K_β subunits speed inactivation of some voltage-gated delayed rectifier K⁺ channels [43, 44], calcium-activated K⁺ channel β subunits increase sensitivity to calcium [45], and ATP-sensitive channels operate when their P domain subunits aggregate with sulfonylurea receptors [46].

The P domain

Pores are the catalytically active sites of ion channels, and identification of P domains was a major step in understanding the structural basis for channel function [47–51]. The P domain residues in Shaker K⁺ channels were found by systematic point mutation in an effort to

Fig. 2a–d Drawings to represent common attributes of inward rectifier K⁺ channels. **a** Whole cell currents in high levels of external KCl (90 mM) in response to a family of voltage pulses from a holding voltage of 0 mV to command voltages of –160 to +50 mV in 30-mV steps. **b** In 20 mM external KCl. **c** Steady-state current/voltage relationships in 90 and 20 mM external KCl. **d** Probable 1P/2TM membrane topology. Scale bars, 1 μA and 5 ms



a

Shaker		I P D A F W W A V V <u>T M T T V G Y G</u> D M T P
TOK P1		Y G N A L Y F C T V S L L T V G L G D I L P
TOK P2		Y F N C I Y F C F L C L L T I G Y G D Y A P
ORK P1		F Y H A F F F A F T V C S T V G Y G N I S P
ORK P2		S S I S L Y Y S Y V T T T T I G F G D Y V P
TPKC P1		L G S S F F F A G T V I T T I G F G N I S P
TPKC P2		A L D A I Y F V V I T L T T I G F G D Y V A
HOHO P1		F T S A L F F A S T V L S T T G Y G H T V P
HOHO P2		F L E S F Y F C F I S L S T I G L G D Y V P



b

Fig. 3a, b P domain amino acid sequences and model for tetrameric pore formation. **a** Residues of the P domains in Shaker, Tok1, d-ORK1, h-TPKC1, and HOHO1 K⁺ channel subunits (Genbank accession numbers M17211, U28005, U55321, AF004711, and U76996, respectively); underlined, the P domain signature sequence residues in Shaker [53]. **b** A drawing to suggest how four P domains may form the pore lining in voltage-gated and inward rectifier K⁺ channels

disrupt high-affinity pore blockade by a scorpion toxin [48]. That residues linking the fifth and sixth transmembrane segments enter the membrane to form the ion conduction pathway was shown by two approaches. First, two binding sites in this region were identified for the

Fig. 4a–d Tok1 K⁺ channels. **a** Currents measured by two electrode voltage clamp in an oocyte injected 2 days previously with 1 ng *TOK1* cRNA and bathed in 2 mM external KCl with voltage pulses of 100 ms from a holding voltage of –80 mV to command voltages of –150 to +45 mV in 15-mV steps. For method details see [11]. **b** Currents measured in 40 mM KCl. **c** steady-state current/voltage relationships for measurements shown in **a** and **b**. **d** proposed 2P/8TM membrane topology. Scale bars, 1 μ A and 25 ms



pore blocker tetraethylammonium ion, one extracellular and one intracellular [49, 50]. Second, a channel carrying transplanted linker residues exhibited pore characteristics of the donor including single-channel conductance and affinity for internal and external tetraethylammonium ion [51]. Later a “signature sequence” of eight highly conserved residues that are critical to K⁺ ion selectivity was identified in the P domain, TxTTxGYG [52, 53]. Figure 3a shows the residues found in the P domain of Shaker as well as those in examples from each of the four known subfamilies of two P domain K⁺ channels.

Site-directed mutagenesis has revealed that K⁺ channel subunits with a single P domain associate as tetrameric aggregates [54, 55]. In these channels the ion conduction pathway is lined by four P domains and portions of the adjacent hydrophobic transmembrane stretches (Fig. 3b) [56–58]. Voltage-gated Na⁺ and Ca²⁺ have a similar fourfold pseudosymmetric pore [1]. These larger proteins carry four homologous domains, each like a complete voltage-gated K⁺ channel subunit with a P domain between every fifth and sixth membrane-spanning segment, and the four P domains fold together to form a central conduction pore. While the three-dimensional structure of intact K⁺ channels has not yet been determined, the K⁺ selectivity filter appears to form through association of pore “loops” like those seen in the bacterial porins [59].

Tok1, novel structure and function

Tok1, was the first K⁺ channel identified to carry two P domains in one continuous polypeptide [11]. The gene was identified on *Saccharomyces cerevisiae* chromosome X by searching with the BLAST algorithm the data collected by the yeast genome sequencing project and available through the National Center for Biotechnology Information (NCBI). The Tok1 gene encodes a protein of 691 amino acids with two P domains and eight probable transmembrane segments (Fig. 4c). When expressed in *Xenopus laevis* oocytes, it produced a current unlike any previously cloned channel but reminiscent of a current described in guard cells of the plant species, *Vicia* [60].


Tok1 is an *outward rectifier* that preferentially passes outward K^+ currents in a fashion coupled to changes in the external K^+ concentration (Fig. 4c). This behavior is similar to that of inward rectifier channels but in the reverse direction (Fig. 2c). It is unlike the behavior of voltage-gated channels, whose gating depends on transmembrane potential in an invariant fashion (Fig. 1c). The macroscopic kinetics of Tok1 currents reveal two components, one rapid, the other time dependent (Fig. 4b) [11, 61, 62].

Much remains to be discovered about Tok1. The mechanism underlying outward rectification is as yet unknown. Tok1 is expressed in the plasma membrane of yeast cells [63] and regulated by protein kinase C and intracellular pH [61], but its functional role *in vivo* is undetermined. Based on the tetrameric structure of single P domain channels, it seems likely that dimers of Tok1 will be shown to form a single conduction pore, but this also requires experimental confirmation.

d-ORK1, a second lineage of two P domain channels with just 4 probable TM segments

d-ORK1 is a K^+ channel with two P domains expressed in the neuromuscular tissues of *Drosophila melanogaster* [10]. The gene was isolated from a cDNA expression library produced from fly larvae based on its ability to complement a strain of yeast defective in K^+ transport. Both the high- and low-affinity K^+ transport proteins (Trk1, Trk2) are disrupted in the yeast cells, limiting their ability to grow in low- K^+ medium. Expression of d-ORK1 in this strain allows the cells to survive in low- K^+ medium and confers the ability to accumulate K^+ ions.

Fig. 5a–d d-ORK1 K^+ channels. **a** Currents measured by two electrode voltage clamp in an oocyte injected 2 days previously with 0.2 ng d-ORK1 cRNA and bathed in 5 mM external KCl with voltage pulses of 100 ms from a holding voltage of -80 mV to command voltages of -150 to $+60$ mV in steps of 15 mV. For method details see [10]. **b** Currents measured in 50 mM KCl. **c** Steady-state current/voltage relationship in **a** and **b**. **d** Proposed 2P/4TM membrane topology. Scale bars, 2 μ A and 25 ms



Based on its nucleotide sequence and glycosylation pattern, the channel protein appears to contain two P domains and only four transmembrane segments [10].

When expressed in *Xenopus* oocytes, d-ORK1 behaves as an open rectifier (or leak) K^+ channel, that is, currents behave as predicted for an open K^+ -selective pore in an electric field. Currents are instantaneous with changes in voltage and rectify when transmembrane K^+ concentrations are unequal (Fig. 5a). These attributes are similar to those of the unidentified leak K^+ current underlying resting membrane potential in myelinated nerves of vertebrates [64].

d-ORK1, like Tok1, is still poorly understood. How do complete channels form? How are they regulated? Do d-ORK1 channels in fact mediate cellular resting potential? To address this question agents that act on the channel are being identified in rapid, large-scale screens exploiting channel-dependent growth of yeast cells (see [65]). Identified agents will later be used to evaluate the channels' role *in vivo*. Indeed, the ability to functionally express d-ORK1 in yeast cells allows application of a wide array of powerful genetic tools to study its structure, regulation, and function [66].

h-TPKC1, an outward rectifier from human brain related to d-ORK1

A degenerate PCR approach (using primers based on the P domain sequences of d-ORK1 and several putative two P domain channels in the genome of *C. elegans*) were used to clone h-TPKC1, so named as an abbreviation for two P domain K^+ channel, from a human brain cDNA library [13]. That h-TPKC1 encodes a functional 2P/4TM K^+ channel was deduced from its expression in *Saccharomyces cerevisiae* cells defective for K^+ uptake which survived on low- K^+ medium as a result. Evaluation of the channel when expressed in *Xenopus* oocytes using two electrode voltage clamp revealed a largely instantaneous, noninactivating, K^+ -selective outward current when measured at physiological levels of external K^+ (approx. 5 mM). As bath K^+ levels increased, the current reversal potential shifted toward E_K , in accordance with

the Nernst equation, and only a modest inward potassium current was detected (data not shown). These properties are held in common with those of Tok1, indicating that outwardly rectifying channel behavior does not require a 2P/8TM structure. A mouse homolog of h-TPKC1 was cloned while this work was in progress and is called mTREK1 [67]. The channels' are highly similar (85% identical) and exhibit common biophysical properties. Despite their similarity, their tissue and brain region specific pattern of expression differ substantially. Northern blotting has revealed that the strongest h-TPKC1 signal corresponds to a 3.8-kb brain transcript found at low steady-state levels in skeletal muscle, small intestine, and colon but undetectable in lung, kidney, and heart. In contrast, substantial mTREK1 mRNA is present in brain, lung, kidney, and heart.

HOHO1, another human homolog of d-ORK1

A search of the NCBI database of expressed sequence tags with the nucleotide sequence for d-ORK1 repeatedly identified a homologous sequence in humans. We isolated HOHO1 (for human *ORK-homologous open reading frame*), from human brain cDNA (NCBI accession no. U76996). The protein has 337 amino acids and, as with d-ORK1, a probable 2P/4TM topology. HOHO1 is identical in its coding region to a human kidney protein, TWIK1 [12], and similar to a murine clone, mTWIK1 [68]; the TWIKs are reported to behave as inwardly rectifying K⁺ channels when expressed in *Xenopus* oocytes. We have observed no currents by two-electrode voltage clamp when studying oocytes injected with 1–10 ng cRNA encoding HOHO1 in constructs that yield high expression of Tok1, d-ORK1 and h-TPKC1 [10, 11, 13]. This suggests that HOHO1/TWIK1 may need to associate with other channel subunits, is only a partial clone, or is subject to regulatory influences that vary between oocyte preparations.

Conclusions

Since the cloning of Shaker in 1988, over 100 K⁺ channels with a single pore-forming P domain on each subunit have been identified [6]. This has revealed the molecular basis for many K⁺ currents essential to normal physiology and allowed elucidation of the mechanism underlying a number of human diseases [24, 25, 30, 69, 70]. Now, a new superfamily of K⁺ channels containing two P domains has emerged. Members of the new superfamily appear likely to be as numerous as their single P domain cousins. In the 2 years since the cloning of TOKI [11] three additional subtypes of two P domain channels have been identified which carry four rather than eight probable transmembrane segments. Database searching reveals this 2P/4TM subfamily to be common in the genomes of many organisms, indeed just the ge-

nome of the nematode *C. elegans* shows over 20 predicted open reading frames with this pattern [71, 72].

These findings herald four changes in the field of K⁺-selective ion channels. First, increasingly rich databases for genomic and expressed nucleotide sequences are changing the way in which genes for ion channels are identified, cloned, and studied. Second, it is now clear that K⁺-selective ion channels can be formed by association of protein subunits with either one or two P domains; how their structures are similar and how unique remains to be revealed. Third, the functional repertoire of cloned K⁺ channels is growing; although two P domain channels share attributes common to known K⁺ channels, such as high selectivity for potassium, they also exhibit novel phenotypes, such as outward rectifier and open (or leak) K⁺ currents. Fourth, powerful new methods for exploring the molecular basis for function of K⁺ channels from animals are now feasible through expression of K⁺ channels in yeast cells [73]. It is premature to conjecture the roles two P domain channels will play in human health and disease. However, by determining the place of two P domain channels in normal physiology we may discover new targets for the development of therapeutic agents [73] and uncover clues to the molecular basis for disorders that result from ion channel dysfunction [15].

Acknowledgements S.A.N.G. is supported in part by grants from the NIH-NIGMS and the Donaghue Foundation; N.I. is supported by a grant from U.S.-Israel Binational Agricultural Research and Development Fund (BARD), and K. W. W. by a grant from the American Heart Association.

References

- Hille B (1992) Ionic channels of excitable membranes, 2nd edn. Sunderland, Sinauer
- Miller C (1987) How ion channel proteins work. In: Kaczmarek L, Levitan I (eds) Neuromodulation. Oxford, Oxford University Press, pp 39–63
- Papazian DM, Schwarz TL, Tempel BL, Jan YN, Jan LY (1987) Cloning of genomic and complementary DNA from Shaker, a putative potassium channel gene from *Drosophila*. *Science* 237:749–753
- Pongs O, Kecskemethy N, Muller R, Krah-Jentgens I, Baumann A, Kiltz HH, Canal I, Llamazares S, Ferrus A (1988) Shaker encodes a family of putative potassium channel proteins in the nervous system of *Drosophila*. *EMBO J* 7: 1087–1096
- Iverson LE, Tanouye MA, Lester HA, Davidson N, Rudy B (1988) A-type potassium channels expressed from Shaker locus cDNA. *Proc Natl Acad Sci USA* 85:5723–5727
- Chandy KG, Gutman GA (1993) Nomenclature for mammalian potassium channel genes. *Trends Pharmacol Sci* 14:434
- Ho K, Nichols CG, Lederer WJ, Lytton J, Vassilev PM, Kanazirska MV, Hebert SC (1993) Cloning and expression of an inwardly rectifying ATP-regulated potassium channel. *Nature* 362:31–38
- Kubo Y, Reuveny E, Slesinger PA, Jan YN, Jan LY (1993) Primary structure and functional expression of a rat G-protein-coupled muscarinic potassium channel. *Nature* 364:802–806
- Canessa CM, Schild L, Buell G, Thorens B, Gautschi I, Horisberger JD, Rossier BC (1994) Amiloride-sensitive epithelial Na⁺ channel is made of three homologous subunits. *Nature* 367:463–467

10. Goldstein SAN, Price LA, Rosenthal DN, Pausch MH (1996) ORK1, a potassium-selective leak channel with two pore domains cloned from *Drosophila melanogaster* by expression in *Saccharomyces cerevisiae*. Proc Natl Acad Sci 93:13256–13261
11. Ketchum KA, Joiner WJ, Sellers AJ, Kaczmarek LK, Goldstein SAN (1995) A new family of outwardly-rectifying potassium channel proteins with two pore domains in tandem. Nature 376:690–695
12. Lesage F, Guillemare E, Fink M, Duprat F, Lazdunski M, Romey G, Barhanin J (1996) TWIK-1, a ubiquitous human weakly inward rectifying K⁺ channel with a novel structure. EMBO 15:1004–1011
13. Price LA, Hellings SE, Hayashi JH, Pausch MH (1997) Functional expression of the human brain outwardly rectifying two-pore potassium channel, hTPKC1, in *Saccharomyces cerevisiae* and *Xenopus laevis* cells. (submitted)
14. Yellen G (1984) Ionic permeation and blockade in Ca²⁺-activated K⁺ channels of bovine chromaffin cells. J Gen Physiol 84:157–186
15. Goldstein SAN (1996) Ion channels: structural basis for function and disease. Semin Perinatol 20:520–530
16. Hodgkin AL, Huxley AF (1952) A quantitative description of membrane current and its application to conduction and excitation in nerve. J Physiol (Lond) 117:500–544
17. Rudy B (1988) Diversity and ubiquity of K channels. Neuroscience 25:729–749
18. Hoshi T, Zagotta WN, Aldrich RW (1990) Biophysical and molecular mechanisms of Shaker potassium channel inactivation. Science 250:533–538
19. Ruppersberg JP, Schroter KH, Sakmann B, Stocker M, Sewing S, Pongs O (1990) Heteromultimeric channels formed by rat brain potassium-channel proteins. Nature 345:535–537
20. Isacoff EY, Jan YN, Jan LY (1990) Evidence for the formation of heteromultimeric potassium channels in *Xenopus* oocytes. Nature 345:530–534
21. Chen M-L, Hoshi T, Wu C-F (1996) Heteromultimeric interactions among K⁺ channel subunits from Shaker and eag families in *xenopus* oocytes. Neuron 17:535–542
22. Wang KW, Goldstein SAN (1995) Subunit composition of minK potassium channels. Neuron 14:1303–1309
23. Tai K-K, Wang K-W, Goldstein SAN (1997) MinK potassium channels are heteromultimeric complexes. J Biol Chem 272:1654–1658
24. Sanguinetti MC, Curran ME, Zou A, Shen J, Spector PS, Atkinson DL, Keating MT (1996) Coassembly Of K(V)Lqt1 and minK (Isk) proteins to form cardiac I-Ks potassium channel. Nature 384:80–83
25. Barhanin J, Lesage F, Guillemare E, Fink M, Lazdunski M, Romey G (1996) K(V)LQT1 and IsK (minK) proteins associate to form the I(Ks) cardiac potassium current. Nature 384:78–80
26. McDonald TV, Yu Z, Ming Z, Palma E, Meyers MB, Wang KW, SAN Goldstein, GI Fishman (1997) A minK-HERG complex regulates the cardiac potassium current IKr. Nature 388:289–292
27. Wang, K-W, Tai K-K, Goldstein SAN (1996) MinK residues line a potassium channel pore. Neuron 16:571–577
28. Wang Q et al (1996) Positional cloning of a novel potassium channel gene: KVLQT1 mutations cause cardiac arrhythmias. Nat Genet 12:17–23
29. Roden DM, Lazzara R, Rosen M, Schwartz PJ, Towbin J, Vincent GM (1996) Multiple mechanisms in the long-QT syndrome – current knowledge, gaps, and future directions. Circulation 94:1996–2012
30. Sanguinetti MC, Jiang C, Curran ME, Keating MT (1995) A mechanistic link between an inherited and an acquired cardiac arrhythmia: HERG encodes the IKr potassium channel. Cell 81:299–307
31. Sanguinetti MC, Curran ME, Spector PS, Keating MT (1996) Spectrum of HERG K⁺-channel dysfunction in an inherited cardiac arrhythmia. Proc Natl Acad Sci USA 93:2208–2212
32. Suzuki M, Takahashi K, Ikeda M, Hayakawa H, Ogawa A, Kawaguchi Y, Sakai O (1994) Cloning of a pH-sensitive K⁺ channel possessing two transmembrane segments. Nature 367:642–645
33. Sutcliffe MJ, Stanfield PR (1994) Pore region of K⁺ channel RACTK1. Nature 369:616
34. Miller AG, Aldrich RW (1996) Conversion of a delayed rectifier K⁺ channel to a voltage-gated inward rectifier K⁺ channel by three amino acid substitutions. Neuron 16:853–858
35. Smith PL, Baukowitz T, Yellen G (1996) The inward rectification mechanism of the HERG cardiac potassium channel. Nature 379:833–836
36. Cao Y, Ward JM, Kelly WB, Ichida AM, Gaber RF, Anderson JA, Uozumi N, Schroeder JI, Crawford NM (1995) Multiple genes, tissue specificity, and expression-dependent modulation contribute to the functional diversity of potassium channels in *Arabidopsis thaliana*. Plant Physiol 109:1093–1106
37. Butler A, Tsunoda S, McCobb DP, Wei A, Salkoff L (1993) mSlo, a complex mouse gene encoding “maxi” calcium-activated potassium channels. Science 261:221–224
38. Kohler M, Hirschberg B, Bond CT, Kinzie JM, Marrion NV, Maylie J, Adelman JP (1996) Small-conductance, calcium-activated potassium channels from mammalian brain. Science 273:1709–1714
39. Nichols CG, Shyng SL, Nestorowicz A, Glaser B, Clement JPT, Gonzalez G, Aguilar-Bryan L, Permutt MA, Bryan J (1996) Adenosine diphosphate as an intracellular regulator of insulin secretion. Science 272:1785–1787
40. Tucker SJ, Gribble FM, Zhao C, Trapp S, Ashcroft FM (1997) Truncation of Kir6.2 produces ATP-sensitive K⁺ channels in the absence of the sulphonylurea receptor. Nature 387:179–183
41. Krapivinsky G, Gordon EA, Wickman K, Velimirovic B, Krapivinsky L, Clapham DE (1995) The G-protein-gated atrial K⁺ channel IKACH is a heteromultimer of two inwardly rectifying K(+) channel proteins. Nature 374:135–141
42. Isom LL, De Jongh KS, Catterall WA (1994) Auxiliary subunits of voltage-gated ion channels. Neuron 12:1183–1194
43. Rettig J, Heinemann SH, Wunder F, Lorra C, Parcej DN, Dolly JO, Pongs O (1994) Inactivation properties of voltage-gated K⁺ channels altered by presence of beta-subunit. Nature 369:289–294
44. Sewing S, Roeper J, Pongs O (1996) Kv beta 1 subunit binding specific for shaker-related potassium channel alpha subunits. Neuron 16:455–63
45. Knaus HG, Folander K, Garcia CM, Garcia ML, Kaczorowski GJ, Smith M, R Swanson (1994) Primary sequence and immunological characterization of beta-subunit of high conductance Ca(2+)-activated K⁺ channel from smooth muscle. J Biol Chem 269:17274–17278
46. Inagaki N, Gono T, Clement JP IV, Namba N, Inazawa J, Gonzalez G, Aguilar-Bryan L, Seino S, Bryan J (1995) Reconstitution of IKATP: an inward rectifier subunit plus the sulfonylurea receptor. Science 270:1160–1170
47. Guy HR, Seetharamulu P (1986) Molecular model of the action potential sodium channel. Proc Natl Acad Sci USA 83:508–512
48. MacKinnon R, Miller C (1989) Mutant potassium channels with altered binding of charybdotoxin, a pore-blocking peptide inhibitor. Science 245:1382–1385
49. MacKinnon R, Yellen G (1990) Mutations affecting TEA blockade and ion permeation in voltage-activated K⁺ channels. Science 250:276–279
50. Yellen G, Jurman ME, Abramson T, MacKinnon R (1991) Mutations affecting internal TEA blockade identify the probable pore-forming region of a K⁺ channel. Science 251:939–942
51. Hartmann HA, Kirsch GE, Drewe JA, Tagliatela M, Joho RH, Brown AM (1991) Exchange of conduction pathways between two related K⁺ channels. Science 251:942–944
52. Heginbotham L, Abramson T, MacKinnon R (1992) A functional connection between the pores of distantly related ion

- channels as revealed by mutant K⁺ channels. *Science* 258: 1152–1155
53. Heginbotham L, Lu Z, Abramson T, MacKinnon R (1994) Mutations in the K⁺ channel signature sequence. *Biophys J* 66:1061–1067
 54. MacKinnon R (1991) Determination of the subunit stoichiometry of a voltage-activated potassium channel. *Nature* 350: 232–235
 55. Yang J, Jan YN, Jan LY (1995) Determination of the subunit stoichiometry of an inwardly rectifying potassium channel. *Neuron* 15:1441–1447
 56. Goldstein SAN (1996) A structural vignette common to voltage sensors and conduction pores: canaliculi. *Neuron* 16: 717–122
 57. Choi KL, Mossman C, Aube J, Yellen G (1993) The internal quaternary ammonium receptor site of Shaker potassium channels. *Neuron* 10:533–41
 58. Lopez GA, Jan YN, Jan LY (1994) Evidence that the S6 segment of the Shaker voltage-gated K⁺ channel comprises part of the pore. *Nature* 367:179–182
 59. MacKinnon R (1995) Pore loops: an emerging theme in ion channel structure. *Neuron* 14:889–982
 60. Blatt MR (1991) Ion channel gating in plants: physiological implications and integration for stomatal function. *J Membr Biol* 124:95–112
 61. Lesage F, Guillemare E, Fink M, Duprat F, Lazdunski M, Romey G, Barhanin J (1996) A pH-sensitive yeast outward rectifier K⁺ channel with two pore domains and novel gating properties. *J Biol Chem* 271:4183–4187
 62. Vergani P, Miosga T, Jarvis SM, Blatt MR (1997) Extracellular K⁺ and Ba²⁺ mediate voltage-dependent inactivation of the outward-rectifying K⁺ channel encoded by the yeast gene TOK1. *FEBS Lett* 405:337–344
 63. Zhou XL, Vaillant B, Loukin SH, Kung C, Saimi Y (1995) YKC1 encodes the depolarization-activated K⁺ channel in the plasma membrane of yeast. *FEBS Lett* 373:170–176
 64. Baker M, Bostock H, Grafe P, Martius P (1987) Function and distribution of three types of rectifying channel in rat spinal root myelinated axons. *J Physiol* 383:45–67
 65. Kirsch DR (1993) Development of improved cell-based assays and screens in *Saccharomyces* through the combination of molecular and classical genetics. *Curr Opin Biotechnol* 4:543–552
 66. Uozumi N, Gassmann W, Cao Y, Schroeder JI (1995) Identification of strong modifications in cation selectivity in an Arabidopsis inward rectifying potassium channel by mutant selection in yeast. *J Biol Chem* 270:24276–24281
 67. Fink M, Duprat F, Lesage F, Reyes R, Romey G, Heurteaux C, Lazdunski M (1996) Cloning, functional expression and brain localization of a novel unconventional outward rectifier K⁺ channel. *EMBO* 15:6854–6862
 68. Lesage F, Lauritzen I, Duprat F, Reyes R, Fink M, Heurteaux C, Lazdunski M (1997) The structure, function and distribution of the mouse TWIK-1 K⁺ channel. *FEBS Lett* 402:28–32
 69. Thomas PM, Cote GJ, Wohlk N, Haddad B, Mathew PM, Rabl W, Aguilar-Bryan L, Gagel RF, Bryan J (1995) Mutations in the sulfonylurea receptor gene in familial persistent hyperinsulinemic hypoglycemia of infancy. *Science* 268:426–429
 70. Vincent A, Roberts M, Willison H, Lang B, Newsom-Davis J (1995) Autoantibodies, neurotoxins and the nervous system. *J Physiol Paris* 89:129–136
 71. Salkoff L, Jegla T (1995) Surfing the DNA databases for K⁺ channels nets yet more diversity. *Neuron* 15:489–492
 72. Wei A, Jegla T, Salkoff L (1996) Eight potassium channel families revealed by the *C-Elegans* Genome Project. *Neuropharmacology* 35:805–829
 73. Goldstein SAN, Colatsky TJ (1996) Ion channels: too complex for rational drug design? *Neuron* 16:913–919