

# UC Irvine

## UC Irvine Previously Published Works

### Title

YeiE Regulates Motility and Gut Colonization in Salmonella enterica Serotype Typhimurium

### Permalink

<https://escholarship.org/uc/item/48c3b3q8>

### Journal

mBio, 12(3)

### ISSN

2161-2129

### Authors

Westerman, TL

McClelland, M

Elfenbein, JR

### Publication Date

2021-06-29

### DOI

10.1128/mbio.03680-20

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



# YeiE Regulates Motility and Gut Colonization in *Salmonella enterica* Serotype Typhimurium

T. L. Westerman,<sup>a,b</sup> M. McClelland,<sup>c</sup>  J. R. Efenbein<sup>a,b,d</sup>

<sup>a</sup>Department of Pathobiological Sciences, University of Wisconsin—Madison, Madison, Wisconsin, USA

<sup>b</sup>Department of Clinical Sciences, North Carolina State University, Raleigh, North Carolina, USA

<sup>c</sup>Department of Microbiology and Molecular Genetics, University of California, Irvine, Irvine, California, USA

<sup>d</sup>Food Research Institute, University of Wisconsin—Madison, Madison, Wisconsin, USA

**ABSTRACT** Regulation of flagellum biosynthesis is a hierarchical process that is tightly controlled to allow for efficient tuning of flagellar expression. Flagellum-mediated motility directs *Salmonella enterica* serovar Typhimurium toward the epithelial surface to enhance gut colonization, but flagella are potent activators of innate immune signaling, so fine-tuning flagellar expression is necessary for immune avoidance. In this work, we evaluate the role of the LysR transcriptional regulator YeiE in regulating flagellum-mediated motility. We show that *yeiE* is necessary and sufficient for swimming motility. A  $\Delta yeiE$  mutant is defective for gut colonization in both the calf ligated ileal loop model and the murine colitis model due to its lack of motility. Expression of flagellar class 2 and 3 but not class 1 genes is reduced in the  $\Delta yeiE$  mutant. We linked the motility dysregulation of the  $\Delta yeiE$  mutant to repression of the anti-FlhD<sub>4</sub>C<sub>2</sub> factor STM1697. Together, our results indicate that YeiE promotes virulence by enhancing cell motility, thereby providing a new regulatory control point for flagellar expression in *Salmonella* Typhimurium.

**IMPORTANCE** The ability to finely tune virulence factor gene expression is required for bacterial pathogens to successfully colonize a host. Flagellum-mediated motility is critical for many gut pathogens to establish productive infections. However, flagella activate the immune system, leading to bacterial clearance; therefore, tight control of flagellar gene expression enhances bacterial fitness in the host. Here, we demonstrate that the transcriptional regulator YeiE acts as a control point for flagellar gene expression and is required for *Salmonella* Typhimurium to establish a productive infection in mammals. The expression of an inhibitor of flagellar biogenesis is repressed in the absence of *yeiE*. Our work adds a new layer to the tightly controlled cascade regulating control of flagellar gene expression to facilitate the fitness of an enteric pathogen.

**KEYWORDS** *Salmonella*, flagellar gene regulation, gastrointestinal infection, host-pathogen interactions, transcriptional regulation

Flagellum-mediated motility is a key virulence determinant facilitating gut colonization by many organisms, including nontyphoidal salmonellae (1, 2). Flagellum-mediated motility allows movement through the intestinal mucus layer toward the epithelium, allowing *Salmonella* to scan epithelial cells for permissive entry sites (1, 3). Upon contact with the epithelium, *Salmonella* invades intestinal cells, induces a neutrophilic inflammatory response, and replicates intracellularly, processes that require secretion of effector proteins through the two type 3 secretion systems (4). During early intracellular growth, *Salmonella enterica* serovar Typhimurium downregulates the expression of flagella to avoid immune activation induced by sensing intracellular flagellin (5). However, flagella are expressed during late intracellular infection in preparation for escape from the intracellular

**Citation** Westerman TL, McClelland M, Efenbein JR. 2021. YeiE regulates motility and gut colonization in *Salmonella enterica* serotype Typhimurium. mBio 12:e03680-20. <https://doi.org/10.1128/mBio.03680-20>.

**Invited Editor** Cagla Tukul, Temple University

**Editor** N. Luisa Hiller, Carnegie Mellon University

**Copyright** © 2021 Westerman et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to J. R. Efenbein, [jelfenbein@wisc.edu](mailto:jelfenbein@wisc.edu).

**Received** 30 December 2020

**Accepted** 29 April 2021

**Published** 8 June 2021

environment (6, 7). The dynamic expression of flagellar genes requires tight control to facilitate gut colonization while avoiding inappropriate stimulation of the innate immune response.

Flagellar biosynthesis is regulated by a transcriptional hierarchy comprised of genes in three classes. Class 1 genes encode the heterodimeric master transcriptional regulator FlhD<sub>4</sub>C<sub>2</sub>, required for expression of class 2 and 3 genes that encode the flagellar apparatus, motor force-generating elements, chemotaxis proteins, and numerous regulatory proteins (8–15). Global transcriptional regulators integrate environmental signals at the single class 1 promoter to control expression of *flhDC* (16). Small RNAs and the RNA binding protein CsrA exert posttranscriptional control of *flhDC* (17, 18). Further control of flagellar biosynthesis is mediated by regulation of FlhD<sub>4</sub>C<sub>2</sub> function. Two proteins, each containing EAL-like domains, YdiV and STM1697 (STM14\_2047), inhibit motility using different mechanisms to regulate FlhD<sub>4</sub>C<sub>2</sub> function (19–22). YdiV binds FlhD to prevent FlhD<sub>4</sub>C<sub>2</sub> from binding DNA and targets it for proteolysis (21, 23), whereas STM1697 restricts FlhD<sub>4</sub>C<sub>2</sub> from recruiting  $\sigma^{70}$  RNA polymerase to promoters of class 2 flagellar genes (20). Together, the extensive transcriptional, posttranscriptional, and functional control of the flagellar master regulator enables tight control over the initiation of flagellar biosynthesis.


In prior work, we found a motility defect for a *Salmonella* Typhimurium mutant in the putative LysR-type transcriptional regulator (LTTR) *yeiE* (STM2201/STM14\_2717) (24). LTTRs comprise the largest family of transcriptional regulators among prokaryotes and can regulate local or global gene expression in response to small-molecule ligands (25, 26). There are at least 44 annotated LTTRs encoded in the *Salmonella* Typhimurium genome (27–29). Although YeiE is poorly characterized in *S. Typhimurium*, it is highly conserved across related organisms. In *Escherichia coli*, there are no genes with clear roles that regulate flagellar biosynthesis in the YeiE regulon and no motility defect identified for a  $\Delta yeiE$  mutant (30, 31). However, YeiE homologs improve the *in vivo* fitness of other bacterial pathogens, including *Cronobacter sakazakii* (gpESA\_01081), *Pseudomonas aeruginosa* (PA3398/*finR*), and *Vibrio cholerae* (VC2324/*tehR*) (32–34). These data indicate an important role for YeiE in the pathogenesis of numerous organisms.

Despite extensive study of the regulation of flagellar biogenesis, the role of YeiE in flagellar regulation has not previously been described. The purpose of this study was to determine the role of *yeiE* in the regulation of *S. Typhimurium* flagellum-mediated motility and gastrointestinal colonization. We hypothesized that *yeiE* impacts flagellar gene expression and that dysregulated motility results in reduced gut colonization in animal models. Our work demonstrates the critical role of *yeiE* in the complex pathway regulating flagellum-mediated motility in *Salmonella* Typhimurium.

## RESULTS

***yeiE* is required for swimming motility.** Flagellum-mediated motility is a potent agonist of the neutrophil respiratory burst (35). In a screen of *Salmonella* Typhimurium mutants, we found that a  $\Delta yeiE$  mutant elicited a decreased neutrophil respiratory burst due to its lack of motility (24). To definitively link the observed motility defect to the disruption of *yeiE*, we assessed the swimming motility of the complemented  $\Delta yeiE$  mutant. Complementation in *trans* restored the swimming motility of the  $\Delta yeiE$  mutant to levels greater than that of the wild-type (WT) organism (Fig. 1A and B). The  $\Delta yeiE$  mutant grew normally in both rich and minimal media (Fig. 1C and D), ruling out any potential effects of abnormal growth on swimming motility. These data demonstrate that *yeiE* plays an important role in *S. Typhimurium* swimming motility.

***yeiE* is required for gastrointestinal colonization.** Since flagellum-mediated motility is required for efficient colonization of the gut, we hypothesized that the  $\Delta yeiE$  mutant would have a colonization defect in animal models of enteric salmonellosis. We used the calf ligated ileal loop model of infection because calves are natural hosts for *Salmonella*, infections occur in the presence of an intact microbiota, and pathology is consistent with that of enteric salmonellosis in humans (36). In competitive infections with the virulent WT organism, we found that the  $\Delta yeiE$  mutant was defective for colonization




**FIG 1** *yeiE* is required for swimming motility. (A) Normalized overnight cultures of the WT (HA420) and the  $\Delta motA$  mutant (JE1202),  $\Delta yeiE$  mutant (JE973),  $\Delta yeiE$  mutant with an empty vector ( $\Delta yeiE + V$ ; JE1511), and the  $\Delta yeiE$  mutant with a complementing plasmid ( $\Delta yeiE + CP$ ; JE1513) were spotted onto swimming plates. Diameters of cell spread were measured 4 h postinoculation. Each assay was performed in 3 replicates on 3 independent occasions. Bars represent means  $\pm$  standard errors of the means (SEM). Statistical significance was determined by Student's *t* test (*P* of <0.05). \* indicates a significant difference between the WT and mutant, and † indicates a significant difference between the indicated mutants. (B) Representative photographs of swimming plates 5 h postinoculation from one experiment. Growth curves of the WT (HA420) and the  $\Delta yeiE$  mutant (JE973) in rich (C) and minimal (D) media.

of the intestinal mucus layer and tissue but not for survival within luminal fluid (Fig. 2A). The lack of a defect in the fluid compartment suggests no effect of the  $\Delta yeiE$  mutation on fitness in the inflamed gut, and the failure of the  $\Delta yeiE$  mutant to penetrate the mucus layer and colonize intestinal tissue is consistent with a defect in flagellum-mediated motility.

Next, we used the murine colitis model to investigate the mechanism of the  $\Delta yeiE$  mutant gut colonization defect. In the murine colitis model, antibiotic treatment disrupts the gut microbiota prior to *Salmonella* infection, allowing development of a neutrophilic typhlitis (37). As with our findings from the calf model, the  $\Delta yeiE$  mutant had a colonization defect in the murine cecum and large intestine (Fig. 2B). The colonization defect was reversed by complementation in *trans* in both the large intestine and the cecum (Fig. 2B), definitively linking *yeiE* to the observed gut colonization defect.


We hypothesized that the motility defect of the  $\Delta yeiE$  mutant was the likely cause of the observed gut colonization defect. To investigate the role flagellum-mediated motility played in the  $\Delta yeiE$  gut colonization defect, we compared the competitive index (CI) of the aflagellated  $\Delta fliC \Delta fljB$  mutant with that of the  $\Delta yeiE$  mutant. We found that the  $\Delta yeiE$  mutant and the  $\Delta fliC \Delta fljB$  mutant had similar colonization defects in competition with the WT organism (Fig. 3). To establish whether abnormal motility was the reason for the fitness defect of the  $\Delta yeiE$  mutant, we performed a competitive infection experiment with the  $\Delta fliC \Delta fljB$  mutant and a  $\Delta fliC \Delta fljB \Delta yeiE$  mutant. We found no significant alteration in fitness of the  $\Delta fliC \Delta fljB \Delta yeiE$  mutant from that of




**FIG 2** *yeiE* is required for efficient gut colonization. (A) Ligated ileal loops from five 3- to 6-week-old calves were inoculated with  $\sim 10^9$  CFU of an equivalent mixture of the WT (HA420) and  $\Delta yeiE$  mutant (JE973). Loops were harvested 12 h after inoculation, and luminal fluid, mucus, and intestinal tissue were processed for enumeration of CFU. (B) Six C57BL/6J mice were treated with streptomycin prior to infection with  $\sim 10^8$  CFU of an equivalent mixture of the WT and  $\Delta yeiE$  mutant bearing the empty plasmid (JE1511) or the complemented mutant (JE1513) by gavage. Mice were euthanized 72 h after infection and the cecum and large intestine (LI) processed for enumeration of CFU. The competitive index (CI) was determined by dividing the ratio of WT cells to mutant cells in each tissue compartment by the ratio in the inoculum. Each data point represents a single animal, with median and interquartile ranges indicated. Statistical significance was determined by Student's *t* test ( $P < 0.05$ ). \* indicates a significant difference in competitive indexes, and † indicates a significant difference between groups.

the  $\Delta fljC \Delta fljB$  mutant (Fig. 3). Together, these data demonstrate that the  $\Delta yeiE$  mutant poorly colonizes the mammalian intestine, likely as a result of defective flagellum-mediated motility.

**Role of YeiE in flagellar regulation.** Although the *E. coli* YeiE regulon does not include genes responsible for flagellar biogenesis (30), our data demonstrate a role for *yeiE* in *S. Typhimurium* flagellum-mediated motility. We hypothesized that YeiE influences the expression of flagellar genes. To investigate the level at which YeiE affects the transcriptional hierarchy of flagellar biogenesis, we measured the mRNA expression of a subset of genes from each of the three flagellar regulatory classes in the WT and  $\Delta yeiE$  mutant. Expression of the class 1 gene *flhD* was not affected by deletion of *yeiE*



**FIG 3** The  $\Delta yeiE$  mutant gut colonization defect is linked to defective flagellar motility. The CI between the indicated strains in cecum from the murine colitis model was determined as described in the legend for Fig. 2. Statistical significance was determined by Student's *t* test ( $P < 0.05$ ). \* indicates a significant difference in competitive indexes, and † indicates a significant difference between groups.




**FIG 4** Flagellar class 2 and 3 genes are downregulated in the  $\Delta yeiE$  mutant. Relative expression of the indicated genes from the  $\Delta yeiE$  mutant compared with that of the WT in late exponential growth in rich medium. Each data point represents the mean from triplicate samples from one biological replicate; means  $\pm$  standard deviations (SD) are indicated. Statistical significance was determined by Student's *t* test ( $P < 0.05$ ). \* indicates a significant difference in relative levels of expression of the given gene.

(Fig. 4A). In contrast, expression of genes from both class 2 and class 3 was significantly downregulated in the  $\Delta yeiE$  mutant (Fig. 4A).

There is regulatory cross talk between the expression of flagellar genes and those borne in *Salmonella* pathogenicity island 1 (SPI-1), encoding type 3 secretion system 1 and effector proteins needed for the invasion of epithelial cells (38–40). To rule out an effect of *yeiE* on SPI-1 gene expression, we measured the mRNA expression of three SPI-1 regulators in the WT and  $\Delta yeiE$  mutant. We observed no difference in expression of *hilA*, *hilC*, or *hilD* in the  $\Delta yeiE$  mutant (Fig. 4B), suggesting that *yeiE* does not impact SPI-1 gene expression. These data are consistent with our prior work demonstrating no role for *yeiE* in the activation of a promoter for a type 3 secretion system 1 apparatus protein (24). These findings suggest that YeiE promotes the expression of flagellar class 2 genes through an SPI-1-independent process.

**Characterization of the  $\Delta yeiE$  suppressor mutant.** In prior work evaluating the role of *S. Typhimurium* genes in the stimulation of the neutrophil respiratory burst, we found that the effects of the  $\Delta yeiE$  mutation in a strain obtained from the kanamycin-resistant *Salmonella* Typhimurium ATCC 14028s single-gene-deletion library ( $\Delta yeiE^{lib}$  strain) differed from those of the  $\Delta yeiE$  mutation in a clean genetic background (24, 41). To establish whether the differential effects were due to a motility difference between the  $\Delta yeiE$  mutants, we measured the motility of the two  $\Delta yeiE$  mutants on semisolid agar. The  $\Delta yeiE$  mutant in the clean genetic background was amotile, whereas the  $\Delta yeiE^{lib}$  mutant was hypermotile compared to the WT organism (Fig. 5). Furthermore, the complemented  $\Delta yeiE^{lib}$  mutant remained hypermotile (Fig. 5), suggesting that the hypermotility is not due to disruption of *yeiE* in the  $\Delta yeiE^{lib}$  mutant. Therefore, we hypothesized that a suppressor mutation was the likely cause of the observed hypermotility of the  $\Delta yeiE^{lib}$  mutant.

We performed whole-genome sequencing of the amotile  $\Delta yeiE$  mutant and the hypermotile  $\Delta yeiE^{lib}$  suppressor mutant to establish the identity of the suppressor mutation. In a comparison with the published ATCC 14028s genome sequence (42), we identified three single-nucleotide variants (SNVs) in the amotile  $\Delta yeiE$  mutant and three SNVs in the  $\Delta yeiE^{lib}$  suppressor mutant that were different between the two genomes (see Tables S1 and S2 in the supplemental material). In the amotile mutant, we found an SNV in *gyrA* (260T>C) which confers nalidixic acid resistance and is consistent with the phenotypic nalidixic acid resistance in the amotile  $\Delta yeiE$  mutant (Table 1). We also




**FIG 5** The  $\Delta yeiE^{lib}$  mutant is hypermotile. Normalized overnight cultures of the WT (HA420),  $\Delta motA$  mutant (JE1202),  $\Delta yeiE$  mutant (JE973),  $\Delta yeiE^{lib}$  mutant (JE1681),  $\Delta yeiE^{lib}$  mutant with the empty vector ( $\Delta yeiE^{lib} + V$ ; JE2014), and complemented  $\Delta yeiE^{lib}$  mutant ( $\Delta yeiE^{lib} + CP$ ; JE2016) were spotted onto swimming plates. Statistical significance was determined by Student's *t* test ( $P < 0.05$ ). \* indicates a significant difference between the WT and mutant. (B) Representative photographs of swimming plates from one experiment.

identified an SNV in *eehH* on the 94-kb virulence plasmid and a single-nucleotide deletion in *cyoB* leading to a frameshift resulting in a predicted truncation of the protein from 664 amino acids to 278 amino acids. CyoB encodes cytochrome *bo* ubiquinol oxidase subunit I, which helps generate the proton-motive force (43). Since the proton-motive force drives flagellum-mediated motility, the predicted CyoB truncation may play a contributing role in its decreased motility (44). However, complementation in *trans* reverses the  $\Delta yeiE$  mutant motility defect (Fig. 1), suggesting that CyoB truncation is not the likely cause of the motility defect.

Three SNVs were identified in the  $\Delta yeiE^{lib}$  suppressor mutant (Tables S1 and S2). An SNV identified in *hrpA* (248G>T) resulted in a threonine-to-asparagine amino acid sequence change that is unlikely to alter protein function due to replacement of one polar amino acid for another. Two other SNVs in the  $\Delta yeiE^{lib}$  mutant were in *hupA* and *STM1697*. Published work demonstrated that deletion of *hupA* causes an ~10% motility reduction compared with the motility of its wild-type parent (45). The *STM1697* mutation (565G>A) resulted in a stop codon after amino acid 188, causing early termination of the protein. Deletion of *STM1697* caused hypermotility (19). Since the  $\Delta yeiE^{lib}$  mutant

**TABLE 1** Bacterial strains and plasmids

Strain or plasmid	Genotype or description	Reference or source
<b>Strains</b>		
HA420	ATCC 14028.s (spontaneous Nal <sup>r</sup> )	59
JE973	HA420 $\Delta yeiE::kan$ (Nal <sup>r</sup> Kan <sup>r</sup> )	24
JE1202	HA420 $\Delta motA::kan$ (Nal <sup>r</sup> Kan <sup>r</sup> )	35
JE1511	JE973 carrying pWSK29 (Nal <sup>r</sup> Kan <sup>r</sup> Amp <sup>r</sup> )	This study
JE1513	JE973 carrying pWSK29:: <i>yeiE</i> (Nal <sup>r</sup> Kan <sup>r</sup> Amp <sup>r</sup> )	This study
JE1681	14028.s $\Delta yeiE::kan$ (Kan <sup>r</sup> )	41
JE1389	HA420 $\Delta yeiE::frt$ (Nal <sup>r</sup> )	This study
JE1699	HA420 $\Delta STM1697::cm$ (Nal <sup>r</sup> Cm <sup>r</sup> )	This study
JE1907	JE973 $\Delta STM1697::cm$ (Nal <sup>r</sup> Kan <sup>r</sup> Cm <sup>r</sup> )	This study
JE1915	HA420 $\Delta fliC::kan \Delta fljB::cm$ (Nal <sup>r</sup> Kan <sup>r</sup> Cm <sup>r</sup> )	This study
JE1919	HA420 $\Delta fliC::frt \Delta fljB::frt$ (Nal <sup>r</sup> )	This study
JE1921	HA420 $\Delta fliC::frt \Delta fljB::frt \Delta yeiE::kan$ (Nal <sup>r</sup> Kan <sup>r</sup> )	This study
JE2014	JE1681 carrying pWSK29 (Kan <sup>r</sup> Amp <sup>r</sup> )	This study
JE2016	JE1681 carrying pWSK29:: <i>yeiE</i> (Kan <sup>r</sup> Amp <sup>r</sup> )	This study
<b>Plasmids</b>		
pCP20	<i>flp</i> recombinase, Amp <sup>r</sup>	51
pWSK29	Cloning vector, Amp <sup>r</sup>	53
pWSK29:: <i>yeiE</i>	pWSK29:: <i>yeiE</i> Amp <sup>r</sup>	This study



**FIG 6** YeiE interacts with STM1697 to alter cell motility. (A) Normalized overnight cultures of the WT (HA420),  $\Delta motA$  mutant (JE1202),  $\Delta yeiE$  mutant (JE973),  $\Delta STM1697$  mutant (JE1699), and  $\Delta yeiE \Delta STM1697$  mutant (JE1907) were spotted onto swimming plates. Data analysis was as described in the legend of Fig. 1. (B) Representative photographs of swimming plates from one experiment. (C) Gene expression from the  $\Delta yeiE$  mutant compared with that of the WT as described in the legend of Fig. 4. Each data point represents the mean from triplicate samples from one biological replicate; means  $\pm$  SD are indicated.

is hypermotile, the SNV in *STM1697* was considered the most likely cause of the observed phenotype.

**The association between YeiE and STM1697.** *STM1697* inhibits cell motility by preventing RNA polymerase recruitment to FlhD<sub>4</sub>C<sub>2</sub>, leading to repressed flagellar class 2 and 3 gene expression (20). Similarly, flagellar class 2 and 3 genes are downregulated by deletion of *yeiE* without an effect on class 1 gene expression (Fig. 4). Therefore, we hypothesized that there was an interaction between *STM1697* and *yeiE*. We found that the  $\Delta STM1697$  mutant is hypermotile on semisolid agar (Fig. 6A and B), consistent with published reports (19). Deletion of *STM1697* exhibited a dominant effect on the  $\Delta yeiE$  mutant, resulting in the hypermotility of the  $\Delta yeiE \Delta STM1697$  mutant (Fig. 6A and B). These data are similar to our observations of hypermotility for the  $\Delta yeiE^{iib}$  suppressor mutant and suggest that YeiE may repress the expression of the *STM1697* FlhD<sub>4</sub>C<sub>2</sub> repressor.


To test whether YeiE influences the expression of *STM1697*, we determined the relative expression of *STM1697* in the WT and the  $\Delta yeiE$  mutant. We found that the relative expression of *STM1697* is upregulated in the  $\Delta yeiE$  mutant (Fig. 6C). Since YdiV is a related EAL domain-containing protein which also inhibits FlhD<sub>4</sub>C<sub>2</sub> function, we measured the effects of deletion of *yeiE* on the expression of *ydiV*. We found no significant alteration from *ydiV* expression in the  $\Delta yeiE$  mutant under the conditions tested (Fig. 6C). These data demonstrate that YeiE regulates motility by repressing the expression of the FlhD<sub>4</sub>C<sub>2</sub> functional repressor *STM1697* (Fig. 7).

## DISCUSSION

The putative LysR-type transcriptional regulator YeiE is critical for *Salmonella* Typhimurium flagellum-mediated motility and gut colonization. We demonstrate that flagellar class 2 gene expression is reduced in a  $\Delta yeiE$  mutant. Furthermore, we link the positive effect of YeiE on cell motility with *STM1697*, an anti-FlhD<sub>4</sub>C<sub>2</sub> factor, and propose that YeiE is a repressor of *STM1697* (Fig. 7).

*E. coli* and *S. Typhimurium* share a core genome, with approximately 70% of their genetic material conserved between the two species (29). *S. Typhimurium* YeiE (STM2201/STM14\_2717) has 89% amino acid identity and 92% similarity with its *E. coli* homolog, suggesting high functional similarity (46). YeiE is both an activator and a repressor of transcription, binds DNA upstream and downstream of the promoter, and can occupy the same sequence as RNA polymerase (30). YeiE has more than 100 predicted binding sites in the *E. coli* MG1655 genome, with target genes enriched in functional groups including energy production, amino acid and inorganic ion transport and metabolism, and iron transport (30). The *E. coli* *yeiE* regulon also includes numerous genes with roles in the regulation of transcription and translation and genes with





**FIG 7** Proposed mechanism for YeiE regulation of cell motility. YeiE acts as a repressor of *STM1697* to promote flagellum-mediated motility. The dotted line indicates an unknown mechanism of repression. RNAP, RNA polymerase.

unknown or poorly characterized functions (30). There were no genes with a direct relationship to flagellar biogenesis in the *E. coli* YeiE regulon and no motility defects in a  $\Delta yeiE$  mutant (30, 31). Unlike in *E. coli*, the *S. Typhimurium*  $\Delta yeiE$  mutant has a severe motility defect that we linked to dysregulation of the FlhD<sub>4</sub>C<sub>2</sub> inhibitor *STM1697*. *STM1697* restricts bacterial motility and adds a layer of flagellar regulation that promotes *Salmonella* intracellular survival and evasion of the host immune system (20). There is no ortholog of *STM1697* in the *E. coli* MG1655 genome (19) or in the genomes of several thousand other *E. coli* and *Shigella* strains (data not shown), explaining the different phenotypes observed for  $\Delta yeiE$  mutants of *S. Typhimurium* and *E. coli*. There are multiple different mechanisms by which YeiE may influence *STM1697* expression. YeiE may directly repress *STM1697* expression by binding to its promoter to inhibit transcription. YeiE may indirectly influence *STM1697* expression by regulating the expression of an activator or inhibitor of *STM1697* transcription. Further work is needed to establish the full YeiE regulon of *S. Typhimurium* and to establish how YeiE represses *STM1697* expression in *S. Typhimurium*.

We observed a significant gut colonization defect for the *S. Typhimurium*  $\Delta yeiE$  mutant that we linked to defective motility. Since YeiE is likely to have numerous regulatory targets within the *S. Typhimurium* genome, it is possible that YeiE also regulates other genes that influence *S. Typhimurium* interactions with the host. For example, *yeiE* responds to DNA damage in *E. coli*, suggesting that it may be activated during exposure to oxidative and nitrosative stresses encountered by *S. Typhimurium* during infection (47). Furthermore, iron import genes are included in the *E. coli* *yeiE* regulon, suggesting a potential role for *yeiE* in iron homeostasis during *S. Typhimurium* host colonization (30). Although there are numerous potential mechanisms by which *yeiE* may influence pathogenesis, elimination of the flagellar filament restored virulence to the  $\Delta yeiE$  mutant, suggesting that the regulation of flagellum-mediated motility is the primary YeiE target facilitating *S. Typhimurium* colonization during acute enterocolitis. To our knowledge, this is the first demonstration of a role for YeiE in *S. Typhimurium* pathogenesis.

The related organisms *Yersinia enterocolitica* and *Cronobacter sakazaki* each encode homologs of both *yeiE* and *STM1697*. The *Cronobacter sakazakii* *yeiE* homolog (*gpESA\_01081*)

shares 84% amino acid identity with *S. Typhimurium* YeiE and is required for enterocyte invasion, biofilm formation, neutrophil recruitment, and virulence in neonatal rats (32, 48). The *Yersinia enterocolitica* *yeiE* homolog (*rscR*) shares 75% identity with *S. Typhimurium* YeiE and limits systemic dissemination of the organism (48, 49). The effects of the *C. sakazaki* and *Y. enterocolitica* *yeiE* homologs (*gpESA\_01081* and *rscR*, respectively) on motility have not yet been characterized. We hypothesize that the role of the *yeiE* homologs in virulence is due, in part, to regulation of the *STM1697* homolog in each of these organisms.

Evaluation of the  $\Delta yeiE^{lib}$  suppressor mutant led to the identification of *STM1697* as a target for YeiE motility regulation. *STM1697* is an anti-FlhD<sub>4</sub>C<sub>2</sub> factor that represses cell motility by binding FlhD, thereby preventing RNA polymerase recruitment to the FlhD<sub>4</sub>C<sub>2</sub>-DNA complex (20). The C terminus of *STM1697* is critical for its function, as truncation of the 235-amino-acid protein at residue 192 results in a nonfunctional protein and hypermotility (19). Although the crystal structure of the *STM1697*-FlhD complex suggests that Tyr180 is the residue closest to the C terminus required for stable *STM1697*-FlhD interaction (20), our data demonstrate an SNV in *STM1697* causing a stop codon after amino acid 188, and deletion of *STM1697* resulted in the same hypermotile phenotype, confirming prior reports that the C terminus is critical for *STM1697* function (see Table S2 in the supplemental material). These data, in combination with prior work (19), suggest that one or more residues located even closer to the C terminus may be required for *STM1697* stability or function *in vivo*.

*Salmonella* uses flagellum-mediated motility to interact with host cells, driving successful gut colonization; however, it is critical for *Salmonella* to repress flagellin expression when residing within host cells to evade immune detection (2, 4, 7, 50). Our data suggest that YeiE functions as a regulatory control point to promote flagellar motility by inhibiting the anti-FlhD<sub>4</sub>C<sub>2</sub> factor *STM1697*. The promotility effect of *STM1697* inhibition by YeiE likely facilitates initial gut colonization, whereas *STM1697*-mediated repression of flagellin aids immune evasion once *Salmonella* is located intracellularly (20). The regulatory functions of LysR regulators are influenced by small-molecule coinducers that sense environmental conditions (25, 26). Further molecular characterization of YeiE and the identification of its coinducer will help to delineate how this LysR regulator exerts control over cell motility during infection.

We definitively link *yeiE* with *S. Typhimurium* motility and demonstrate its requirement for gut colonization using two animal models of enterocolitis. We propose that YeiE serves as a control point for flagellar regulation through inhibition of the anti-FlhD<sub>4</sub>C<sub>2</sub> factor *STM1697*, although additional work is needed to establish the mechanism by which YeiE influences *STM1697* expression. Tight control of flagellar biogenesis is critical to facilitate pathogenesis. Whereas the flagellum-based motility state is needed for gut colonization, elimination of flagellins is critical to evade immune detection. Our work demonstrates that YeiE serves as a key regulatory control point for flagellar biogenesis to facilitate *S. Typhimurium* enteropathogenesis.

## MATERIALS AND METHODS

**Bacterial strains and growth conditions.** All bacterial strains are derivatives of *Salmonella enterica* serotype Typhimurium ATCC 14028s (Table 1). Mutations were moved into a clean genetic background by P22 transduction, and antibiotic cassettes were removed by *flp*-mediated recombination as previously described (51, 52). Bacteria were grown on Luria-Bertani (LB)-Miller agar or in LB broth at 37°C with agitation (225 rpm) unless otherwise noted. Medium was supplemented with the following antibiotics, as appropriate: nalidixic acid (50 mg/liter), chloramphenicol (20 mg/liter), kanamycin (50 mg/liter), and carbenicillin (100 mg/liter).

**Complementing plasmid construction.** Genomic DNA was isolated from the wild-type organism using the GenElute bacterial genomic DNA kit (Sigma-Aldrich). A 1.5-kb PCR product of *yeiE*, including its native promoter, was generated from genomic DNA by PCR using Q5 polymerase (New England Biolabs), with an annealing temperature of 72°C and an extension time of 40s for 30 cycles. Restriction sites for endonucleases were incorporated into the following primers to facilitate cloning: *STM2201*BamH1Fwd, 5'-GTCCGATCCTGCCTGCCAGACCAAGA-3', and *STM2201*Kpn1Rev, 5'-GTCCGGTACCTGCGCGTTATAAGAGACCT-3'. The expected size of the PCR product was confirmed by agarose gel electrophoresis. The PCR product was digested with restriction endonucleases BamHI and KpnI (New England Biolabs) and purified with the QIAquick PCR purification kit (Qiagen). The insert was cloned into the pWSK29 vector, sequentially digested with BamHI and KpnI (53). Ligation

was performed overnight at 14°C with T4 DNA ligase (New England Biolabs). The resulting construct was transformed into DH5 $\alpha$  *Escherichia coli* by heat shock. Transformants were obtained by selection on LB agar with carbenicillin and X-gal (5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside; 40  $\mu$ g/ml). Plasmids were isolated (QIAprep miniprep kit; Qiagen), and the correct insert size was confirmed by agarose gel electrophoresis of digested plasmids. The insert sequence was confirmed by Sanger sequencing (Eton Bioscience). The complementing plasmid (CP) was transformed into restriction-deficient modification-positive *S. Typhimurium* LB5000 by electroporation, and transformants were isolated by selection on LB agar with carbenicillin (54). The complementing plasmid and empty vector were isolated and then transformed into the indicated mutants by electroporation. The resulting strains were purified by streaking them twice for single colonies on LB with carbenicillin and stored in glycerol stocks at –80°C.

**Growth curves.** Overnight cultures were diluted 1:100 into 50 ml of LB or M9 minimal medium and grown at 37°C with agitation (225 rpm) for 24 h. M9 minimal medium (48 mM Na<sub>2</sub>HPO<sub>4</sub>, 22 mM KH<sub>2</sub>PO<sub>4</sub>, 9 mM NaCl, 19 mM NH<sub>4</sub>Cl, 0.1 mM CaCl<sub>2</sub>, and 2 mM MgSO<sub>4</sub>) was supplemented with 0.2% (wt/vol) dextrose as a carbon source (55). Samples were taken hourly for 6 h and once at 24 h, serially diluted, and plated for enumeration of CFU per milliliter.

**Bacterial motility assays.** Swimming assays were performed on semisolid agar as previously described (56). Swimming motility was assayed on plates containing 0.3% Difco Bacto agar (LB-Miller base, 25 g/liter). Overnight cultures were grown at 37°C with agitation, and the cell concentration was normalized by optical density. Bacterial strains were spotted (3  $\mu$ l) onto swimming plates and incubated at 37°C. The widest diameter of each colony was measured in two intersecting planes after 4 h of incubation. Images of swimming plates were obtained on ChemiDoc MP (Bio-Rad) after 5 h of incubation. Each assay was performed in triplicate on three independent occasions.

**Calf infections.** Calf infections were approved by the North Carolina State University Institutional Animal Care and Use Committee (protocol 15-047-B). Calves were obtained from the university teaching herd and transferred to individual AALAC-approved housing within 8 h following birth. Calves were administered a commercial colostrum replacer (AgriLabs Colostrx CR) by an esophageal feeder upon arrival and 2 h following initial administration. Calves were fed milk replacer twice daily and provided hay and water *ad libitum*. At 1 day of age, adequate passive transfer of immunity was estimated by measuring serum total solids with a refractometer. Fecal cultures were performed at least twice weekly using selective media to ensure that calves were not shedding *Salmonella* prior to experimental infection (57).

In preparation for the ligated ileal loop surgery, bacteria were grown overnight at 37°C with agitation in LB broth. Overnight cultures were subcultured 1:100 into LB broth and incubated for 3 to 4 h at 37°C with agitation. Bacteria were washed twice in sterile LB broth, and the mutant and WT were mixed in a 1:1 ratio based on optical density (600 nm). Loops were inoculated with approximately 10<sup>9</sup> CFU of the mixture in 3 ml LB broth.

Ligated ileal loop surgery was performed on 5 bull calves (3 Jersey and 2 Holstein) aged 3 to 6 weeks, as previously described (57). At 12 h postinfection, the luminal contents, mucus overlying the epithelium, and epithelial tissue were harvested, collected in phosphate-buffered saline (PBS), homogenized, serially diluted, and plated to determine numbers of CFU. The competitive index (CI) between the WT and mutant was determined as the ratio of WT to mutant bacteria after infection normalized to the ratio in the inoculum.

**Mouse infections.** Mouse infections were approved by the University of Wisconsin—Madison Institutional Animal Care and Use Committee (protocol no. V006255). The acute murine colitis model was used as previously described, with 10- to 12-week-old female C57BL/6J mice obtained from Jackson Laboratories (strain 000664) (37). Mice were administered 20 mg streptomycin in sterile water by oral gavage 24 h prior to infection. Overnight bacterial cultures were washed in PBS, and mice were infected with approximately 10<sup>8</sup> CFU of a 1:1 mixture of the two competing strains by oral gavage. Mice were euthanized 72 h postinfection. Organs were harvested, homogenized, serially diluted, and plated for enumeration of CFU. The competitive index was determined as for calf infections.

**Genome sequencing and analysis.** Genomic DNA from the amotile (JE973) and hypermotile (JE1681) *ΔyeiE* mutants was isolated (GenElute bacterial genomic DNA kit; Sigma-Aldrich) and submitted to the North Carolina State University Genomic Sciences Laboratory. Genomic DNA (gDNA) quality was analyzed using TapeStation (Agilent), and library preparation and whole-genome shotgun sequencing was performed using a MiSeq platform (PE300; Illumina). Raw sequencing data were provided as demultiplexed .fastq files. Variants were found using the CLC Genomics Workbench (Qiagen) resequencing analysis module. Reads were first mapped to the published ATCC 14028s genome (42) (NCBI accession no. [NC\\_016856.1](https://.ncbi.nlm.nih.gov/nuccore/NC_016856.1)), followed by local realignment. Variants were then detected using the fixed-ploidy variant detection tool. Variants that were supported by reads in both strands with more than 50% frequency were further investigated.

**Gene expression analyses.** Overnight cultures of the WT and *ΔyeiE* mutant were diluted 1:100 into LB broth and grown at 37°C with agitation for 3.5 h. Total RNA was isolated using TRIzol (Invitrogen) according to the manufacturer's instructions. RNA quantity and integrity were determined using a Qubit 4 fluorometer (Invitrogen), and samples with an RNA IQ value of >7.5 (Qubit RNA IQ assay; Invitrogen) were considered of good quality and were used for downstream applications. Removal of contaminating DNA was performed using TURBO DNase (Invitrogen) according to the manufacturer's instructions.

Reverse transcription (RT) of RNA to cDNA was performed by random-hexamer-dependent amplification using TaqMan reverse transcription reagents according to the manufacturer's instructions (Invitrogen). Real-time PCR was performed using probe-based 5' nuclease assays (IDT PrimeTime quantitative PCR [qPCR]). Primer and probe sets (see Table S3 in the supplemental material) were designed using the IDT

PrimerQuest tool (<https://www.idtdna.com/SciTools>) and incorporated 6-carboxyfluorescein (FAM)/ZEN™ (IDT trademarked dark quencher)/lowa black fluorescent quencher (IBFQ) doubly quenched probes (Integrated DNA Technologies). Cycling parameters were an initial 95°C polymerase activation step for 2 min, followed by 40 cycles of 95°C for 15 s and 60°C for 1 min, performed on a StepOnePlus real-time PCR system (Applied Biosystems). Amplification efficiency was determined to be 90 to 100% for all primer/probe sets (Table S4). All qPCR products were confirmed to have a single product of the expected size by agarose gel electrophoresis. All real-time PCR assays were performed in three technical repeats using cDNA from at least 3 biological replicates. The change in threshold cycle ( $\Delta C_t$ ) for each gene of interest was determined for each strain, using *rpoD* as the reference gene for normalization (58). The relative expression of each gene of interest was determined by comparing the  $\Delta C_t$  of the mutant to that of the WT using the comparative threshold ( $\Delta\Delta C_t$ ) method. All quantitative RT-PCR data are available in Table S4.

**Data analysis.** Statistical significance was determined by a two-tailed Student *t* test, with significance set at a *P* of <0.05. Analyses were performed using GraphPad Prism version 8.0.

**Data availability.** The raw DNA sequences were deposited in GenBank (BioProject accession no. PRJNA704982).

## SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

**TABLE S1**, XLSX file, 0.01 MB.

**TABLE S2**, XLSX file, 0.01 MB.

**TABLE S3**, XLSX file, 0.01 MB.

**TABLE S4**, XLSX file, 0.1 MB.

## ACKNOWLEDGMENTS

We thank M. Mitsu Suyemoto, Craig Radi, Steffen Porwollik, and WeiPing Chu for assistance with experiments associated with this work.

This work was supported by startup funds from the University of Wisconsin—Madison (to J.R.E.). J.R.E. was supported, in part, by USDA-NIFA grant 2018-67017-27632 and NIH grant K08AI108794. T.L.W. was supported by grant 5T32OD01113. M.M. was supported, in part, by U.S. Department of Agriculture, National Institute of Food and Agriculture grants 2015-67017-23360 and 2017-67015-26085, NIFA Hatch grant CA-d-PLS-2327-H, NIFA-BARD award 2017-67017-26180, and NIH grant AI139557.

## REFERENCES

1. Stecher B, Hapfelmeier S, Muller C, Kremer M, Stallmach T, Hardt WD. 2004. Flagella and chemotaxis are required for efficient induction of *Salmonella enterica* serovar Typhimurium colitis in streptomycin-pretreated mice. *Infect Immun* 72:4138–4150. <https://doi.org/10.1128/IAI.72.7.4138-4150.2004>.
2. Schmitt CK, Ikeda JS, Darnell SC, Watson PR, Bispham J, Wallis TS, Weinstein DL, Metcalf ES, O'Brien AD. 2001. Absence of all components of the flagellar export and synthesis machinery differentially alters virulence of *Salmonella enterica* serovar Typhimurium in models of typhoid fever, survival in macrophages, tissue culture invasiveness, and calf enterocolitis. *Infect Immun* 69:5619–5625. <https://doi.org/10.1128/iai.69.9.5619-5625.2001>.
3. Dibb-Fuller MP, Allen-Vercoe E, Thorns CJ, Woodward MJ. 1999. Fimbriae- and flagella-mediated association with and invasion of cultured epithelial cells by *Salmonella enteritidis*. *Microbiology* 145:1023–1031. <https://doi.org/10.1099/13500872-145-5-1023>.
4. Zhang S, Kingsley RA, Santos RL, Andrews-Polymeris H, Raffatellu M, Figueiredo J, Nunes J, Tsolis RM, Adams LG, Bäumlér AJ. 2003. Molecular pathogenesis of *Salmonella enterica* serotype Typhimurium-induced diarrhea. *Infect Immun* 71:1–12. <https://doi.org/10.1128/iai.71.1.1-12.2003>.
5. Miao EA, Alpuche-Aranda CM, Dors M, Clark AE, Bader MW, Miller SJ, Aderem A. 2006. Cytoplasmic flagellin activates caspase-1 and secretion of interleukin 1 $\beta$  via Ipaf. *Nat Immunol* 7:569–575. <https://doi.org/10.1038/ni1344>.
6. Hautefort I, Thompson A, Eriksson-Ygberg S, Parker ML, Lucchini S, Danino V, Bongaerts RJ, Ahmad N, Rhen M, Hinton JC. 2008. During infection of epithelial cells *Salmonella enterica* serovar Typhimurium undergoes a time-dependent transcriptional adaptation that results in simultaneous expression of three type 3 secretion systems. *Cell Microbiol* 10:958–984. <https://doi.org/10.1111/j.1462-5822.2007.01099.x>.
7. Ibarra JA, Knodler LA, Sturdevant DE, Virtaneva K, Carmody AB, Fischer ER, Porcella SF, Steele-Mortimer O. 2010. Induction of *Salmonella* pathogenicity island 1 under different growth conditions can affect *Salmonella* host cell interactions in vitro. *Microbiology (Reading)* 156:1120–1133. <https://doi.org/10.1099/mic.0.032896-0>.
8. Kutsukake K, Ohya Y, Iino T. 1990. Transcriptional analysis of the flagellar regulon of *Salmonella typhimurium*. *J Bacteriol* 172:741–747. <https://doi.org/10.1128/jb.172.2.741-747.1990>.
9. Yanagihara S, Iyoda S, Ohnishi K, Iino T, Kutsukake K. 1999. Structure and transcriptional control of the flagellar master operon of *Salmonella typhimurium*. *Genes Genet Syst* 74:105–111. <https://doi.org/10.1266/ggs.74.105>.
10. Liu X, Matsumura P. 1994. The FlhD/FlhC complex, a transcriptional activator of the *Escherichia coli* flagellar class II operons. *J Bacteriol* 176:7345–7351. <https://doi.org/10.1128/jb.176.23.7345-7351.1994>.
11. Ohnishi K, Kutsukake K, Suzuki H, Iino T. 1990. Gene FlIA encodes an alternative sigma factor specific for flagellar operons in *Salmonella-Typhimurium*. *Mol Gen Genet* 221:139–147. <https://doi.org/10.1007/BF00261713>.
12. Kutsukake K, Ikebe T, Yamamoto S. 1999. Two novel regulatory genes, flIT and flIZ, in the flagellar regulon of *Salmonella*. *Genes Genet Syst* 74:287–292. <https://doi.org/10.1266/ggs.74.287>.
13. Yokoseki T, Kutsukake K, Ohnishi K, Iino T. 1995. Functional analysis of the flagellar genes in the flID operon of *Salmonella typhimurium*. *Microbiology* 141:1715–1722. <https://doi.org/10.1099/13500872-141-7-1715>.
14. Ohnishi K, Kutsukake K, Suzuki H, Iino T. 1992. A novel transcriptional regulation mechanism in the flagellar regulon of *Salmonella typhimurium*: an antisigma factor inhibits the activity of the flagellum-specific sigma factor, sigma F. *Mol Microbiol* 6:3149–3157. <https://doi.org/10.1111/j.1365-2958.1992.tb01771.x>.
15. Liu X, Matsumura P. 1995. An alternative sigma factor controls transcription of flagellar class-III operons in *Escherichia coli*: gene sequence, overproduction, purification and characterization. *Gene* 164:81–84. [https://doi.org/10.1016/0378-1119\(95\)00480-t](https://doi.org/10.1016/0378-1119(95)00480-t).
16. Soutourina O, Kolb A, Krin E, Laurent-Winter C, Rimsky S, Danchin A, Bertin P. 1999. Multiple control of flagellum biosynthesis in *Escherichia*

- coli: role of H-NS protein and the cyclic AMP-catabolite activator protein complex in transcription of the flhDC master operon. *J Bacteriol* 181:7500–7508. <https://doi.org/10.1128/JB.181.24.7500-7508.1999>.
17. De Lay N, Gottesman S. 2012. A complex network of small non-coding RNAs regulate motility in *Escherichia coli*. *Mol Microbiol* 86:524–538. <https://doi.org/10.1111/j.1365-2958.2012.08209.x>.
  18. Wei BL, Brun-Zinkernagel AM, Simecka JW, Pruss BM, Babitzke P, Romeo T. 2001. Positive regulation of motility and flhDC expression by the RNA-binding protein CsrA of *Escherichia coli*. *Mol Microbiol* 40:245–256. <https://doi.org/10.1046/j.1365-2958.2001.02380.x>.
  19. Ahmad I, Wigren E, Le Guyon S, Vekkei S, Blanka A, El Mouali Y, Anwar N, Chuah ML, Lunsdorf H, Frank R, Rhen M, Liang ZX, Lindqvist Y, Romling U. 2013. The EAL-like protein STM1697 regulates virulence phenotypes, motility and biofilm formation in *Salmonella typhimurium*. *Mol Microbiol* 90:1216–1232. <https://doi.org/10.1111/mmi.12428>.
  20. Li B, Yue Y, Yuan Z, Zhang F, Li P, Song N, Lin W, Liu Y, Yang Y, Li Z, Gu L. 2017. *Salmonella* STM1697 coordinates flagella biogenesis and virulence by restricting flagellar master protein FlhD4C2 from recruiting RNA polymerase. *Nucleic Acids Res* 45:9976–9989. <https://doi.org/10.1093/nar/gkx656>.
  21. Li B, Li N, Wang F, Guo L, Huang Y, Liu X, Wei T, Zhu D, Liu C, Pan H, Xu S, Wang H-W, Gu L. 2012. Structural insight of a concentration-dependent mechanism by which YdiV inhibits *Escherichia coli* flagellum biogenesis and motility. *Nucleic Acids Res* 40:11073–11085. <https://doi.org/10.1093/nar/gks869>.
  22. Wada T, Morizane T, Abo T, Tominaga A, Inoue-Tanaka K, Kutsukake K. 2011. EAL domain protein YdiV acts as an anti-FlhD(4)C(2) factor responsible for nutritional control of the flagellar regulon in *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 193:1600–1611. <https://doi.org/10.1128/JB.01494-10>.
  23. Takaya A, Erhardt M, Karata K, Winterberg K, Yamamoto T, Hughes KT. 2012. YdiV: a dual function protein that targets FlhDC for ClpXP-dependent degradation by promoting release of DNA-bound FlhDC complex. *Mol Microbiol* 83:1268–1284. <https://doi.org/10.1111/j.1365-2958.2012.08007.x>.
  24. Westerman TL, Sheats MK, Eifenbein JR. 5 April 2021. Sulfate import in *Salmonella Typhimurium* impacts bacterial aggregation and the respiratory burst in human neutrophils. *Infect Immun* <https://doi.org/10.1128/IAI.00701-20>.
  25. Schell MA. 1993. Molecular biology of the LysR family of transcriptional regulators. *Annu Rev Microbiol* 47:597–626. <https://doi.org/10.1146/annurev.mi.47.100193.003121>.
  26. Maddocks SE, Oyston PC. 2008. Structure and function of the LysR-type transcriptional regulator (LTTR) family proteins. *Microbiology (Reading)* 154:3609–3623. <https://doi.org/10.1099/mic.0.2008/022772-0>.
  27. Pareja E, Pareja-Tobes P, Manrique M, Pareja-Tobes E, Bonal J, Tobes R. 2006. ExtraTrain: a database of extragenic regions and transcriptional information in prokaryotic organisms. *BMC Microbiol* 6:29. <https://doi.org/10.1186/1471-2180-6-29>.
  28. Lahiri A, Das P, Chakravorty D. 2009. *Salmonella Typhimurium*: insight into the multi-faceted role of the LysR-type transcriptional regulators in *Salmonella*. *Int J Biochem Cell Biol* 41:2129–2133. <https://doi.org/10.1016/j.biocel.2009.05.006>.
  29. McClelland M, Sanderson KE, Spieth J, Clifton SW, Latreille P, Courtney L, Porwollik S, Ali J, Dante M, Du F, Hou S, Layman D, Leonard S, Nguyen C, Scott K, Holmes A, Grewal N, Mulvaney E, Ryan E, Sun H, Florea L, Miller W, Stoneking T, Nhan M, Waterston R, Wilson RK. 2001. Complete genome sequence of *Salmonella enterica* serovar Typhimurium LT2. *Nature* 413:852–856. <https://doi.org/10.1038/35101614>.
  30. Gao Y, Yurkovich JT, Seo SW, Kabimoldayev I, Dräger A, Chen K, Sastry AV, Fang X, Mih N, Yang L, Eichner J, Cho B-K, Kim D, Pålsson BO. 2018. Systematic discovery of uncharacterized transcription factors in *Escherichia coli* K-12 MG1655. *Nucleic Acids Res* 46:10682–10696. <https://doi.org/10.1093/nar/gky752>.
  31. Inoue T, Shingaki R, Hirose S, Waki K, Mori H, Fukui K. 2007. Genome-wide screening of genes required for swarming motility in *Escherichia coli* K-12. *J Bacteriol* 189:950–957. <https://doi.org/10.1128/JB.01294-06>.
  32. Choi Y, Kim KP, Kim K, Choi J, Shin H, Kang DH, Ryu S. 2012. Possible roles of LysR-type transcriptional regulator (LTTR) homolog as a global regulator in *Cronobacter sakazakii* ATCC 29544. *Int J Med Microbiol* 302:270–275. <https://doi.org/10.1016/j.ijmm.2012.06.001>.
  33. Boonma S, Romsang A, Duang-Nkem J, Atichartpongkul S, Trinachartvanit W, Vattanaviboon P, Mongkolsuk S. 2017. The FinR-regulated essential gene *fprA*, encoding ferredoxin NADP+ reductase: roles in superoxide-mediated stress protection and virulence of *Pseudomonas aeruginosa*. *PLoS One* 12:e0172071. <https://doi.org/10.1371/journal.pone.0172071>.
  34. Pei B, Wang Y, Katzianer DS, Wang H, Wu H, Zhong Z, Zhu J. 2013. Role of a TehA homolog in *Vibrio cholerae* C6706 antibiotic resistance and intestinal colonization. *Can J Microbiol* 59:136–139. <https://doi.org/10.1139/cjm-2012-0673>.
  35. Westerman TL, Bogomolnaya L, Andrews-Polymenis HL, Sheats MK, Eifenbein JR. 2018. The *Salmonella* type-3 secretion system-1 and flagellar motility influence the neutrophil respiratory burst. *PLoS One* 13:e0203698. <https://doi.org/10.1371/journal.pone.0203698>.
  36. Santos RL, Zhang S, Tsois RM, Bäumlner AJ, Adams LG. 2002. Morphologic and molecular characterization of *Salmonella typhimurium* infection in neonatal calves. *Vet Pathol* 39:200–215. <https://doi.org/10.1354/vp.39-2-200>.
  37. Barthel M, Hapfelmeier S, Quintanilla-Martinez L, Kremer M, Rohde M, Hogardt M, Pfeiffer K, Russmann H, Hardt WD. 2003. Pretreatment of mice with streptomycin provides a *Salmonella enterica* serovar Typhimurium colitis model that allows analysis of both pathogen and host. *Infect Immun* 71:2839–2858. <https://doi.org/10.1128/IAI.71.5.2839-2858.2003>.
  38. Chubiz JEC, Golubeva YA, Lin D, Miller LD, Schlauch JM. 2010. Fliz regulates expression of the *Salmonella* pathogenicity island 1 invasion locus by controlling HiiD protein activity in *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 192:6261–6270. <https://doi.org/10.1128/JB.00635-10>.
  39. Ellermeier CD, Schlauch JM. 2003. RtsA and RtsB coordinately regulate expression of the invasion and flagellar genes in *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 185:5096–5108. <https://doi.org/10.1128/jb.185.17.5096-5108.2003>.
  40. Singer HM, Kühne C, Deditius JA, Hughes KT, Erhardt M. 2014. The *Salmonella* Spi1 virulence regulatory protein HiiD directly activates transcription of the flagellar master operon flhDC. *J Bacteriol* 196:1448–1457. <https://doi.org/10.1128/JB.01438-13>.
  41. Porwollik S, Santiviago CA, Cheng P, Long F, Desai P, Fredlund J, Srikumar S, Silva CA, Chu W, Chen X, Canals R, Reynolds MM, Bogomolnaya L, Shields C, Cui P, Guo J, Zheng Y, Endicott-Yazdani T, Yang H-J, Maple A, Ragoza Y, Blondel CJ, Valenzuela C, Andrews-Polymenis H, McClelland M. 2014. Defined single-gene and multi-gene deletion mutant collections in *Salmonella enterica* sv Typhimurium. *PLoS One* 9:e99820. <https://doi.org/10.1371/journal.pone.0099820>.
  42. Jarvik T, Smillie C, Groisman EA, Ochman H. 2010. Short-term signatures of evolutionary change in the *Salmonella enterica* serovar Typhimurium 14028 genome. *J Bacteriol* 192:560–567. <https://doi.org/10.1128/JB.01233-09>.
  43. Nakamura H, Saiki K, Mogi T, Anraku Y. 1997. Assignment and functional roles of the *cyo*ABCDE gene products required for the *Escherichia coli* bo-type quinol oxidase. *J Biochem* 122:415–421. <https://doi.org/10.1093/oxfordjournals.jbchem.a021769>.
  44. Manson MD, Tedesco P, Berg HC, Harold FM, Van der Drift C. 1977. A protonmotive force drives bacterial flagella. *Proc Natl Acad Sci U S A* 74:3060–3064. <https://doi.org/10.1073/pnas.74.7.3060>.
  45. Mangan MW, Lucchini S, T OC, Fitzgerald S, Hinton JCD, Dorman CJ. 2011. Nucleoid-associated protein HU controls three regulons that coordinate virulence, response to stress and general physiology in *Salmonella enterica* serovar Typhimurium. *Microbiology (Reading)* 157:1075–1087. <https://doi.org/10.1099/mic.0.046359-0>.
  46. Stothard P. 2000. The sequence manipulation suite: JavaScript programs for analyzing and formatting protein and DNA sequences. *Biotechniques* 28:1102–1104. <https://doi.org/10.2144/00286ir01>.
  47. Hong J, Ahn JM, Kim BC, Gu MB. 2009. Construction of a functional network for common DNA damage responses in *Escherichia coli*. *Genomics* 93:514–524. <https://doi.org/10.1016/j.ygeno.2009.01.010>.
  48. Altschul SF, Wootton JC, Gertz EM, Agarwala R, Morgulis A, Schaffer AA, Yu YK. 2005. Protein database searches using compositionally adjusted substitution matrices. *FEBS J* 272:5101–5109. <https://doi.org/10.1111/j.1742-4658.2005.04945.x>.
  49. Nelson KM, Young GM, Miller VL. 2001. Identification of a locus involved in systemic dissemination of *Yersinia enterocolitica*. *Infect Immun* 69:6201–6208. <https://doi.org/10.1128/IAI.69.10.6201-6208.2001>.
  50. Eriksson S, Lucchini S, Thompson A, Rhen M, Hinton JC. 2003. Unravelling the biology of macrophage infection by gene expression profiling of intracellular *Salmonella enterica*. *Mol Microbiol* 47:103–118. <https://doi.org/10.1046/j.1365-2958.2003.03313.x>.
  51. Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A* 7:6640–6645. <https://doi.org/10.1073/pnas.120163297>.

52. Sternberg NL, Maurer R. 1991. Bacteriophage-mediated generalized transduction in *Escherichia coli* and *Salmonella typhimurium*. *Methods Enzymol* 204:18–43. [https://doi.org/10.1016/0076-6879\(91\)04004-8](https://doi.org/10.1016/0076-6879(91)04004-8).
53. Wang RF, Kushner SR. 1991. Construction of versatile low-copy-number vectors for cloning, sequencing and gene expression in *Escherichia coli*. *Gene* 100:195–199. [https://doi.org/10.1016/0378-1119\(91\)90366-J](https://doi.org/10.1016/0378-1119(91)90366-J).
54. Bullas LR, Ryu JI. 1983. *Salmonella typhimurium* LT2 strains which are  $r^- m^+$  for all three chromosomally located systems of DNA restriction and modification. *J Bacteriol* 156:471–474. <https://doi.org/10.1128/JB.156.1.471-474.1983>.
55. Sambrook J, Fritsch E, Maniatis T. 1989. *Molecular cloning: a laboratory manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
56. Bogomolnaya LM, Aldrich L, Ragoza Y, Talamantes M, Andrews KD, McClelland M, Andrews-Polymenis HL. 2014. Identification of novel factors involved in modulating motility of *Salmonella enterica* serotype Typhimurium. *PLoS One* 9:e111513. <https://doi.org/10.1371/journal.pone.0111513>.
57. Elfenbein JR, Endicott-Yazdani T, Porwollik S, Bogomolnaya LM, Cheng P, Guo J, Zheng Y, Yang HJ, Talamantes M, Shields C, Maple A, Ragoza Y, Deatley K, Tatsch T, Cui P, Andrews KD, McClelland M, Lawhon SD, Andrews-Polymenis H. 2013. Novel determinants of intestinal colonization of *Salmonella enterica* serotype Typhimurium identified in bovine enteric infection. *Infect Immun* 81:4311–4320. <https://doi.org/10.1128/AI.00874-13>.
58. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods* 25:402–408. <https://doi.org/10.1006/meth.2001.1262>.
59. Bogomolnaya LM, Santiviago CA, Yang HJ, Baumberg AJ, Andrews-Polymenis HL. 2008. ‘Form variation’ of the O12 antigen is critical for persistence of *Salmonella Typhimurium* in the murine intestine. *Mol Microbiol* 70:1105–1119. <https://doi.org/10.1111/j.1365-2958.2008.06461.x>.