

# UC San Diego

## UC San Diego Previously Published Works

### Title

Non-invasive cervical vagus nerve stimulation effects on reaction time and valence image anticipation response

### Permalink

<https://escholarship.org/uc/item/49n622rw>

### Journal

Brain Stimulation, 15(4)

### ISSN

1935-861X

### Authors

Lerman, Imanuel

Klaming, Ruth

Spadoni, Andrea

et al.

### Publication Date

2022-07-01

### DOI

10.1016/j.brs.2022.06.006

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed



Published in final edited form as:

*Brain Stimul.* 2022 ; 15(4): 946–956. doi:10.1016/j.brs.2022.06.006.

## Non-invasive Cervical Vagus Nerve Stimulation Effects on Reaction Time and Valence Image Anticipation Response

Immanuel Lerman<sup>1,2,3,#,\*</sup>, Ruth Klaming<sup>1,4</sup>, Andrea Spadoni<sup>1,4</sup>, Dewleen G Baker<sup>1,4</sup>, Alan N Simmons<sup>1,4</sup>

<sup>1</sup>VA Center of Excellence for Stress and Mental Health, VA San Diego Healthcare System, La Jolla, CA, United States of America

<sup>2</sup>Department of Anesthesiology, Center for Pain Medicine, University of California San Diego School of Medicine, La Jolla, CA, United States of America

<sup>3</sup>Department of Electrical and Computer Engineering, University of California San Diego, La Jolla, CA, United States of America

<sup>4</sup>Department of Psychiatry University of California San Diego School of Medicine, La Jolla, CA, United States of America

### Abstract

**Background:** Norepinephrine (NE) driven noninvasive vagus nerve stimulation (nVNS), which improves attention and reduces reaction time, augments learning. Equally important, endogenous NE mediated arousal is highly dependent on the valence (positive or negative) of the exogenous stimulus. But to date, no study has measured valence specific effects of nVNS on both functional magnetic resonance imaging (fMRI) anticipation task response and reaction time in healthy individuals. Therefore, the aim of this pilot study was to assess whether nVNS vs sham modulates valence cortical anticipation task response and reaction time in a normative sample.

**Methods:** Participants received right sided transcutaneous cervical nVNS (N = 12) or sham (N = 12) stimulation during a 3T fMRI scan. Subjects first performed a continuous performance task (CPT) and then a cued anticipation task to images of positively and negatively valenced events during fMRI. Reaction times to cues and Blood oxygen level dependent (BOLD) response were examined over phase to identify effects of nVNS/sham over time.

**Results:** nVNS reduced reaction time for all valenced image anticipation trials. With the fMRI anticipation task, we observed a valence-specific effect; nVNS increased responsivity to images

\*Corresponding author: Immanuel Lerman, MD, MSc, ilterman@ucsd.edu.

#Current Address: VA San Diego, 3350 La Jolla Village Dr, San Diego, CA 92161 (MC116A), United States of America  
IL Conceptualized the study. IL, and ANS carried out the analyses. IL, and ANS wrote the manuscript, IL, ANS, RK, AS, DGB edited the manuscript and provided valuable feedback. IL is funded in part by Veterans Affairs, Career Development Program - Panel I (RRD8) #1IK2RX002920-01A1 and the VA Center for Stress and Mental Health.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

with negative valence and decreased responsivity to images with positive valence, whereas sham showed an inverse valence response.

**Conclusions:** nVNS was linked to reduced reaction time during the anticipation task. In tandem, nVNS consistently enhanced responsivity to negatively valenced images and diminished responsivity to positively valenced images, suggesting specific nVNS driven endogenous neurotransmitter signaling may contribute.

### Keywords

Norepinephrine; Locus Coeruleus; Attention; Arousal; Anticipation; Cervical Noninvasive Vagal Nerve Stimulation; functional magnetic resonance imaging

### Introduction:

Vagus nerve stimulation (VNS) modulates human brain activity, producing enhanced arousal and attention effects that are linked to augmented reaction time and learning [1–4]. Afferent vagal fibers largely terminate in the nucleus of the solitary tract (NTS) [5–7]. Anatomical tracer studies show NTS projection fibers innervate brainstem targets including the rostral ventromedial medulla, parabrachial nucleus, raphe nuclei and locus coeruleus (LC) [8]. Locus coeruleus activation was consistently shown in preclinical vagus nerve stimulation [9–11], while temporally dependent LC activation (with auricular vagus nerve stimulation) has been measured (maxima occurring within 6–7 min post stimulation) using functional imaging [1, 12–15]. Increases in LC norepinephrine (NE) concentrations activate the prefrontal cortex as well as a broad network of cortical and subcortical circuitry projections throughout the brain [16, 17]. In addition to LC-NE circuit projection effects, high adrenergic receptor concentrations (responsive to volume transmission NE release) are localized to nodes critical in the Salience Network (SN), (including anterior insula, amygdala, hippocampus, anterior cingulate) that drive the SN network specific activation [18]. LC-NE induced changes do not necessarily directly impact decision making or reward anticipation, but rather facilitate the dynamic reorganization of neural networks designed to quickly adapt to a changing environment [19]. As such, persistent LC activated neurons electrotonically couple enabling synchronic phasic bursting that is associated with task related, novel, threat based or unpredictable stimuli [20–22]. *Additive enhancement in LC activated NE signaling directly results from increases in negatively valenced “threatening” environmental stimuli. This activated NE signaling results in:* amplified alertness and attention, enhanced visual processing speed, augmented reaction time, that jointly reinforces learning [23–27]. Put simply, as negatively valenced threat or stress increases, the human neural response (i.e. requisitely executed via LC-NE signaling) is to react quickly, thus avoiding immediate and imminent harm [23]. In conjunction, the threat-based stimuli that induces alertness and attention coactivates complex cortico- limbic regulatory pathways, thereby continuously calibrating necessary responsivity through specific anticipation circuitry [28–31].

Altered threat anticipation of future aversive events is a core feature of anxiety disorders [32, 33]. In line with this maxim, generalized anxiety disorder (GAD), and posttraumatic stress disorder (PTSD) can be conceptualized as altered learning states characterized by

exaggerated prediction error due to an overgeneralization, followed by an exaggerated reaction to uncontrollable or unpredictable stressors [34]. Advanced functional brain imaging protocols can quantify the neural network that categorizes subjects with general anxiety and trauma disorders (i.e., GAD and PTSD) response during negative anticipation [35, 36].

Neuroimaging affective anticipation paradigms that model threat anticipation by measuring brain response during a cued period of expectancy to negatively relative to positively-valenced images demonstrate specific node activations including: the anterior insula, the anterior cingulate cortex (ACC), the amygdala, as well as the lateral (IPFC) and medial prefrontal cortex (mPFC) [37]. The anterior insula, highly responsive in anticipation paradigms, has efferent connections with ventral frontal brain regions such as the ACC as well as other nodes, (i.e., orbital frontal cortex (OFC) and periamygdaloid areas). Conventionally, the ventral subdivision of the ACC is known to play a significant role in regulatory emotional and physiological processing (BA-24, 32). Its ventral region has reciprocal projections to the anterior insula [38, 39] and the amygdala [40–43], exerting top-down regulation on these structures [42, 43]. Consequently, the ventral ACC is involved in fear conditioning [43–45], in the pathophysiology of anxiety disorders [45, 46], in cognition that is self-relevant [47–53], and in error processing [47–49, 52]. Given the importance of the anterior insula and the ACC in the integration of physiological and psychological processes, changes in their activity are established neural biomarkers for the efficacy of pharmacotherapies. But to date, bio-electronic medicine-based therapies (i.e., nVNS, Transmagnetic Stimulation, Transcranial Electrical Stimulation, and Deep Brain Stimulation) have been largely understudied in threat anticipation. Given the presumed nVNS NE neurochemical substrate, and the well-established neural nodes responsive in threat anticipation paradigms, we proposed to compare nVNS vs sham on both: 1) task performance reaction time, and 2) cortical activation patterns during an image anticipation task, utilizing valence specific cues (negative/positive).

## MATERIALS AND METHODS

### Subjects

A total of 24 mentally and physically healthy, right-handed subjects (10 females and 14 males; age range 18–54 years; mean  $\pm$  SD:  $27 \pm 8.4$  years) were recruited and enrolled in the study (Table 1). A subset of the cohort was included from a prior study [54]. Exclusion criteria were prior surgery or abnormal anatomy in the anterior cervical neck region, history of neurologic disease, implanted neurostimulator or cardiac pacemaker, cardiovascular disease or carotid artery disease, metallic implants that are not MRI-safe as well as any psychiatric disorder as defined by the Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DMS-IV) Axis I. The study was approved by the University of California San Diego, Institutional Review Board (IRB#150943). All subjects provided written informed consent prior to participation.

## Stimulation Paradigm

Subjects were randomized to receive either right sided transcutaneous cervical nVNS (n = 12) or sham (n = 12) stimulation during a functional magnetic resonance imaging (fMRI) scan. They were blinded to which condition they received. A pair of nonferromagnetic stainless-steel surface electrodes (1-cm diameter) were placed on the neck and secured with an adjustable Velcro strap collar, applied to either the right anterior cervical area (overlying the carotid artery) for active nVNS, or the right lateral cervical area (posterior to sternocleidomastoid) for the sham treatment. The surface electrodes were connected to the battery-powered stimulation unit by a 6-m shielded, grounded cable. Both the sham and nVNS devices delivered 1-ms duration bursts of 5 sinusoidal wave pulses at 5000 Hz with a repetition rate of 25 Hz, and a continuous train duration of 2 minutes. In both the nVNS and sham treatments, a computational fixed, initial 30-second ramp-up period (to peak voltage) was followed by 90 seconds of peak stimulation. In the nVNS treatment, the voltage was increased to 24 V, whereas in the sham stimulation it was increased to 4.5 V. Sham and nVNS devices were identical in appearance and produced similar sensations on the skin. In our previous work (pilot extension of parent study) [54] subjects were unable to differentiate sham from active nVNS. With the sham stimulation, subjects generally experience greater discomfort (because of stimulation of neck muscles), with maximal tolerable amplitude typically being only 2–8 V [2, 55]. Therefore, we employed a maximum sham voltage of 4.5 V to the far lateral neck position [2, 55]. Prior work suggests that use of the far lateral neck position with low voltage does not result in stimulation of the vagus nerve [2, 55]. Active transcutaneous cervical (20 V) nVNS is predicted to result in an electric field penetrance required to activate the vagus nerve, when the device is positioned directly over the carotid artery [56].

## Neuroimaging task

Approximately forty minutes after either nVNS or sham stimulation, subjects performed a validated valenced image anticipation task that combined a Continuous Performance Task (CPT) with interspersed presentation of affective stimuli (see Fig 1) [35, 57, 58]. During the CPT, subjects pressed the ‘FIRST (index finger)’ or ‘SECOND (middle finger)’ button on a hand-held button box in response to a blue “CIRCLE” or a “SQUARE”, respectively. Simultaneously, a 250 ms long 500-Hz tone was presented every 2 s. Subjects were instructed that when the shape turned green, accompanied by a 250-Hz (low pitch) tone, a “pleasant” positively valenced image would appear, whereas when the shape turned red, accompanied by a 1000-Hz (high pitch) tone, an “unpleasant” negatively valenced image would appear. Trials with green or red shapes represented anticipation periods. Picture stimuli included 17 positive and 17 negative valenced images. Positively valenced images from the International Affective Picture System (IAPS) and proprietary negatively valenced images were used [59], and evenly distributed across the 9 min 40 sec task duration. Response accuracy and response reaction time (ms) were obtained for the CPT, anticipation of a positive image (API), and anticipation of a negative image (ANI). To examine the behavioral effect of anticipation, we examined the difference between behavioral measures during the API and ANI. No subject response was required during affective image presentation. The valenced image anticipation task was conducted during a single fMRI scan sensitive to BOLD contrast using a 3T General Electric Discovery

MR 750 scanner (GE Healthcare). Images were collected with an eight-channel head coil (T2\*-weighted echoplanar imaging; repetition time, 1500 milliseconds; flip angle = 80, echo time, 32 milliseconds;  $64 \times 64$  matrix; field of view 240mm; thirty 4-mm axial sections; 386 scans). During the same experimental session, a high-resolution T1- weighted image (spoiled gradient recalled; flip angle =12, repetition time, 8 milliseconds; echo time, 3 milliseconds; 168 sections; field of view, 256 mm;  $256 \times 256$  matrix 1-mm<sup>3</sup> voxels) was obtained for anatomical reference.

Imaging analysis was conducted using a combination of AFNI [60] and ANTsR (<https://github.com/ANTsX/ANTsR>). T1 images were inhomogeneity-corrected to enhance registration using the N4 bias correction method [61]. Each subject's T1 was brain extracted, diffeomorphically registered to the 1mm mni\_icbm152\_nlin template, and segmented using ANTS [62, 63]. For preprocessing of the functional images, the first three EPI volumes were removed to allow for scanner equilibration. Next, EPI images were despiked, slice-time corrected, outliers were censored and N4 biased corrected. EPI images were then temporally whitened to reduce BOLD-signal temporal autocorrelation, motion corrected, detrended, the four top noise components were removed using CompCor [64], and data was spatially smoothed using Perona-Malik anisotropic diffusion [65]. Corrected EPI images were coregistered to the individual standardized T1 image using SyNbold. Forward and inverse registration transformations were concatenated and applied in a single step to minimize excess blurring of images. DVARs (>2.5) and framewise displacement (mean FD >.3mm) were identified if subjects moved however no subjects were removed. Hemodynamics were modelled (AFNI: 3dDeconvolve/TENT) during the task for valence (positive and negative anticipation blocks) and phase (phase 1 and phase 2 correspond to first half and second half of the valenced image anticipation task respectively). This two-phase model was chosen for the imaging data to retain a minimum of 8 per phase. Neural responses were measured for a duration of: 1) 6 seconds during anticipation 2) 2 seconds during the image presentation and, 3) approximately 8 seconds post image presentation; that in aggregate allow ample time for the API and ANI hemodynamic response.

To identify group and task effects on the hemodynamic response, hemodynamics for API and ANI were subjected to linear mixed-effects analysis using the R statistical package. A linear mixed-effects analysis for Group (sham and nVNS) by anticipation valence (positive and negative) by phase (phase 1 and phase 2) as fixed effects and subject as random effects, to identify task main effects. An additional linear mixed-effects analysis for Group (sham and nVNS) by anticipation valence (positive and negative) by trial (over the duration of the whole task) as fixed effects and subject as random effects, to identify task main effects. Multivoxel multiple comparisons were performed using Monte Carlo simulations (3dClustSim [bug fixed] with three-parameter noise modeling) in order to reduce potential false positives. To achieve cluster-wise threshold of  $p < 0.000001$ , a per-voxel thresholds of  $p < .000001$  were set along with a minimum of 4 voxels per cluster. As an exploratory measure valence specific neural response (AUC) to valence specific reaction time (in phase 1 and phase 2 and from phase 1 to phase 2) correlation analyses was carried out. Statistical analyses were performed with R version 3.4.3.

## Results:

### Participant demographics and psychiatric assessments

Age, sex and race were similar between the nVNS and the sham groups (Table I). Subjects were healthy; anxiety, depression, or posttraumatic stress disorder (PTSD), as measured by the Beck Anxiety Index (BAI), Beck Depression Inventory 2 (BDI-2), or the PTSD Check List–Civilian version (PCL-C) were below threshold cutoffs. Accordingly, no significant difference in mean scores between groups was noted for these measures. Two subjects failed the initial screen and were excluded from the study; one potential participant had a preexisting arrhythmia disorder (Wolf-Parkinson-White syndrome) and another had dental braces (Table I). The total sample used for analysis 12 subjects in sham and 12 in the VNS group, after exclusion of the 2 subjects who failed screening, and 6 others (3 in each group) due to technical problems with the task or scan.

### 3.2. Continuous Image Task Performance Accuracy and Reaction Time

As expected, all participants' accuracy on the continuous performance task was high (Mean=98.7%). It did not significantly differ across conditions (Continuous Task/No Anticipation: Mean=99.1%, Positive/Affective Control Anticipation: Mean=97.3%, Negative Anticipation: Mean=99.1% ( $p>.05$ ). Accuracy rates (correct control box button push) were also similar between groups ( $p>.05$ ) when examined across and between phase ( $p>.05$ ). There was a significant difference in reaction time between groups over time ( $p=.01$ ) with the nVNS group showing greater reduction in reaction time over task duration (i.e., over phases 1–2) for CPT, positive and negative image trials (see Fig 2 and Supplemental Table 1), when compared with the sham group. Similarly, there was a significant difference in reaction time between groups over time ( $p=.0009$ ) with the nVNS group showing greater reduction in reaction time over task duration (i.e., over trial) for CPT, positive and negative image trials (see Supplemental Table 2) when compared with the sham group.

#### Imaging results

**Regions of Interest (ROI):** The results of the whole brain analysis revealed a broad set of neural regions that demonstrate Group\*Valence\*Phase regions of interest. During the image anticipation task (independent of positive or negative imagery), 20 regions ( $p<.000001$ ) met cluster thresholds (with 4 voxel minimum) in the group\*valence\*phase main effect derivative of the LME analyses (Table 2). Significant clusters  $>9$  voxel in Group\*Valence\*Phase analysis (separated into 1<sup>st</sup>/2<sup>nd</sup> phase) demonstrates activation in the left precentral gyrus, Broadman Areas (BA) 44 and BA6, left anterior insula BA13, left anterior cingulate BA24 and BA32, and left post central gyrus BA3 ( $p<.000001$ ). Notable clusters  $>4$  voxels in Group\*Valence\*Phase analysis (separated into 1<sup>st</sup>/2<sup>nd</sup> phase) demonstrates activation in the left precentral gyrus (BA 3, 6), left orbital gyrus (BA 11), and left superior frontal gyrus (BA 6). Of note 80% (16/20) clusters identified were localized contralateral (left sided) to the nVNS (right sided). During the valence anticipation task, Group\*Valence\*Phase analyses (all listed ROI) of positive imagery demonstrate nVNS initial phase 1 greater response followed by consistent low response during phase 2; in converse, sham stimulation response was low during phase 1 but increased significantly during phase 2 (Fig 3). During the valence anticipation task, Group\*Valence\*Phase analyses



(all listed ROI) of negative imagery demonstrate an increase in response over task in the nVNS group (initial phase 1 low response followed by consistent greater response during phase 2); in converse, the sham group showed a decrease in response over task (response was greatest during phase 1 but decreased during phase 2) (Fig 3). Valence specific neural response (AUC) to valence specific reaction time (in phase 1 and phase 2 and for phase 1 transition to phase 2) correlation analyses did not demonstrate any significant correlations with Bonferroni correction for multiple comparisons (Supplemental Table 3, 4).

## Discussion:

Given the known alerting effects of nVNS, our study goal was to directly assess the effects of cervical nVNS vs sham on: (1) the task performance (i.e., reaction time) and (2) the coinciding neural response during an affective anticipation task. In this healthy control cohort, results revealed a stronger task performance (i.e., greater reduction in reaction time) in the nVNS group compared to the sham group during the valence anticipation task. Coordinate cortical neuroimaging revealed a valence-specific neuromodulatory effect; nVNS increased responsivity to images with negative valence and decreased responsivity to images with positive valence), whereas sham showed an inverse valence response. In aggregate nVNS reliably reduced reaction time independent of valence, while for the first time to our knowledge this is the first report of nVNS valence specific effects on the neural substrates thought to subserve alertness, attention, and anticipation.

## Potential Mechanisms of Enhancing Reaction Time and Alerting Effects Following nVNS

In this cohort, (nVNS vs sham), we observed consistent reduction in reaction time over phase. Considerable prior work provides the premise for our findings. Noradrenergic mechanisms can be modulated by stimulation (VNS) that primarily occurs through activation of the LC. This effect is supported by preclinical [10, 11, 66–68] as well as consistent clinical reports of nVNS mediated: 1) increases in salivary alpha-amylase (sAA) [69, 70], 2) pupillary change [71–73], and 3) alteration of visual evoked potential amplitude [69, 70, 74]. Multiple fMRI studies suggest nVNS results in activation of the LC [14, 15, 75] and other brainstem nuclei important in autonomic regulation [54]; in aggregate they suggest LC activation impacts nVNS effects, however change in additional neurotransmitters (GABAergic, Serotonergic, Cholinergic) may contribute as well [74]. Vagus nerve stimulation GABAergic effects are observed in the contralateral cortex to stimulation site (that correspond to ipsilateral hand) [76–78]. In our cohort, we observed a reduction of right-hand reaction time response in the right sided nVNS group. Therefore, if highly contributory, we would have expected a GABAergic mediated increase in (slowed) right hand reaction time as has been prior reported [79]. In as much, we postulate that the observed effect is likely compound, due to multiple neurotransmitters that are known to effect reaction time (Noradrenergic, Serotonergic, Cholinergic) [80–82]. Increases in LC-NE selectively activate SN nodes due to NE volume transmission amongst high density adrenergic receptors found in these regions [16, 17, 29, 83–87]. In support of this notion, nVNS reliably enhances visual search processing speed [3, 69, 87], that verify VNS as an alerting visual enhancement tool. Recent work from McIntire and colleagues (2021) support the tenet that cervical nVNS can improve attention, visual processing performance and



reaction time [4] that mirror LC-NE signaling attention effects of common stimulant drugs, such as Modafinil [88, 89] and Dextromethamphetamine [88]. Taken together, the present findings further support nVNS NE effects on reaction time, as described by other groups [3, 4, 70, 90]. In this cohort, we were likely underpowered to detect a group by valence specific reduction in the behavioral measure (i.e., reduction in reaction time see Supplemental Table 1, 2). However, we postulate that the nVNS evinced valence specific (negatively valenced image) neural (increases in cortical anticipatory node response) may also produce tangible behavioral (reduction in reaction time) effects (if adequately powered in a larger study), that are likely to be driven by additive threat and nVNS actuated LC-NE signaling. Further study, (in both pre-clinical and clinical models), is needed to determine whether negative alerting stimuli result in additive nVNS LC-NE signaling that reduce reaction time; *it may identify and translate an optimal arousal state that additively enhance nVNS effects on attention and alertness*. Moreso, preclinical and clinical work is required that may specify nVNS neurotransmitter (i.e., Noradrenergic, Serotonergic, Cholinergic) contribution to reduction in reaction time.

### Effect of nVNS on Valence Image Anticipation Task

The top six clusters with a 9 voxel minimum separated by group and phase included: two in the left precentral gyrus, left superior frontal gyrus, left anterior insula, left anterior cingulate, and left post central gyrus (Fig 3). The left anterior cingulate and the left anterior insula are known SN anticipatory nodes (Fig 3 K–T) [32–36, 91–93]. The other brain areas include premotor regions that may relate to the increased task attention discussed above [94, 95]. They are important in the Ventral Attention Network (VAN), also known to be preferentially activated by LC-NE signaling (Fig 3 A–M) [28, 85, 87]. In these areas, nVNS study subjects demonstrate increased activation over time (phase-1 to phase-2) during negative valence image anticipation, but in contrast, demonstrate decreased activation over time during positive valence image anticipation, (Fig 3). Remarkably, the sham group response was the reciprocal pattern. Specifically, sham subjects demonstrated a decrease in brain activation over time during negative valence image anticipation and an increase in activation for positive valence image anticipation (Fig 3). As a core tenant, exogenous stimuli that present an immediate environmental threat incrementally increase LC-NE release in turn augmenting alertness and attention [25–27]. Consistently in clinical and neuroimaging studies, negatively valenced images provoke higher arousal than positively valenced images [96–98]. Aversive stimuli conditioning, known to result in arousal, are dependent on LC-NE signaling that counteract learning adaptation [24]. Other studies have demonstrated that healthy adults have enhanced activity in insular cortex during the anticipation of aversive events, [91–93, 99–108] and that individuals with anxiety demonstrate exaggerated anterior insula activity and altered activity in anterior cingulate during emotional processing [35, 109–111]. Both high density LC-NE and noradrenergic receptor concentrations found in the right hemisphere, (including right anterior insula) [28, 29, 85–87] may contribute to greater right anterior insula anticipatory response to negatively valenced stimuli in healthy controls [91, 93, 101, 104, 105, 107, 112] and in subjects prone to anxiety disorders [35, 110]. In addition to commonly reported right anterior insula activation, bilateral [99, 103, 106, 107, 110, 113] and left anterior insula response activation [92, 102] are also reported in anticipation response to negatively valenced stimuli. In a

large (N=965 individuals) meta-analysis [114], bilateral anterior insula uniformly activate in anticipation to negatively valenced stimuli. In that study, the non-uniform population included both healthy and anxious individuals, that may have contributed to their bilateral anterior insula finding. Future studies and meta-analyses that covary the population (healthy and anxious or PTSD), may further support lateralized activation in aversive anticipation (right vs left anterior insula activation). Comparable to the anterior insula, anterior cingulate activation is universally reported: left, [92, 102, 106], right [115] and bilaterally [99, 103, 112, 113, 116] in aversive anticipation paradigms; likewise meta-analysis reports of bilateral anterior cingulate activation may indicate non-uniform population effects (i.e., inclusion of healthy and anxious individuals) in that study [114]. The observed (in our cohort) left anterior insula and anterior cingulate areas otherwise known to highly relate to cardiac parasympathetic responsivity, [117–119], potentially point to nVNS modulation of a valence specific autonomic arousal set point specific to our healthy control cohort. In our healthy control cohort, right sided cervical nVNS was carried out on all subjects. Canonical pre-clinical work by Randich and colleagues, (1990) [120], consistently instantiate VNS to LC projection cortical effects are predominantly contralateral to the site of stimulation, possibly explaining our observed left side (anterior insula, anterior cingulate, frontal cortex) effects. However, other as yet unstudied mechanisms may contribute in mental health populations (anxiety prone, GAD, PTSD) and therefore are not generalizable. Taken together, we hypothesize that in the nVNS group the intrinsic arousal associated with negative image anticipation met an additive threshold reciprocally activating SN nodes including the anterior insula and anterior cingulate. To our knowledge, the present study is the first to suggest that cervical noninvasive VNS improved reaction time by enhancing attentional networks during the valenced image (i.e., affective) anticipation task.

## Limitations

There are limitations that need to be addressed. First, we do not have a direct measurement of vagus nerve activity. Prior work utilizing the same cervical nVNS device have found neural evidence supporting stimulation of the vagus with similar stimulation-specific sensory evoked potentials as implantable devices [121–123] and changes consistent with increases in p300b amplitude indicative of noradrenaline specific effects (auricular vagus nerve stimulation) [69, 70, 123], while one study failed to do so [124]. Future studies are planned to objectively acquire fMRI scan evoked potential measures and peripheral and central noradrenaline biomarkers (salivary alpha amylase, plasma and cerebrospinal fluid norepinephrine) that may evince parameter specific change in noradrenaline pre-to-post nVNS. Second, we do not have direct measures of skin impedance during the stimulation as the procedures were carried out in the fMRI suite; therefore, we did not calculate the current imparted at the anterior cervical skin. Future studies will calculate current and as well as complete finite element modeling to identify current density at the cervical vagus nerve. Although the target of this stimulator is the vagal nerve, cutaneous afferents to the skin and innervation of the underlying tissue may have been stimulated as well. Baseline reaction times (prior to sham or nVNS) were not measured. Future work will incorporate a within subject design that allows for measures of pre-to-post intervention reaction time. We do not have direct measure of blindness between sham and active stimulation. Future within subject study will measure pre-to-post blindness. Although substantial data support

the anticipation protocol employed in this study (>300 subjects across studies) [58, 125–132]; color (red vs green) and tone (250 Hz vs 1000Hz) effects of the anticipation stimuli may have contributed. And, although the hemodynamic response measured includes the anticipation and image presentation, future work may disambiguate the which components, i.e., color, tone, or image contribute to the observed group by phase effects. This study was carried out in healthy control subjects and as yet cannot be extrapolated to disease state populations. As this was a pilot study, the small sample size was modest and may not adequately represent a larger population, thus results should be considered preliminary. However, the positive findings observed in this small cohort of healthy control subjects were robust and significant, warranting further investigation in a larger cohort, as well as in patients with mental health disorders (Posttraumatic Stress, Depression and Generalized Anxiety Disorder) or cognitive deficits (due to traumatic brain injury, stroke, dementia, etc.). Further, the valence image anticipation task was administered 40 minutes after nVNS; dose response and duration of effects were not assessed. Prior work demonstrates LC activation maximal response at 7 minutes post nVNS stimulation. The lack of LC activation response in the current cohort is likely due to the long post stimulation scan window, i.e., 40 min post stimulation. Additional work is needed to address these methodological considerations and to evaluate whether nVNS has only short-term effects or whether longer-term improvements, through noradrenergic driven facilitation of arousal-based learning acquisition and or consolidation, can be achieved. Accumulating work from Bremner et al., (2021) provide substantial evidence of nVNS improved learning in clinical populations with PTSD over a three-day period [133–136]. The study task, a self-motivated personalized traumatic script, when coupled with nVNS likely increased perceived control contributing to enhanced extinction learning; the authors observed a reduced SN (including anterior insula, anterior cingulate) response [136]. In contrast to the self-motivated personalized traumatic script task, valence anticipation paradigms provide a model of reduced perceived control that demonstrates a clear link between anxiety proneness and exaggerated anterior insula activity [35, 108]. These task paradigm differences (self-motivated vs reduced perceived control) likely contribute to the inverse effects observed in our studies [133–136]. Although promising, more work is clearly needed that may identify optimal arousal levels titrated for disease specific (i.e., PTSD, Generalized Anxiety Disorder, Anorexia) enhanced learning.

## Conclusion

This study revealed two major findings. First, when compared to sham, the nVNS group reaction time was consistently reduced for all (CPT, positive and negative) image anticipation trials. Second, we observed a valence specific nVNS neural effect; nVNS uniformly increased responsivity to negatively valenced images and decreased responsivity to positively valenced images, consistent with an extant nVNS literature suggesting that endogenous NE signaling may contribute. Future studies will be necessary to determine if the observed nVNS effects can be specifically calibrated in healthy control and mental health populations.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Bibliography:

1. Frangos E, Ellrich J, Komisaruk BR: Non-invasive access to the vagus nerve central projections via electrical stimulation of the external ear: fMRI evidence in humans. *Brain stimulation* 2015, 8(3):624–636. [PubMed: 25573069]
2. Frangos E, Komisaruk BR: Access to Vagal Projections via Cutaneous Electrical Stimulation of the Neck: fMRI Evidence in Healthy Humans. *Brain Stimul* 2017, 10(1):19–27. [PubMed: 28104084]
3. McIntire L, Goodyear C, McKinley A: Peripheral Nerve Stimulation to Augment Human Analyst Performance. In: 2019 IEEE Research and Applications of Photonics in Defense Conference (RAPID): 2019: IEEE; 2019: 1–3.
4. McIntire LK, McKinley RA, Goodyear C, McIntire JP, Brown RD: Cervical transcutaneous vagal nerve stimulation (ctVNS) improves human cognitive performance under sleep deprivation stress. *Communications biology* 2021, 4(1):1–9. [PubMed: 33398033]
5. Fornai F, Ruffoli R, Giorgi FS, Paparelli A: The role of locus coeruleus in the antiepileptic activity induced by vagus nerve stimulation. *European Journal of Neuroscience* 2011, 33(12):2169–2178. [PubMed: 21535457]
6. Berger A, Vespa S, Dricot L, Dumoulin M, Iachim E, Doguet P, Vandewalle G, El Tahry R: How is the norepinephrine system involved in the antiepileptic effects of Vagus Nerve Stimulation (VNS)? *Frontiers in neuroscience* 2021:1655.
7. Rutecki P: Anatomical, physiological, and theoretical basis for the antiepileptic effect of vagus nerve stimulation. *Epilepsia* 1990, 31:S1–S6.
8. Berthoud H-R, Neuhuber WL: Functional and chemical anatomy of the afferent vagal system. *Autonomic Neuroscience* 2000, 85(1–3):1–17. [PubMed: 11189015]
9. Hulseley DR, Riley JR, Loerwald KW, Rennaker II RL, Kilgard MP, Hays SA: Parametric characterization of neural activity in the locus coeruleus in response to vagus nerve stimulation. *Experimental neurology* 2017, 289:21–30. [PubMed: 27988257]
10. Dorr AE, Debonnel G: Effect of vagus nerve stimulation on serotonergic and noradrenergic transmission. *Journal of Pharmacology and Experimental Therapeutics* 2006, 318(2):890–898. [PubMed: 16690723]
11. Groves DA, Bowman EM, Brown VJ: Recordings from the rat locus coeruleus during acute vagal nerve stimulation in the anaesthetised rat. *Neuroscience letters* 2005, 379(3):174–179. [PubMed: 15843058]
12. Garcia RG, Lin RL, Lee J, Kim J, Barbieri R, Sclocco R, Wasan AD, Edwards RR, Rosen BR, Hadjikhani N: Modulation of brainstem activity and connectivity by respiratory-gated auricular vagal afferent nerve stimulation (RAVANS) in migraine patients. *Pain* 2017, 158(8):1461. [PubMed: 28541256]
13. Sclocco R, Garcia RG, Kettner NW, Fisher HP, Isenburg K, Makarovskiy M, Stowell JA, Goldstein J, Barbieri R, Napadow V: Stimulus frequency modulates brainstem response to respiratory-gated transcutaneous auricular vagus nerve stimulation. *Brain stimulation* 2020, 13(4):970–978. [PubMed: 32380448]
14. Yakunina N, Kim SS, Nam EC: Optimization of transcutaneous vagus nerve stimulation using functional MRI. *Neuromodulation: technology at the neural interface* 2017, 20(3):290–300. [PubMed: 27898202]
15. Kraus T, Kiess O, Hösl K, Terekhin P, Kornhuber J, Forster C: CNS BOLD fMRI effects of sham-controlled transcutaneous electrical nerve stimulation in the left outer auditory canal—a pilot study. *Brain stimulation* 2013, 6(5):798–804. [PubMed: 23453934]
16. Florin-Lechner SM, Druhan JP, Aston-Jones G, Valentino RJ: Enhanced norepinephrine release in prefrontal cortex with burst stimulation of the locus coeruleus. *Brain research* 1996, 742(1–2):89–97. [PubMed: 9117425]
17. Devoto P, Flore G, Saba P, Fa M, Gessa GL: Stimulation of the locus coeruleus elicits noradrenaline and dopamine release in the medial prefrontal and parietal cortex. *Journal of neurochemistry* 2005, 92(2):368–374. [PubMed: 15663484]
18. Zerbi V, Floriou-Servou A, Markicevic M, Vermeiren Y, Sturman O, Privitera M, von Ziegler L, Ferrari KD, Weber B, De Deyn PP: Rapid reconfiguration of the functional connectome

- after chemogenetic locus coeruleus activation. *Neuron* 2019, 103(4):702–718. e705. [PubMed: 31227310]
19. Bouret S, Sara SJ: Network reset: a simplified overarching theory of locus coeruleus noradrenaline function. *Trends in neurosciences* 2005, 28(11):574–582. [PubMed: 16165227]
  20. Berridge CW, Waterhouse BD: The locus coeruleus–noradrenergic system: modulation of behavioral state and state-dependent cognitive processes. *Brain research reviews* 2003, 42(1):33–84. [PubMed: 12668290]
  21. McEwen BS: Stress, adaptation, and disease: Allostasis and allostatic load. *Annals of the New York academy of sciences* 1998, 840(1):33–44. [PubMed: 9629234]
  22. Howells FM, Stein DJ, Russell VA: Synergistic tonic and phasic activity of the locus coeruleus norepinephrine (LC-NE) arousal system is required for optimal attentional performance. *Metabolic brain disease* 2012, 27(3):267–274. [PubMed: 22399276]
  23. Ross JA, Van Bockstaele EJ: The locus coeruleus-norepinephrine system in stress and arousal: unraveling historical, current, and future perspectives. *Frontiers in Psychiatry* 2021, 11:1581.
  24. Díaz-Mataix L, Piper WT, Schiff HC, Roberts CH, Campese VD, Sears RM, LeDoux JE: Characterization of the amplificatory effect of norepinephrine in the acquisition of Pavlovian threat associations. *Learning & Memory* 2017, 24(9):432–439. [PubMed: 28814469]
  25. Rufener KS, Geyer U, Janitzky K, Heinze HJ, Zaehle T: Modulating auditory selective attention by non-invasive brain stimulation: Differential effects of transcutaneous vagal nerve stimulation and transcranial random noise stimulation. *European Journal of Neuroscience* 2018, 48(6):2301–2309. [PubMed: 30144194]
  26. Rizzo P, Beelke M, De Carli F, Canovaro P, Nobili L, Robert A, Tanganelli P, Regesta G, Ferrillo F: Chronic vagus nerve stimulation improves alertness and reduces rapid eye movement sleep in patients affected by refractory epilepsy. *Sleep* 2003, 26(5):607–611. [PubMed: 12938816]
  27. Sara SJ, Bouret S: Orienting and reorienting: the locus coeruleus mediates cognition through arousal. *Neuron* 2012, 76(1):130–141. [PubMed: 23040811]
  28. Corbetta M, Shulman GL: Control of goal-directed and stimulus-driven attention in the brain. *Nature reviews neuroscience* 2002, 3(3):201–215. [PubMed: 11994752]
  29. Corbetta M, Patel G, Shulman GL: The reorienting system of the human brain: from environment to theory of mind. *Neuron* 2008, 58(3):306–324. [PubMed: 18466742]
  30. Mohanty A, Sussman TJ: Top-down modulation of attention by emotion. *Frontiers in Human Neuroscience* 2013, 7:102. [PubMed: 23554590]
  31. Herrmann MJ, Boehme S, Becker MP, Tupak SV, Guhn A, Schmidt B, Brinkmann L, Straube T: Phasic and sustained brain responses in the amygdala and the bed nucleus of the stria terminalis during threat anticipation. *Human brain mapping* 2016, 37(3):1091–1102. [PubMed: 26678871]
  32. Grillon C: Models and mechanisms of anxiety: evidence from startle studies. *Psychopharmacology* 2008, 199(3):421–437. [PubMed: 18058089]
  33. Eysenck M: *Anxiety and cognition: A unified theory*: Psychology Press; 2014.
  34. Mineka S, Zinbarg R: A contemporary learning theory perspective on the etiology of anxiety disorders: it's not what you thought it was. *American psychologist* 2006, 61(1):10. [PubMed: 16435973]
  35. Simmons AN, Stein MB, Strigo IA, Arce E, Hitchcock C, Paulus MP: Anxiety positive subjects show altered processing in the anterior insula during anticipation of negative stimuli. *Human brain mapping* 2011, 32(11):1836–1846. [PubMed: 21181800]
  36. Simmons AN, Flagan TM, Wittmann M, Strigo IA, Matthews SC, Donovan H, Lohr JB, Paulus MP: The effects of temporal unpredictability in anticipation of negative events in combat veterans with PTSD. *Journal of affective disorders* 2013, 146(3):426–432. [PubMed: 22910447]
  37. Tanovic E, Pruessner L, Joormann J: Attention and anticipation in response to varying levels of uncertain threat: An ERP study. *Cognitive, Affective, & Behavioral Neuroscience* 2018, 18(6):1207–1220.
  38. Dupont S, Bouilleret V, Hasboun D, Semah F, Baulac M: Functional anatomy of the insula: new insights from imaging. *Surgical and Radiologic Anatomy* 2003, 25(2):113–119. [PubMed: 12819943]

39. Augustine JR: Circuitry and functional aspects of the insular lobe in primates including humans. *Brain research reviews* 1996, 22(3):229–244. [PubMed: 8957561]
40. Öngür D, An X, Price J: Prefrontal cortical projections to the hypothalamus in macaque monkeys. *Journal of comparative neurology* 1998, 401(4):480–505. [PubMed: 9826274]
41. Phan KL, Britton JC, Taylor SF, Fig LM, Liberzon I: Corticolimbic blood flow during nontraumatic emotional processing in posttraumatic stress disorder. *Archives of General Psychiatry* 2006, 63(2):184–192. [PubMed: 16461862]
42. Likhtik E, Pelletier JG, Paz R, Paré D: Prefrontal control of the amygdala. *Journal of Neuroscience* 2005, 25(32):7429–7437. [PubMed: 16093394]
43. Quirk GJ, Likhtik E, Pelletier JG, Paré D: Stimulation of medial prefrontal cortex decreases the responsiveness of central amygdala output neurons. *Journal of Neuroscience* 2003, 23(25):8800–8807. [PubMed: 14507980]
44. Sotres-Bayon F, Cain CK, LeDoux JE: Brain mechanisms of fear extinction: historical perspectives on the contribution of prefrontal cortex. *Biological psychiatry* 2006, 60(4):329–336. [PubMed: 16412988]
45. Phan KL, Taylor SF, Welsh RC, Decker LR, Noll DC, Nichols TE, Britton JC, Liberzon I: Activation of the medial prefrontal cortex and extended amygdala by individual ratings of emotional arousal: a fMRI study. *Biological psychiatry* 2003, 53(3):211–215. [PubMed: 12559653]
46. Britton JC, Phan KL, Taylor SF, Fig LM, Liberzon I: Corticolimbic blood flow in posttraumatic stress disorder during script-driven imagery. *Biological psychiatry* 2005, 57(8):832–840. [PubMed: 15820703]
47. Bremner JD, Vermetten E, Vythilingam M, Afzal N, Schmahl C, Elzinga B, Charney DS: Neural correlates of the classic color and emotional stroop in women with abuse-related posttraumatic stress disorder. *Biological psychiatry* 2004, 55(6):612–620. [PubMed: 15013830]
48. Whalen PJ, Bush G, McNally RJ, Wilhelm S, McInerney SC, Jenike MA, Rauch SL: The emotional counting Stroop paradigm: a functional magnetic resonance imaging probe of the anterior cingulate affective division. *Biological psychiatry* 1998, 44(12):1219–1228. [PubMed: 9861465]
49. Bush G, Luu P, Posner MI: Cognitive and emotional influences in anterior cingulate cortex. *Trends in cognitive sciences* 2000, 4(6):215–222. [PubMed: 10827444]
50. Amodio DM, Frith CD: Meeting of minds: the medial frontal cortex and social cognition. *Nature reviews neuroscience* 2006, 7(4):268–277. [PubMed: 16552413]
51. Wicker B, Ruby P, Royet J-P, Fonlupt P: A relation between rest and the self in the brain? *Brain Research Reviews* 2003, 43(2):224–230. [PubMed: 14572916]
52. Northoff G, Bermpohl F: Cortical midline structures and the self. *Trends in cognitive sciences* 2004, 8(3):102–107. [PubMed: 15301749]
53. Keenan JP, Wheeler MA, Gallup GG Jr, Pascual-Leone A: Self-recognition and the right prefrontal cortex. *Trends in cognitive sciences* 2000, 4(9):338–344. [PubMed: 10962615]
54. Lerman I, Davis B, Huang M, Huang C, Sorkin L, Proudfoot J, Zhong E, Kimball D, Rao R, Simon B: Noninvasive vagus nerve stimulation alters neural response and physiological autonomic tone to noxious thermal challenge. *PLoS one* 2019, 14(2):e0201212. [PubMed: 30759089]
55. Lewine JD, Paulson K, Bangera N, Simon BJ: Exploration of the Impact of Brief Noninvasive Vagal Nerve Stimulation on EEG and Event-Related Potentials. *Neuromodulation: Technology at the Neural Interface* 2018.
56. Mourdoukoutas AP, Truong DQ, Adair DK, Simon BJ, Bikson M: High-Resolution Multi-Scale Computational Model for Non-Invasive Cervical Vagus Nerve Stimulation. *Neuromodulation* 2017.
57. Simmons A, Strigo IA, Matthews SC, Paulus MP, Stein MB: Initial evidence of a failure to activate right anterior insula during affective set-shifting in PTSD. *Psychosomatic Medicine* 2009, 71(4):373. [PubMed: 19398499]
58. Aupperle RL, Ravindran L, Tankersley D, Flagan T, Stein NR, Simmons AN, Stein MB, Paulus MP: Pregabalin influences insula and amygdala activation during anticipation of emotional images. *Neuropsychopharmacology* 2011, 36(7):1466–1477. [PubMed: 21430645]



59. Lang P, Bradley M, Cuthbert B: International affective picture system (IAPS): affective ratings of pictures and instruction manual. University of Florida, Gainesville. In.: Tech Rep A-8; 2008.
60. Cox RW: AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical research* 1996, 29(3):162–173. [PubMed: 8812068]
61. Tustison NJ, Avants BB, Cook PA, Zheng Y, Egan A, Yushkevich PA, Gee JC: N4ITK: improved N3 bias correction. *IEEE transactions on medical imaging* 2010, 29(6):1310–1320. [PubMed: 20378467]
62. Avants BB, Tustison NJ, Wu J, Cook PA, Gee JC: An open source multivariate framework for n-tissue segmentation with evaluation on public data. *Neuroinformatics* 2011, 9(4):381–400. [PubMed: 21373993]
63. Klein S, Staring M, Murphy K, Viergever MA, Pluim JP: Elastix: a toolbox for intensity-based medical image registration. *IEEE transactions on medical imaging* 2009, 29(1):196–205. [PubMed: 19923044]
64. Behzadi Y, Restom K, Liao J, Liu TT: A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage* 2007, 37(1):90–101. [PubMed: 17560126]
65. Nam H, Lee D, Lee JD, Park H-J: A method for anisotropic spatial smoothing of functional magnetic resonance images using distance transformation of a structural image. *Physics in Medicine & Biology* 2011, 56(15):5063. [PubMed: 21772082]
66. Raedt R, Clinckers R, Mollet L, Vonck K, El Tahry R, Wyckhuys T, De Herdt V, Carrette E, Wadman W, Michotte Y: Increased hippocampal noradrenaline is a biomarker for efficacy of vagus nerve stimulation in a limbic seizure model. *Journal of neurochemistry* 2011, 117(3):461–469. [PubMed: 21323924]
67. Mridha Z, de Gee JW, Shi Y, Alkashgari R, Williams J, Suminski A, Ward MP, Zhang W, McGinley MJ: Graded recruitment of pupil-linked neuromodulation by parametric stimulation of the vagus nerve. *Nature communications* 2021, 12(1):1–14.
68. Hassert D, Miyashita T, Williams C: The effects of peripheral vagal nerve stimulation at a memory-modulating intensity on norepinephrine output in the basolateral amygdala. *Behavioral neuroscience* 2004, 118(1):79. [PubMed: 14979784]
69. Warren CM, Tona KD, Ouwkerk L, Van Paridon J, Poletiek F, van Steenberg H, Bosch JA, Nieuwenhuis S: The neuromodulatory and hormonal effects of transcutaneous vagus nerve stimulation as evidenced by salivary alpha amylase, salivary cortisol, pupil diameter, and the P3 event-related potential. *Brain stimulation* 2019, 12(3):635–642. [PubMed: 30591360]
70. Ventura-Bort C, Wirkner J, Genheimer H, Wendt J, Hamm AO, Weymar M: Effects of transcutaneous vagus nerve stimulation (tVNS) on the P300 and alpha-amylase level: a pilot study. *Frontiers in human neuroscience* 2018, 12:202. [PubMed: 29977196]
71. Jodoin VD, Lespérance P, Nguyen DK, Fournier-Gosselin M-P, Richer F: Effects of vagus nerve stimulation on pupillary function. *International Journal of Psychophysiology* 2015, 98(3):455–459. [PubMed: 26437126]
72. Urbin MA, Lafe CW, Simpson TW, Wittenberg GF, Chandrasekaran B, Weber DJ: Electrical stimulation of the external ear acutely activates noradrenergic mechanisms in humans. *Brain Stimulation* 2021.
73. Sharon O, Fahoum F, Nir Y: Transcutaneous vagus nerve stimulation in humans induces pupil dilation and attenuates alpha oscillations. *Journal of Neuroscience* 2021, 41(2):320–330. [PubMed: 33214317]
74. Lewine JD, Paulson K, Banger N, Simon BJ: Exploration of the impact of brief noninvasive vagal nerve stimulation on EEG and event-related potentials. *Neuromodulation: Technology at the Neural Interface* 2019, 22(5):564–572. [PubMed: 30288866]
75. Frangos E, Komisaruk BR: Access to vagal projections via cutaneous electrical stimulation of the neck: fMRI evidence in healthy humans. *Brain stimulation* 2017, 10(1):19–27. [PubMed: 28104084]
76. Capone F, Assenza G, Di Pino G, Musumeci G, Ranieri F, Florio L, Barbato C, Di Lazzaro V: The effect of transcutaneous vagus nerve stimulation on cortical excitability. *Journal of Neural Transmission* 2015, 122(5):679–685. [PubMed: 25182412]



77. Marrosu F, Serra A, Maleci A, Puligheddu M, Biggio G, Piga M: Correlation between GABAA receptor density and vagus nerve stimulation in individuals with drug-resistant partial epilepsy. *Epilepsy research* 2003, 55(1–2):59–70. [PubMed: 12948617]
78. Keute M, Ruhnau P, Heinze H-J, Zaehle T: Behavioral and electrophysiological evidence for GABAergic modulation through transcutaneous vagus nerve stimulation. *Clinical Neurophysiology* 2018, 129(9):1789–1795. [PubMed: 29981954]
79. Stagg CJ, Bachtiar V, Johansen-Berg H: The role of GABA in human motor learning. *Current Biology* 2011, 21(6):480–484. [PubMed: 21376596]
80. Kuo H-I, Qi F-X, Paulus W, Kuo M-F, Nitsche MA: Noradrenergic Enhancement of Motor Learning, Attention, and Working Memory in Humans. *International Journal of Neuropsychopharmacology* 2021, 24(6):490–498. [PubMed: 33617635]
81. Furey ML, Pietrini P, Haxby JV, Drevets WC: Selective effects of cholinergic modulation on task performance during selective attention. *Neuropsychopharmacology* 2008, 33(4):913–923. [PubMed: 17534379]
82. Hasbroucq T, Rihet P, Blin O, Possamai C-A: Serotonin and human information processing: fluvoxamine can improve reaction time performance. *Neuroscience letters* 1997, 229(3):204–208. [PubMed: 9237494]
83. Sackeim HA, Keilp JG, Rush AJ, George MS, Marangell LB, Dormer JS, Burt T, Lisanby SH, Husain M, Cullum CM: The effects of vagus nerve stimulation on cognitive performance in patients with treatment-resistant depression. *Cognitive and Behavioral Neurology* 2001, 14(1):53–62.
84. Jacobs HI, Riphagen JM, Razat CM, Wiese S, Sack AT: Transcutaneous vagus nerve stimulation boosts associative memory in older individuals. *Neurobiology of Aging* 2015, 36(5):1860–1867. [PubMed: 25805212]
85. Morrison JH, Foote SL: Noradrenergic and serotonergic innervation of cortical, thalamic, and tectal visual structures in Old and New World monkeys. *Journal of Comparative Neurology* 1986, 243(1):117–138. [PubMed: 3950077]
86. ASTON-JONES G, Foote SL, Bloom FE: Anatomy and physiology of locus coeruleus neurons: functional implications. *Frontiers of clinical neuroscience* 1984, 2:92–116.
87. Pardo JV, Fox PT, Raichle ME: Localization of a human system for sustained attention by positron emission tomography. *Nature* 1991, 349(6304):61–64. [PubMed: 1985266]
88. Killgore WD, Kahn-Greene ET, Grugle NL, Killgore DB, Balkin TJ: Sustaining executive functions during sleep deprivation: a comparison of caffeine, dextroamphetamine, and modafinil. *Sleep* 2009, 32(2):205–216. [PubMed: 19238808]
89. Walsh JK, Randazzo AC, Stone KL, Schweitzer PK: Modafinil improves alertness, vigilance, and executive function during simulated night shifts. *Sleep* 2004, 27(3):434–439. [PubMed: 15164895]
90. Sun L, Peräkylä J, Holm K, Haapasalo J, Lehtimäki K, Ogawa KH, Peltola J, Hartikainen KM: Vagus nerve stimulation improves working memory performance. *Journal of Clinical and Experimental Neuropsychology* 2017, 39(10):954–964. [PubMed: 28492363]
91. Ploghaus A, Tracey I, Gati JS, Clare S, Menon RS, Matthews PM, Rawlins JNP: Dissociating pain from its anticipation in the human brain. *science* 1999, 284(5422):1979–1981. [PubMed: 10373114]
92. Chua P, Krams M, Toni I, Passingham R, Dolan R: A functional anatomy of anticipatory anxiety. *Neuroimage* 1999, 9(6):563–571. [PubMed: 10334900]
93. Simmons A, Matthews SC, Stein MB, Paulus MP: Anticipation of emotionally aversive visual stimuli activates right insula. *Neuroreport* 2004, 15(14):2261–2265. [PubMed: 15371746]
94. Hackley SA, Langner R, Rolke B, Erb M, Grodd W, Ulrich R: Separation of phasic arousal and expectancy effects in a speeded reaction time task via fMRI. *Psychophysiology* 2009, 46(1):163–171. [PubMed: 18992072]
95. Nagai Y, Critchley HD, Featherstone E, Fenwick PB, Trimble M, Dolan RJ: Brain activity relating to the contingent negative variation: an fMRI investigation. *Neuroimage* 2004, 21(4):1232–1241. [PubMed: 15050551]

96. Phan KL, Wager T, Taylor SF, Liberzon I: Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. *Neuroimage* 2002, 16(2):331–348. [PubMed: 12030820]
97. Lang PJ, Greenwald MK, Bradley MM, Hamm AO: Looking at pictures: Affective, facial, visceral, and behavioral reactions. *Psychophysiology* 1993, 30(3):261–273. [PubMed: 8497555]
98. Kimble MO, Fleming K, Bandy C, Kim J, Zambetti A: Eye tracking and visual attention to threatening stimuli in veterans of the Iraq war. *Journal of anxiety disorders* 2010, 24(3):293–299. [PubMed: 20138463]
99. Nitschke JB, Sarinopoulos I, Mackiewicz KL, Schaefer HS, Davidson RJ: Functional neuroanatomy of aversion and its anticipation. *Neuroimage* 2006, 29(1):106–116. [PubMed: 16181793]
100. Carlson JM, Greenberg T, Rubin D, Mujica-Parodi LR: Feeling anxious: anticipatory amygdalo-insular response predicts the feeling of anxious anticipation. *Social cognitive and affective neuroscience* 2011, 6(1):74–81. [PubMed: 20207692]
101. Samanez-Larkin GR, Hollon NG, Carstensen LL, Knutson B: Individual differences in insular sensitivity during loss anticipation predict avoidance learning. *Psychological science* 2008, 19(4):320–323. [PubMed: 18399882]
102. Alvarez RP, Biggs A, Chen G, Pine DS, Grillon C: Contextual fear conditioning in humans: cortical-hippocampal and amygdala contributions. *Journal of Neuroscience* 2008, 28(24):6211–6219. [PubMed: 18550763]
103. Marschner A, Kalisch R, Vervliet B, Vansteenwegen D, Büchel C: Dissociable roles for the hippocampus and the amygdala in human cued versus context fear conditioning. *Journal of Neuroscience* 2008, 28(36):9030–9036. [PubMed: 18768697]
104. Carlsson K, Andersson J, Petrovic P, Petersson KM, Öhman A, Ingvar M: Predictability modulates the affective and sensory-discriminative neural processing of pain. *Neuroimage* 2006, 32(4):1804–1814. [PubMed: 16861005]
105. Shankman SA, Gorka SM, Nelson BD, Fitzgerald DA, Phan KL, O'Daly O: Anterior insula responds to temporally unpredictable aversiveness: an fMRI study. *Neuroreport* 2014, 25(8):596. [PubMed: 24614364]
106. Grupe DW, Oathes DJ, Nitschke JB: Dissecting the anticipation of aversion reveals dissociable neural networks. *Cerebral cortex* 2013, 23(8):1874–1883. [PubMed: 22763169]
107. Somerville LH, Wagner DD, Wig GS, Moran JM, Whalen PJ, Kelley WM: Interactions between transient and sustained neural signals support the generation and regulation of anxious emotion. *Cerebral cortex* 2013, 23(1):49–60. [PubMed: 22250290]
108. Alvarez R, Kirlic N, Misaki M, Bodurka J, Rhudy J, Paulus M, Drevets W: Increased anterior insula activity in anxious individuals is linked to diminished perceived control. *Translational psychiatry* 2015, 5(6):e591–e591. [PubMed: 26125154]
109. Simmons A, Strigo I, Matthews SC, Paulus MP, Stein MB: Anticipation of aversive visual stimuli is associated with increased insula activation in anxiety-prone subjects. *Biological psychiatry* 2006, 60(4):402–409. [PubMed: 16919527]
110. Stein MB, Simmons AN, Feinstein JS, Paulus MP: Increased amygdala and insula activation during emotion processing in anxiety-prone subjects. *American Journal of Psychiatry* 2007, 164(2):318–327. [PubMed: 17267796]
111. Paulus MP, Feinstein JS, Simmons A, Stein MB: Anterior cingulate activation in high trait anxious subjects is related to altered error processing during decision making. *Biological Psychiatry* 2004, 55(12):1179–1187. [PubMed: 15184037]
112. Alvarez RP, Chen G, Bodurka J, Kaplan R, Grillon C: Phasic and sustained fear in humans elicits distinct patterns of brain activity. *Neuroimage* 2011, 55(1):389–400. [PubMed: 21111828]
113. Greenberg T, Carlson JM, Rubin D, Cha J, Mujica-Parodi L: Anticipation of high arousal aversive and positive movie clips engages common and distinct neural substrates. *Social cognitive and affective neuroscience* 2015, 10(4):605–611. [PubMed: 24984958]
114. Andrzejewski JA, Greenberg T, Carlson JM: Neural correlates of aversive anticipation: An activation likelihood estimate meta-analysis across multiple sensory modalities. *Cognitive, Affective, & Behavioral Neuroscience* 2019, 19(6):1379–1390.

115. Sege CT, Bradley MM, Weymar M, Lang PJ: A direct comparison of appetitive and aversive anticipation: Overlapping and distinct neural activation. *Behavioural brain research* 2017, 326:96–102. [PubMed: 28267576]
116. Schlund MW, Brewer AT, Richman DM, Magee SK, Dymond S: Not so bad: avoidance and aversive discounting modulate threat appraisal in anterior cingulate and medial prefrontal cortex. *Frontiers in behavioral neuroscience* 2015, 9:142. [PubMed: 26113813]
117. Matthews SC, Paulus MP, Simmons AN, Nelesen RA, Dimsdale JE: Functional subdivisions within anterior cingulate cortex and their relationship to autonomic nervous system function. *Neuroimage* 2004, 22(3):1151–1156. [PubMed: 15219587]
118. Lane R, Reiman E, Ahern G, Thayer J: 21. Activity in medial prefrontal cortex correlates with vagal component of heart rate variability during emotion. *Brain and cognition* 2001, 47(1–2):97–100.
119. Craig A: Forebrain emotional asymmetry: a neuroanatomical basis? *Trends in cognitive sciences* 2005, 9(12):566–571. [PubMed: 16275155]
120. Randich A, Ren K, Gebhart G: Electrical stimulation of cervical vagal afferents. II. Central relays for behavioral antinociception and arterial blood pressure decreases. *Journal of neurophysiology* 1990, 64(4):1115–1124. [PubMed: 2258737]
121. Nonis R, D’Ostilio K, Schoenen J, Magis D: Evidence of activation of vagal afferents by non-invasive vagus nerve stimulation: an electrophysiological study in healthy volunteers. *Cephalalgia* 2017, 37(13):1285–1293. [PubMed: 28648089]
122. Fallgatter A, Neuhauser B, Herrmann M, Ehlis A-C, Wagener A, Scheuerpflug P, Reiners K, Riederer P: Far field potentials from the brain stem after transcutaneous vagus nerve stimulation. *Journal of neural transmission* 2003, 110(12):1437–1443. [PubMed: 14666414]
123. Polak T, Markulin F, Ehlis A-C, Langer J, Ringel TM, Fallgatter AJ: Far field potentials from brain stem after transcutaneous vagus nerve stimulation: optimization of stimulation and recording parameters. *Journal of neural transmission* 2009, 116(10):1237–1242. [PubMed: 19728032]
124. D’Agostini M, Burger AM, Villca Ponce G, Claes S, von Leupoldt A, Van Diest I: No evidence for a modulating effect of continuous transcutaneous auricular vagus nerve stimulation on markers of noradrenergic activity. *Psychophysiology* 2022:e13984. [PubMed: 34990045]
125. Simmons AN, Norman SB, Spadoni AD, Strigo IA: Neurosubstrates of remission following prolonged exposure therapy in veterans with posttraumatic stress disorder. *Psychotherapy and psychosomatics* 2013, 82(6):382–389. [PubMed: 24061484]
126. Berner LA, Simmons AN, Wierenga CE, Bischoff-Grethe A, Paulus MP, Bailer UF, Kaye WH: Altered anticipation and processing of aversive interoceptive experience among women remitted from bulimia nervosa. *Neuropsychopharmacology* 2019, 44(7):1265–1273. [PubMed: 30840983]
127. Arce E, Simmons AN, Lovero KL, Stein MB, Paulus MP: Escitalopram effects on insula and amygdala BOLD activation during emotional processing. *Psychopharmacology* 2008, 196(4):661–672. [PubMed: 18058090]
128. Strigo IA, Matthews SC, Simmons AN: Right anterior insula hypoactivity during anticipation of homeostatic shifts in major depressive disorder. *Psychosomatic medicine* 2010, 72(3):316. [PubMed: 20100882]
129. Lovero KL, Simmons AN, Aron JL, Paulus MP: Anterior insular cortex anticipates impending stimulus significance. *Neuroimage* 2009, 45(3):976–983. [PubMed: 19280711]
130. Aupperle RL, Allard CB, Grimes EM, Simmons AN, Flagan T, Behrooznia M, Cissell SH, Twamley EW, Thorp SR, Norman SB: Dorsolateral prefrontal cortex activation during emotional anticipation and neuropsychological performance in posttraumatic stress disorder. *Archives of general psychiatry* 2012, 69(4):360–371. [PubMed: 22474105]
131. Brown GG, Ostrowitzki S, Stein MB, von Kienlin M, Liu TT, Simmons A, Wierenga C, Stein OY, Bruns A, Bischoff-Grethe A: Temporal profile of brain response to alprazolam in patients with generalized anxiety disorder. *Psychiatry Research: Neuroimaging* 2015, 233(3):394–401.
132. Harlé KM, Simmons AN, Norman SB, Spadoni AD: Neural affective mechanisms associated with treatment responsiveness in veterans with PTSD and comorbid alcohol use disorder. *Psychiatry Research: Neuroimaging* 2020, 305:111172. [PubMed: 32927371]

133. Gurel NZ, Wittbrodt MT, Jung H, Shandhi MMH, Driggers EG, Ladd SL, Huang M, Ko Y-A, Shallenberger L, Beckwith J: Transcutaneous cervical vagal nerve stimulation reduces sympathetic responses to stress in posttraumatic stress disorder: A double-blind, randomized, sham controlled trial. *Neurobiology of stress* 2020, 13:100264. [PubMed: 33344717]
134. Gurel NZ, Huang M, Wittbrodt MT, Jung H, Ladd SL, Shandhi MMH, Ko Y-A, Shallenberger L, Nye JA, Pearce B: Quantifying acute physiological biomarkers of transcutaneous cervical vagal nerve stimulation in the context of psychological stress. *Brain stimulation* 2020, 13(1):47–59. [PubMed: 31439323]
135. Bremner JD, Gurel NZ, Wittbrodt MT, Shandhi MH, Rapaport MH, Nye JA, Pearce BD, Vaccarino V, Shah AJ, Park J: Application of noninvasive vagal nerve stimulation to stress-related psychiatric disorders. *Journal of personalized medicine* 2020, 10(3):119. [PubMed: 32916852]
136. Wittbrodt MT, Gurel NZ, Nye JA, Ladd S, Shandhi MMH, Huang M, Shah AJ, Pearce BD, Alam ZS, Rapaport MH: Non-invasive vagal nerve stimulation decreases brain activity during trauma scripts. *Brain stimulation* 2020, 13(5):1333–1348. [PubMed: 32659483]


**Highlights:**

nVNS effects, (increase in arousal and reduction in reaction time) is in part ascribed to (LC-NE) cortical signaling.

In this study, nVNS (in contrast to sham) reduced reaction time across all valenced anticipation tasks.


In contrast to sham, nVNS: 1) increased responsivity to images with negative valence and 2) decreased responsivity to images with positive valence.

nVNS additive response during the negatively valenced (arousing) image anticipation tasks may have additively reduced reaction time across all valenced anticipation tasks.



**Figure 1.**


Noninvasive Vagus Nerve Stimulation Study Design. Subjects were screened and randomized to either the sham treatment or nVNS group. Sham stimulation was carried out posteriolateral to the sternocleidomastoid. In the nVNS group, stimulation occurred anteromedial to the sternocleidomastoid and lateral to the trachea. 40 min after completion of stimulation (sham or nVNS), subjects participated in the functional magnetic resonance imaging anticipation task. This task combines a continuous performance task (CPT) with the interspersed presentation of affective stimuli. Subjects were asked to press a button based on a type of shape (circle or square). They were instructed that a green shape accompanied by a low pitch tone indicated a positive image would appear (6 sec). In contrast, a red shape accompanied by a high pitch tone signaled an impending negative image (6 sec). Images displayed (2 sec) in this paradigm were taken from the International Affective Picture System (IAPS) [59].



**Figure 2.**

Reaction time across phases (1–2) demonstrate a group\*phase ( $p=.01$ ) significant effect; the nVNS group showed greater reduction in reaction time over the duration (phase 1–2) of the image task when compared to sham for Continuous Performance Task (CPT), as well as for the positive and negative imagery. Reaction time for all image tasks over all phases (i.e., CPT, positive and negative, for phase 1–2) improved in the nVNS group. (+/- is reported as Standard Error, CPT=Continuous Performance Task, POS=positive image task, NEG=negative image task, nVNS=Non-invasive Vagus Nerve Stimulation, Sham=sham device, Phase 1=1<sup>st</sup> 1/2<sup>nd</sup> of task duration Phase 2=2/2<sup>nd</sup> of task duration).





**Figure 3.**

Group\*Valence\*Phase analysis separated into 1<sup>st</sup> (solid line) and 2<sup>nd</sup> phase (dashed line) demonstrate left precentral gyrus, Brodmann Areas (BA) 44, (A), and BA6 (D, H), left post central gyrus BA3 (K), left anterior insula BA13 (O) and left anterior cingulate BA32 (R), ( $p < .000001$ ). During the valence anticipation task, Group\*Valence\*Phase analyses of negative imagery (red box) demonstrate nVNS phase 1 low response followed by consistent greater response during phase 2; in converse, sham stimulation response was greatest during phase 1, but decreased during phase 2 (B, E, I, L, P, S). During the valence anticipation task, Group\*Valence\*Phase analyses of positive imagery (green box) demonstrate nVNS initial

phase 1 greater response followed by consistent low response during phase 2; in converse, sham stimulation response was low during phase 1, but increased significantly during phase 2 (C, F, J, M, Q, T) for all ROI.

**Table 1.**

Demographic and affective measurement in nVNS and Sham groups.

	nVNS	Sham	
Age (years)	25 (18, 31)	28 (18, 54)	0.410 <sup>a</sup>
Sex	8M:4F [66%, 33%]	7M:5F [58%: 41%]	0.673 <sup>b</sup>
Race			
Asian	4 [33%]	4 [33%]	
Black	0 [0%]	1 [8.3%]	0.428 <sup>b</sup>
White	7 [58%]	7 [58%]	
Other	1 [8.3%]	0 [0%]	
BAI	2.41 (0.0, 13.0)	2.58(0.0, 12.0)	0.79 <sup>a</sup>
BDI-2	3.25 (0.0, 17.0)	2.91 (0.0, 14.0)	0.97 <sup>a</sup>
PCL-C	18.5 (17.0, 28.0)	19.75 (17.0, 28.0)	0.19 <sup>a</sup>

BAI = Beck Anxiety Inventory; BDI-2 = Beck Depression Inventory 2; PCL-C = Posttraumatic Stress Disorder Check List–Civilian version.

<sup>a</sup>= Mann Whitney U statistical test.<sup>b</sup>= Log-Likelihood.

() connote data ranges, [] connote percentile of overall group.

**Table 2.**

Whole-brain linear mixed effects results.

Voxels	x	y	z	F-Value	Region	BA
32	-52	17	9	33.413	Left Precentral Gyrus (44)	44
11	-58	0	32	32.404	Left Precentral Gyrus	6
10	-40	18	-7	28.586	Left Anterior Insula, Inferior Frontal Gyrus	13, 47
10	-61	-16	29	30.671	Left Postcentral Gyrus	3
10	-2	23	32	34.006	Left Cingulate Gyrus	24, 32
9	-4	-5	69	33.424	Left Superior Frontal Gyrus	6
8	39	-58	-38	28.673	Right Tuber	NA
6	-12	38	-28	30.781	Left Orbital Gyrus	11
6	-45	9	39	34.136	Left Middle Frontal Gyrus	9
6	-53	-13	45	27.827	Left Precentral Gyrus	3
6	5	11	69	28.536	Right Superior Frontal Gyrus	6
5	-25	56	28	31.947	Left Superior Frontal Gyrus	10
5	-42	-11	59	31.193	Left Precentral Gyrus	6
4	-26	-42	-38	30.217	Left Cerebellar Tonsil	NA
4	-64	-26	-12	40.139	Left Middle Temporal Gyrus	21
4	-47	37	-12	27.946	Left Inferior Frontal Gyrus	47
4	20	41	33	27.842	Right Superior Frontal Gyrus	9
4	-5	2	61	27.306	Left Superior Frontal Gyrus	6
4	-15	18	61	28.665	Left Superior Frontal Gyrus	6
4	8	-22	82	30.788	Right Paracentral Lobule	NA

Note: Abbreviations: Broadman Area = BA. *F*-value reflects peak, *F*-Value within the cluster. Voxel-size = 4 mm<sup>3</sup>. Cluster-thresholded,  $p < .000001$ ; *F*-Value threshold was 24.22, with whole-brain cluster-corrected  $< 0.000001$  ( $k > 4$  voxels).