

# UC Santa Cruz

## UC Santa Cruz Previously Published Works

### Title

Integration of pro-inflammatory cytokines, 12-lipoxygenase and NOX-1 in pancreatic islet beta cell dysfunction

### Permalink

<https://escholarship.org/uc/item/4b43h7xz>

### Journal

Molecular and Cellular Endocrinology, 358(1)

### ISSN

0303-7207

### Authors

Weaver, Jessica R  
Holman, Theodore R  
Imai, Yumi  
et al.

### Publication Date

2012-07-01

### DOI

10.1016/j.mce.2012.03.004

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

**Integration of Pro-Inflammatory Cytokines, 12-Lipoxygenase and NOX-1 in  
Pancreatic Islet Beta Cell Dysfunction.**

**Jessica R. Weaver<sup>o‡</sup>, Theodore Holman<sup>a</sup>, Ajit Jadhav<sup>b</sup>, Victor Kenyon<sup>a</sup>, David  
Maloney<sup>b</sup>, Jerry Nadler<sup>o‡</sup>, Ganesha Rai<sup>b</sup>, Anton Simenov<sup>b</sup> and David A. Taylor-  
Fishwick<sup>o‡§¶</sup>**

**<sup>o</sup>Strelitz Diabetes Center, <sup>‡</sup>Departments of Medicine, <sup>§</sup>Microbiology & Molecular  
Cell Biology, Eastern Virginia Medical School, Norfolk, VA.**

**<sup>a</sup>Department of Biochemistry, University of California Santa Cruz, Santa Cruz, CA.**

**<sup>b</sup>NIH Chemical Genomics Center, Bethesda, MD.**

**¶ Corresponding Author:**

**David A Taylor-Fishwick, PhD**

**LH 2128**

**Eastern Virginia Medical School**

**700 W Olney Road**

**Norfolk, VA 23501**

**Email           Taylord@evms.edu**

**FAX             757 446 5803**

**Phone          757 446 7359**

## ABSTRACT

Elevated cellular reactive species, which can be produced by diabetic serum conditions such as increased inflammatory cytokines, contribute to islet beta cell dysfunction and cell death. Cellular pathways that result in beta cell oxidative stress are poorly resolved. In this study, stimulation of human donor islets, primary mouse islets or homogeneous beta cell lines with a cocktail of inflammatory cytokines ( $\text{TNF}\alpha$ ,  $\text{IL-1}\beta$ ,  $\text{INF}\gamma$ ) significantly ( $p < 0.05$ ) induced NADPH oxidase-1 (NOX-1) gene expression. This pro-inflammatory cytokine cocktail concomitantly induced loss of islet glucose stimulated insulin response ( $p < 0.05$ ), elevated expression of MCP-1 ( $p < 0.01$ ), increased cellular reactive oxygen species (ROS) and induced cell death. Inhibitors of NADPH oxidase, apocynin and diphenylene iodonium, blocked ROS generation ( $p < 0.01$ ) and induction of MCP-1 ( $p < 0.05$ ) by pro-inflammatory cytokines in beta cells. It has previously been reported that pro-inflammatory cytokine stimulation induces 12-lipoxygenase (12-LO) expression in human islets. 12-hydroxyeicosatetraenoic acid (12-HETE), a product of 12-LO activity, stimulated NOX-1 expression in human islets ( $p < 0.05$ ). Additionally, a new selective inhibitor of 12-LO blocked pro-inflammatory cytokine-induction of NOX-1 in INS-1 beta cells ( $p < 0.01$ ). Inhibition NOX-1 was not seen with a structural analogue with no 12-LO inhibitory activity. Importantly, islets from human type 2 diabetic donors have an elevated expression of NOX-1 ( $p < 0.05$ ). This study reports the initial description of an integrated pathway in beta cells that links pro-inflammatory-induced beta cell dysfunction with 12-lipoxygenase and NADPH oxidase activation. Inhibitors of this pathway may provide a new therapeutic strategy for diabetes.

**KEYWORDS:**

**NADPH Oxidase, 12-Lipoxygenase, Cytokines, Diabetes, Islet beta cell, Inhibitors.**

## 1. INTRODUCTION

Oxidative Stress in the pancreatic beta cell is recognized as a pathogenic step associated with a loss of beta cell function. Pancreatic beta cells express a lower activity of free-radical detoxifying enzymes (eg, catalase, superoxide dismutase, glutathione peroxidase) when compared to other tissues (Grankvist, Marklund et al. 1981; Lenzen, Drinkgern et al. 1996; Tiedge, Lortz et al. 1997; Modak, Datar et al. 2007). Beta cells are also inefficient in rectifying oxidative damage to DNA (Modak, Parab et al. 2009). Thus, under conditions of sustained activation of intracellular reactive species, islets are readily overwhelmed and undergo oxidative stress (Lenzen 2008). Under oxidative stress conditions, the elevated reactive oxygen species (ROS), in addition to oxidizing proteins, lipids and DNA, also activate stress-sensitive second messengers such as p38MAPK, JNK (Purves, Middlemas et al. 2001) and PKC (Koya and King 1998). While a transient increase in ROS generation is a required second messenger for glucose stimulated insulin secretion in the beta cell (Goldstein, Mahadev et al. 2005; Pi, Bai et al. 2007; Morgan, Rebelato et al. 2009; Newsholme, Morgan et al. 2009), chronic activation of ROS is destructive to the function and survival of the beta cell.

Serum conditions associated with the diabetic state, increased pro-inflammatory cytokines, high free fatty acids (FFA) and elevated glucose levels are all potent inducers of high levels of cellular ROS (Janciauskiene and Ahren 2000; Oliveira, Verlengia et al. 2003; Cunningham, McClenaghan et al. 2005; Inoguchi and Nawata 2005; Nakayama,

Inoguchi et al. 2005; Uchizono, Takeya et al. 2006; Morgan, Oliveira-Emilio et al. 2007; Michalska, Wolf et al. 2010). Chronic elevation of pro-inflammatory cytokines is an established feature of type 1 diabetes (Eizirik and Mandrup-Poulsen 2001; Jorns, Gunther et al. 2005), and in recent studies low-grade chronic inflammation and increase in serum pro-inflammatory cytokines is recognized as a key feature of type 2 diabetes (Catalan, Gomez-Ambrosi et al. 2007; Steinberg 2007; Tilg and Moschen 2008; Al-Maskari, Al-Shukaili et al. 2010; Igoillo-Esteve, Marselli et al. 2010; Kang, Song et al. 2010; Su, Pei et al. 2010). Within the beta cell, cellular sources of ROS include induced mitochondrial stress (reviewed (Newsholme, Haber et al. 2007)), induced endoplasmic reticulum stress (reviewed (Volchuk and Ron 2010)) and potentially NADPH oxidase activation.

NADPH oxidases are proteins that transfer electrons across biological membranes. Their function is the generation of ROS, superoxide and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). The phagocyte NADPH oxidase was the first identified example of an enzyme system where ROS generation was the primary function rather than a byproduct, as seen in mitochondria and other cell components. Activation of phagocyte NADPH oxidase occurs through a complex series of protein interactions. Genome sequencing has subsequently identified a family of NOX proteins that form distinct NADPH oxidase complexes. NOX-1 is one of five homologues of the core catalytic component subunit of phagocyte NADPH oxidase (reviewed (Bedard and Krause 2007)). NOX isoforms and subunits of the functional complex have been identified in pancreatic beta cells supporting a role of NADPH oxidase in beta cell function/dysfunction (Cheng, Cao et al.

2001; Uchizono, Takeya et al. 2006; Kowluru 2010). Global inhibition of NADPH oxidase conferred protection to beta cell dysfunction induced by cytokines or fatty acids (Michalska, Wolf et al. 2010). Most recently we have shown that decreased NOX-1 expression is associated with beta cell protection in a transgenic model (Chang, Weaver et al. 2010).

In non-beta cell systems, activators of NOX activity include 12-lipoxygenase (12-LO), a lipid metabolizing enzyme. 12-LO converts arachidonic acid to lipid mediators including 12-hydroxyeicosatetraenoic acid (12-HETE). Platelet oxidation, generated by NADPH oxidase, is downstream of 12-LO (Nardi, Feinmark et al. 2004). Exposure of cultured neurons to methylisothiazolinone induced 12-LO which increased ERK activation and NADPH oxidase activation (Du, McLaughlin et al. 2002), linking 12-LO to ERK activation of NOX. Further, induction of monocyte chemotactic protein (MCP)-1 in peritoneal macrophages by 12-S-HETE (a product of 12-LO activity) was blocked by inhibitors of NADPH oxidase activity (apocynin and diphenyleneiodonium chloride) (Wen, Gu et al. 2008). In pancreatic islets, MCP-1 expression is inversely correlated with islet graft function (Marzorati, Antonioli et al. 2006). Additionally, proliferation and migration of colon adenocarcinoma cells is mediated by NOX-1 (de Carvalho, Sadok et al. 2008; Sadok, Bourgarel-Rey et al. 2008) that is stimulated by 12-LO or arachidonic acid (the substrate for 12-LO activity); inhibition of 12-LO decreased arachidonic acid induced NOX-1 expression and cell migration. However an interaction between 12-LO and NOX-1 has not been described in pancreatic beta cells.

12-Lipoxygenase activity has been linked with the development in T1DM. 12-LO knock out mice were resistant to the induction of diabetes by low dose streptozotocin (Bleich, Chen et al. 1999) and the 12-LO knock out mice lacked cytokine-induced conversion of arachidonic acid to 12-HETE, implying that 12-HETE generation was cytotoxic to beta cells (Bleich, Chen et al. 1999). The role of 12-LO as a key mediator in the development of autoimmune diabetes is supported by studies of 12-LO deletion in the type 1 diabetes (T1DM) mouse model, NOD mice. Knock out of leukocyte 12-LO in NOD mice resulted in a significant reduction (2.5% vs >60%) in the development of diabetes (McDuffie, Maybee et al. 2008). A direct role of pro-inflammatory cytokines in activating 12-LO activity is supported by cytokine-induced production of 12-HETE in both islets and beta-cell lines (Bleich, Chen et al. 1995; Chen, Yang et al. 2005). Addition of 12-HETE to human islets decreased islet viability and function (Ma, Nunemaker et al. 2010).

In this study, we describe activation of NOX-1 by pro-inflammatory cytokines in islets and beta cell lines. This upregulation in NOX-1 is linked to 12-LO activity and reproduced in islets from diabetic donors. The data described integrate pro-inflammatory cytokine induced beta cell dysfunction with 12-LO activation and NOX-1-induced ROS in a unifying pathway.

## **2. MATERIALS AND METHODS**

2.1 Institutional approvals were obtained for all procedures and resources described.



2.2 *Reagents and Cell lines:* Human donor islets were obtained from integrated islet distribution project (<http://iidp.coh.org>) and cultured in CMRL media (Mediatech, Manassas, VA). Mouse islets were freshly isolated by common bile duct cannulation and collagenase digestion. Islets were hand picked prior to use. Beta cell lines, INS-1 and  $\beta$ TC3 were cultured as described (Scharfmann, Tazi et al. 1993). Active inhibitors of 12-lipoxygenase and inactive structural analogues were kindly provided by NIH Chemical Genomics Center, Bethesda, MD. Other reagents were commercially sourced, human and mouse cytokines (R&D Systems, Minneapolis, MN), Apocynin, Diphenylene Iodonium Chloride (DPI), (Sigma Aldrich, St Louis, MO), 12-hydroxyeicosatetraenoic acid (12-HETE, Enzo Life Science, Farmingdale, NY).

2.3 *Cell stimulation:* Cells or islets were treated with (TNF $\alpha$  1ng/ml, IL-1 $\beta$  0.5ng/ml, IFN $\gamma$  10ng/ml) individually or in combination (triple cytokines) for 4 or 24 hours. Recovered cell pellets were analyzed for gene expression by real-time quantitative PCR as previously described (Ma, Nunemaker et al. 2010). Oxidative stress PCR array was performed following manufacturers instructions (SABiosceinces, Valencia, CA). All reactions were done in triplicate. Primer sequences used were; Human NOX-1 forward 5'-CAC AAG AAA AAT CCT TGG GTC AA-3'; Human NOX-1 reverse 5'-GAC AGC AGA TTG CGA CAC ACA-3'; Nox1 forward 5'-CGC TCC CAG CAG AAG GTC GTG ATT ACC AAG G-3'; Nox1 reverse 5'-GGA GTG ACC CCA ATC CCT GCC CCA ACC A-3'; MCP-1 forward, 5'-CTT CTG GGC CTG CTG TTC A-3'; MCP-1 reverse 5'-CCA GCC TAC TCA TTG GGA TCA-3'; Actin forward, 5'-AGG TCA TCA CTA TTG GCA ACG A-3'; Actin reverse, 5'-CAC TTC ATG ATG GAT TGA ATG TAG TT-3'; GAPDH

forward 5'-TCA CCA CCA TGG AG-3'; GAPDH reverse 5'-GCT AAG CAG TTG GT-3'. Taqman primers were used for caspase 3, human GAPDH and human 12-Lipoxygenase (Applied Biosystems, Carlsbad, CA).

2.4 *Detection of Reactive Species and Apoptosis:* Treated INS-1 cells were washed in PBS and placed in PBS containing 10 $\mu$ M 6-carboxy-2',7'-dichlorodihydrofluorescein diacetate, di(acetoxymethyl ester) (DCF-DA, Invitrogen, Carlsbad, CA) for 30 mins at 37°C. Following a PBS wash and re-incubation in PBS for 1 hr at 37°C, fluorescence was measured on a SpectraMax M5 plate reader (Molecular Devices, Sunnyvale, CA); emission wavelength 480 nm; excitation wavelength 530 nm. Islet apoptosis was screened microscopically. Islets were washed in PBS and incubated in ice cold PBS containing 1 $\mu$ g/ml Propidium Iodide and 0.1 $\mu$ M YO-PRO-1 (Invitrogen) for 30 min. Fluorescent signal was determined on an AxioObserver microscope (Carl Zeiss Inc.) using defined settings. Quantitation of fluorescent signal was achieved with Image J ([rsbweb.nih.gov/ij/](http://rsbweb.nih.gov/ij/)).

2.5 *Data and analysis:* All experiments were performed a minimum of three times. Data shown are mean  $\pm$  SEM. Data was analyzed using Prism 4.0 (GraphPad Software, Inc., La Jolla, CA). Significance was calculated using student two-tail t-tests with 95% confidence and defined as  $p < 0.05$ .

### 3. RESULTS AND DISCUSSION

Treatment of human donor islets with a cocktail of pro-inflammatory cytokines ( $\text{TNF}\alpha$ ,  $\text{IL-1}\beta$ ,  $\text{IFN}\gamma$ ) for 24 hours induced cell death as detected microscopically (Supplemental Figure 1A/B) and validated by increased gene expression of the apoptosis gene Caspase-3 (Supplemental Figure 1C). Disruption of islet function by pro-inflammatory cytokines (PICs) was further confirmed by a loss of the glucose-stimulated-insulin-response (Supplemental Figure 1D). PCR array analyses were performed with focus on oxidative stress pathways to determine intracellular pathways that are initiated by PICs leading to beta cell dysfunction (data not shown). An increase in expression of NADPH oxidase NOX-1 was identified and subsequently validated in quantitative real time PCR analysis (Figure 1). Stimulation with a cocktail of pro-inflammatory cytokines ( $\text{TNF}\alpha$ ,  $\text{IL-1}\beta$  and  $\text{IFN}\gamma$ ), at relevant concentrations, resulted in a significant ( $p<0.05$ ) increase in expression for NOX-1. A significant ( $p<0.05$ ) fold elevation in NOX-1 expression was observed in human donor islets, in freshly isolated mouse islets and also in homogeneous beta cell lines, mouse  $\beta\text{TC3}$  cells and rat INS-1 cells following pro-inflammatory cytokine stimulation. Other components of the functional NOX-1 complex have been described in beta cells (Cheng, Cao et al. 2001; Uchizono, Takeya et al. 2006; Kowluru 2010). Thus, elevated NOX-1 expression by PICs could result in a sustained increase in ROS generated via the NADPH oxidase complex. The induction of NOX-1 by triple pro-inflammatory cytokine combination was not reproduced by individual cytokine stimulation (Supplemental Figure 1E) suggesting an interaction between the triple cytokine stimulation occurs in beta cells to induce NOX-1 expression. Triple cytokine stimulation also induced markers of beta cell dysfunction, such as apoptotic markers, loss of glucose-stimulated insulin response and expression of MCP-

1, to levels that are not reproduced by individual cytokine stimulation (JW and DATF data not shown).

Elevated expression of MCP-1 is a recognized marker of beta cell dysfunction. Indeed, for human islet transplantation protocols, MCP-1 expression is inversely correlated with islet graft function (Marzorati, Antonioli et al. 2006). We therefore used MCP-1 expression as a marker of PIC-induced islet dysfunction to determine the importance of the NADPH oxidase in beta cell damage. Highly selective inhibitors of NOX isoforms and NADPH oxidase are not available; however, extensive literature on the use of apocynin and diphenylene iodonium chloride (DPI) as inhibitors of NADPH oxidase are published (Lambeth, Krause et al. 2008). Stimulation of the INS-1 beta cell line with triple pro-inflammatory cytokines produced a marked elevation in MCP-1 expression, consistent with induction of beta cell dysfunction (Figure 2). Treatment with apocynin or DPI significantly ( $p < 0.05$ ) reduced MCP-1 expression induced by pro-inflammatory cytokines. The inhibition was dose dependent. These data suggest that pro-inflammatory cytokines induce ROS production in beta cells through a NADPH oxidase mechanism that leads to beta cell dysfunction. Apocynin and DPI reduced beta cell intracellular ROS production that was induced by PICs (Supplement Figure 2).

Considering non-beta cell cellular models (Du, McLaughlin et al. 2002; Nardi, Feinmark et al. 2004; de Carvalho, Sadok et al. 2008; Sadok, Bourgarel-Rey et al. 2008; Wen, Gu et al. 2008), upstream activators of NADPH oxidase activity include 12-Lipoxygenase, an enzyme that has been implicated in the pathogenesis of diabetes development

(Bleich, Chen et al. 1999; McDuffie, Maybee et al. 2008) . Studies of a genetic deletion of 12-LO associate 12-LO activity with beta cell destruction and onset of diabetes. Further, 12-HETE, a lipid product of 12-LO activity directly induced beta cell dysfunction (Ma, Nunemaker et al. 2010). Induction of MCP-1 in peritoneal macrophages by 12-S-HETE was blocked by inhibitors of NADPH oxidase activity (apocynin or diphenyleneiodonium chloride) (Wen, Gu et al. 2008). These data indicate 12-LO activates NOX activity in a pathway leading to upregulation of MCP-1. However, this link has not previously been made in islet beta cells in relation to diabetes pathogenesis. To test this association, human donor islets were stimulated directly with 12-HETE, a bioactive lipid product of 12-LO activity. Treatment of human islets with 12-HETE (1nM) induced a significant 13-fold increase in NOX-1 expression ( $p < 0.05$ ) (Figure 3A). Further, a recently described selective inhibitor of 12-LO, cmpd #1 ( $IC_{50}$  12-LO 0.8 $\mu$ M) (Kenyon, Rai et al. 2011) was used to confirm the association between pro-inflammatory cytokines, 12-LO activity and NOX-1 expression. In the homogeneous beta cell line INS-1, the expression of NOX-1 induced by pro-inflammatory cytokines was significantly blocked by 5 $\mu$ M of the selective 12-LO inhibitor cmpd #1, and not by a structurally related molecule (Supplemental Figure 3) that is inactive ( $IC_{50} > 70\mu$ M) in 12-LO inhibition (cpmd #E Figure 3B).

Translating the clinical relevance of these findings to human diabetic patients, islets from human type 2 diabetic donors were compared with non-diabetic donors. Islet dysfunction was confirmed both by increased gene expression of MCP-1 ( $p < 0.05$ ) and additionally by the presence of a defective first phase insulin release in a perfusion

glucose-stimulated-insulin-secretion test (data not shown). Unstimulated gene expression of NOX-1 from 6 representative non-diabetic donors (non DM) and five representative type 2 diabetes donors (T2DM) was determined. Plotted in Figure 4 are the normalized (to GAPDH) inverse Ct values for the two donor groups. A significantly higher expression of NOX-1 was detected in islets from type 2 diabetic donors ( $p < 0.05$ ). Further, supporting an interaction between 12-LO and NOX-1 expression in human diabetes, the protein expression for 12-LO was elevated in the T2DM group in Western blot analysis relative to the housekeeping protein, actin (data not shown). T2DM is associated with an increase in serum pro-inflammatory cytokines (Catalan, Gomez-Ambrosi et al. 2007; Steinberg 2007; Tilg and Moschen 2008; Al-Maskari, Al-Shukaili et al. 2010; Igoillo-Esteve, Marselli et al. 2010; Kang, Song et al. 2010; Su, Pei et al. 2010).

In summary, this study describes for the first time in islet beta cells a linked association in pro-inflammatory cytokine-induced beta cell dysfunction between 12-lipoxygenase, NADPH oxidase NOX-1 and reactive oxygen species. Inhibition of this pathway is a candidate target to preserve and protect beta cell mass in diabetes. Development of selective NOX-1 inhibitors could offer a new therapeutic strategy in diabetes.

#### **4. REFERENCES**

Al-Maskari, M., Al-Shukaili, A. and Al-Mammari, A. (2010). Pro-inflammatory cytokines in Omani type 2 diabetic patients presenting anxiety and depression. *Iran J Immunol.* 7(2): 124-129.

Bedard, K. and Krause, K. H. (2007). The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev.* 87(1): 245-313.

Bleich, D., Chen, S., Gu, J. L., Thomas, L., Scott, S., Gonzales, N., Natarajan, R. and Nadler, J. L. (1995). Interleukin-1 beta regulates the expression of a leukocyte type of 12-lipoxygenase in rat islets and RIN m5F cells. *Endocrinology.* 136(12): 5736-5744.

Bleich, D., Chen, S., Zipser, B., Sun, D., Funk, C. D. and Nadler, J. L. (1999). Resistance to type 1 diabetes induction in 12-lipoxygenase knockout mice. *J Clin Invest.* 103(10): 1431-1436.

Catalan, V., Gomez-Ambrosi, J., Ramirez, B., Rotellar, F., Pastor, C., Silva, C., Rodriguez, A., Gil, M. J., Cienfuegos, J. A. and Fruhbeck, G. (2007). Proinflammatory cytokines in obesity: impact of type 2 diabetes mellitus and gastric bypass. *Obes Surg.* 17(11): 1464-1474.

Chang, T. J., Weaver, J., Bowman, A., Leone, K., Raab, R., Vinik, A. I., Pittenger, G. L. and Taylor-Fishwick, D. A. (2010). Targeted expression of Islet Neogenesis Associated Protein to beta cells enhances glucose tolerance and confers resistance to streptozotocin-induced hyperglycemia. *Mol Cell Endocrinol.*

Chen, M., Yang, Z. D., Smith, K. M., Carter, J. D. and Nadler, J. L. (2005). Activation of 12-lipoxygenase in proinflammatory cytokine-mediated beta cell toxicity. *Diabetologia.* 48(3): 486-495.

Cheng, G., Cao, Z., Xu, X., van Meir, E. G. and Lambeth, J. D. (2001). Homologs of gp91phox: cloning and tissue expression of Nox3, Nox4, and Nox5. *Gene.* 269(1-2): 131-140.

Cunningham, G. A., McClenaghan, N. H., Flatt, P. R. and Newsholme, P. (2005). L-Alanine induces changes in metabolic and signal transduction gene expression in a clonal rat pancreatic beta-cell line and protects from pro-inflammatory cytokine-induced apoptosis. *Clin Sci (Lond).* 109(5): 447-455.

de Carvalho, D. D., Sadok, A., Bourgarel-Rey, V., Gattacceca, F., Penel, C., Lehmann, M. and Kovacic, H. (2008). Nox1 downstream of 12-lipoxygenase controls cell proliferation but not cell spreading of colon cancer cells. *Int J Cancer.* 122(8): 1757-1764.

Du, S., McLaughlin, B., Pal, S. and Aizenman, E. (2002). In vitro neurotoxicity of methylisothiazolinone, a commonly used industrial and household biocide, proceeds via a zinc and extracellular signal-regulated kinase mitogen-activated protein kinase-dependent pathway. *J Neurosci.* 22(17): 7408-7416.

Eizirik, D. L. and Mandrup-Poulsen, T. (2001). A choice of death--the signal-transduction of immune-mediated beta-cell apoptosis. *Diabetologia.* 44(12): 2115-2133.

Goldstein, B. J., Mahadev, K. and Wu, X. (2005). Redox paradox: insulin action is facilitated by insulin-stimulated reactive oxygen species with multiple potential signaling targets. *Diabetes.* 54(2): 311-321.

Grankvist, K., Marklund, S. L. and Taljedal, I. B. (1981). CuZn-superoxide dismutase, Mn-superoxide dismutase, catalase and glutathione peroxidase in pancreatic islets and other tissues in the mouse. *Biochem J.* 199(2): 393-398.

Igoillo-Esteve, M., Marselli, L., Cunha, D. A., Ladriere, L., Ortis, F., Grieco, F. A., Dotta, F., Weir, G. C., Marchetti, P., Eizirik, D. L. and Cnop, M. (2010). Palmitate induces a pro-inflammatory response in human pancreatic islets that mimics CCL2 expression by beta cells in type 2 diabetes. *Diabetologia.* 53(7): 1395-1405.

Inoguchi, T. and Nawata, H. (2005). NAD(P)H oxidase activation: a potential target mechanism for diabetic vascular complications, progressive beta-cell dysfunction and metabolic syndrome. *Curr Drug Targets.* 6(4): 495-501.

Janciauskiene, S. and Ahren, B. (2000). Fibrillar islet amyloid polypeptide differentially affects oxidative mechanisms and lipoprotein uptake in correlation with cytotoxicity in two insulin-producing cell lines. *Biochem Biophys Res Commun.* 267(2): 619-625.

Jorns, A., Gunther, A., Hedrich, H. J., Wedekind, D., Tiedge, M. and Lenzen, S. (2005). Immune cell infiltration, cytokine expression, and beta-cell apoptosis during the development of type 1 diabetes in the spontaneously diabetic LEW.1AR1/Ztm-iddm rat. *Diabetes.* 54(7): 2041-2052.

Kang, Y. S., Song, H. K., Lee, M. H., Ko, G. J. and Cha, D. R. (2010). Plasma concentration of visfatin is a new surrogate marker of systemic inflammation in type 2 diabetic patients. *Diabetes Res Clin Pract.* 89(2): 141-149.

Kenyon, V., Rai, G., Jadhav, A., Schultz, L., Armstrong, M., Jameson, J. B., 2nd, Perry, S., Joshi, N., Bougie, J. M., Leister, W., Taylor-Fishwick, D. A., Nadler, J. L., Holinstat, M., Simeonov, A., Maloney, D. J. and Holman, T. R. (2011). Discovery of potent and selective inhibitors of human platelet-type 12- lipoxygenase. *J Med Chem.* 54(15): 5485-5497.

Kowluru, A. (2010). Small G proteins in islet beta-cell function. *Endocr Rev.* 31(1): 52-78.

Koya, D. and King, G. L. (1998). Protein kinase C activation and the development of diabetic complications. *Diabetes.* 47(6): 859-866.

Lambeth, J. D., Krause, K. H. and Clark, R. A. (2008). NOX enzymes as novel targets for drug development. *Semin Immunopathol.* 30(3): 339-363.

Lenzen, S. (2008). Oxidative stress: the vulnerable beta-cell. *Biochem Soc Trans.* 36(Pt 3): 343-347.

Lenzen, S., Drinkgern, J. and Tiedge, M. (1996). Low antioxidant enzyme gene expression in pancreatic islets compared with various other mouse tissues. *Free Radic Biol Med.* 20(3): 463-466.



Ma, K., Nunemaker, C. S., Wu, R., Chakrabarti, S. K., Taylor-Fishwick, D. A. and Nadler, J. L. (2010). 12-Lipoxygenase Products Reduce Insulin Secretion and  $\beta$ -Cell Viability in Human Islets. *J Clin Endocrinol Metab.* 95(2): 887-893.

Marzorati, S., Antonioli, B., Nano, R., Maffi, P., Piemonti, L., Giliola, C., Secchi, A., Lakey, J. R. and Bertuzzi, F. (2006). Culture medium modulates proinflammatory conditions of human pancreatic islets before transplantation. *Am J Transplant.* 6(11): 2791-2795.

McDuffie, M., Maybee, N. A., Keller, S. R., Stevens, B. K., Garmey, J. C., Morris, M. A., Kropf, E., Rival, C., Ma, K., Carter, J. D., Tersey, S. A., Nunemaker, C. S. and Nadler, J. L. (2008). Nonobese diabetic (NOD) mice congenic for a targeted deletion of 12/15-lipoxygenase are protected from autoimmune diabetes. *Diabetes.* 57(1): 199-208.

Michalska, M., Wolf, G., Walther, R. and Newsholme, P. (2010). The effects of pharmacologic inhibition of NADPH oxidase or iNOS on pro-inflammatory cytokine, palmitic acid or H<sub>2</sub>O<sub>2</sub>-induced mouse islet or clonal pancreatic beta cell dysfunction. *Biosci Rep.*

Modak, M. A., Datar, S. P., Bhonde, R. R. and Ghaskadbi, S. S. (2007). Differential susceptibility of chick and mouse islets to streptozotocin and its co-relation with islet antioxidant status. *J Comp Physiol B.* 177(2): 247-257.

Modak, M. A., Parab, P. B. and Ghaskadbi, S. S. (2009). Pancreatic islets are very poor in rectifying oxidative DNA damage. *Pancreas.* 38(1): 23-29.

Morgan, D., Oliveira-Emilio, H. R., Keane, D., Hirata, A. E., Santos da Rocha, M., Bordin, S., Curi, R., Newsholme, P. and Carpinelli, A. R. (2007). Glucose, palmitate and pro-inflammatory cytokines modulate production and activity of a phagocyte-like NADPH oxidase in rat pancreatic islets and a clonal beta cell line. *Diabetologia.* 50(2): 359-369.

Morgan, D., Rebelato, E., Abdulkader, F., Graciano, M. F., Oliveira-Emilio, H. R., Hirata, A. E., Rocha, M. S., Bordin, S., Curi, R. and Carpinelli, A. R. (2009). Association of NAD(P)H oxidase with glucose-induced insulin secretion by pancreatic beta-cells. *Endocrinology.* 150(5): 2197-2201.

Nakayama, M., Inoguchi, T., Sonta, T., Maeda, Y., Sasaki, S., Sawada, F., Tsubouchi, H., Sonoda, N., Kobayashi, K., Sumimoto, H. and Nawata, H. (2005). Increased expression of NAD(P)H oxidase in islets of animal models of Type 2 diabetes and its improvement by an AT1 receptor antagonist. *Biochem Biophys Res Commun.* 332(4): 927-933.

Nardi, M., Feinmark, S. J., Hu, L., Li, Z. and Karpatkin, S. (2004). Complement-independent Ab-induced peroxide lysis of platelets requires 12-lipoxygenase and a platelet NADPH oxidase pathway. *J Clin Invest.* 113(7): 973-980.

Newsholme, P., Haber, E. P., Hirabara, S. M., Rebelato, E. L., Procopio, J., Morgan, D., Oliveira-Emilio, H. C., Carpinelli, A. R. and Curi, R. (2007). Diabetes associated cell stress and

dysfunction: role of mitochondrial and non-mitochondrial ROS production and activity. *J Physiol.* 583(Pt 1): 9-24.

Newsholme, P., Morgan, D., Rebelato, E., Oliveira-Emilio, H. C., Procopio, J., Curi, R. and Carpinelli, A. (2009). Insights into the critical role of NADPH oxidase(s) in the normal and dysregulated pancreatic beta cell. *Diabetologia.* 52(12): 2489-2498.

Oliveira, H. R., Verlengia, R., Carvalho, C. R., Britto, L. R., Curi, R. and Carpinelli, A. R. (2003). Pancreatic beta-cells express phagocyte-like NAD(P)H oxidase. *Diabetes.* 52(6): 1457-1463.

Pi, J., Bai, Y., Zhang, Q., Wong, V., Floering, L. M., Daniel, K., Reece, J. M., Deeney, J. T., Andersen, M. E., Corkey, B. E. and Collins, S. (2007). Reactive oxygen species as a signal in glucose-stimulated insulin secretion. *Diabetes.* 56(7): 1783-1791.

Purves, T., Middlemas, A., Agthong, S., Jude, E. B., Boulton, A. J., Fernyhough, P. and Tomlinson, D. R. (2001). A role for mitogen-activated protein kinases in the etiology of diabetic neuropathy. *FASEB J.* 15(13): 2508-2514.

Sadok, A., Bourgarel-Rey, V., Gattacceca, F., Penel, C., Lehmann, M. and Kovacic, H. (2008). Nox1-dependent superoxide production controls colon adenocarcinoma cell migration. *Biochim Biophys Acta.* 1783(1): 23-33.

Scharfmann, R., Tazi, A., Polak, M., Kanaka, C. and Czernichow, P. (1993). Expression of functional nerve growth factor receptors in pancreatic beta-cell lines and fetal rat islets in primary culture. *Diabetes.* 42(12): 1829-1836.

Steinberg, G. R. (2007). Inflammation in obesity is the common link between defects in fatty acid metabolism and insulin resistance. *Cell Cycle.* 6(8): 888-894.

Su, S. C., Pei, D., Hsieh, C. H., Hsiao, F. C., Wu, C. Z. and Hung, Y. J. (2010). Circulating pro-inflammatory cytokines and adiponectin in young men with type 2 diabetes. *Acta Diabetol.*

Tiedge, M., Lortz, S., Drinkgern, J. and Lenzen, S. (1997). Relation between antioxidant enzyme gene expression and antioxidative defense status of insulin-producing cells. *Diabetes.* 46(11): 1733-1742.

Tilg, H. and Moschen, A. R. (2008). Inflammatory mechanisms in the regulation of insulin resistance. *Mol Med.* 14(3-4): 222-231.

Uchizono, Y., Takeya, R., Iwase, M., Sasaki, N., Oku, M., Imoto, H., Iida, M. and Sumimoto, H. (2006). Expression of isoforms of NADPH oxidase components in rat pancreatic islets. *Life Sci.* 80(2): 133-139.

Volchuk, A. and Ron, D. (2010). The endoplasmic reticulum stress response in the pancreatic beta-cell. *Diabetes Obes Metab.* 12 Suppl 2: 48-57.

Wen, Y., Gu, J., Vandenhoff, G. E., Liu, X. and Nadler, J. L. (2008). Role of 12/15-lipoxygenase in the expression of MCP-1 in mouse macrophages. *Am J Physiol Heart Circ Physiol.* 294(4): H1933-1938.

## 5. ACKNOWLEDGEMENTS:

We are grateful to the technical expertise of Kendall Leone who performed mouse islet isolation, Norine Kuhn for GSIS and to Dr. Yumi Imai for support analysis of T2DM donor islets. Human donor islets were supplied by the integrated islet distribution program (<http://iidp.coh.org>). This research was funded in part by JDRF (DATF, TH, JN), NIH R01 *DK55240* (DATF, JN), NIH RO1 *GM5606* (TH) and CDMRP *PR093521* (DATF).

## 6. FIGURE LEGENDS

**FIGURE 1:** Pro-inflammatory cytokines induce expression of NOX-1 in islets and beta cell lines. Expression of NOX-1 gene was determined by qRT-PCR in unstimulated (untreated) cells and cells stimulated with a cocktail of TNF $\alpha$ , IL-1 $\beta$  and IFN $\gamma$  (PIC treated). Data shows fold change in NOX-1 over control cells from three or greater experiments; \* p<0.05.

**FIGURE 2:** NADPH Oxidase inhibitors, Apocynin and DPI block expression of MCP-1 induced by pro-inflammatory cytokines. Expression of MCP-1 in INS-1 cells was determined by qRT-PCR following stimulation with a cocktail of TNF $\alpha$ , IL-1 $\beta$  and IFN $\gamma$  (PICs) with or without apocynin or DPI at the concentrations shown. Graph shows fold

change in MCP-1 expression relative to untreated cells. Data are from three experiments; \* $p < 0.05$ , \*\* $p < 0.01$ .

**FIGURE 3:** NOX-1 expression in islets and beta cells is induced by 12-Lipoxygenase.

A), Nox-1 gene expression of human donor islets treated with 1nM 12-HETE. Graph shows fold change in qRT-PCR NOX-1 expression relative to untreated islets. B), Pro-inflammatory cytokine stimulation of INS-1 cells in the presence or absence of 5 $\mu$ M Cmpd#1 (a selective 12-LO-inhibitor) or 5 $\mu$ M CmpdE (a structurally related molecule that is not active in 12-LO inhibition). Graph shows percent of cytokine response. Data are from  $\geq 3$  experiments; \*  $p < 0.05$ , \*\*  $p < 0.01$ .

**FIGURE 4:** NOX-1 expression is increased in islets from type 2 diabetic donors. Basal expression of NOX-1 was determined by qRT-PCR in unstimulated human donor islets, from non diabetic (non DM) or type 2 diabetic (T2DM). Graph shows inverse Ct value. Data are from six non diabetics and five type 2 diabetic donors; \* $p < 0.05$ .

**SUPPLEMENTAL FIGURE 1:** Pro-inflammatory cytokines induce beta cell death and

dysfunction. B), Human islets stimulated with a cocktail of  $TNF\alpha$ ,  $IL-1\beta$  and  $IFN\gamma$  (PIC) for 24 hours undergo apoptosis and necrosis (green, red and yellow fluorescence) unlike A), islets incubated in media alone for 24 hours. C), Induction of caspase-3 in mouse and human primary islets following treatment with PICs. Graph shows fold increase in qRT-PCR relative to untreated (control). D), Insulin secretion in human donor islets following 24 hours in media only (control) or media plus  $TNF\alpha$ ,  $IL-1\beta$  and

IFN $\gamma$  (cytokines) in response to 3mM (Low) or 18mM (High) glucose. E), NOX-1 expression in INS-1 cells following four hour stimulation with IL-1 $\beta$ , TNF $\alpha$ , IFN $\gamma$  alone or in combination (PICs). Graph shows fold increase in NOX-1 relative to untreated cells determined by qRT-PCR. Data are from  $\geq 3$  experiments; \*p<0.05, \*\* p<0.01.

**SUPPLEMENTAL FIGURE 2:** ROS production induced pro-inflammatory cytokines is inhibited by apocynin and DPI. INS-1 cells were stimulated with a cocktail of TNF $\alpha$ , IL-1 $\beta$  and IFN $\gamma$  (PIC) in the presence of 2mM apocynin (APO) or 10 $\mu$ M DPI. As a positive control, cells were also treated with 1mM H<sub>2</sub>O<sub>2</sub>. Graph shows fluorescence readout of DCF-DA expressed as a percentage of untreated cells (control). Data are from four experiments; \*\*p<0.01.

**SUPPLEMENTAL FIGURE 3:** Chemical structures of compound 1 and compound E.