

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

THE K- PROTON AND NEUTRON INTERACTION FROM 0.6 Bev TO 3.5 Bev

### Permalink

<https://escholarship.org/uc/item/4bc3v0m6>

### Authors

Cook, V.  
Cork, B.  
Hoang, T.F.  
et al.

### Publication Date

1960-08-01

UCRL 8736

cy 2

UNIVERSITY OF  
CALIFORNIA

*Ernest O. Lawrence*

*Radiation  
Laboratory*

FREE ENERGY FUNCTIONS FOR SOME  $\text{MO}_2$  OXIDES

BERKELEY, CALIFORNIA

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-8736  
Chemistry Distribution

*cy 2*

UNIVERSITY OF CALIFORNIA  
Lawrence Radiation Laboratory  
Berkeley, California

Contract No. W-7405-eng-48

FREE ENERGY FUNCTIONS FOR SOME  $MO_2$  OXIDES

M. S. Chandrasekharaiah and L. Brewer

April 1959

Printed for the U.S. Atomic Energy Commission

Printed in USA. Price 50 cents. Available from the  
Office of Technical Services  
U. S. Department of Commerce  
Washington 25, D. C.

FREE ENERGY FUNCTIONS FOR SOME  $MO_2$  OXIDES

M. S. Chandrasekharaiah and L. Brewer  
Lawrence Radiation Laboratory and College of Chemistry  
University of California, Berkeley 4, California

April 1959

## ABSTRACT

The free energy functions for nine transition metal dioxides have been estimated by considering them as linear symmetrical molecules.

Unlike the diatomic oxide case,<sup>1,2</sup> there are very few spectroscopic data available for the gaseous triatomic oxides of metals. Hence in calculating the thermodynamic functions for these dioxides, almost all of the necessary molecular constants have to be estimated. In the first instance, the free energy functions (fef) for nine transition metal oxides are calculated on the basis of a linear symmetrical molecular approximation.

(1) Electronic Free Energy Functions

Since the experimentally observed electronic states for any of these molecules are not known, an ionic model similar to one assumed in the calculating of fef for monoxides,<sup>3</sup> is assumed. Thus, the molecular electronic states are equated to the energy levels of  $M^{+4}$  ions as given by Moore.<sup>4</sup> Employing these energy levels of ions, the electronic free energy functions were calculated. However, no  $M^{+4}$  ion energy levels for Hf, Ta or W are available and the electronic free energy functions for them have been obtained by extrapolating the values of other molecules in the corresponding periodic groups.

(2) Rotational and Vibrational Free Energy Functions

In computing the rotational contribution to the fef, the interatomic distances of oxygen atom to central metal atom were approximated to the internuclear distance of the corresponding monoxides. This approximation implies that the bond order in the two classes of oxides are the same which may not

strictly be true. But the difference in the bond orders in these two classes introduces fairly small changes in the  $r_e$  values. However, this approximation of assuming the same bond distances in both dioxides and monoxides introduces a small correction in  $f_{ef}$  which is much smaller than the other uncertainties in calculating  $f_{ef}$ . Moreover, our knowledge of internuclear distances in the monoxides is not accurate to warrant any corrections. Walsh<sup>5</sup> has concluded that triatomic molecules with a total of 18 valence electrons would be bent as is observed for dioxides of the sulfur family. However, we have assumed that this would not be true with some of the electrons in d orbitals.

One of the biggest uncertainties introduced in calculating the  $f_{ef}$  for  $MO_2$  oxides is in estimating the bending frequencies of vibration. These frequencies are generally small and are doubly degenerate. Hence the contribution of bending vibrational mode of molecule to  $f_{ef}$  is considerable and any estimation of this frequency introduces a large uncertainty.

In our calculation, we have assumed that the stretching force constants of dioxides are the same as the force constants of the corresponding monoxides. Bending force constants are thus calculated assuming a constant ratio for the bending force constant to stretching force constant and this ratio was considered the same for all molecules as for  $CO_2$ . Thus all the three fundamental frequencies of vibrations were evaluated using valence force method.

The total free energy functions and the estimated frequencies of vibration for gaseous  $TiO_2$ ,  $VO_2$ ,  $CrO_2$ ,  $ZrO_2$ ,  $NbO_2$ ,  $MoO_2$ ,  $HfO_2$ ,  $TaO_2$  and  $WO_2$  are given in Table 1.

Table 1

Vibrational frequencies and $-\left(\frac{F^O - H^O}{T}\right)$ for $MO_2$ molecules									
	$\omega_1 \text{ cm}^{-1}$	$\omega_2 \text{ cm}^{-1}$	$\omega_3 \text{ cm}^{-1}$	298	1000	1500	2000	2500	3000
$TiO_2$	873	292	1125		60.5	65.5	69.3	72.3	74.9
$VO_2$	882	291	1125		64.2	69.4	73.0	76.4	79.0
$CrO_2$	785	259	996		66.0	71.4	75.4	78.6	81.2
$ZrO_2$	863	260	1000		62.4	67.5	71.2	74.3	76.8
$NbO_2$	870	262	1009		65.3	70.6	74.5	77.8	80.4
$MoO_2$	776	232	895		66.8	72.3	76.4	79.7	82.4
$HfO_2^*$	860	244	935		64.3	69.4	73.2	76.2	78.8
$TaO_2^*$	864	244	935		67.4	72.5	76.3	79.4	81.9
$WO_2^*$	675	216	835		62.7	74.0	78.0	80.9	83.4

\*electronic contributions are extrapolated values from values of other members of the same group.



REFERENCES

1. G. Herzberg, Spectra of diatomic molecules, Van Nostrand, New York, 1950.
2. B. Rosen, Données Spectroscopiques Concernant Les Molécules Diatomiques (Hermann and Cie, Paris, 1951).
3. L. Brewer and M. S. Chandrasekharaiah, Free energy functions for gaseous monoxides, University of California Radiation Laboratory Report UCRL-8713, April, 1959.
4. C. Moore, Atomic Energy Levels, National Bureau of Standards Circular 467, Vols I, II, and III. 1949-58.
5. A. D. Walsh, J. Chem. Soc. 1953, 2266.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.