

UC Davis

UC Davis Previously Published Works

Title

M2 Muscarinic Receptor-Dependent Contractions of Airway Smooth Muscle are Inhibited by Activation of β -Adrenoceptors.

Permalink

<https://escholarship.org/uc/item/4bk8j3jb>

Journal

Function, 3(6)

Authors

Alkawadri, Tuleen

Wong, Pei

Fong, Zhihui

et al.

Publication Date

2022


DOI

10.1093/function/zqac050

Peer reviewed

RESEARCH ARTICLE

M2 Muscarinic Receptor-Dependent Contractions of Airway Smooth Muscle are Inhibited by Activation of β -Adrenoceptors

Tuleen Alkawadri¹, Pei Yee Wong¹, Zhihui Fong¹, Fionnuala T Lundy², Lorcan P McGarvey², Mark A Hollywood¹, Keith D Thornbury¹, Gerard P Sergeant ^{1,*}

¹ Smooth Muscle Research Centre, Dundalk Institute of Technology, Dublin Road, Dundalk, Co. Louth A91 K584, Ireland and ²The Wellcome-Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Belfast, BT9 7BL, Northern Ireland


*Address correspondence to G.P.S. (e-mail: gerard.sergeant@dkit.ie)

Abstract

Beta-adrenoceptor (β -AR) agonists inhibit cholinergic contractions of airway smooth muscle (ASM), but the underlying mechanisms are unclear. ASM cells express M3 and M2 muscarinic receptors, but the bronchoconstrictor effects of acetylcholine are believed to result from activation of M3Rs, while the role of the M2Rs is confined to offsetting β -AR-dependent relaxations. However, a profound M2R-mediated hypersensitization of M3R-dependent contractions of ASM was recently reported, indicating an important role for M2Rs in cholinergic contractions of ASM. Here, we investigated if M2R-dependent contractions of murine bronchial rings were inhibited by activation of β -ARs. M2R-dependent contractions were apparent at low frequency (2Hz) electric field stimulation (EFS) and short (10s) stimulus intervals. The β 1-AR agonist, denopamine inhibited EFS-evoked contractions of ASM induced by reduction in stimulus interval from 100 to 10 s and was more effective at inhibiting contractions evoked by EFS at 2 than 20 Hz. Denopamine also abolished carbachol-evoked contractions that were resistant to the M3R antagonist 4-DAMP, similar to the effects of the M2R antagonists, methoctramine and AFDX-116. The inhibitory effects of denopamine on EFS-evoked contractions of ASM were smaller in preparations taken from M2R^{-/-} mice, compared to wild-type (WT) controls. In contrast, inhibitory effects of the β 3-AR agonist, BRL37344, on EFS-evoked contractions of detrusor strips taken from M2R^{-/-} mice were greater than WT controls. These data suggest that M2R-dependent contractions of ASM were inhibited by activation of β 1-ARs and that genetic ablation of M2Rs decreased the efficacy of β -AR agonists on cholinergic contractions.

Submitted: 26 August 2022; Revised: 19 September 2022; Accepted: 21 September 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of American Physiological Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.



Key words: airways; cholinergic; COPD; smooth muscle; muscarinic; bronchodilation

Introduction

β 2-adrenoceptor (β 2-AR) agonists and muscarinic receptor (MR) antagonists are the mainstay bronchodilator treatments used to prevent, or alleviate the symptoms of, obstructive lung conditions such as COPD and asthma.¹ β 2-AR agonists target β 2-ARs on airway smooth muscle (ASM) cells to induce ASM relaxation and reduce airway resistance,² whereas MR antagonists prevent the bronchoconstrictor effects of acetylcholine (ACh) released from parasympathetic nerves.³ There is also evidence of “cross-talk” between the beta-adrenoceptor (β -AR) and MR signalling pathways in ASM cells,^{4–7} and it has been shown that β -AR agonists inhibit contractile responses induced by MR agonists.^{8–11} β -ARs are coupled to Gs-proteins, which activate adenylate cyclase to elevate cytosolic cAMP levels and stimulate protein kinase A (PKA), while the bronchoconstrictor effects of ACh are thought to result from activation of postjunctional M3Rs, even though they are outnumbered by M2Rs by a ratio of 4:1 in most species.^{12–14} The role of the postjunctional M2Rs in ASM is less clear.

The inhibitory effects of β -AR agonists on ASM contraction are potentiated by M2R antagonists, presumably by preventing activation of G_i-proteins that decrease adenylate cyclase activity and would counteract the effects of β -AR activation.^{6,10,12,15–18} Therefore, it is thought that the main role of M2Rs in ASM is to provide a functional antagonism to the inhibitory effects of β -AR activation, rather than directly contributing to contractions induced by neurally released ACh. Nevertheless, studies on small airways of M3R knockout mice showed that cholinergic responses were ~60% lower than wild-type (WT) controls, but were absent in mice that had both M2 and M3Rs knocked out, consistent with a direct role for M2Rs in cholinergic contractions of ASM.¹⁹ In addition, a profound M2R-mediated hypersensitization of M3R-dependent contractions of murine ASM was reported by Alkawadri et al.,²⁰ indicating that activation of postjunctional M2Rs could make a greater contribution to cholinergic nerve-mediated contractions of ASM than previously realized.²¹

β -AR agonists are thought to exert their inhibitory effects on cholinergic contractions of ASM via inhibition of M3R signalling pathways. However, it has been noted that this assertion is based on sensitivity of known M3R-dependent pathways to PKA, rather than direct evidence showing that M3R signalling is affected by β -ARs in ASM cells.^{4,5} The finding that M2Rs make a direct contribution to ACh-dependent contractions of ASM raises the possibility that β -AR agonists could affect M2R-dependent contractions. In the present study, we found that the β -AR agonist denopamine inhibited M2R-dependent potentiation of electric field stimulation (EFS)-evoked contractions of ASM and abolished 4-DAMP-resistant contractions induced by CCh. Furthermore, the inhibitory effects of denopamine on EFS-induced contractions were reduced, rather than enhanced, in M2R null mice. These data suggest that M2R-dependent contractions of ASM are regulated by activation of β -ARs.

Material and Methods

Tissue Dissection

All procedures were carried out in accordance with current EU legislation and with the approval of Dundalk Institute of Technology Animal Use and Care Committee. Male and female C57BL/6 WT and B6N.129S4(Cg)-Chrm2^{tm1jwe}/J (M2R knock out) mice aged 10–16 wk old were humanely killed by intraperitoneal injection of pentobarbitone (100 mg/kg) and the lungs were removed and placed in oxygenated Krebs solution. M2R knock out (M2R KO) mice were purchased from the Jackson Laboratory (United States). Homozygote M2R KO mice were generated from the breeding of heterozygotes and were identified by genotyping using PCR as per JAX protocols and reagents. The bronchial tree was exposed by sharp dissection under a microscope to remove surrounding blood vessels and lung tissue.²² The left and right bronchi were removed and cut into rings and placed in Krebs solution.

Isometric Tension Recordings

Rings (1–2 mm) from the right and left main bronchi were mounted in water-jacketed organ baths, perfused with warmed Krebs solution, adjusted to 5 mN tension, and equilibrated for 40 min. Isometric contractions, were recorded using a Myobath system, and data acquired using DataTrax2 software (World Precision Instruments). Mean contraction amplitude was measured by averaging peak contraction amplitude of 10 electric-field stimulation (EFS)-induced contractions during each parameter, or before and during drug addition (when they had their maximal effect). Drugs were added directly to the organ bath, where they were diluted in Krebs solution to their final concentration. Electrical field stimulation was used to excite transmural nerves and was applied via two platinum electrode wires (5 mm length, 2.5 mm apart) by a MultiStim system-D330 stimulator (Digitimer Ltd, England), which delivered trains of pulses (pulse amplitude 20 V, nominal; pulse width 0.3 ms; 1 or 10 s duration) at frequencies of 2, 4, or 20 Hz, at intervals of 10 or 100 s. Experiments, which compared responses in M2R KO mice with WT controls were performed at the same time.

Drugs and Solutions

Carbachol, methoctramine, and BRL-37344 (Sigma Aldrich), indomethacin (Abcam), denopamine (Santa Cruz), and α,β -methylene ATP (Cayman) were dissolved in DMSO, ethanol, or distilled water as appropriate. Krebs solution was composed of: (mM) 120 NaCl, 5.9 KCl, 25 NaHCO₃, 1.2 NaH₂PO₄·2H₂O, 5.5 glucose, 1.2 MgCl₂, and 2.5 CaCl₂. pH was adjusted to 7.4 by bubbling the solution with 95% O₂, –5% CO₂.




Figure 1. Effects of the β_1 -AR agonist denopamine on EFS-evoked contractions of murine bronchial rings. (A) is a representative isometric tension recording showing the effect of denopamine ($3 \mu\text{M}$) on EFS-evoked contractions (2 Hz) that were augmented by a reduction in stimulus interval from 10 to 100 s. Contraction amplitude was reduced to levels observed prior to reduction in stimulus interval. (B) is a summary bar chart showing the effect of denopamine on mean contraction amplitude, before and during the presence of denopamine ($n = 15$, $N = 9$, and **** $P < .0001$ ANOVA).

Data Analysis and Statistics

Experimental series were obtained from four or more animals; n refers to the number of tissue strips studied and N to the number of animals. Data were analyzed using Prism software (GraphPad). Summary data are presented as mean \pm SEM. Statistical comparisons were performed on original (non-normalized) data using either Student's paired t -test or, if three experimental groups were compared, ANOVA followed by Bonferroni post hoc test, with $P < .05$ considered statistically significant.

Results

Alkawadri et al. demonstrated that the amplitude of EFS-evoked contractions of murine ASM were enhanced by a reduction in the stimulus interval from 100 to 10 s.²⁰ This effect was inhibited by the M2R antagonists, methoctramine, and AFDX-116, and was absent in preparations taken from M2R null mice. Application of the M3R antagonist abolished the entire response.²⁰ These data indicated that M3R-dependent contractions of ASM were potentiated by activation of postjunctional M2Rs. The purpose of the present study was to investigate if the M2R-dependent component of cholinergic contractions of murine ASM was affected by activation of β -ARs. Previous studies found that the inhibitory effects of β -AR agonists in murine ASM were brought about by activation of β_1 -ARs, and not β_2 -ARs as is the case in humans.^{23,24} For example, Birrell et al. showed that relaxations of murine trachea induced by the nonselective β -adrenoceptor agonist, isoprenaline, were inhibited by

genetic ablation of β_1 -AR, but not β_2 -ARs.²⁴ Therefore, we examined if denopamine, a selective β_1 -AR agonist, inhibited the M2R-dependent potentiation of EFS-evoked contractions of ASM induced by reduction in stimulus interval. Figure 1(A) shows a representative trace demonstrating that reducing the stimulus interval from 100 to 10 s enhanced contraction amplitude, and that this effect was reversed by denopamine ($3 \mu\text{M}$). In 15 preparations, mean contraction amplitude increased from 0.8 ± 0.07 to 2.1 ± 0.2 mN, following reduction in stimulus interval and decreased to 0.8 ± 0.1 mN ($P < .0001$, ANOVA, $n = 15$, and $N = 9$) when denopamine was added.

M2R-dependent potentiation of ASM contractions was evident at 2 Hz EFS, but not 20 Hz.²⁰ Therefore, to further examine if β -AR activation affected the M2R-component of the response we compared the effects of denopamine on contractions evoked at both frequencies. The representative traces in Figure 2(A) and (B) show the effect of cumulative addition of denopamine, over the concentration range 0.03 – $30 \mu\text{M}$, on contractions induced by EFS at 2 and 20 Hz, respectively. It was clear that contractions evoked by EFS at 2 Hz were more sensitive to denopamine than those at 20 Hz. This is borne out in the summary concentration-effect curves in Figure 2(C), which indicate that denopamine inhibited responses evoked by 2 Hz EFS with an IC_{50} of $1.2 \mu\text{M}$ (95% CI: 0.86 – $1.73 \mu\text{M}$, $n = 17$, and $N = 9$) versus $48 \mu\text{M}$ (95% CI: 27.2 – $86.4 \mu\text{M}$, $n = 8$, and $N = 5$) for responses evoked by 20 Hz EFS.

The representative traces in Figure 3(A) and (B) show that CCh-evoked contractions of ASM were reduced by the selective M3R antagonist, 4-DAMP (3 nM), resulting in a series of




Figure 2. Comparison of the effects of increasing concentrations of denopamine on contractions of murine bronchial rings evoked by EFS at 2 and 20 Hz. (A) and (B) are representative isometric tension recordings showing the effect of cumulative addition of denopamine (30 nM to 30 μM) on contractions by EFS at 2 and 20 Hz, respectively (10 s intervals). (C) Summary concentration-effect curves for denopamine on contractions evoked by EFS at 2 Hz (circles, $\text{IC}_{50} = 1.2 \mu\text{M}$, $n = 17$, and $N = 9$) and 20 Hz (triangles, $\text{IC}_{50} = 48 \mu\text{M}$, $n = 8$, and $N = 5$), respectively.

oscillatory contractions. These 4-DAMP-resistant contractions were abolished by the selective M2R antagonists methoctramine (100 nM, **Figure 3A**) and AFDX-116 (300 nM, **Figure 3B**), respectively. Summary data in **Figure 3(C)** show that 4-DAMP reduced the amplitude of CCh-induced contractions by 72% and that the amplitude of the residual contractions were reduced to less than 1% of control in the presence of 4-DAMP and methoctramine ($n = 9$, $N = 9$, $P < .01$, ANOVA). Similar results were achieved with AFDX-116 ($n = 6$, $N = 4$, $P < .01$, ANOVA, **Figure 3D**). Therefore, it was apparent that the 4-DAMP-insensitive contractions, evoked by 300 nM CCh, resulted from activation of M2Rs and we next examined if these contractions were sensitive to denopamine treatment.

The representative trace in **Figure 4(A)** shows that the oscillatory contractions that remained in the presence of 4-DAMP (3 nM) were abolished by denopamine (3 μM), indicating that the M2R component of the response was abolished by activation of β_1 -ARs. Summary data in **Figure 4(C)** indicate that

4-DAMP reduced the contractions by 68% ($n = 7$, $N = 6$, and $P < .001$) and that subsequent addition of denopamine (3 μM) abolished the remaining contractions ($P < .01$) and reduced resting tone below that recorded prior to CCh application. In contrast, when denopamine was added first, before 4-DAMP, it reduced the amplitude of the CCh response by 46% ($P < .001$, $n = 16$, and $N = 8$). Subsequent addition of 4-DAMP greatly reduced the amplitude of the denopamine-resistant contraction (**Figures 4B** and **D**; $P < .0001$). Therefore, denopamine abolished 4-DAMP-resistant contractions of ASM, but a large 4-DAMP-sensitive component of the CCh response remained when denopamine was added first. This indicates that denopamine abolished the M2R-component of the CCh response, but that a large M3R-component was unaffected by denopamine.

Previous studies indicated that β -AR agonists inhibited cholinergic contractions of ASM via PKA-mediated inhibition of M3R-dependent signalling pathways.⁴ In addition, it has




Figure 3. Effects of M2R antagonists methoctramine and AFDX-116 on CCh-evoked contractions of murine bronchial rings in the presence of 4-DAMP. (A) and (B) are isometric tension recording showing that methoctramine (100 nM; A) and AFDX-116 (300 nM; B) abolished CCh-induced contractions that remained in the presence of 4-DAMP (3 nM). (C) is a summary bar chart showing mean contraction amplitude (measured by area under the curve) before and during the presence of 4-DAMP and 4-DAMP plus methoctramine ($n = 9$, $N = 9$, $**P < .01$, ANOVA). (D) is a summary bar chart showing mean contraction amplitude before and during the presence of 4-DAMP, and 4-DAMP plus AFDX-116 ($n = 6$, $N = 4$, $**P < .01$, $*P < .05$, ANOVA).

been suggested that postjunctional M2Rs are not functionally involved in cholinergic contractions of ASM, but instead provide a functional antagonism to β -AR activation.^{15–18} If this model were correct, then blockade, or removal, of M2Rs would be expected to enhance the inhibitory effects of β -AR agonists, since the opposing effects of M2R activation would be lost. We investigated if this was the case in ASM by comparing the effects of denopamine on cholinergic nerve-evoked contractions taken from WT and M2R null mice. Representative traces showing the effect of denopamine on EFS-evoked contractions of ASM from WT and M2R null mice are shown in Figure 5(A) and (B) and corresponding summary data, are provided in Figure 5(C) and (D), respectively. In 9 WT mice, denopamine reduced mean contraction amplitude by 62% from 2.1 ± 0.2 to 0.8 ± 0.1 mN ($P < .0001$, and $n = 15$), whereas in M2R null preparations contractions were only reduced by 40.9% (from 2.2 ± 0.2 to 1.3 ± 0.1 mN, $P < .01$, $n = 9$, $N = 5$). Therefore, the effects of denopamine on EFS responses were reduced, rather than enhanced, in preparations lacking M2Rs. These data are at odds with the current models, which suggest that only role of M2Rs in ASM is to provide a functional antagonism to β -AR activation.

Functional antagonism between M2R and β -AR activation has been observed in studies of detrusor smooth muscle.²⁵ Therefore, we examined if the β 3-AR agonist, BRL37344 (100 nM) induced larger inhibitory effects on EFS-evoked contractions in detrusor strips from M2R null mice, compared to WT. These experiments were performed in the presence of α,β -methylene

ATP ($10 \mu\text{M}$) to desensitize P2X1 receptors and isolate the cholinergic component of the EFS response. In contrast to the effects of denopamine on ASM, BRL37344 (100 nM) induced greater inhibitory effects in bladder strips taken from M2R null mice (Figure 6B) compared to WT mice (Figure 6A). Summary data in Figure 6(C) and (D) show that BRL37344 reduced mean contraction amplitude by 18.7% in WT mice (5.6 ± 0.8 to 4.5 ± 0.6 mN, $n = 16$, $N = 8$) compared to 32.2% (7 ± 0.5 to 4.8 ± 0.4 mN, $n = 16$, and $N = 8$) in M2R KOs.

Discussion

Airway diameter is regulated by contraction of ASM and enhanced ASM contraction leads to increased airway resistance, a feature of obstructive airway conditions such as COPD.^{26,27} Increased cholinergic tone is the main reversible component of airway obstruction in COPD,^{28–31} therefore, it is of interest to understand the mechanisms underlying cholinergic contractions of ASM and how they are targeted by β -AR agonists. Most studies have determined that cholinergic tone of ASM results from activation of postjunctional M3Rs on ASM cells and is regulated by activation of autoinhibitory M2Rs located on parasympathetic nerves that limit ACh output.^{32,33} The contribution of postjunctional M2Rs to cholinergic tone of ASM is less clear and most studies conclude that their contribution to ACh-induced contractions is negligible, but that their activation can counteract inhibitory effects of β -AR agonists.^{10,12,14,16,18}




Figure 4. Effects of denopamine on CCh-evoked contractions of murine bronchial rings, before and during the presence of 4-DAMP. (A) is an isometric tension recording showing that denopamine (3 μM) abolished CCh-induced contractions that remained in the presence of 4-DAMP (3 nM). (B) is an isometric tension recording showing that denopamine (3 μM) reduced the amplitude of CCh-induced contractions of ASM and that the residual contractions were inhibited by 4-DAMP (3 nM). (C) is a summary bar chart showing mean contraction amplitude (measured by area under the curve) before and during the presence of 4-DAMP and 4-DAMP plus denopamine ($n = 7$, $N = 6$, $***P < .001$, and $**P < .01$, ANOVA). (D) is a summary bar chart showing mean contraction amplitude before and during the presence of denopamine, and denopamine plus 4-DAMP ($n = 16$, $N = 8$, $***P < .001$, and $****P < .0001$, ANOVA).

The findings of the present study demonstrate that activation of β_1 -ARs reduces the amplitude of cholinergic contractions of ASM and that this effect involves inhibition of M2R-dependent responses. In addition, we found that the inhibitory effects of the β_1 -AR agonist denopamine on contractions of ASM induced by cholinergic stimuli in M2R KO mice were reduced and not enhanced. Therefore, contrary to previous studies, we suggest that M2Rs are functionally involved in cholinergic contractions of ASM and that inhibition of this response contributes to the inhibitory effects of β -AR agonists on the bronchoconstrictor effects of ACh. Consequently, our understanding of the mechanisms responsible for contraction of ASM by ACh and its regulation by β -AR agonists should be reviewed.

Alkawadri et al.²⁰ demonstrated that the amplitude of M3R-dependent contractions of ASM were augmented by activation of M2Rs. M2R-dependent contractions were revealed by EFS at low frequency (2 Hz) and short stimulus intervals (10 s).²⁰ Here, we show that the inhibitory effects of M2R antagonists on these responses, were mimicked by application of the β_1 -AR agonist, denopamine. In addition, denopamine also inhibited 4-DAMP-resistant contractions of ASM evoked by CCh, consistent with the idea that M2R-dependent contractions of ASM are inhibited by β -AR activation. Previous studies suggested that there was

cross-talk between β -AR and M3R signalling pathways in ASM cells,⁷ but our results suggest that there may also be an interaction between β -AR and M2R pathways.

The degree to which inhibition of M2R-dependent responses contributes to the overall inhibitory effect of β -AR agonists on cholinergic contractions of ASM is likely to depend on a combination of the relative contribution of M2R and M3Rs to the cholinergic response and on the relative sensitivity of the M3 and M2R components to β -AR activation. The former is highly dependent on the stimulus parameters and it seems likely from other studies that the M2R component is more prevalent at low ACh concentrations.^{34,35} With regards to the relative sensitivity of the M2 and M3R responses, it was interesting to note that a large 4-DAMP-sensitive contraction remained in the presence of denopamine, suggesting that the M3R-dependent contractions was not fully blocked by β -ARs activation. In contrast, the 4-DAMP-resistant component of the CCh response was abolished by denopamine, suggesting that the entire M2R response was ablated by activation of β -ARs. Similarly, contractions evoked by 20 Hz EFS, known to involve M3, but not M2Rs, displayed a weaker sensitivity to denopamine than those induced by 2 Hz EFS, which contain a larger M2R-dependent component. Therefore, it is plausible that, at least in murine




Figure 5. Comparison of the effects of denopamine on EFS-evoked contractions of ASM taken from WT and M2R null mice. (A) and (B) are representative isometric tension recordings showing the effect of 3 μ M denopamine on EFS-evoked contractions (2 Hz), taken from WT and M2R null mice, respectively. (C) and (D) are summary bar charts showing the effect of denopamine (3 μ M) on mean contraction amplitude in WT ($n = 15$, $N = 9$, and **** $P < .0001$, and two-tailed Student's t -test) and M2R null mice ($n = 9$, $N = 5$, and ** $P < .01$, and two-tailed Student's t -test).

ASM, M2R-dependent signalling pathways have a greater sensitivity to activation of the β -AR/cAMP pathway than the M3R pathways.

The cellular mechanisms underlying M2R-dependent contractions of ASM have not been fully elucidated. However, Semenov et al. reported that M2R signalling in tracheal smooth muscle leads to depolarization and recruitment of voltage-operated Ca^{2+} channels, and that these responses are functionally opposed by activation of large conductance Ca^{2+} -activated K^{+} (BK) channels.³⁴ It has also been shown that activation of β -ARs in ASM leads to BK channel activation.^{36,37} Therefore, it is conceivable that β -AR agonists inhibit M2R-dependent responses by such a mechanism, leading to membrane hyperpolarization and reduced activation of voltage-operated Ca^{2+} channels. However, further investigation would be required to determine if this is the case.

The results of the present study support the concept of functional antagonism between activation of β -AR and M2Rs in the bladder, as reported previously,^{25,38} but not in the airways. It is unclear why this difference exists between both tissues, but it seems likely that a greater functional role for M2Rs in cholinergic contractions of ASM is a factor. If M2R responses are targeted by β -AR agonists to a greater degree than the M3R component, then tissues with more involvement of M2Rs in the cholinergic response would be expected to display a greater sensitivity to β -AR agonists. Consequently, ablation of M2Rs would result in reduced efficacy of β -AR agonists in these tissues. Our results are in agreement with findings from Matsui et al.,³⁸ which showed that the inhibitory effects of isoprenaline on contractions of murine bladder induced by the cholinergic agonist oxotremorine-M were enhanced in M2R KO mice, but were unchanged in tracheal smooth muscle. They reasoned that

part of the isoprenaline response in the airways was independent of cAMP, and was thus unaffected by reduction in cAMP levels following M2R activation. However, this explanation does not account for the large potentiation of β -AR effects induced by M2R antagonists in ASM, such as those reported by Brown et al.,¹⁸ which found that isoproterenol-induced relaxations of rat and human small airways were significantly increased by AFDX-116 and gallamine. Therefore, it appears that the effects of β -AR agonists on cholinergic responses in ASM are enhanced by M2R antagonists^{10,12,18} but are unaffected, or reduced, when M2Rs are genetically ablated.³⁸ It is possible that this difference arises from compensatory changes in gene expression that may occur in M2R KO mice. However, based on the data presented here, this appears to be only a feature of ASM, and not bladder smooth muscle, and further investigation of this issue is required. It should also be noted that some studies suggest that enhanced effects of β -AR agonists in the presence of M2R antagonists, result from reduced levels of cholinergic tone and are not necessarily due to blockade of M2Rs.³⁹ In the present study, it was notable that even though the mean amplitude of EFS-induced contractions was lower in the M2R knock-outs compared to WT preparations, the effects of denopamine were reduced, rather than enhanced. Therefore, the lack of functional antagonism observed in the present study does not appear to be due to a difference in the level of tone.

The therapeutic effects of anticholinergics, such as tiotropium, for asthma and COPD, are primarily thought to arise from blockade of M3Rs.⁴ Indeed, anticholinergics with a higher kinetic selectivity for M3Rs over M2Rs are preferred as they limit unwanted effects arising from inhibition of autoinhibitory M2Rs that would enhance ACh output.^{40,41} However, this logic fails to recognize the possibility that, under specific




Figure 6. Comparison of the effects of the β -AR agonist BRL37344 on EFS-evoked contractions of detrusor strips taken from WT and M2R null mice. (A) and (B) are representative isometric tension recordings showing the effect of 100 nM BRL37344 on EFS-evoked contractions (4 Hz) of detrusor strips taken from WT and M2R null mice, respectively. (C) and (D) are summary bar charts showing the effect of BRL37344 on mean contraction amplitude in WT ($n = 16$, $N = 8$, and **** $P < .0001$, two-tailed Student's paired t-test) and M2R null mice ($n = 16$, $N = 8$, and **** $P < .0001$, and two-tailed Student's paired t-test).

conditions, activation of M2Rs may contribute to the bronchoconstrictor effects of ACh.²⁰ The results of the present study demonstrate that β -AR agonists inhibit the M2R-dependent component of the cholinergic response and it is possible that this may contribute to the therapeutic effects of these agents. If so, this may also contribute to the superior improvements in pulmonary function in patients taking a combination of β -AR agonists and M2R-sparing anticholinergics, than when either are administered alone.^{42,43} It is clear that further experimentation using human samples would be required to discern if this is the case. In addition, it should be noted that such experiments would require use of selective β 2, rather than β 1, adrenoceptor agonists.

In summary, we demonstrate that M2R-dependent contractions of ASM are inhibited by activation of β -ARs and that genetic ablation of M2Rs decreases the efficacy of the β -AR agonist denopamine on cholinergic contractions.

Acknowledgments

The authors are grateful for technical support provided by Billie McIlveen.

Authors' Contributions

T.A. completed experiments on ASM, drafted figures, performed statistical analyses, contributed to study design and data interpretation, and manuscript review. Z.F. and P.Y.W. completed

experiments on bladder, drafted figures, and performed statistical analyses. M.H., F.L., L.Mc.G., and K.T. contributed to study design, data interpretation, and manuscript review. G.S. wrote the paper, drafted figures, contributed to study design and data interpretation, and manuscript review.

Funding

This work was supported by the BREATH project, funded by the European Union Interreg VA Health and Life Science Programme. INT-VA/045. Z.F. was funded by the Irish Research Council (GOIPG/2016/1300) and P.Y.W. is supported by Landscape funding from the HEA and is in receipt of the HEA's Covid19 Relief funding for Researchers scheme.

Conflict of Interest Statement

None declared.

Data Availability

The data underlying this article will be shared on reasonable request to the corresponding author.

References

- Matera MG, Page CP, Calzetta L, Rogliani P, Cazzola M. Pharmacology and therapeutics of bronchodilators revisited. *Pharmacol Rev* 2020;**72**(1):218–252.
- Billington CK, Penn RB, Hall IP. β_2 -agonists. *Handb Exp Pharmacol* 2017;**237**:23–40.
- Gazzola M, Maillhot-Larouche S, Beucher C, Bossé Y. The underlying physiological mechanisms whereby anticholinergics alleviate asthma. *Can J Physiol Pharmacol* 2018;**96**(5):433–441.
- Pera T, Penn RB. Crosstalk between beta-2-adrenoceptor and muscarinic acetylcholine receptors in the airway. *Curr Opin Pharmacol* 2014;**16**:72–81.
- Panettieri Jr RA. Bronchodilators, receptors and cross-talk: together is better?. *Postgrad Med* 2015;**127**(7):771–780.
- Kume H, Nishiyama O, Isoya T, Higashimoto Y, Tohda Y, Noda Y. Involvement of allosteric effect and K_{Ca} channels in crosstalk between β_2 -adrenergic and muscarinic M_2 receptors in airway smooth muscle. *Int J Mol Sci* 2018;**19**(7):1999.
- Cazzola M, Calzetta L, Matera MG. Long-acting muscarinic antagonists and small airways in asthma: which link?. *Allergy* 2021;**76**(7):1990–2001.
- Lin CH, Yang CM, Chen CM, Ko FN, Teng CM. Pharmacological characteristics of BDTI, a new isoquinoline-derived beta 2-adrenoceptor agonist, in canine trachea and rat heart. *Pharmacology* 1996;**53**(1):19–27.
- Calzetta L, Rogliani P, Page C, Rinaldi B, Cazzola M, Matera MG. Pharmacological characterization of the interaction between tiotropium bromide and olodaterol on human bronchi and small airways. *Pulm Pharmacol Ther* 2019;**56**:39–50.
- Sarria B, Naline E, Zhang Y,, et al. Muscarinic M_2 receptors in acetylcholine-isoproterenol functional antagonism in human isolated bronchus. *Am J Physiol Lung Cell Mol Physiol* 2002; **283**(5):L1125–L1132.
- Maarsingh H, Oldenburger A, Han B,, et al. Effects of (a combination of) the beta $_2$ -adrenoceptor agonist indacaterol and the muscarinic receptor antagonist glycopyrrolate on intrapulmonary airway constriction. *Cells* 2021;**10**(5):1237.
- Fernandes LB, Fryer AD, Hirshman CA. M_2 muscarinic receptors inhibit isoproterenol-induced relaxation of canine airway smooth muscle. *J Pharmacol Exp Ther* 1992;**262**(1):119–26.
- Haddad EB, Mak JC, Hislop A, Haworth SG, Barnes PJ. Characterization of muscarinic receptor subtypes in pig airways: radioligand binding and northern blotting studies. *Am J Physiol* 1994;**266**(6 Pt 1):L642–8.
- Roffel AF, Meurs H, Elzinga C R, Zaagsma J. Characterization of the muscarinic receptor subtype involved in phosphoinositide metabolism in bovine tracheal smooth muscle. *Br J Pharmacol* 1990;**99**(2):293–296.
- Schramm CM, Arjona N C, Grunstein M M. Role of muscarinic M_2 receptors in regulating beta-adrenergic responsiveness in maturing rabbit airway smooth muscle. *Am J Physiol* 1995;**269**(6 Pt 1):L783–90.
- Fayon M, Dumas De La Roque E, Berger P,, et al. Increased relaxation of immature airways to beta2-adrenoceptor agonists is related to attenuated expression of postjunctional smooth muscle muscarinic M_2 receptors. *J Appl Physiol* 2005. **98**(4):1526–1533.
- Watson N, Eglen RM. Effects of muscarinic M_2 and M_3 receptor stimulation and antagonism on responses to isoprenaline of guinea-pig trachea in vitro. *Br J Pharmacol* 1994;**112**(1):179–187.
- Brown SM, Koarai A, Sturton RG, Nicholson AG, Barnes PJ, Donnelly LE. A role for $M(2)$ and $M(3)$ muscarinic receptors in the contraction of rat and human small airways. *Eur J Pharmacol* 2013;**702**(1–3):109–115.
- Struckmann N, Schwering S, Wiegand S,, et al. Role of muscarinic receptor subtypes in the constriction of peripheral airways: studies on receptor-deficient mice. *Mol Pharmacol* 2003;**64**(6):1444–1451.
- Alkawadri T, McGarvey LP, Mullins ND, Hollywood MA, Thornbury KD, Sergeant GP. Contribution of postjunctional M_2 muscarinic receptors to cholinergic nerve-mediated contractions of murine airway smooth muscle. *Function* 2021;**3**(1):zqab053.
- Ward SM. Postjunctional M_2 muscarinic receptors augment neurally mediated cholinergic contractions of murine airway smooth muscle. *Function* 2022;**3**(1):zqab067.
- Bradley E, Large RJ, Bihun VV,, et al. Inhibitory effects of openers of large-conductance Ca^{2+} -activated K^+ channels on agonist-induced phasic contractions in rabbit and mouse bronchial smooth muscle. *Am J Physiol Cell Physiol* 2018;**315**(6):C818–C829.
- Henry PJ, Goldie RG. Beta 1-adrenoceptors mediate smooth muscle relaxation in mouse isolated trachea. *Br J Pharmacol* 1990;**99**(1):131–135.
- Birrell MA, Bonvini SJ, Wortley MA,, et al. The role of adenylyl cyclase isoform 6 in β -adrenoceptor signalling in murine airways. *Br J Pharmacol* 2015;**172**(1):131–141.
- Ehlert FJ, Ahn S, Pak KJ,, et al. Neuronally released acetylcholine acts on the M_2 muscarinic receptor to oppose the relaxant effect of isoproterenol on cholinergic contractions in mouse urinary bladder. *J Pharmacol Exp Ther* 2007;**322**(2):631–637.
- Agustí A, Hogg JC. Update on the pathogenesis of chronic obstructive pulmonary disease. *N Engl J Med* 2019;**381**(13):1248–1256.
- Kume H. Role of airway smooth muscle in inflammation related to asthma and COPD. *Adv Exp Med Biol* 2021;**1303**:139–172.
- Gross NJ, Skorodin MS. Role of the parasympathetic system in airway obstruction due to emphysema. *N Engl J Med* 1984;**311**(7):421–425.
- Barnes PJ. The role of anticholinergics in chronic obstructive pulmonary disease. *Am J Med* 2004;**117**(12A):24S–32S.
- Gosens R, Zaagsma J, Meurs H, Halayko A. Muscarinic receptor signaling in the pathophysiology of asthma and COPD. *Respir Res* 2006;**7**(1):73.
- Meurs H, Oenema TA, Kistemaker LE, Gosens R. A new perspective on muscarinic receptor antagonism in obstructive airways diseases. *Curr Opin Pharmacol* 2013;**13**(3): 316–323.
- Fryer AD, Jacoby DB. Muscarinic receptors and control of airway smooth muscle. *Am J Respir Crit Care Med* 1998;**158**(5 Pt 3):S154–S160. Supplement.2.
- Belmonte KE. Cholinergic pathways in the lungs and anticholinergic therapy for chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 2005;**2**(4):297–304
- Semenov I, Wang B, Herlihy JT, Brenner R. BK channel β_1 subunits regulate airway contraction secondary to M_2 muscarinic acetylcholine receptor mediated depolarization. *J Physiol* 2011;**589**(7 Pt 7):1803–1817.
- Unno T, Komori S, Ohashi H. Inhibitory effect of muscarinic receptor activation on Ca^{2+} channel current in smooth muscle cells of guinea-pig ileum. *J Physiol* 1995;**484**(3):567–581. (Pt 3)

36. Kume H, Takai A, Tokuno H, Tomita T. Regulation of Ca²⁺-dependent K⁺-channel activity in tracheal myocytes by phosphorylation. *Nature* 1989;**341**(6238):152–154.
37. Savaria D, Lanoue C, Cadieux A, Rousseau E. Large conducting potassium channel reconstituted from airway smooth muscle. *J Physiol* 1992;**262**(3 Pt 1):L327–36.
38. Matsui M, Griffin MT, Shehnaz D, Taketo MM, Ehlert FJ. Increased relaxant action of forskolin and isoproterenol against muscarinic agonist-induced contractions in smooth muscle from M2 receptor knockout mice. *J Pharmacol Exp Ther* 2003;**305**(1):106–113.
39. Roffel AF, Meurs H, Elzinga CR, Zaagsma J. No evidence for a role of muscarinic M2 receptors in functional antagonism in bovine trachea. *Br J Pharmacol* 1995;**115**(4):665–671.
40. Spina D. Current and novel bronchodilators in respiratory disease. *Curr Opin Pulm Med* 2014;**20**(1):73–86.
41. Ora J, Coppola A, Cazzola M, Calzetta L, Rogliani P. Long-acting muscarinic antagonists under investigation to treat chronic obstructive pulmonary disease. *J Exp Pharmacol* 2020;**12**:559–574.
42. Petite SE. Role of long-acting muscarinic antagonist/long-acting β 2-agonist therapy in chronic obstructive pulmonary disease. *Ann Pharmacother* 2017;**51**(8):696–705.
43. Calzetta L, Rogliani P, Ora J, Puxeddu E, Cazzola M, Matera MG. LABA/LAMA combination in COPD: a meta-analysis on the duration of treatment. *Eur Respir Rev* 2017;**26**(143):160043.