

UC Irvine

UC Irvine Previously Published Works

Title

Draft Genome Sequence of Salmonella enterica subsp. enterica Serovar Blockley Strain CRJJGF_00147 (Phylum Gammaproteobacteria)

Permalink

<https://escholarship.org/uc/item/4cs37252>

Journal

Microbiology Resource Announcements, 4(5)

ISSN

2576-098X

Authors

Gupta, Sushim K
McMillan, Elizabeth A
Jackson, Charlene R
et al.

Publication Date

2016-10-27

DOI

10.1128/genomea.00954-16

Peer reviewed

Draft Genome Sequence of *Salmonella enterica* subsp. *enterica* Serovar Blockley Strain CRJJGF_00147 (Phylum *Gammaproteobacteria*)

Sushim K. Gupta,^a Elizabeth A. McMillan,^{a,b} Charlene R. Jackson,^a Prerak T. Desai,^c Steffen Porwollik,^c Michael McClelland,^c Lari M. Hiott,^a Shaheen B. Humayoun,^a Jonathan G. Frye^a

Bacterial Epidemiology and Antimicrobial Resistance Research Unit, USDA-ARS, Athens, Georgia, USA^a; Department of Microbiology, University of Georgia, Athens, Georgia, USA^b; Department of Microbiology and Molecular Genetics, University of California Irvine, Irvine, California, USA^c

Here, we report a 4.72-Mbp draft genome sequence of *Salmonella enterica* subsp. *enterica* serovar Blockley strain CRJJGF_00147, isolated from chicken rinse in 2009.

Received 14 July 2016 Accepted 19 July 2016 Published 8 September 2016

Citation Gupta SK, McMillan EA, Jackson CR, Desai PT, Porwollik S, McClelland M, Hiott LM, Humayoun SB, Frye JG. 2016. Draft genome sequence of *Salmonella enterica* subsp. *enterica* serovar Blockley strain CRJJGF_00147 (phylum *Gammaproteobacteria*). *Genome Announc* 4(5):e00954-16. doi:10.1128/genomeA.00954-16.

Copyright © 2016 Gupta et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jonathan G. Frye, Jonathan.Frye@ars.usda.gov.

Salmonella enterica subsp. *enterica* serovar Blockley was first isolated from the stool from a 60-year-old patient in 1955 (1). An S. Blockley outbreak was reported for the first time from frozen unpasteurized egg yolks used in the preparation of ice cream (2). S. Blockley is a rarely isolated serotype in most countries; however, S. Blockley has been isolated from agricultural and clinical samples periodically worldwide (3–7). Here, we announce the draft genome sequence of S. Blockley, isolated from chicken rinse in 2009.

Standard microbiology techniques were applied to isolate *Salmonella* strains from food animals. The isolates were serotyped using SMART typing (8), and reads were used to determine antigenic formula to predict the serotype using SeqSero (9), which predicted the antigenic formula of 8:k:1,5, designated Blockley. Using pulsed-field gel electrophoresis (PFGE), as described by PulseNet (10), the isolate was assigned PFGE pattern JBGX01.0001. Susceptibility testing for the strain was performed using broth microdilution plates for the Sensititre semiautomated antimicrobial susceptibility system (Trek Diagnostic Systems, Inc., Westlake, OH). The results were interpreted according to Clinical and Laboratory Standards Institute (CLSI) guidelines (11).

Genomic DNA was isolated from an overnight-grown culture using the GenElute bacterial genomic DNA kit (Sigma-Aldrich, St. Louis, MO). DNA libraries were constructed using Nextera-XT DNA preparation kit, and paired-end sequencing was performed on the Illumina HiSeq 2500 (Illumina, Inc., San Diego, CA) using a 500-cycle MiSeq reagent kit. A total of 2,599,790 reads were generated. Reads were *de novo* assembled using Velvet (12), which assembled to 154 contigs ≥ 200 bp. The combined length of contigs was 4,722,084 bases, with a G+C content of 52.11% and N_{50} value of 76.5 kb. The contigs were ordered with Mauve (13) using

Salmonella LT2 as a reference genome, and coding sequences were predicted with Prodigal (14). A total of 4,399 coding sequences (≥ 50 amino acids) were predicted within the genome. Signal peptide, clustered regularly interspaced short palindromic repeat (CRISPR) elements, and prophages were predicted using SignalP (15), CRISPRFinder (16), and PHAST (17), respectively. We identified signal peptides in 429 coding sequences (CDSs), two CRISPR loci, and two intact phages, Gifsy2 (accession no. NC_010393) and Salmon_SEN1 (accession no. NC_029003), in the analyzed contigs. The isolate was susceptible to all tested antibiotics, although a cryptic *aac6-Iy* gene was identified through ARG-ANNOT (18). The information generated from genome sequencing will improve our understanding of the role of genomic regions and how they may lead to *Salmonella* pathogenicity.

Accession number(s). The genome sequence of *Salmonella enterica* subsp. *enterica* serovar Blockley strain CRJJGF_00147 has been deposited in the GenBank database (NCBI) under the accession no. [JQXY00000000](https://doi.org/10.1101/090000). This paper describes the first version of the genome.

ACKNOWLEDGMENTS

J.G.F. and C.R.J. were supported by USDA project numbers 6040-32000-006-00 and 6040-32000-009-00, and a grant from the American Meat Institute Foundation. M.M. was supported in part by NIH grants R01AI052237, AI039557, AI052237, AI073971, AI075093, AI077645, and AI083646, USDA grants 2009-03579 and 2011-67017-30127, the Binational Agricultural Research and Development Fund, and a grant from the Center for Produce Safety.

We thank Calvin Williams for all the technical support.

The mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

FUNDING INFORMATION

This work, including the efforts of Charlene R. Jackson and Jonathan Gray Frye, was funded by American Meat Institute Foundation. This work, including the efforts of Michael McClelland, was funded by HHS | National Institutes of Health (NIH) (R01AI052237, AI039577, AI053327, AI073971, and AI083646). This work, including the efforts of Charlene R. Jackson and Jonathan Gray Frye, was funded by U.S. Department of Agriculture (USDA) (6040-32000-006-00 and 6040-32000-009-00). This work, including the efforts of Michael McClelland, was funded by U.S. Department of Agriculture (USDA) (2009-03579 and 2011-67107-30127). This work, including the efforts of Michael McClelland, was funded by United States - Israel Binational Agricultural Research and Development Fund (BARD). This work, including the efforts of Michael McClelland, was funded by Center for Produce Safety (CPS).

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- Friedman S, Wassermann MM, Saphra I. 1955. A new *Salmonella* type: *Salmonella* Blockley. *J Bacteriol* 70:354–355.
- Morse LJ, Rubenstein AD. 1967. A foodborne institutional outbreak of enteritis due to *Salmonella* Blockley. *JAMA* 202:939–940.
- Limawongpranee S, Hayashidani H, Okatani AT, Hirota C, Kaneko K, Ogawa M. 1999. Contamination of *Salmonella* Blockley in the environment of a poultry farm. *Avian Dis* 43:302–309. <http://dx.doi.org/10.2307/1592621>.
- Fell G, Hamouda O, Lindner R, Rehmet S, Liesegang A, Prager R, Gericke B, Petersen L. 2000. An outbreak of *Salmonella* Blockley infections following smoked eel consumption in Germany. *Epidemiol Infect* 125:9–12. <http://dx.doi.org/10.1017/S0950268899004069>.
- Thai TH, Lan NT, Hirai T, Yamaguchi R. 2012. Antimicrobial resistance in *Salmonella* serovars isolated from meat shops at the markets in North Vietnam. *Foodborne Pathog Dis* 9:986–991. <http://dx.doi.org/10.1089/fpd.2011.1121>.
- Gonose T, Smith AM, Keddy KH, Sooka A, Howell V, Jacobs CA, Haffeejee S, Govender P. 2012. Human infections due to *Salmonella* Blockley, a rare serotype in South Africa: a case report. *BMC Res Notes* 5:562. <http://dx.doi.org/10.1186/1756-0500-5-562>.
- Molla B, Berhanu A, Muckle A, Cole L, Wilkie E, Kleer J, Hildebrandt G. 2006. Multidrug resistance and distribution of *Salmonella* serovars in slaughtered pigs. *J Vet Med B Infect Dis Vet Public Health* 53:28–33. <http://dx.doi.org/10.1111/j.1439-0450.2006.00900.x>.
- Leader BT, Frye JG, Hu J, Fedorka-Cray PJ, Boyle DS. 2009. High-throughput molecular determination of *Salmonella enterica* serovars by use of multiplex PCR and capillary electrophoresis analysis. *J Clin Microbiol* 47:1290–1299. <http://dx.doi.org/10.1128/JCM.02095-08>.
- Zhang S, Yin Y, Jones MB, Zhang Z, Deatherage Kaiser BL, Dinsmore BA, Fitzgerald C, Fields PI, Deng X. 2015. *Salmonella* serotype determination utilizing high-throughput genome sequencing data. *J Clin Microbiol* 53:1685–1692. <http://dx.doi.org/10.1128/JCM.00323-15>.
- Gerner-Smidt P, Hise K, Kincaid J, Hunter S, Rolando S, Hyytiä-Trees E, Ribot EM, Swaminathan B, Pulsenet Taskforce. 2006. PulseNet USA: a five-year update. *Foodborne Pathog Dis* 3:9–19. <http://dx.doi.org/10.1089/fpd.2006.3.9>.
- CLSI. 2015. Performance standards for antimicrobial susceptibility testing: 25th informational supplement. CLSI document M100–S25. Clinical and Laboratory Standards Institute, Wayne, PA.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
- Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. 2009. Reordering contigs of draft genomes using the Mauve aligner. *Bioinformatics* 25:2071–2073. <http://dx.doi.org/10.1093/bioinformatics/btp356>.
- Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <http://dx.doi.org/10.1186/1471-2105-11-119>.
- Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* 8:785–786. <http://dx.doi.org/10.1038/nmeth.1701>.
- Grissa I, Vergnaud G, POURCEL C. 2007. CRISPRFinder: a Web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35:W52–W57. <http://dx.doi.org/10.1093/nar/gkm360>.
- Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. *Nucleic Acids Res* 39:W347–W352. <http://dx.doi.org/10.1093/nar/gkr485>.
- Gupta SK, Padmanabhan BR, Diene SM, Lopez-Rojas R, Kempf M, Landraud L, Rolain JM. 2014. ARG-ANNOT, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes. *Antimicrob Agents Chemother* 58:212–220. <http://dx.doi.org/10.1128/AAC.01310-13>.