

Lawrence Berkeley National Laboratory

Recent Work

Title

CHEMICAL EVIDENCE FOR MAIN-CHAIN SCISSION AS A MAJOR DECOMPOSITION MODE IN THE RADIOLYSIS OF SOLID PEPTIDES

Permalink

<https://escholarship.org/uc/item/4h51d739>

Authors

Garrison, Warren M.

Jayko, Michael E.

Weeks, Boyd M.

et al.

Publication Date

1966-12-01

University of California
Ernest O. Lawrence
Radiation Laboratory

CHEMICAL EVIDENCE FOR MAIN-CHAIN SCISSION AS A MAJOR DECOMPOSITION
MODE IN THE RADIOLYSIS OF SOLID PEPTIDES

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Journal of Physical Chemistry

UCRL-17307
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

CHEMICAL EVIDENCE FOR MAIN-CHAIN SCISSION AS A MAJOR DECOMPOSITION
MODE IN THE RADIOLYSIS OF SOLID PEPTIDES

Warren M. Garrison, Michael E. Jayko, Boyd M. Weeks,
Harvey A. Sokol, and Winifred Bennett-Corniea

December 1966

CHEMICAL EVIDENCE FOR MAIN-CHAIN SCISSION AS A MAJOR DECOMPOSITION
MODE IN THE RADIOLYSIS OF SOLID PEPTIDES

Warren M. Garrison, Michael E. Jayko, Boyd M. Weeks,
Harvey A. Sokol, and Winifred Bennett-Corniea

Lawrence Radiation Laboratory
University of California
Berkeley, California


December 1966

We find that a major chemical effect of ionizing radiation on peptide derivatives of the aliphatic α -amino acids is the formation of labile "amide-like" products which are readily degraded to yield ammonia on mild hydrolysis. The N-acetyl derivatives of alanine, valine and leucine, for example, give $G(\text{NH}_3)$ values of 3.8, 3.0 and 2.7 (molecules/100 ev) respectively on irradiation with Co^{60} γ -rays in vacuo; the irradiated solids were dissolved in water in the absence of oxygen and hydrolyzed for 90 minutes in 2 N HCl to effect a quantitative liberation of ammonia which was then assayed after the method of Conway.¹


A detailed study of other reaction products from N-acetylalanine has been made to obtain evidence as to the mechanism of the radiation-induced degradation. Organic acids were assayed by gas-phase chromatography² and by partition chromatography.^{3,4} In certain runs, the acetylalanine was "spiked" with $\text{CH}_3\text{CONHCH}(\text{CH}_2\text{C}^{14}\text{OOH})$ to facilitate product identification; the radiometric techniques have been described.⁴ Carbonyl products were determined as the 2, 4-dinitrophenylhydrazone derivatives.^{4,5} A colorimetric method⁶ was used to set a limit on the yield of lactic acid. Gaseous products were pumped-off following

complete dissolution of an irradiated sample in degassed water on the vacuum line. Gases were analyzed by mass spectrometry. A Varian EPR spectrometer, model E-3, was used to measure the production of spin centers in the irradiated solids.


The evidence is that the observed radiolytic degradation of acetylalanine cannot be represented simply in terms of the rearrangement




since we find $G(\text{acrylic acid}) < 0.2$. The possibility that the N, O shift




occurs in high yield and leads to ammonia formation through subsequent hydrolysis of the labile imino ester



is negated by the fact that $G(\text{lactic acid}) \leq 0.2$. Similarly, we conclude that the production of dehydropeptide via




is relatively unimportant in view of the fact that we find $G(\text{pyruvic acid}) \approx 0.4$; the dehydropeptides are hydrolyzed quantitatively to yield carbonyl and ammonia





under the conditions of hydrolysis employed here.⁷ Additional evidence that the dehydrogenation reaction 4 occurs in low yield is given by the gas-yield data which show $G(\text{H}_2) = 0.4$.⁸

We do find that propionic acid is produced as a major product, $G(\text{propionic acid}) \approx 1.2$. It is clear that main-chain scission occurs and the simplest formulation in accord with present observations is given by:




followed by the abstraction reactions




to give amide, fatty acid and the long-lived α -carbon radicals which have been observed in irradiated peptides at room temperatures through EPR spectroscopy.^{9,10} On dissolution in water (oxygen-free), the α -carbon radicals, $\text{RCONHC}(\text{CH}_3)\text{R}$, undergo dimerization to yield the α, α -diaminosuccinic acid derivative.¹¹ The formulation of reactions 6, 7, 8 is intended only to convey the nature of the overall reaction stoichiometry.¹² Ionic or "hot" radical intermediates presumably are involved since cage effects in the condensed medium would lead to preferential radical-recombination.

We estimate from EPR measurements that the yield of the long-lived α -carbon radicals is roughly $G \sim 3$. This value and the propionic acid yield value of $G=1.2$ are somewhat lower than would be predicted from the reaction sequence 6, 7, 8 on the basis of $G(\text{amide}) \sim 3.8$. The apparent discrepancy may be understood if a fraction of the $\text{CH}(\text{CH}_3)\text{R}$ radicals are removed through radical-radical interactions in the spur-regions of high radical concentration. That the "amide" type of radiolytic degradation of the peptide chain is not confined to the N-acylamino acid configuration is shown by the fact that the G values for ammonia and propionic acid from poly-DL-alanine are almost identical with those obtained with N-acetylalanine.


It has been suggested that the long-lived α -carbon radicals are produced in peptide radiolysis through side-chain cleavage,^{13,14} e.g.,



followed by



where H formed in reaction 9 may have excess kinetic energy. However the low $G(H_2)$ values observed in the γ -radiolysis of simple peptides, polypeptides and protein,⁸ does not support the concept that reaction 10 represents a major source of α -carbon radicals. On the other hand, reaction 9 followed by



(or the equivalent heterolytic steps involving e^- and H^+) would satisfy the requirement of main-chain scission. Note that the stoichiometry of reactions 9, 11 followed by 8 is equivalent to that given by the reaction sequence 6, 7, 8.

Detailed reports of these studies will appear in forthcoming publications. Acknowledgement is due to ^{now} Miss Sibyl A. Cole for assistance in some of the measurements.

This work was performed under the auspices of the U. S. Atomic Energy Commission.

References and Footnotes

- (1) E. J. Conway, Microdiffusion Analysis (Crosby Lockwood and Son, Ltd., London, 1962) p. 90.
- (2) E. M. Emery and W. E. Koerner, Anal. Chem. 33, 146 (1961).
- (3) C. S. Marvel and R. Rands, J. Am. Chem. Soc., 72, 2642 (1950).
- (4) B. M. Weeks, S. A. Cole and W. M. Garrison, J. Phys. Chem. 69, 4131 (1965).
- (5) G. R. A. Johnson and G. Scholes, Ind. Eng. Chem., Anal. Ed. 79, 217 (1954).
- (6) F. Feigl, Spot Tests In Organic Analysis (Elsevier Publishing Co., New York, 1956) p. 346.
- (7) J. P. Greenstein and M. Winitz, Chemistry of Amino Acids (John Wiley and Sons, Inc., New York, New York, 1961) p. 843.
- (8) Hydrogen yields from compounds containing the peptide bond are uniformly low. We have found initial $G(H_2)$ values of 0.45, 0.85 and 0.22 for poly-alanine, nylon, and gelatin respectively. See, M. E. Jayko and W. M. Garrison, Abstract 465, Abstract of Papers, Third International Congress of Radiation Research, Cortina, Italy, June 26 to July 2, 1966.
- (9) R. C. Drew and W. Gordy, Radiation Res. 18, 552 (1963).
- (10) N. C. Box, H. G. Freund and K. Lilga, in "Free Radicals In Biological Systems" ed. by M. Blois, et al., (Academic Press, New York, 1961).
- (11) W. M. Garrison and B. M. Weeks, Radiation Res. 17, 341 (1962).
- (12) The propionic acid yield from acetylalanine is directly proportional to dosage up to 10^{21} ev/gm and the dose-yield plot extrapolates through the origin. We conclude from this that propionic acid is not produced through the reduction by e^- or H of acrylic acid formed in reactions akin to 1. However we cannot rule out the possibility that acrylic acid "steady-states"

at a concentration level below that detectable by our present analytical techniques.

- (13) R. Braams, Nature 200, 152 (1966).
- (14) P. Riesz, F. N. White, Jr., and H. Kon, J. Am. Chem. Soc. 88, 872 (1966).

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

[Faint, illegible text covering the majority of the page]

