

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

REGGE POLES AND THE PHASE OF THE FORWARD SCATTERING AMPLITUDE

### Permalink

<https://escholarship.org/uc/item/4jb351qv>

### Authors

Phillips, Roger J.N.  
Rarita, William.

### Publication Date

1965-02-02

University of California  
Ernest O. Lawrence  
Radiation Laboratory

REGGE POLES AND THE PHASE  
OF THE FORWARD SCATTERING AMPLITUDE

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

Berkeley, California

UNIVERSITY OF CALIFORNIA  
Lawrence Radiation Laboratory  
Berkeley, California  
AEC Contract No. W-7405-eng-48

REGGE POLES AND THE PHASE  
OF THE FORWARD SCATTERING AMPLITUDE

Roger J. N. Phillips and William Rarita

February 2, 1965

## REGGE POLES AND THE PHASE OF THE FORWARD SCATTERING AMPLITUDE\*

Roger J. N. Phillips<sup>†</sup>Lawrence Radiation Laboratory  
University of California  
Berkeley, California

and

William Rarita

752 Grizzly Peak Boulevard  
Berkeley, California

February 2, 1965

There is currently great interest in the phase of forward elastic scattering amplitudes at high energy. In a simple diffraction picture, one expects a purely imaginary amplitude. However, recent Coulomb interference measurements<sup>1-6</sup> in pp and np scattering have shown that the real part is substantial, of order 20 to 30%, up to 26 GeV/c. Various dispersion relation calculations<sup>7-9</sup> have confirmed that something of this order may be expected, but they do not explain what mechanism produces it.

The present note points out that simple Regge pole models predict a substantial real part at the energies in question, in surprisingly good agreement with experiment. We employ the latest  $\pi N$ ,  $NN$ , and  $\bar{N}N$  total cross section data<sup>5,10</sup> from 6 to 26 GeV/c, to fix the parameters of the forward scattering amplitudes, taking three Regge poles for  $\pi N$  and four for  $NN$ . The ratio of the real to the imaginary part of the forward scattering amplitude is illustrated and compared with experiment in Figs. 1 and 2.

Let us define and normalize the spin-averaged forward elastic amplitude  $A(0)$  for each process, such that the optical theorem reads

$$\sigma_T = \text{Im } A(0) \quad (1)$$

where  $\sigma_T$  is the corresponding total cross section. Then in a high-energy approximation each Regge pole contributes to  $A(0)$  a term of the form<sup>11</sup>

$$A_i(0) = B_i \left\{ \left[ 1 \pm \exp(-i\pi\alpha_i) \right] / \sin \pi\alpha_i \right\} (E/E_0)^{\alpha_i-1} \quad (2)$$

Here  $i$  labels the Regge pole,  $\alpha_i$  is its trajectory at squared momentum transfer  $t = 0$ ,  $B_i$  is a real coefficient measured in millibarns,  $E$  is the total laboratory-system energy of the bombarding particle, and  $E_0$  is an arbitrary scale parameter which we choose to be 1 GeV. Note that the phase of  $A_i(0)$  is determined by  $\alpha_i$ , through the "signature factor" (in braces), and is therefore directly related to the energy dependence.

For  $\pi N$  scattering at least three Regge poles are needed. The Pomeranchuk pole  $P$  gives the asymptotic limit; a second vacuum pole  $P'$  and the  $\rho$  pole give the differences of the  $\pi^{\pm}p$  amplitudes from the asymptotic limit and from each other. The signature  $\pm$  in Eqs. (2) is  $+$  and  $P$  and  $P'$ ,  $-$  for  $\rho$ . Let us take the coefficients  $B_i$  above to refer to  $\pi^{\mp}p$  elastic scattering (for  $\pi^{\pm}p$ ,  $B_i$  changes sign). Since the fit to data is not very sensitive to the precise values of the  $\alpha_i$ , we fix them at suitable values  $\alpha_P = 1$ ,  $\alpha_{P'} = 0.5$ ,  $\alpha_\rho = 0.6$ . Then by a least-squares fit to the total cross sections<sup>10</sup> the  $B_i$  are determined:  $B_P = 19.9$  mb,  $B_{P'} = 18.1$  mb, and  $B_\rho = 2.4$  mb.

The predicted ratios  $\text{Re } A(0)/\text{Im } A(0)$  for  $\pi^{\pm}p$  scattering are shown in Fig. 1. They agree with the experimental determination in sign, in magnitude, and in giving a larger value for  $\pi^{\pm}p$  than for  $\pi^{\mp}p$ ; however, the experimental uncertainties are rather large. The dispersion calculation

of reference 9 agrees closely with our  $\pi^- p$  curve but gives a  $\pi^+ p$  curve displaced upwards.

For NN and  $\bar{N}N$  scattering, at least two more poles are usually invoked. One is the negative-signature  $\omega$  pole (which is supposed to include any contribution from the  $\phi$  pole, lying near, with the same quantum numbers). The other is the R pole, proposed by Pignotti.<sup>12</sup> However, the experimental uncertainties in the data we use are such that the R contribution that is determined is not significantly different from zero, and we ignore this term. Let us take the coefficients  $B_i$  in Eq. (2) to refer to  $\bar{p}p$  elastic scattering; for  $\bar{n}p$  the  $\rho$  term changes sign; for  $pp$  the  $\omega$  and  $\rho$  terms change sign; for  $np$  the  $\omega$  term changes sign. We fix  $\alpha_\omega = 0.5$ , with  $\alpha_p$ ,  $\alpha_{p'}$ , and  $\alpha_\rho$  as before; then by a least-squares fit to the total cross sections<sup>5,10</sup> we determine  $B_p = 36.2$  mb,  $B_{p'} = 33.8$  mb,  $B_\rho = 1.0$  mb, and  $B_\omega = 21.0$  mb.

The predicted ratios  $\text{Re } A(0)/\text{Im } A(0)$  for  $pp$  and  $\bar{p}p$  scattering are shown in Fig. 2. The experimental points refer only to  $pp$ . Below 10 GeV/c there is a marked divergence between prediction and experiment; the former becomes steadily more negative while the latter (in a region not illustrated) finally becomes positive below 1.5 GeV/c. Above 10 GeV/c, however, the agreement with experiment is surprisingly good. There have been several dispersion calculations which agree roughly with one another and with experiment. Söding's calculation,<sup>7</sup> for example, gives roughly 70% of our  $pp$  values (above 10 GeV/c); it also agrees rather closely with our  $\bar{p}p$  values.

Regge pole models are designed for high energies. As the energy is lowered, the various correction terms play more and more important roles.

We may expect the real part of the forward scattering amplitude to be especially sensitive to these corrections, since it is in a sense a correction term itself--coming wholly from the secondary trajectories. Thus a divergence of the prediction from experiment is to be expected at lower energies.

The parameters of our Regge poles have been fixed by total cross sections only. A complete Regge pole model should also fit elastic angular distributions; this seems to present no serious difficulty,<sup>13,14</sup> but the best fit to this wider range of data generally gives slightly different parameters. However, in the cases we have studied<sup>14</sup> the change is small; the curves in Figs. 1 and 2 should be little altered by a more complete fit to data.

#### ACKNOWLEDGMENT

We thank Professor Geoffrey F. Chew for useful discussions and Dr. David L. Judd for the hospitality of the Theoretical Physics Group at the Lawrence Radiation Laboratory, where this work was done.

FOOTNOTES AND REFERENCES

- \* This work was done under the auspices of the U.S. Atomic Energy Commission.
- † Permanent Address: A. E. R. E., Harwell, Berks., England.
1. S. J. Lindenbaum, rapporteur report to the 1964 Conference on High Energy Physics at Dubna, U.S.S.R.
  2. E. Lohrmann, H. Meyer, and H. Winzeler, *Physics Letters* 13, 78 (1964).
  3. L. Kirillova, L. Khristov, V. Nikitin, M. Shafranov, L. Strunov, V. Sviridov, Z. Korbel, L. Rob, P. Markov, Kh. Tchernev, T. Todorov, and Z. Zlateva, *Phys. Letters* 13, 93 (1964).
  4. K. J. Foley, R. S. Gilmore, R. S. Jones, S. J. Lindenbaum, W. A. Love, S. Ozaki, E. H. Willen, R. Yamada, and L. C. L. Yuan, *Phys. Rev. Letters* 14, 74 (1965).
  5. G. Bellettini, G. Cocconi, A. N. Diddens, E. Lillethun, J. Pahl, J. P. Scanlon, J. Walters, A. M. Wetherell, and P. Zanella, *Phys. Letters* 14, 164 (1965).
  6. A. E. Taylor, A. Ashmore, W. S. Chapman, D. F. Falla, W. H. Range, D. B. Scott, A. Astbury, F. Capocci, and T. G. Walker, *Phys. Letters* 14, 54 (1965).
  7. P. Soding, *Phys. Letters* 8, 285 (1964).
  8. I. I. Levintov and G. M. Adelson-Velsky, *Phys. Letters* 13, 185 (1964).
  9. V. S. Barashenkov and V. I. Dedyu, Dubna preprint R-1598 (1964), submitted to *Nucl. Phys.*
  10. W. Galbraith, E. W. Jenkins, T. F. Kycia, B. A. Leontic, R. H. Phillips, A. L. Reed, and R. Rubinstein, report presented to 1964 Conference on High Energy Physics at Dubna, U.S.S.R.

11. See, for instance, S. Frautschi, M. Gell-Mann, and F. Zachariasen, Phys. Rev. 126, 2204 (1962).
12. A. Pignotti, Phys. Rev. 134, B630 (1964).
13. T. O. Binford and B. R. Desai, High Energy Elastic Scattering at Low Momentum Transfers, University of Wisconsin preprint, 1964.
14. R. J. N. Phillips and W. Rarita, unpublished calculations.

FIGURE CAPTIONS

Fig. 1. The ratio of the real to the imaginary part of the forward amplitude for  $\pi^+p$  and  $\pi^-p$  scattering. The curves are Regge pole predictions. The data are from reference 1: the inner error flags are statistical; the outer ones are estimated limits of systematic error.

Fig. 2. The ratio of the real to the imaginary part of the forward spin-averaged amplitude, for  $pp$  and  $\bar{p}p$  scattering. The curves are Regge pole predictions. The data refer only to  $pp$  scattering. Where double error flags are shown, the inner ones are statistical and the outer ones are estimated limits of systematic error.




Fig. 1




Fig. 2

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

2004  
2003  
2002