

UCSF

UC San Francisco Previously Published Works

Title

Physical intimate partner violence and prenatal oral health experiences in the United States.

Permalink

<https://escholarship.org/uc/item/4k04h0dn>

Journal

BMC Oral Health, 23(1)

Authors

Testa, Alexander
Lee, Jacqueline
Jackson, Dylan
et al.

Publication Date

2023-10-12

DOI

10.1186/s12903-023-03491-0

Peer reviewed

RESEARCH

Open Access



Physical intimate partner violence and prenatal oral health experiences in the United States

Alexander Testa¹, Jacqueline G. Lee², Dylan B. Jackson³, Rahma Mungia⁴, Kyle T. Ganson⁵ and Jason M. Nagata^{6*}

Abstract

Background Intimate partner violence (IPV) is a significant public health issue, and when experienced during pregnancy, IPV substantially harms maternal health. Still, limited research has examined how IPV may influence prenatal oral health and dental care utilization. This study investigates the relationship between IPV during pregnancy and women's oral health experiences.

Data Data are from 31 states from 2016–2019 in the United States that participated in the Pregnancy Risk Assessment Monitoring System ($N=85,289$)—a population-based surveillance system of live births conducted annually by the Centers for Disease Control and Prevention and state health departments. Multivariable logistic regression analyses were used to examine the association between physical IPV during pregnancy (measured by being pushed, hit, slapped, kicked, choked, or physically hurt any other way by a current or ex-husband/partner) and various oral health experiences.

Findings Women who experienced prenatal physical IPV reported worse oral health experiences during pregnancy, including being more likely to report not knowing it was important to care for their teeth, not talking about dental health with a provider, needing to see a dentist for a problem, going to see a dentist for a problem, as well as having more unmet dental care needs.

Conclusions Together, these findings indicate that women who experience physical IPV during pregnancy have lower knowledge of prenatal oral health care, more oral health problems, and greater unmet dental care needs. Given the risk of IPV and oral health problems for maternal and infant health, the study findings point to greater attention toward the oral health needs of IPV-exposed pregnant women.

Keywords Intimate partner violence, Pregnancy, Oral health, Maternal health, PRAMS

*Correspondence:

Jason M. Nagata
jason.nagata@ucsf.edu

¹ School of Public Health, University of Texas Health Science Center at Houston, Houston, TX, USA

² Department of Criminal Justice, Boise State University, Boise, ID, USA

³ Johns Hopkins Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA

⁴ School of Dentistry, University of Texas Health Science Center at San Antonio, San Antonio, TX, USA

⁵ Factor-Inwentash Faculty of Social Work, University of Toronto, Toronto, Canada

⁶ Department of Pediatrics, University of California, San Francisco, San Francisco, CA, USA



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Physical intimate partner violence (IPV)—physical aggression or abuse by a romantic partner—is a traumatic life event that can harm health and well-being [1]. According to estimates, approximately one-in-five biological women in the United States experienced severe physical violence from an intimate partner in their lifetime [2]. Physical IPV can be especially detrimental when it is experienced during pregnancy. While there are no precise estimates of the cost of physical IPV during pregnancy, estimates from the United States that the lifetime cost of intimate partner violence defined as contact sexual violence, physical violence, or stalking victimization results in a lifetime population economic burden of nearly \$3.6 trillion (2014 US\$), including 2.1 trillion in medical costs, \$1.3 trillion in lost productivity among victims and perpetrators, \$73 billion in criminal justice spending, and \$62 billion (2%) in other costs, including victim property loss or damage [3].

Estimates suggest that over 300,000 pregnant women in the United States experience any IPV (i.e., physical, emotional, sexual, and/or psychological) annually [4] (out of approximately 6 million births) [5]. Importantly, research has connected IPV to adverse health consequences for maternal and infant health, including physical injury, preterm birth, and maternal and neonatal death [6–9]. Even so, oral health is one key but overlooked area that IPV may harm. Oral health is strongly connected to general health and quality of life [10], and poor oral health is associated with other systemic diseases due to the shared link with inflammation [11]. During pregnancy, women are at heightened risk of oral health problems as changes in diet and hormonal fluctuations can contribute to an increased risk of gingivitis and periodontal disease [12–14]. While the mechanisms are not fully understood, oral health problems during pregnancy have been found to negatively impact birth outcomes, including preterm birth and low birth weight [15–17]. Indeed, scholars have noted that “maternal oral disease during pregnancy is a significant public health issue due to its prevalence and life course connections with adverse pregnancy/birth outcomes, early childhood caries, and chronic diseases.” [18] Due to this growing evidence, recent public health efforts have focused on increasing awareness of the importance of oral health care during pregnancy and improving access to preventative and problem-related dental care for pregnant women [18, 19]. However, women who experience physical IPV may face challenges accessing health care services and be at risk for oral health problems.

Even though research demonstrates IPV is associated with physical health problems [1] and access to health care services in general [20], there is limited

prior research on the influence of IPV on the oral health experiences of women, especially during pregnancy. For instance, one study found that women who experienced IPV during the prenatal period had elevated rates of oral health problems during pregnancy, including problems that stem from physical violence, including painful gingivae, toothache, and an injury to the mouth, teeth, or gingivae [21]. However, there is a lack of research on a broad range of oral health experiences that might be influenced by IPV, including general oral health problems, oral health literacy, dental care utilization, and unmet dental care needs, despite reasons to suspect that IPV might serve as a risk factor that undermines oral health and oral health care utilization during pregnancy.

First, studies of pregnant women who experience abuse often report that their partners attempt to isolate them from support systems [22–24]. In such cases, an abusive partner may actively prevent a woman from accessing health care services, including dental care [20]. Moreover, women may not want to deal with the trouble of this process, particularly if they feel that preventative dental care is not that critical, given the host of other issues they are dealing with. Second, and relatedly an abusive partner may interfere with health care visits or treatment to prevent detection of the abuse by a medical provider [25]. One recent study found that women who experienced IPV during pregnancy were less likely to receive adequate prenatal care and faced greater barriers to prenatal care services [26]. However, no research has assessed the relationship between IPV and patterns of dental care utilization. Third, oral health care is often costly, and dental care requires more out-of-pocket spending than other medical expenses [27, 28]. A physically abusive partner may exercise economic dependence that may discourage pregnant women from pursuing oral health care and impact the quality of dental care when received [22, 24, 29]. In turn, if women experiencing IPV are not receiving regular preventive care, this can lead to additional problems with oral health. These include more oral health problems due to the lack of preventive treatment, as well as less knowledge about the importance of proper oral health care due to infrequent interactions with dental care professionals. In addition, aside from oral health problems that may emerge from a lack of preventive care, IPV can increase oral health problems through traumatic injury from physical violence [30–32]. Finally, if women who experience IPV during pregnancy have more oral health care needs, but IPV also reduces the likelihood of receiving dental care services., It is also likely that women who experience IPV will be less likely to receive oral health treatment for a problem when needed.

Given the lack of research on IPV and oral health during pregnancy, the current study draws on state

surveillance system data of recent mothers to investigate the association between prenatal physical IPV and a range of oral health experiences during pregnancy.

Methods

Data

The data for this study is from the Pregnancy Risk Assessment Monitoring System (PRAMS). The PRAMS is a population-based surveillance system of live births conducted annually by the Centers for Disease Control and Prevention and state health departments. Each participating site (i.e., states and territories) uses birth certificate records to collect a stratified systematic sample of approximately 100 to 250 mothers who delivered a live birth. The PRAMS data are derived from three primary sources: (1) birth certificates, (2) state and territory vital record systems, and (3) survey responses via a questionnaire. Surveys are distributed to mothers in a series of three mailings made 2–4 months after birth. Non-responders are followed up with a series of 15 phone calls made throughout a 2-to-3-week following the last mailing attempt. Sites are included in PRAMS if a minimum response rate has been met, set at 55% in 2015–2017 and 50% since 2018.

The PRAMS survey comprises different questionnaire types that determine the final sample in the current study. A core survey is distributed to all participating sites, which includes questions about the following topics: attitudes about pregnancy, preconception care, prenatal care, breastfeeding, cigarette and alcohol use, health insurance coverage, physical abuse, infant health care, and contraception use. A subset of sites also administered questions from a pretested list developed by the CDC or individual states. This latter questionnaire includes questions about oral health experiences. Therefore, the current study uses data from the 31 states from 2016–2019 with available data on oral health experiences and the other variables used in the analysis ($N=85,289$ recent mothers). The full list of states and years included in the study are reported in Appendix A. Additional information about the PRAMS survey and methodology is available in Shulman et al. [33]. The CDC institutional review board approved the use of the PRAMS data for this study as part of the external researcher data sharing agreement. Additional information on access and use of the PRAMS data can be found at: <https://www.cdc.gov/prams/prams-data/researchers.htm>.

Independent variable

Physical Intimate Partner Violence is measured using survey questions asking respondents whether they experienced physical abuse during pregnancy. Specifically, women were asked two questions: (1) “During your most

recent pregnancy, did your husband or partner push, hit, slap, kick, choke, or physically hurt you in any other way?” and (2) “During your most recent pregnancy, did an ex-husband or ex-partner push, hit, slap, kick, choke, or physically hurt you in any other way?” Using these questions, we created a binary variable where women who responded affirmatively to either of these questions were coded as experiencing physical IPV (1), and those who answered no to both were coded as not experiencing physical IPV (0).

Dependent variables

The six dependent variables come from a series of questions regarding the mother’s self-reported oral health experiences during pregnancy. These questions encompass all oral health and dental care questions included in the PRAMS survey. Consistent with prior research, all are coded in the direction of risk [34, 35]. *Didn’t Know Importance of Oral Health Care* is measured using a question asking, “Did you know it was important to care for your teeth and gums during your pregnancy?” (1=no, 0=yes). *Didn’t Talk about Dental Health with Provider* is measured from a survey question asking: “Did a dental or other health care worker talk with you about how to care for your teeth and gums?” (1=no, 0=yes). *No Dental Prophylaxis* is based on a question asking, “During your most recent pregnancy, did you have your teeth cleaned by a dentist or dental hygienist?” (1=no, 0=yes). *Needed to see Dentist for a Problem* is measured from an item asking, “Did you need to see a dentist for a problem?” (1=yes, 0=no). *Visited Dentist for a Problem* is measured from a question asking, “Did you go to see a dentist or dental clinic about a problem?” (1=yes, 0=no). Finally, *Visited a Dentist, Conditional on Needing to See a Dentist* is a binary variable based on a subsample of respondents who responded affirmatively to the question inquiring about needing to see a dentist for a problem but who also replied “no” when asked if they visited a dentist for a problem.

Control variables

Control variables are included to account for the demographic and socioeconomic characteristics of the mother. Variables are selected based on prior research with the PRAMS data assessing IPV and health and dental care outcomes [26, 26, 34]. Socio-demographic control variables include *mother’s age* (<24, 25–29, 30–34, and 35 or older), *mother’s race/ethnicity* (White, Hispanic, Black, Native American, Asian or Other), *currently married* (1=currently married; 0=not currently married). To account for socioeconomic differences and household size that might influence oral health and dental care, we include control variables for *educational attainment*

(less than high school, high school graduate, some college, college graduate), *household income* (\leq \$16,000, \$16,0001-\$40,000, \$40,001-\$85,000, >\$85,000), *number of financial dependents* (range 0–7), and *number of prior births*(0, 1, 2, 3 or more) [36, 37]. Pre-pregnancy *body mass index*(underweight, normal weight, overweight, obese) is included as a proxy for health and because of a possible connection between obesity and oral health [38, 39]. Given the importance of dental insurance for oral health and dental care, we control for whether a mother reported having *no dental insurance*, which is measured using a question asking, “Did you have insurance to cover dental care during your pregnancy” (1=no, 0=yes) [37, 40]. Finally, control variables are also included for the state of residence and year of birth.

Statistical analysis

The analysis is performed in three stages. First, the descriptive statistics for the analytic sample are presented. Next, the bivariate association between oral health experiences and IPV is examined using a two-tailed t-test. Third, multiple logistic regression analyses adjusting for control variables are performed. All analyses were adjusted for survey weights and strata information using the SVY command in Stata Version 17. Missing data were handled using listwise deletion, considering that the sample size is relatively large and that listwise deletion may be more robust to violations of the Missing at Random (MAR) assumption than imputation [41].

Results

Table 1 presents the weighted descriptive statistics for the analytic sample (N=85,289). Overall, 1.8% (n=1,706) of respondents reported physical IPV during their most recent pregnancy. Across oral health experience measures, 11.9% of the sample reported not knowing it was important to care for their teeth (n=12,353), 47.5% did not talk about dental health with a provider (n=40,712), 51.5% did not have dental prophylaxis (n=44,592), 18.2% reported needing to see a dentist for a problem (n=44,592), 13.8% reported having visited a dentist for a problem (n=16,100), and among those who reported needing to see a dentist for a problem (N=16,100), 68.3% reported having visited a dentist for a problem (n=10,887). Appendix B provides the summary statistics stratified by IPV exposure.

Next, to assess how these patterns vary by physical IPV exposure during pregnancy, Fig. 1 displays the dependent variables stratified by IPV status. The results of two-tailed t-tests demonstrate that women who experienced physical IPV during pregnancy reported worse oral health experiences across all

Table 1 Summary statistics from pregnancy risk assessment monitoring system, 2016–2019 (N=85,289)

Variables	%/ Mean (SD)
<i>Oral Health Experiences</i>	
Didn't Know Important to Care for Teeth	11.9%
Didn't Talk about Dental Health with Provider	47.5%
No Dental Prophylaxis	51.5%
Needed to see Dentist for a Problem	18.2%
Visited Dentist for Problem	13.8%
Visited a Dentist Needing to see a Dentist	68.3%
Physical Intimate Partner Violence	1.8%
<i>Maternal Age</i>	
< 24	20.3%
25–29	29.1%
30–34	31.0%
35 +	19.6%
<i>Maternal Race/Ethnicity</i>	
White	63.0%
Hispanic	15.0%
Black	13.3%
Other Race/Ethnicity	8.6%
<i>Maternal Educational Attainment</i>	
Less than High School	9.2%
High School Graduate	22.8%
Some College	27.1%
College Graduate	40.9%
Married	65.3%
<i>Number of Prior Births</i>	
0	39.1%
1	33.7%
2	16.1%
3 +	11.1%
<i>Body Mass Index</i>	
Underweight	3.1%
Normal Weight	44.1%
Overweight	24.8%
Obese	28.0%
<i>Household Income</i>	
\leq \$16,000	17.4%
\$16,000, \$40,000	22.6%
\$40,001 – \$85,000	30.1%
> \$85,000	29.9%
Number of Dependents	2.94 (1.37)
No Dental Insurance	19.7%

Visited a Dentist | Needing to see a Dentist available for 16,100 respondents who reported needing to see a dentist for a problem

Abbreviations: SD Standard deviation

measures. Specifically, IPV exposure was associated with reporting not knowing it was important to care for teeth during pregnancy (19.3% vs. 11.8%, $p < 0.001$),




Fig. 1 Dental health measures by physical intimate partner violence. *Note:* two-sample test for proportions between no IPV and IPV are statistically significant at the $\alpha = .01$ level for all variables above

not talking about dental health with a provider (61.0% vs. 47.3%, $p < 0.001$), no dental prophylaxis (63.5% vs. 51.3%, $p < 0.001$), needing to see a dentist for a problem (36.0% vs. 17.9%, $p < 0.001$), having gone to see a dentist for a problem (23.3% vs. 13.6%, $p < 0.001$), and having unmet dental care needs (68.7% vs. 57.5%, $p < 0.01$).

Table 2 presents the oral health measures regressed on IPV, adjusting for covariates. Findings show that women who experienced IPV during pregnancy were significantly more likely to report not knowing the importance of oral health care (Odds Ratio [OR]=1.403, 95% Confidence Interval [CI]=1.138, 1.729), not having an oral health discussion with their provider (OR=1.451, 95% CI=1.217, 1.729), needing to see a dentist for a problem (OR=1.731, 95% CI=1.445, 2.072) and having visited a dental care provider for a problem (OR=1.429, 95% CI=1.164, 1.756). Finally, restricting to the subsample of women who reported needing to see a dentist for a problem ($n=16,100$), the results in Model 6 show that IPV exposure had a marginally significant but negative, association with visiting a dentist for a problem. Overall, IPV-exposed women were about 25% less likely to see a dentist for a problem conditional on needing to see a dentist for a problem (OR=0.746, CI=0.555, 1.004), indicating that IPV-exposed women have more unmet dental care needs.

Discussion

This study aimed to examine the relationship between a mother's experiences with physical IPV during pregnancy and various oral health experiences. The findings

revealed four general patterns in the relationship between IPV and oral health. First, IPV was associated with greater odds of not knowing it was important to care for oral health during pregnancy and not talking with an oral health provider about oral health during pregnancy. This indicates that IPV is related to lower oral health knowledge during pregnancy. While the exact mechanisms for this relationship are unclear, it may be that because physical IPV is associated with less frequent interactions with dental care professionals and with prenatal care in general [26], women who experience physical IPV may receive less health-related consultation and therefore have less knowledge about proper health behaviors during pregnancy. Second, women who experienced IPV during pregnancy were more likely to report needing to see a dentist for a problem and going to see a dentist for a problem indicating that IPV is associated with more oral health problems. This finding is consistent with other research demonstrating a link between IPV and a series of oral health problems, including physical injury to the maxillofacial area that would require dental care [21, 30, 42–45].

Third, IPV was related to being less likely to see an oral health care provider for a problem, conditional on needing oral health care for a problem, highlighting that IPV is related to more unmet oral health care needs. Accordingly, this finding suggests that despite the greater overall need for oral health services, physical IPV may pose several barriers that prevent adequate access to dental care services for women in need [20, 26]. A useful direction for future research would be to identify the potential barriers that prevent women in need of dental care services

Table 2 Logistic regression of oral health experiences during pregnancy on physical intimate partner violence and other covariates

Variables	Model 1: Didn't Know Important to Care for Teeth		Model 2: Didn't Talk about Dental Health		Model 3: No Dental Prophylaxis		Model 4: Needed to see Dentist for a Problem		Model 5: Went to see Dentist for a Problem		Model 6: Visited a Dentist Needing to see a Dentist	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Physical IPV	1.403**	(1.138, 1.729)	1.451***	(1.217, 1.729)	1.012	(0.849, 1.207)	1.731***	(1.445, 2.072)	1.429***	(1.164, 1.756)	0.746†	(0.555, 1.004)
Control Variables	✓		✓		✓		✓		✓		✓	
Observations	85,289		85,289		85,289		85,289		85,289		16,100	

Control variables include maternal age, maternal race/ethnicity, maternal educational attainment, marital status, number of prior births, pre-pregnancy body mass index, household income, number of financial dependents, does not have dental insurance, state of residence, and year of birth

*** $p < 0.001$

** $p < 0.01$

* $p < 0.05$

† $p < .10$

from accessing such services, including cost, psychological control by the abuser, low trust in the health care system, low self-efficacy, or being unable to find or access dental care services because of transportation issues, getting time off work, or finding childcare [20, 26]. Finally, while the results showed that in a bivariate model, IPV was related to a lower likelihood of receiving a dental cleaning during pregnancy, the results of the multiple regression analysis adjusting for control variables found no statistical difference in the odds of receiving dental prophylaxis during pregnancy among women with and without IPV exposure.

Limitations and future directions

Before discussing the implications of these results, it is important to highlight some limitations that can be expanded upon in future research. First, the measure of IPV used in this study refers to physical violence, and therefore the study findings are not necessarily generalizable to other forms of IPV, including emotional, psychological, or sexual abuse. Second, because the measure of IPV is focused on the violence that occurs in the contexts during pregnancy, the results cannot speak to the influence of IPV at earlier points in life on oral health experiences during pregnancy. Third, the binary nature of the independent variable captures the presence of whether IPV occurred. However, we lack details on the frequency, duration, and severity of IPV and when the IPV occurred during pregnancy. Fourth, while the prevalence of physical IPV during in the current study (1.8%) is consistent with other PRAMS research, this figure is likely lower than estimates in the general population, considering (a) the focus is only on physical IPV compared to other forms of IPV (emotional, sexual, or psychological), (b) the focus is only on physical IPV during pregnancy, compared to lifetime, and (c) the focus on women who recently gave birth means that our sample is younger than a general population sample [21, 26, 46]. Fifth, the questions about oral health experiences during pregnancy lack information on specific types of oral health problems such as dental caries, periodontitis, or traumatic dental injuries. Future research that examines the relationship between IPV and more detailed measures of dental health issues would be valuable. Sixth, the IPV and oral health experiences measures are self-reported and can be subject to recall or social desirability biases. Seventh, because of social desirability issues, women being concerned about the repercussions of marking a response affirmative of abuse, and coercive control preventing survey responses

in the context of ongoing IPV, women experiencing IPV may be underrepresented in this survey. Finally, because the PRAMS is a cross-sectional survey, this study cannot establish a causal association between IPV and oral health experiences during pregnancy.

Public health implications

Considering these limitations, the results hold important implications for improving oral health experiences among women experiencing IPV. The general pattern detected in this study is that IPV exposure is associated with (1) less knowledge about oral health during pregnancy, (2) greater oral health care needs, and (3) more unmet dental care needs. Therefore, these findings highlight the need for better oral healthcare access among women who have experienced IPV. One way to expand access to dental care knowledge is by providing information on proper oral health during pregnancy as a component of prenatal care. Such information can include information about standard oral health guidelines during pregnancy and locations of local affordable dental care providers. Second, expanding oral health treatment to domestic violence shelters can be a useful way to reach IPV-exposed women who are out of the reach of the traditional medical care system [47]. Finally, another important implication of this study is increasing dental care providers' knowledge about IPV. Because many IPV-related injuries are to the head, neck, and face, dental care providers play an important role in detecting IPV and providing resources to aid IPV victims [48–51]. Even so, studies show that a sizeable portion of dental providers do not feel they have sufficient training to assist IPV victims or screen for IPV properly [51, 52]. Thus, expanding educational offerings on IPV in dental schools and continuing education programs would be beneficial [51]. Indeed, evidence suggests domestic violence education can improve the dental care provided to survivors of domestic violence [47].

Conclusions

The current study showed that women who experienced physical IPV during pregnancy also exhibited more oral health problems, lower oral health knowledge, and greater unmet oral health care needs. Considering the profound risks that both IPV and oral health problems can pose for maternal and child health, the findings point to the need for efforts to reduce the prevalence of IPV during pregnancy and expand oral health care services to IPV-exposed populations.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12903-023-03491-0>.

Additional file 1: Appendix A. List of Sites and Years in Analytic Sample.
Appendix B. Summary Statistics from Pregnancy Risk Assessment Monitoring System, 2016-2019 Stratified by Physical IPV

Acknowledgements

The authors wish to acknowledge the CDC PRAMS Working Group: Tammie Yellidell, MPH (AL), Kathy Perham-Hester, MS, MPH (AK), Letitia de Graft-Johnson, DrPH, MHSA (AR), Ashley Juhl, MSPH (CO), Jennifer Morin, MPH (CT), George Yocher, MS (DE), Tara Hylton, MPH (FL), Florence A. Kanu, PhD, MPH (GA), Matt Shim, PhD, MPH (HI), Julie Doetsch, MA (IL), Jennifer Pham (IA), Tracey D. Jewell, MPH (KY), Rosaria Trichilo, MPH (LA), Tom Patenaude, MPH (ME), Laurie Kettinger, MS (MD), Hafsatou Diop, MD, MPH (MA), Peterson Haak (MI), Mira Grice Sheff, PhD, MS (MN), Brenda Hughes, MPPA (MS), Venkata Garikapaty, PhD (MO), Emily Healy, MS (MT), Jessica Seberger (NE), David J. Laflamme, PhD, MPH (NH), Sharon Smith Cooley, MPH (NJ), Sarah Schrock, MPH (NM), Anne Radigan (NY), Pricila Mullachery, MPH (New York City), Kathleen Jones-Vessey, MS (NC), Grace Njau, MPH (ND), Ayesha Lampkins, MPH, CHES (OK), Cate Wilcox, MPH (OR), Sara Thuma, MPH (PA), Karine Tolentino Monteiro, MPH (RI), Kristin Simpson, MSW, MPA (SC), Ransom Wyse, MPH, MPH (TN), Tanya Guthrie, PhD (TX), Nicole Stone, MPH (UT), Peggy Brozicevic (VT), Kenesha Smith, MSPH (VA), Linda Lohdefinck (WA), Melissa Baker, MA (WV), Fiona Weeks, MSPH (WI), Lorie Chesnut, PhD (WY), *CDC PRAMS Team, Women's Health and Fertility Branch, Division of Reproductive Health.*

Authors' contributions

AT conceptualized and designed the study, carried out the analyses, drafted portions of the initial manuscript, reviewed and revised the manuscript, and approved of the final manuscript submitted; JL assisted in the design of the study, drafted portions of the initial manuscript, reviewed and revised the manuscript, and approved of the final manuscript submitted; DJ assisted in the design of the study, drafted portions of the initial manuscript, reviewed and revised the manuscript, and approved of the final manuscript submitted; RM drafted portions of the initial manuscript, reviewed and revised the manuscript, and approved of the final manuscript submitted; KG drafted portions of the initial manuscript, reviewed and revised the manuscript, and approved of the final manuscript submitted; JM drafted portions of the initial manuscript, reviewed and revised the manuscript, and approved of the final manuscript submitted.

Funding

J.M.N. was funded by National Institutes of Health (K08HL159350), the American Heart Association Career Development Award (CDA34760281), and the Doris Duke Charitable Foundation (2022056).

Availability of data and materials

The datasets generated and/or analyzed during the current study are not publicly available due to the nature of Pregnancy Risk Assessment Monitoring System not being publicly available. Data used in this study can be requested at <https://www.cdc.gov/prams/index.htm>. Queries about the data can be directed to Alexander Testa: alexander.testa@uth.tmc.edu.

Declarations

Ethics approval and consent to participate

The study was performed in accordance with the Declaration of Helsinki. The study was approved by the Centers for Disease Control and Prevention in accordance with the data usage agreement for the Pregnancy Risk Assessment Monitoring System. All participants provided informed consent.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 3 July 2023 Accepted: 4 October 2023

Published online: 12 October 2023

References

- Plichta SB. Intimate partner violence and physical health consequences: policy and practice implications. *J Interpers Violence*. 2004;19(11):1296–323. <https://doi.org/10.1177/0886260504269685>.
- Breiding MJ. Prevalence and characteristics of sexual violence, stalking, and intimate partner violence victimization—national intimate partner and sexual violence survey, United States, 2011. *Morb Mortal Wkly Rep Surveill Summ Wash DC* 2002. 2014;63(8):1–18.
- Peterson C, Kearns MC, McIntosh WL, et al. Lifetime economic burden of intimate partner violence among U.S. Adults *Am J Prev Med*. 2018;55(4):433–44. <https://doi.org/10.1016/j.amepre.2018.04.049>.
- American College of Obstetricians and Gynecologists. ACOG Committee Opinion No. 518: Intimate partner violence. *Obstet Gynecol*. 2012;119(2 Pt 1):412–7. <https://doi.org/10.1097/AOG.0b013e318249ff74>.
- Curtin SC, Abma JC, Ventura SJ. Pregnancy Rates for U.S. Women Continue to Drop. *NCHS Data Brief*. 2013;136:1–8.
- Sharps PW, Laughon K, Giangrande SK. Intimate partner violence and the childbearing year: maternal and infant health consequences. *Trauma Violence Abuse*. 2007;8(2):105–16. <https://doi.org/10.1177/1524838007302594>.
- Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes: a meta-analysis. *JAMA*. 2006;295(15):1809–23. <https://doi.org/10.1001/jama.295.15.1809>.
- Coker AL, Sanderson M, Dong B. Partner violence during pregnancy and risk of adverse pregnancy outcomes. *Paediatr Perinat Epidemiol*. 2004;18(4):260–9. <https://doi.org/10.1111/j.1365-3016.2004.00569.x>.
- Yost NP, Bloom SL, McIntire DD, Leveno KJ. A prospective observational study of domestic violence during pregnancy. *Obstet Gynecol*. 2005;106(1):61. <https://doi.org/10.1097/01.AOG.0000164468.06070.2a>.
- McGrath C, Bedi R. The association between dental anxiety and oral health-related quality of life in Britain. *Community Dent Oral Epidemiol*. 2004;32(1):67–72. <https://doi.org/10.1111/j.1600-0528.2004.00119.x>.
- Haumschild MS, Haumschild RJ. The importance of oral health in long-term care. *J Am Med Dir Assoc*. 2009;10(9):667–71. <https://doi.org/10.1016/j.jamda.2009.01.002>.
- Kloetzel MK, Huebner CE, Milgrom P. Referrals for dental care during pregnancy. *J Midwifery Womens Health*. 2011;56(2):110–7. <https://doi.org/10.1111/j.1542-2011.2010.00022.x>.
- Lachat MF, Solnik AL, Nana AD, Citron TL. Periodontal disease in pregnancy: review of the evidence and prevention strategies. *J Perinat Neonatal Nurs*. 2011;25(4):312–9. <https://doi.org/10.1097/JPN.0b013e31821072e4>.
- Lieff S, Boggess KA, Murtha AP, et al. The Oral Conditions and Pregnancy Study: Periodontal Status of a Cohort of Pregnant Women. *J Periodontol*. 2004;75(1):116–26. <https://doi.org/10.1902/jop.2004.75.1.116>.
- Bobetsis YA, Graziani F, Gürsoy M, Madianos PN. Periodontal disease and adverse pregnancy outcomes. *Periodontol* 2000. 2020;83(1):154–74. <https://doi.org/10.1111/prd.12294>.
- Corbella S, Taschieri S, Del Fabbro M, Francetti L, Weinstein R, Ferrazzi E. Adverse pregnancy outcomes and periodontitis: A systematic review and meta-analysis exploring potential association. *Quintessence Int*. 2016;47(3):193–204. <https://doi.org/10.3290/j.qi.a34980>.
- Iheozor-Ejirofor Z, Middleton P, Esposito M, Glenny AM. Treating periodontal disease for preventing adverse birth outcomes in pregnant women. *Cochrane Database Syst Rev*. 2017;(6). <https://doi.org/10.1002/14651858.CD005297.pub3>
- Vamos CA, Thompson EL, Avendano M, Daley EM, Quinonez RB, Boggess K. Oral health promotion interventions during pregnancy: a systematic review. *Community Dent Oral Epidemiol*. 2015;43(5):385–96. <https://doi.org/10.1111/cdoe.12167>.
- Hartnett E, Haber J, Krainovich-Miller B, Bella A, Vasilyeva A, Lange KJ. Oral health in pregnancy. *J Obstet Gynecol Neonatal Nurs*. 2016;45(4):565–73. <https://doi.org/10.1016/j.jogn.2016.04.005>.
- Wilson K, Silberberg M, Brown A, Yaggy S. Health needs and barriers to healthcare of women who have experienced intimate partner violence.

- J Womens Health. 2007;16:1485–98. <https://doi.org/10.1089/jwh.2007.0385>.
21. Testa A, Lee J, Neumann A, Jackson DB. Physical intimate partner violence and oral health problems during pregnancy. *J Am Dent Assoc*. Published online January 25, 2023. <https://doi.org/10.1016/j.adaj.2022.11.019>
 22. Bacchus L, Mezey G, Bewley S. A qualitative exploration of the nature of domestic violence in pregnancy. *Violence Women*. 2006;12(6):588–604. <https://doi.org/10.1177/1077801206289131>.
 23. Noel NL, Yam M. Domestic violence: the pregnant battered woman. *Nurs Clin North Am*. 1992;27(4):871–84. [https://doi.org/10.1016/S0029-6465\(22\)02816-X](https://doi.org/10.1016/S0029-6465(22)02816-X).
 24. Sales P, Murphy S. Surviving violence: pregnancy and drug use. *J Drug Issues*. 2000;30(4):695–723. <https://doi.org/10.1177/002204260003000403>.
 25. McCloskey LA, Williams CM, Lichter E, Gerber M, Ganz ML, Sege R. Abused women disclose partner interference with health care: an unrecognized form of battering. *J Gen Intern Med*. 2007;22(8):1067–72. <https://doi.org/10.1007/s11606-007-0199-z>.
 26. Testa A, Lee J, Semenza DC, Jackson DB, Ganson KT, Nagata JM. Intimate partner violence and barriers to prenatal care. *Soc Sci Med*. 2023;320:115700. <https://doi.org/10.1016/j.socscimed.2023.115700>.
 27. Vujicic M, Buchmueller T, Klein R. Dental care presents the highest level of financial barriers, compared to other types of health care services. *Health Aff (Millwood)*. 2016;35(12):2176–82. <https://doi.org/10.1377/hlthaff.2016.0800>.
 28. Institute of Medicine. Improving Access to Oral Health Care for Vulnerable and Underserved Populations. National Academies Press; 2011:13116. <https://doi.org/10.17226/13116>
 29. Brownridge DA, Taillieu TL, Tyler KA, Tiwari A, Chan KL, Santos SC. Pregnancy and intimate partner violence: risk factors, severity, and health effects. *Violence Women*. 2011;17(7):858–81. <https://doi.org/10.1177/1077801211412547>.
 30. Caldas IM, Grams AC, Afonso A, Magalhães T. Oral injuries in victims involving intimate partner violence. *Forensic Sci Int*. 2012;221(1):102–5. <https://doi.org/10.1016/j.forsciint.2012.04.010>.
 31. Glendor U. Aetiology and risk factors related to traumatic dental injuries – a review of the literature. *Dent Traumatol*. 2009;25(1):19–31. <https://doi.org/10.1111/j.1600-9657.2008.00694.x>.
 32. Nelms AP, Gutmann ME, Solomon ES, DeWald JP, Campbell PR. What victims of domestic violence need from the dental profession. *J Dent Educ*. 2009;73(4):490–8. <https://doi.org/10.1002/j.0022-0337.2009.73.4.tb04720.x>.
 33. Shulman HB, D'Angelo DV, Harrison L, Smith RA, Warner L. The pregnancy risk assessment monitoring system (PRAMS): overview of design and methodology. *Am J Public Health*. 2018;108(10):1305–13. <https://doi.org/10.2105/AJPH.2018.304563>.
 34. Testa A, Ganson KT, Jackson DB, Bojorquez-Ramirez P, Weiser SD, Nagata JM. Food insecurity and oral health care experiences during pregnancy. *J Am Dent Assoc*. 2022;153(6):503–10. <https://doi.org/10.1016/j.adaj.2021.12.010>.
 35. Testa A, de Gimeno RuizPorras D. The association of employment status and unwanted job loss with maternal oral health experiences: findings from the pregnancy risk assessment monitoring system. *BMC Oral Health*. 2023;23(1):168. <https://doi.org/10.1186/s12903-023-02869-4>.
 36. Cha A, Cohen R. Dental Care Utilization Among Adults Aged 18–64: United States, 2019 and 2020. National Center for Health Statistics (U.S.); 2022. <https://doi.org/10.15620/cdc:115597>
 37. National Institute of Dental and Craniofacial Research. Oral Health in America: Advances and Challenges. Bethesda, MD: National Institutes of Health; 2021.
 38. Jepsen S, Suvan J, Deschner J. The association of periodontal diseases with metabolic syndrome and obesity. *Periodontol*. 2020;83(1):125–53. <https://doi.org/10.1111/prd.12326>.
 39. Northridge ME, Kumar A, Kaur R. Disparities in access to oral health care. *Annu Rev Public Health*. 2020;41:513–35. <https://doi.org/10.1146/annurev-publhealth-040119-094318>.
 40. Borrell LN, Reynolds JC, Fleming E, Shah PD. Access to dental insurance and oral health inequities in the United States. *Community Dent Oral Epidemiol*. 2023;51(4):615–20. <https://doi.org/10.1111/cdoe.12848>.
 41. Allison PD. Missing Data. Thousand Oaks, CA: SAGE Publications; 2001.
 42. Sheridan DJ, Nash KR. Acute injury patterns of intimate partner violence victims. *Trauma Violence Abuse*. 2007;8(3):281–9. <https://doi.org/10.1177/1524838007303504>.
 43. Ferreira MC, Batista AM, Ferreira F de O, Ramos-Jorge ML, Marques LS. Pattern of oral–maxillofacial trauma stemming from interpersonal physical violence and determinant factors. *Dent Traumatol*. 2014;30(1):15–21. <https://doi.org/10.1111/edt.12047>.
 44. Lam R. Epidemiology and outcomes of traumatic dental injuries: a review of the literature. *Aust Dent J*. 2016;61(S1):4–20. <https://doi.org/10.1111/adj.12395>.
 45. Saddki N, Suhaimi AA, Daud R. Maxillofacial injuries associated with intimate partner violence in women. *BMC Public Health*. 2010;10(1):268. <https://doi.org/10.1186/1471-2458-10-268>.
 46. Alhusen JL, Geller R, Dreisbach C, Constantoulakis L, Siega-Riz AM. Intimate partner violence and gestational weight gain in a population-based sample of perinatal women. *J Obstet Gynecol Neonatal Nurs*. 2017;46(3):390–402. <https://doi.org/10.1016/j.jogn.2016.12.003>.
 47. Abel SN, Bhoopathi V, Herzig K, Godoy MT, Kowal HC, Gerbert B. The impact of an oral health program on domestic violence survivors within community shelters. *J Am Dent Assoc*. 2013;144(12):1372–8. <https://doi.org/10.14219/jada.archive.2013.0073>.
 48. Coulthard P, Feder G, Evans MA, et al. Dentistry responding to domestic violence and abuse: a dental, practice-based intervention and a feasibility study for a cluster randomised trial. *Br Dent J*. 2022;233(11):949–55. <https://doi.org/10.1038/s41415-022-5271-x>.
 49. Femi-Ajao O, Doughty J, Evans MA, et al. Dentistry responding in domestic violence and abuse (DRiDVA) feasibility study: a qualitative evaluation of the implementation experiences of dental professionals. *BMC Oral Health*. 2023;23(1):475. <https://doi.org/10.1186/s12903-023-03059-y>.
 50. Doughty JL, Ferns H, Taylor-Weetman K, et al. Dental Public Health in Action: Delivering a domestic violence and abuse screening and identification training programme in North Staffordshire-based dental practices. *Community Dent Health*. 2023;40(1):3–8. https://doi.org/10.1922/CDH_00153Doughty06.
 51. Parish CL, Pereyra MR, Abel SN, Siegel K, Pollack HA, Metsch LR. Intimate partner violence screening in the dental setting: Results of a nationally representative survey. *J Am Dent Assoc*. 2018;149(2):112–21. <https://doi.org/10.1016/j.adaj.2017.09.003>.
 52. Harris CM, Boyd L, Rainchuso L, Rothman AT. Oral health care providers' knowledge and attitudes about intimate partner violence. *Am Dent Hyg Assoc*. 2016;90(5):283–96.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

