

# UCLA

## UCLA Previously Published Works

**Title**

Surgery of Olfactory Groove Meningiomas: When in Rome, Do as the Romans Do?

**Permalink**

<https://escholarship.org/uc/item/4kg4w2tf>

**Journal**

World Neurosurgery, 83(2)

**ISSN**

1878-8750

**Author**

Schaller, Karl

**Publication Date**

2015-02-01

**DOI**

10.1016/j.wneu.2014.12.004

Peer reviewed

## Imaging Techniques in Spinal Cord Injury

Benjamin M. Ellingson<sup>1-3</sup>, Noriko Salamon<sup>1</sup>, Langston T. Holly<sup>4</sup>

### Key words

- Cord
- Imaging
- Injury
- Spinal

### Abbreviations and Acronyms

**ADC:** Apparent diffusion coefficient  
**CT:** Computed tomography  
**DCE:** Dynamic contrast-enhanced  
**DTI:** Diffusion tensor imaging  
**FA:** Fractional anisotropy  
**fMRI:** Functional MRI  
**MRI:** Magnetic resonance imaging  
**PET:** Positron emission tomography  
**SCI:** Spinal cord injury



From the Departments of <sup>1</sup>Radiological Sciences, <sup>2</sup>Biomedical Physics, <sup>3</sup>Bioengineering, and <sup>4</sup>Neurosurgery, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California, USA

To whom correspondence should be addressed:  
 Langston T. Holly, M.D.

[E-mail: lholly@mednet.ucla.edu]

Citation: *World Neurosurg.* (2014) 82, 6:1351-1358.  
<http://dx.doi.org/10.1016/j.wneu.2012.12.004>

Journal homepage: [www.WORLDNEUROSURGERY.org](http://www.WORLDNEUROSURGERY.org)

Available online: [www.sciencedirect.com](http://www.sciencedirect.com)

1878-8750/\$ - see front matter © 2014 Elsevier Inc.  
 All rights reserved.

### INTRODUCTION

Spinal imaging plays an essential role in the diagnosis, treatment, and rehabilitation of patients with spinal cord injury (SCI). Traditionally, these modalities have consisted of plain radiography (Figure 1A), computed tomography (CT; Figure 1B), and magnetic resonance imaging (MRI; Figure 1C and D). In combination, these techniques provide excellent macrostructural information regarding the classification and magnitude of the osseous and ligamentous injury, which, coupled with the clinical examination, guides patient management.

Despite their critical importance, these modalities offer comparatively less information regarding the microstructural injury to the spinal cord, and correlations between the radiographic and clinical findings largely have been limited. In part this has led to the development of novel

■ **BACKGROUND:** Spinal imaging plays a critical role in the diagnosis, treatment, and rehabilitation of patients with spinal cord injury (SCI). In recent years there has been increasing interest in the development of advanced imaging techniques to provide pertinent microstructural and metabolic information that is not provided by conventional modalities.

■ **METHODS:** This review details the pathophysiological structural changes that accompany SCI, as well as their imaging correlates. The potential clinical applications of novel spinal cord imaging techniques to SCI are presented.

■ **RESULTS:** There are a variety of novel advanced imaging techniques that are principally focused on the microstructural and/or biochemical function of the spinal cord, and can potentially be applied to traumatic SCI, including diffusion tensor imaging, magnetic resonance spectroscopy, positron emission tomography, single-photon emission computed tomography, and functional magnetic resonance imaging. These techniques are presently in various stages of development, including some whose applications are primarily limited to laboratory investigation, whereas others are being actively used in clinical practice.


■ **CONCLUSION:** Advanced imaging of the spinal cord has tremendous potential to provide patient-specific physiological information about the status of cord integrity and health. Advanced spinal cord imaging is still at early stages of development and clinical implementation but is likely to play an increasingly important role in the management of spinal cord health in the foreseeable future.

imaging techniques that are principally focused on the microstructural and/or biochemical function of the spinal cord, including diffusion tensor imaging (DTI), magnetic resonance spectroscopy, positron emission tomography (PET), single-photon emission computed tomography, and functional MRI (fMRI). These techniques are presently in various stages of development, including some whose applications are primarily limited to laboratory investigation, whereas others are being actively used in clinical practice. This review outlines the major structural and vascular changes that are expected to accompany the phases of traumatic injury of the spinal cord, along with the imaging correlates of these physiological changes. The application of conventional and novel imaging techniques to SCI will be discussed.

### PHASES OF SCI

#### Hyper-Acute and Acute Spinal Trauma

The hyper-acute and acute stages of SCI, typically referring from the time of traumatic insult to hours after the initial injury, result in direct mechanical injury as well as other indirect effects including local hypoxia, ischemia, hemorrhage, and edema. Mechanical disruption of neural tissue structure results in immediate death of cells in the region of the insult. The stretching and tearing of large axons results in damage to axonal membranes and an increase in membrane permeability (82). Magnetic resonance spectroscopy clearly demonstrates a decrease in N-acetyl aspartate, a neuronal marker, after traumatic SCI in animal models (69). When diffusion MRI, which is sensitive to the magnitude of water self-diffusion, is used, early axonal



death and the disruption of the cell membranes has been shown to result in elevated apparent diffusion coefficient (ADC) in animal studies (22, 38), numerical simulations (37, 40), and human patients (16, 32, 74, 95). Despite these intriguing observations, hyperacute imaging of SCI is relatively difficult in the clinical scenario because patients often are admitted after this stage of injury.

Some of the earliest imaging changes after SCI are the result of hemorrhages in the central gray matter adjacent to the central canal that spread radially from the central canal into the anterior horns and neighboring white matter regions around the injury epicenter and extend both rostrally and caudally (45, 75, 89). CT is ideally suited for identifying acute hemorrhage after SCI (23, 87) because it shows

hyperdense regions in areas of blood products. MRI techniques that are sensitive to changes in magnetic susceptibility, or susceptibility-weighted images, are also useful for identifying hemorrhagic lesions because of the sensitivity to microscopic magnetic perturbations from iron products after spinal trauma (92). Blood products may also be visible as hyper-intensity on precontrast T<sub>1</sub>-weighted images (36, 59) because the iron within the blood can dramatically shorten tissue relaxation rates.

Compressive or impact-induced spinal trauma often causes restriction of blood flow to the injury site (24, 49, 72). Arterial spin labeling, an MRI technique that uses magnetization tagged blood water in an artery and to quantify blood flow, has been successfully used to quantify blood flow in the mouse spinal cord (16, 25), although similar results in humans have not yet been obtained. Dynamic contrast-enhanced (DCE) MRI, an MR technique that uses a pharmacokinetic model for contrast agent extravasation from the vascular to the extravascular space to quantify blood volume and vascular permeability, also has been used to successfully quantify vascular changes in animal models (7-9, 17), but again analogous studies in human SCI have not yet been performed.

Similarly, the DCE approach has been successfully applied to CT-contrast agents in the spinal cord after injury and demonstrated a decrease in blood flow and volume after acute spinal trauma in animals (44, 54). DCE applied to CT has only recently been shown to be feasible in humans (10); thus, this technique may be useful in the very near future when applied to either CT or MRI data. Other techniques for imaging blood flow can be performed in the spinal cord, including PET imaging using radiolabeled water (H<sub>2</sub>-[<sup>15</sup>O]), single-photon emission computed tomography imaging, xenon-enhanced CT, and phase-contrast MRI; however, these techniques are not yet routinely performed after spinal trauma.

The reduction in blood flow commonly observed often results in a decrease in oxygen tension (24, 48, 83), forcing neural tissue to use anaerobic metabolic pathways in the form of high-energy phosphates, resulting in an overall decrease in metabolism for up to four hours after

the initial insult (76). After a period of anaerobic metabolism lasting from 4 to 24 hours after injury, a shift toward oxidative metabolism occurs in the viable tissue, and the remaining tissue starts to become necrotic (1, 90). PET studies that incorporate [<sup>18</sup>F]-fluoro-2-deoxy-d-glucose to examine glucose metabolism have clearly demonstrated this decrease in metabolic activity after acute trauma in the rat (64). In addition, investigators all report a decrease in the measured ADC in the early onset of spinal stroke, which is indicative of the ischemic events after infarction (4, 58, 60, 73). Similarly, other investigations have shown a decrease in the ADC within the center of the spinal cord in patients with cervical spondylotic myelopathy (4, 21), which was thought to occur as the result of vascular compromise. Although the precise mechanisms responsible for changes in the diffusion characteristics after acute ischemia are still speculative, it is believed that a shift in water from the greater ADC extracellular compartment to the low ADC intracellular compartment is a result of the decrease in the observed ADC, particularly when one uses clinical scanners that can only measure water mobility at long diffusion times.

Edema occurs relatively abruptly after traumatic injury as a direct result of the mechanical disruption of axon cell membranes, damage to local blood vessels, and electrolytic imbalances (6, 36, 43, 52, 70, 84). This damage to cell membranes increases the amount of extracellular, mobile water, resulting in hyperintensity on T2-weighted (20, 46, 94) or fluid-attenuated inversion recovery MRI (22, 38). ADC in the transverse plane as measured with DTI (38), which is thought to be a result of increased extracellular (fast diffusing) volume fraction. In addition, DTI in acute spinal trauma indicates a decrease in ADC parallel to the spinal cord, resulting in an overall decrease in diffusion anisotropy, or fractional anisotropy (FA), in the lesion sites during the period of severe edema and hemorrhage (79). It has been suggested that this decrease in diffusivity parallel to spinal cord orientation may be attributed to metabolic dysfunction as opposed to specific changes in axon morphology (2); however, this is still currently under active investigation.

### Subacute Stage of SCI

After the hyper-acute and acute stages of SCI reactive cells infiltrate near the injury site, axonal degeneration occurs within the regions of injured spinal tracts, and changes in the cellular structure of the neurons occur as a result of change in connectivity. In particular, the initial response to spinal trauma invokes infiltration of inflammatory cells from both the central nervous system and periphery. Activated microglia and astroglia increase in a number of processes (26) relatively proportional to the severity of the injury, and spread from the lesion epicenter into adjacent gray and white matter (65). Activated astrocytes then begin to proliferate and undergo hypertrophy (5), extending long distances into both gray and white matter (3). Polymorphonuclear granulocytes and macrophages are also present in the traumatic cavity in humans (12), as well as Schwann, meningeal, and fibroblast cells (78).

The correlation between these biological changes and structural MRI features is nonspecific, but more advanced MR techniques do show sensitivity to microstructural changes. Regions of local reactive cell infiltration remain hyperintense on T2 as the relative water concentration from edematous processes remain dominant; however, on diffusion MRI, reactive cells such as glia produce collagenous scar tissue that is expected to have a relatively high impact on tissue diffusivity. Schwartz et al. (78) demonstrated that DTI tractography can be used to identify and visualize glial scar orientation. Despite relatively widespread infiltration of reactive cells, the sensitivity of DTI tractography (i.e., DTI eigenvector orientation) appears isolated to regions close to the injury epicenter because microstructural changes must have a sense of “directionality” that is relatively coherent throughout a voxel. Thus, reactive astrocytes may only have an influence on eigenvector orientations close to the injury epicenter, or in relatively close proximity to the forming lesion cavity. The influx of high numbers of astrocytes, microglia, and macrophages are likely to decrease the extracellular volume, which is expected to decrease the overall apparent diffusion coefficient, counteracting the initial increase associated with edema.

Wallerian degeneration (35, 91), or axonal degeneration after traumatic injury,

is also present during the subacute stages of SCI and contributes to changes in both structural and diffusion-weighted MRI, even at locations distant from the injury site. Axon degeneration first manifests as breakdown of the myelin sheath and cytoskeletal proteins such as microtubules and neurofilaments. If the distal segments of damaged axons are not reconnected both structurally and functionally, these axons will die and degeneration will continue anterograde. If not connected, the proximal ends of the damaged axons will produce retraction bulbs in order to cleave leaking axoplasm (19). Experimental data suggest apoptosis and necrosis (53, 67, 71) result in extensive retrograde degeneration following injury in humans (3), starting with swelling of neuronal cell bodies (62, 96), nuclear movement within the soma to an eccentric position, then programmed cell death (85). This degenerative process may even continue in presynaptic cells due to loss of target cell signaling, resulting in a cascade of degeneration and total or partial dysfunction of spinal pathways.

Hyperintensity on T2-weighted images arises along affected tracts during Wallerian degeneration (18, 42, 61), which is thought to largely reflect the increased extracellular water fraction during axonal dieback. T1-weighted images show hypointensity within these regions, although the effect is very subtle. Magnetization transfer MRI, a sequence in which the hydrogen on large macromolecules like myelin are excited with radiofrequency energy that is exchanged with the mobile water pool, also has shown sensitivity to these degenerative processes (22, 77). Diffusion MRI shows an elevated ADC in the transverse plane above baseline levels (22) due to the decrease in boundaries to water diffusion during degeneration. Anterograde degeneration results in disintegration of both the axonal membrane and myelin sheath, which decreases the number and extent of transverse boundaries to diffusion and results in a higher transverse ADC. Retrograde degeneration also shows a similar, but slightly larger, increase in transverse ADC in experimental animal models (3), perhaps the subsequent increase in intracellular space from axonal swelling (34).

Gray matter also undergoes morphological changes, many of which have been implicated in severe neurological symptoms including chronic pain (55) and


spasticity (55). Experimental data have shown that the number of dendrites decreases and the length of the remaining dendritic projections increases in motoneurons after injury (88). In a mouse model of spinal injury, similar morphologic changes in spinal neurons have been reported; in addition, an enlargement of the soma during the subacute stages can also occur (29, 39). Despite these quite dramatic changes in gray matter morphology after injury, imaging characteristics resulting from changes are less profound. No real changes in standard MRI have been noted; however, changes on diffusion MRI have been observed. Spinal cord gray matter has been shown to be anisotropic (4, 40, 41), particularly compared with brain gray matter (50). The eigenvector orientations within spinal gray follow a similar rostral-caudal dominance to white matter, and the

particular eigenvector orientation may follow microstructures and the predominant soma orientation (14, 86, 93). Given the sensitivity of diffusion to gray matter microstructures, it is conceivable that large changes in soma morphology after SCI may influence diffusion measurements. Although not thoroughly investigated, an increase in overall apparent diffusion coefficient and a decrease in diffusion anisotropy are likely in spinal gray matter in the acute and subacute stages of injury because of the increase in soma size and decrease in the number of dendritic projections.

#### Chronic Stage of SCI

Although most of the degenerative processes are stabilized by the chronic stage of injury, which is typically defined from months to years after the initial injury, there is evidence to suggest ongoing

degeneration long after the initial response. Progressive demyelination has been documented during the chronic stages of injury (11, 13, 47, 86) and remyelination, when it does occur, results in axons with significantly thinner myelin sheaths (13). Large diameter axons are preferentially at risk for degeneration during chronic injury (51), which often leads to small, unmyelinated axons in damaged axonal tracts. Cysts formed from reactive astrocytes in the subacute stage can spread longitudinally down the spinal cord from the initial site of injury, leading to widespread changes in spinal cord morphology including necrosis (85). In addition, as spinal tracts are left nonfunctional for long periods of time, significant atrophy of the spinal cord also occurs, causing the remaining axons to be compressed and tightly packed. These structural changes all



**Figure 2.** A 77-year-old woman with chronic SCI caused by advanced cervical spondylosis, resulting in myelomalacia and impaired neurological function. **(A)** T2-weighted sagittal MRI showing anterior compression from the intervertebral disk at the C3-4 and C4-5 levels. **(B)** Fractional anisotropy (i.e., FA) color maps showing the relative degree of diffusion anisotropy (brightness) and primary eigenvector orientation (color). *Blue* indicates superior/inferior orientation; *green*, anterior/posterior orientation; *red*, left/right orientation. A greater degree of diffusion anisotropy is observed in healthy white matter tracts due to diffusion being more restricted perpendicular to axon tract orientation. In levels above and below the site of compression, FA color maps are both bright and blue because of high diffusion anisotropy and a rostral-caudal (superior-inferior) orientation,

respectively. Color maps at the level of compression show reduced brightness and both red and green colors, reflecting loss of diffusion anisotropy and distorted fiber tract orientation. **(C)** DTI tractography, where pseudoaxonal tracts are created based on the information present in the diffusion tensor, shows lack of fiber continuity through the site of compression. This finding reflects disruption in the architecture of the spinal cord such that the tractography algorithm does not propagate fiber tracts through this area. **(D)** ADC, a quantitative measurement of water mobility, is highest within the cerebrospinal fluid within the spinal canal and lower within the spinal tissue. In areas of compression both abnormally high and low ADC can manifest, reflecting the myriad of inflammatory processes and pathogenesis that can alter water diffusivity.

contribute to observed abnormalities on both structural and diffusion MRI (Figure 2). Widespread axonal degeneration results in T2 hyperintensity in affected spinal tracts (69). These tracts also have elevated choline as the result of degenerative changes (27, 30, 31). Diffusion MR characteristics in chronic injury have only recently been explored (37). These studies demonstrate significant changes in diffusion distributions in chronic injury, indicative of expected changes in the spinal cord microstructure. Specifically, ADC measurements taken along the transverse orientation in spinal white matter have shown to be lower than uninjured controls and demonstrate a kurtosis (or flattening) of the diffusion distribution, likely due to loss of large diameter axons because of the known dependence of transverse ADC to axon diameter (6, 36, 43, 52, 66, 70, 81, 84).

### CLINICAL CORRELATION

Conventional MRI is considered the “gold standard” imaging modality in the assessment of spinal cord damage (63) yet has not been shown to consistently correlate with the clinical findings after SCI. Consequently, there has been an increasing interest in identifying noninvasive imaging studies that can provide pertinent microstructural and metabolic information about the injured spinal cord, which is theorized to provide more accurate clinical correlation. The potential utility of such imaging modalities and impact on treatment of SCI patients are manifold: (1) the ability to predict functional outcome which can influence rehabilitation strategies, (2) to have a better understanding of the degree of neurologic impairment related to SCI in patients that have concomitant head injury or are otherwise unexamined, (3) the assessment of neurological injury in children, a patient population in whom the clinical examination is often times unreliable (21), (4) the assessment of spinal cord integrity below the level of injury, and (5) the measurement of cellular changes in response to neural repair or biologic therapies directed towards healing the injured spinal cord.

One such imaging modality, DTI, has been demonstrated to be a more sensitive biomarker of spinal cord damage than standard T2 weighted imaging for both SCI related to cervical spondylosis (15, 33,

68, 80) and traumatic injury (15). For example, Chang et al (68) evaluated 10 chronic SCI patients and 10 healthy controls by using DTI and conventional MRI. Quantitative parameters of DTI (e.g., FA, ADC) were calculated for each cervical spinal cord level, and then DTI tractography parameters were determined for the sub-axial levels, which included the number of “tracts” passing through the injury site, or “connection rate.” Neurologic function was assessed by use of the International Standards for Neurological Classification of Spinal Cord Injury. Results demonstrated that abnormal appearing cervical levels on conventional MRI were not correlated with clinical findings in SCI patients; however, FA correlated with the motor function, as did the number of DTI tractography fibers and connection rates through the injury site. Specifically, on fiber tractography the one American Spinal Injury Association scale A (i.e., complete) patient demonstrated no visible connections crossing the lesion, and minimal connections were observed in three patients without motor function but spared sensory function. In patients with incomplete injury, the overall imaginary fiber numbers were greater than those with complete or motor complete paralysis. In a separate study, Petersen et al. (68) evaluated 19 chronic cervical SCI patients by using DTI, electrophysiological measures (SSEP, MEP), and neurological examination via the American Spinal Injury Association impairment scale. They found that the FA values were decreased compared to healthy control subjects and that the decrease in DTI values correlated with the clinical completeness of the SCI and with the somatosensory-evoked potential amplitudes.

Another novel imaging techniques for SCI is spinal fMRI. This noninvasive modality relies on changes in blood flow and oxygen levels that occur in metabolically active neural tissue that is responsible for a tested neurological function (57). The signal change is related to both the blood oxygen level—dependent contrast and signal enhancement from extracellular water protons caused by an increase in water content in the area. As such, spinal fMRI is able to map out areas of spinal function related to motor and sensory function. Kornelsen and Macky (56) performed spinal fMRI by using an active and passive lower-limb task

paradigm in a cohort of 12 patients with traumatic SCI. They found that neural activity was present in all patients irrespective of the degree of injury and that both active and passive motion elicited activity below the level of injury.

### Future Clinical Applications

Because there are a rapidly increasing number of experimental studies in which authors investigate cellular therapies in animal SCI models, advanced spinal imaging will likely play a critical role in helping to determine the efficacy of these novel treatments. Some potential applications include: (1) serial DTI to initially assess spinal cord integrity and monitor microstructural changes during therapies; (2) perfusion MRI, which may be useful for determining whether blood flow or perfusion has been re-established in the injured cord after therapy; and (3) PET imaging of [<sup>18</sup>F]-fluoro-2-deoxy-d-glucose, which may be useful for monitoring neural repair, as an increase in glucose uptake might be expected in actively healing spinal tissue.

Several experimental studies have supported the potential role of advanced spinal imaging techniques in the assessment of response to cellular therapy. Schwartz et al. (78) disrupted the rubrospinal tract via cervical funiculotomies in adult rats and then transplanted fibroblasts and vitrogen into the surgical lesion. In addition to discerning between injured and un-injured tracts, DTI was also able to identify glial scarring as well as the orientation of the glial processes. Glial scarring is well recognized as a both a physical and cellular barrier to axonal growth after SCI. Thus, DTI could have a potential role in assessing the efficacy of cellular SCI therapies designed to reduce or inhibit glial scar formation. Ellingson et al. (28) used DTI to monitor the effects of spinal cord regeneration after the transplantation of epidermal neural crest stem cell grafts. Their study showed increased diffusion anisotropy and decreased mean diffusivity at the site of injury in treated animals, suggesting structural and functional improvement of the cord.

Despite the promise of novel imaging techniques for SCI such as spinal DTI and fMRI, there are some limitations to these techniques. Presently, these modalities are not practiced at the vast majority of

institutions, so their overall impact will be limited until more widely adopted. In comparison with the brain, the spinal cord is very small and has a greater chance of artificial contamination related to the surrounding osseous and ligamentous structures, particularly in the setting of traumatic injury. These techniques are sensitive to patient and physiological spinal cord movement, and methods such as cardiac gating are required to minimize distortion.

## CONCLUSION

In summary, advanced imaging of the spinal cord has tremendous potential to provide patient-specific physiological information about the status of cord integrity and health. Advanced spinal cord imaging is still at early stages of development and clinical implementation, but is likely to play an increasingly important role in the management of spinal cord health in the foreseeable future.

## REFERENCES

- Anderson DK, Means ED, Waters TR, Spears CJ: Spinal cord energy metabolism following compression trauma to the feline spinal cord. *J Neurosurg* 53:375-380, 1980.
- Andersson PB, Perry VH, Gordon S: The kinetic and morphological characteristics of the macrophage-microglial response to kainic acid-induced neuronal degeneration. *Neuroscience* 42:201-214, 1991.
- Balentine JD: Pathology of experimental spinal cord trauma. *Lab Invest* 39:236-253, 1978.
- Bammer R, Fazekas F, Augustin M, Simbrunner J, Strasser-Fuchs S, Seifert T, Stollberger R, Hartung HP: Diffusion-weighted MR imaging of the spinal cord. *AJNR Am J Neuroradiol* 21: 587-591, 2000.
- Barrett CP, Guth L, Donati EJ, Krikorian JG: Astroglial reaction in gray matter of lumbar segments after midthoracic transection of the adult rat spinal cord. *Exp Neurol* 73:365-377, 1981.
- Betz RR, Gelman AJ, DeFilipp GJ, Mesgarzadeh M, Clancy M, Steel HH: Magnetic resonance imaging (MRI) in the evaluation of spinal cord injured children and adolescents. *Paraplegia* 25:92-99, 1987.
- Bilgen M, Abbe R, Narayana PA: Dynamic contrast-enhanced MRI of experimental spinal cord injury: in vivo serial studies. *Magn Reson Med* 45:614-622, 2001.
- Bilgen M, Dogan B, Narayana PA: In vivo assessment of blood-spinal cord barrier permeability: serial dynamic contrast enhanced MRI of spinal cord injury. *Magn Reson Imaging* 20:337-341, 2002.
- Bilgen M, Narayana PA: A pharmacokinetic model for quantitative evaluation of spinal cord injury with dynamic contrast-enhanced magnetic resonance imaging. *Magn Reson Med* 46:1099-1106, 2001.
- Bisdas S, Rumboldt Z, Surlan K, Koh TS, Deveikis J, Spampinato MV: Perfusion CT measurements in healthy cervical spinal cord: feasibility and repeatability of the study as well as interchangeability of the perfusion estimates using two commercially available software packages. *Eur Radiol* 18:2321-2328, 2008.
- Blakemore WF: Pattern of remyelination in the CNS. *Nature* 249:577-578, 1974.
- Blakemore WF: Remyelination by Schwann cells of axons demyelinated by intraspinal injection of 6-aminocaproic acid in the rat. *J Neurocytol* 4: 745-757, 1975.
- Blight AR, Decrescito V: Morphometric analysis of experimental spinal cord injury in the cat: the relation of injury intensity to survival of myelinated axons. *Neuroscience* 19:321-341, 1986.
- Bunge RP, Puckett WR, Becerra JL, Marcillo A, Quencer RM: Observation on the pathology of human spinal cord injury. A review and classification of 22 new cases with details from a case of chronic cord compression with extensive focal demyelination. *Adv Neurol* 59:75-89, 1993.
- Chang Y, Jung TD, Yoo DS, Hyun JK: Diffusion tensor imaging and fiber tractography of patients with cervical spinal cord injury. *J Neurotrauma* 27: 2033-2040, 2010.
- Cheran S, Shanmuganathan K, Zhuo J, Mirvis SE, Aarabi B, Alexander MT, Gullapalli RP: Correlation of MR diffusion tensor imaging parameters with ASIA motor scores in hemorrhagic and nonhemorrhagic acute spinal cord injury. *J Neurotrauma* 28:1881-1892, 2011.
- Cohen DM, Patel CB, Ahobila-Vajjula P, Sundberg LM, Chacko T, Liu SJ, Narayana PA: Blood-spinal cord barrier permeability in experimental spinal cord injury: dynamic contrast-enhanced MRI. *NMR Biomed* 22:332-341, 2009.
- Cohen-Adad J, El Mendili MM, Lehericy S, Pradat PF, Blanche S, Rossignol S, Benali H: Demyelination and degeneration in the injured human spinal cord detected with diffusion and magnetization transfer MRI. *Neuroimage* 55: 1024-1033, 2011.
- Crowe MJ, Bresnahan JC, Shuman SL, Masters JN, Beattie MS: Apoptosis and delayed degeneration after spinal cord injury in rats and monkeys. *Nat Med* 3:240, 1997.
- Demaerel P, Sunaert S, Wilms G: Sequences and techniques in spinal MR imaging. *JBR-BTR* 86: 221-222, 2003.
- Demir A, Ries M, Moonen CT, Vital J, Dehais J, Arne P, Caille J, Dousset V: Diffusion-weighted MR imaging with apparent diffusion coefficient and apparent diffusion tensor maps in cervical spondylotic myelopathy. *Radiology* 229:37-43, 2003.
- Deo AA, Grill RJ, Hasan K, Narayana PA: In vivo serial diffusion tensor imaging of experimental spinal cord injury. *J Neurosci Res* 83:801-810, 2006.
- Djang WT: Radiology of acute spinal trauma. *Crit Care Clin* 3:495-518, 1987.
- Ducker TB, Saleman M, Perot PL, Balentine D: Experimental spinal cord trauma. I. Correlation of blood flow, tissue oxygen, and neurologic status in the dog. *Surg Neurol* 10:60-63, 1978.
- Duhamel G, Callot V, Cozzone PJ, Kober F: Spinal cord blood flow measurement by arterial spin labeling. *Magn Reson Med* 59:846-854, 2008.
- Dusart I, Schwab ME: Secondary cell death and the inflammatory reaction after dorsal hemisection of the rat spinal cord. *Eur J Neurosci* 6: 712-724, 1994.
- Ellingson BM, Prost RW, Ulmer JL, Schmit BD: Morphology and morphometry in chronic spinal cord injury assessed using diffusion tensor imaging and fuzzy logic. *Conf Proc IEEE Eng Med Biol Soc* 1:1885-1888, 2006.
- Ellingson BM, Schmit BD, Gourab K, Sieber-Blum M, Hu YF, Schmainda KM: Diffusion heterogeneity tensor MRI ( $\alpha$ -Dti): mathematics and initial applications in spinal cord regeneration after trauma. *Biomed Sci Instrum* 45:167-172, 2009.
- Ellingson BM, Ulmer JL, Kurpad SN, Schmit BD: Diffusion tensor magnetic resonance imaging of the neurologically intact human spinal cord. *AJNR Am J Neuroradiol* 29:1279-1284, 2008.
- Ellingson BM, Ulmer JL, Schmit BD: A new technique for imaging the human spinal cord in vivo. *Biomed Sci Instrum* 42:255-260, 2006.
- Ellingson BM, Ulmer JL, Schmit BD: Morphology and morphometry of human chronic spinal cord injury using diffusion tensor imaging and fuzzy logic. *Ann Biomed Eng* 36:224-236, 2008.
- Endo T, Suzuki S, Utsunomiya A, Uenohara H, Tominaga T: Prediction of neurological recovery using apparent diffusion coefficient in cases of incomplete spinal cord injury. *Neurosurgery* 68: 329-336, 2011.
- Facon D, Ozanne A, Fillard P, Lepointre J, Tournoux-Facon C, Ducreux D: MR diffusion tensor imaging and fiber tracking in spinal cord compression. *AJNR Am J Neuroradiol* 26: 1587-1594, 2005.
- Finnerup NB, Gyldensted C, Nielsen E, Kristensen AD, Bach FW, Jensen TS: MRI in chronic spinal cord injury patients with and without central pain. *Neurology* 61:1569-1575, 2003.
- Fishman PS, Kelley JP: The fate of severed corticospinal axons. *Neurology* 34:1161-1167, 1984.

36. Flanders AE, Schaefer DM, Doan HT, Mishkin MM, Gonzalez CF, Northrup BE: Acute cervical spine trauma: correlation of MR imaging findings with degree of neurologic deficit. *Radiology* 177:25-33, 1990.
37. Ford JC, Hackney DB: Numerical model for calculations of apparent diffusion coefficient (ADC) in permeable cylinders—comparison with measured ADC in spinal cord white matter. *Magn Reson Med* 37:387-394, 1997.
38. Ford JC, Hackney DB, Alsop DC, Jara H, Joseph PM, Hand CM, Black P: MRI characterization of diffusion coefficients in a rat spinal cord injury model. *Magn Reson Med* 31:488-494, 1994.
39. Ford JC, Hackney DB, Joseph PM, Phelan M, Alsop DC, Tabor SL, Hand CM, Markowitz RS, Black P: A method for in vivo high resolution MRI of rat spinal cord injury. *Magn Reson Med* 31:218-223, 1994.
40. Ford JC, Hackney DB, Lavi E, Phillips M, Patel U: Dependence of apparent diffusion coefficients on axonal spacing, membrane permeability, and diffusion time in spinal cord white matter. *J Magn Reson Imaging* 8:775-782, 1998.
41. Fraidakis M, Klason T, Cheng H, Olson L, Spenger C: High-resolution MRI of intact and transected rat spinal cord. *Exp Neurol* 153:299-312, 1998.
42. Gareau PJ, Weaver LC, Dekaban GA: In vivo magnetization transfer measurements of experimental spinal cord injury in the rat. *Magn Reson Med* 45:159-163, 2001.
43. Goldberg AL, Rothfus WE, Deeb ZL, Daffner RH, Lupetin AR, Wilberger JE, Prostko ER: The impact of magnetic resonance on the diagnostic evaluation of acute cervicothoracic spinal trauma. *Skeletal Radiol* 17:89-95, 1988.
44. Guha A, Tator CH, Rochon J: Spinal cord blood flow and systemic blood pressure after experimental spinal cord injury in rats. *Stroke* 20:372-377, 1989.
45. Hagg T, Oudega M: Degenerative and spontaneous regenerative processes after spinal cord injury. *J Neurotrauma* 23:264-280, 2006.
46. Hajnal JV, Kasuboski L, deSouza NM, Bydder GM: Magnetic resonance imaging: spinal cord imaging with the turbo-fluid attenuated inversion recovery (FLAIR) pulse sequence. *Clin Radiol* 50:1-5, 1995.
47. Harrison BM, McDonald WI: Remyelination after transient experimental compression of the spinal cord. *Ann Neurol* 1:542-551, 1977.
48. Hayashi N, De La Torre JC, Green BA: Regional spinal cord blood flow and tissue oxygen content after spinal cord trauma. *Surg Forum* 31:461-463, 1980.
49. Holtz A, Nystrom B, Gerdin B: Spinal cord blood flow measured by <sup>14</sup>C-iodoantipyrine autoradiography during and after graded spinal cord compression in rats. *Surg Neurol* 31:350-360, 1989.
50. Inglis BA, Yang L, Wirth ED 3rd, Plant D, Mareci TH: Diffusion anisotropy in excised normal rat spinal cord measured by NMR microscopy. *Magn Reson Imaging* 15:441-450, 1997.
51. Ito T, Oyanagi K, Wakabayashi K, Ikuta F: Traumatic spinal cord injury: a neuropathological study on the longitudinal spreading of the lesions. *Acta Neuropathol (Berl)* 93:13-18, 1997.
52. Kalfas I, Wilberger J, Goldberg A, Prostko ER: Magnetic resonance imaging in acute spinal cord trauma. *Neurosurgery* 23:295-299, 1988.
53. Kalil K, Schneider GE: Retrograde cortical and axonal changes following lesions of the pyramidal tract. *Brain Res* 89:15-27, 1975.
54. Kang CE, Clarkson R, Tator CH, Yeung IW, Shoichet MS: Spinal cord blood flow and blood vessel permeability measured by dynamic computed tomography imaging in rats after localized delivery of fibroblast growth factor. *J Neurotrauma* 27:2041-2053, 2010.
55. Kitzman P: Alterations in axial motoneuron morphology in the spinal cord injured spastic cat. *Exp Neurol* 192:100-108, 2005.
56. Kornelsen J, Mackey S: Potential clinical applications for spinal functional MRI. *Curr Pain Headache Rep* 11:165-170, 2007.
57. Kornelsen J, Stroman PW: Detection of the neuronal activity occurring caudal to the site of spinal cord injury that is elicited during lower limb movement tasks. *Spinal Cord* 45:485-490, 2007.
58. Kuker W, Weller M, Klose U, Krapf H, Dichgans J, Nagele T: Diffusion-weighted MRI of spinal cord infarction—high resolution imaging and time course of diffusion abnormality. *J Neurol* 251:818-824, 2004.
59. Kulkarni MV, McArdle CB, Kopanicky D, Miner M, Cotler HB, Lee KF, Harris JH: Acute spinal cord injury: MR imaging at 1.5 T. *Radiology* 164:837-843, 1987.
60. Mamata H, Jolesz FA, Maier SE: Apparent diffusion coefficient and fractional anisotropy in spinal cord: age and cervical spondylosis-related changes. *J Magn Reson Imaging* 22:38-43, 2005.
61. McCreary CR, Bjarnason TA, Skihar V, Mitchell JR, Yong VW, Dunn JF: Multiexponential T<sub>2</sub> and magnetization transfer MRI of demyelination and remyelination in murine spinal cord. *Neuroimage* 45:1173-1182, 2009.
62. Moriya T, Hassan AZ, Young W, Chesler M: Dynamics of extracellular calcium activity following contusion of the rat spinal cord. *J Neurotrauma* 11:255-263, 1994.
63. Mulcahey MJ, Gaughan J, Betz RR, Johansen KJ: The International Standards for Neurological Classification of Spinal Cord Injury: reliability of data when applied to children and youths. *Spinal Cord* 45:452-459, 2007.
64. Nandoe Tewarie RD, Yu J, Seidel J, Rahiem ST, Hurtado A, Tsui BM, Grotenhuis JA, Pomper MG, Oudega M: Positron emission tomography for serial imaging of the contused adult rat spinal cord. *Mol Imaging* 9:108-116, 2010.
65. Norton WT, Aquino DA, Hozumi I, Chiu FC, Brosnan CF: Quantitative aspects of reactive gliosis: a review. *Neurochem Res* 17:877-885, 1992.
66. Paleologos TS, Fratzoglou MM, Papadopoulos SS, Chatzidakis EE, Gouliamos AD, Kourousis DD: Posttraumatic spinal cord lesions without skeletal or discal and ligamentous abnormalities: the role of MR imaging. *J Spinal Disord* 11:346-349, 1998.
67. Pallini R, Fernandez E, Sbriccoli A: Retrograde degenerations of corticospinal axons following transection of the spinal cord in rats. *J Neurosurg* 68:124-128, 1988.
68. Petersen JA, Wilm BJ, von Meyenburg J, Schubert M, Seifert B, Najafi Y, Dietz V, Kollias S: Chronic cervical spinal cord injury: DTI correlates with clinical and electrophysiological measures. *J Neurotrauma* 29:1556-1566, 2012.
69. Qian J, Herrera JJ, Narayana PA: Neuronal and axonal degeneration in experimental spinal cord injury: in vivo proton magnetic resonance spectroscopy and histology. *J Neurotrauma* 27:599-610, 2010.
70. Quencer RM, Sheldon JJ, Post MJ, Diaz RD, Montalvo BM, Green BA, Eismont FJ: MRI of the chronically injured cervical spinal cord. *AJR Am J Roentgenol* 147:125-132, 1986.
71. Ramon y Cajal S: *Degeneration and Regeneration of the Nervous System*. London: Oxford University Press; 1928.
72. Rivlin AS, Tator CH: Regional spinal cord blood flow in rats after severe cord trauma. *J Neurosurg* 49:470-477, 1978.
73. Sagiuchi T, Iida H, Tachibana S, Kusumi M, Kan S, Fujii K: Diffusion-weighted MRI in anterior spinal artery stroke of the cervical spinal cord. *J Comput Assist Tomogr* 27:410-414, 2003.
74. Sagiuchi T, Tachibana S, Endo M, Hayakawa K: Diffusion-weighted MRI of the cervical cord in acute spinal cord injury with type II odontoid fracture. *J Comput Assist Tomogr* 26:654-656, 2002.
75. Sandler AN, Tator CH: Review of the effect of spinal cord trauma on the vessels and blood flow in the spinal cord. *J Neurosurg* 45:638-646, 1976.
76. Schwab ME, Bartholdi D: Degeneration and regeneration of axons in the lesioned spinal cord. *Physiol Rev* 76:319-370, 1996.
77. Schwartz ED, Chin CL, Shumsky JS, Jawad AF, Brown BK, Wehrli S, Tessler A, Murray M, Hackney DB: Apparent diffusion coefficients in spinal cord transplants and surrounding white matter correlate with degree of axonal dieback after injury in rats. *AJNR Am J Neuroradiol* 26:7-18, 2005.



78. Schwartz ED, Duda JT, Shumsky JS, Cooper ET, Gee JC: Spinal cord diffusion tensor imaging and fiber tracking can identify white matter tract disruption and glial scar orientation following lateral funiculotomy. *J Neurotrauma* 22:1388-1398, 2005.
79. Schwartz ED, Hackney DB: Diffusion-weighted MRI and the evaluation of spinal cord axonal integrity following injury and treatment. *Exp Neurol* 184:570-589, 2003.
80. Shanmuganathan K, Gullapalli RP, Zhuo J, Mirvis SE: Diffusion tensor MR imaging in cervical spine trauma. *AJNR Am J Neuroradiol* 29:655-659, 2008.
81. Shepard MJ, Bracken MB: Magnetic resonance imaging and neurological recovery in acute spinal cord injury: observations from the National Acute Spinal Cord Injury Study 3. *Spinal Cord* 37:833-837, 1999.
82. Shi R, Pryor JD: Pathological changes of isolated spinal cord axons in response to mechanical stretch. *Neuroscience* 110:765-777, 2002.
83. Stokes BT, Garwood M, Walters P: Oxygen fields in specific spinal loci of the canine spinal cord. *Am J Physiol* 240:H761-H766, 1981.
84. Tarr RW, Drolshagen LF, Kerner TC, Allen JH, Partain CL, James AE Jr: MR imaging of recent spinal trauma. *J Comput Assist Tomogr* 11:412-417, 1987.
85. Terae S, Taneichi H, Abumi K: MRI of wallerian degeneration of the injured spinal cord. *J Comput Assist Tomogr* 17:700-703, 1993.
86. Totoiu MO, Keirstead HS: Spinal cord injury is accompanied by chronic progressive demyelination. *J Comp Neurol* 486:373-383, 2005.
87. Trafton PG, Boyd CA Jr: Computed tomography of thoracic and lumbar spine injuries. *J Trauma* 24:506-515, 1984.
88. Uchida K, Baba H, Maezawa Y, Furukawa S, Furukawa N, Imura S: Histological investigation of spinal cord lesions in the spinal hyperostotic mouse (twy/twy): morphological changes in anterior horn cells and immunoreactivity to neurotropic factors. *J Neurol* 245:781-793, 1998.
89. Wagner FC Jr, Dohrmann GJ, Bucy PC: Histopathology of transitory traumatic paraplegia in the monkey. *J Neurosurg* 35:272-276, 1971.
90. Walker JG, Yates RR, Yashon D: Regional canine spinal cord energy state after experimental trauma. *J Neurochem* 33:397-401, 1979.
91. Waller A: Experiments on the section of glossopharyngeal and hypoglossal nerves of the frog and observation of the alternatives produced thereby in the structure of their primitive fibres. *Phil Trans R Soc Lond* 140:423, 1850.
92. Wang M, Dai Y, Han Y, Haacke EM, Dai J, Shi D: Susceptibility weighted imaging in detecting hemorrhage in acute cervical spinal cord injury. *Magn Reson Imaging* 29:365-373, 2011.
93. Waxman SG: Demyelination in spinal cord injury. *J Neurol Sci* 91:1-14, 1989.
94. White SJ, Hajnal JV, Young IR, Bydder GM: Use of fluid-attenuated inversion-recovery pulse sequences for imaging the spinal cord. *Magn Reson Med* 28:153-162, 1992.
95. Yin B, Tang Y, Ye J, Wu Y, Wang P, Huang L, Yang R, Shen H: Sensitivity and specificity of in vivo diffusion-weighted MRI in acute spinal cord injury. *J Clin Neurosci* 17:1173-1179, 2010.
96. Young W, Koreh I: Potassium and calcium changes in injured spinal cords. *Brain Res* 365:42-53, 1986.

*Conflict of interest statement: The authors declare that the article content was composed in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.*

*Received 9 July 2012; accepted 7 December 2012; published online 12 December 2012*

*Citation: World Neurosurg. (2014) 82, 6:1351-1358. <http://dx.doi.org/10.1016/j.wneu.2012.12.004>*

*Journal homepage: [www.WORLDNEUROSURGERY.org](http://www.WORLDNEUROSURGERY.org)*

*Available online: [www.sciencedirect.com](http://www.sciencedirect.com)*

*1878-8750/\$ - see front matter © 2014 Elsevier Inc. All rights reserved.*



A FORUM FOR SIX CONTINENTS

**WORLD  
NEUROSURGERY**  
THE OFFICIAL JOURNAL OF THE WORLD FEDERATION OF NEUROLOGICAL SOCIETIES

CONTACT THE  
EDITORIAL OFFICE

WORLD NEUROSURGERY (ISSN 1878-8750) is published monthly by Elsevier Inc., 360 Park Avenue South, New York, NY 10010-1710.

Editorial Correspondence:

Michael L.J. Apuzzo, M.D., Ph.D. (Hon)  
WORLD NEUROSURGERY Editor-in-Chief  
1640 Marengo Street, Suite 701  
Los Angeles, CA 90033  
Tel: (323) 442-3001, Fax: (323) 987-2828  
E-mail: [worldneurosurgery-L@usc.edu](mailto:worldneurosurgery-L@usc.edu)  
Website: [www.WORLDNEUROSURGERY.org](http://www.WORLDNEUROSURGERY.org)