

# UCSF

## UC San Francisco Previously Published Works

### Title

Self-monitoring and reminder text messages to increase physical activity in colorectal cancer survivors (Smart Pace): a pilot randomized controlled trial

### Permalink

<https://escholarship.org/uc/item/4kw8v2pn>

### Journal

BMC Cancer, 19(1)

### ISSN

1471-2407

### Authors

Van Blarigan, Erin L  
Chan, Hilary  
Van Loon, Katherine  
[et al.](#)

### Publication Date

2019-12-01

### DOI

10.1186/s12885-019-5427-5

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>


Peer reviewed

RESEARCH ARTICLE

Open Access



# Self-monitoring and reminder text messages to increase physical activity in colorectal cancer survivors (Smart Pace): a pilot randomized controlled trial

Erin L. Van Blarigan<sup>1,2\*</sup> , Hilary Chan<sup>3</sup>, Katherine Van Loon<sup>4,5</sup>, Stacey A. Kenfield<sup>2</sup>, June M. Chan<sup>1,2</sup>, Emily Mitchell<sup>4</sup>, Li Zhang<sup>1,4,5</sup>, Alan Paciorek<sup>1,5</sup>, Galen Joseph<sup>6</sup>, Angela Laffan<sup>5</sup>, Chloe E. Atreya<sup>4,5</sup>, Yoshimi Fukuoka<sup>7</sup>, Christine Miaskowski<sup>5,7</sup>, Jeffrey A. Meyerhardt<sup>8</sup> and Alan P. Venook<sup>4,5</sup>

## Abstract

**Background:** Over 1.3 million people live with colorectal cancer in the United States. Physical activity is associated with lower risk of colorectal cancer recurrence and mortality. Interventions are needed to increase physical activity in colorectal cancer survivors.

**Methods:** We conducted a 2-arm non-blinded pilot randomized controlled trial at the University of California, San Francisco among 42 individuals who had completed curative-intent treatment for colorectal cancer to determine the feasibility and acceptability of a 12-week (84 days) physical activity intervention using a Fitbit Flex™ and daily text messages. Participants were randomized 1:1 to receive the intervention with print educational materials or print educational materials alone. We explored the impact of the intervention versus usual care on physical activity using ActiGraph GT3X+ accelerometers pre-/post-intervention.

**Results:** We screened 406 individuals and randomized 42 to intervention ( $n = 21$ ) or control ( $n = 21$ ) groups. During the 12-week study, the intervention arm wore their Fitbits a median of 74 days [88% of days in study period, interquartile range: 23–83 days] and responded to a median of 34 (out of 46) text messages that asked for a reply (interquartile range: 13–38 text messages). Among the 16 intervention participants who completed the feedback survey, the majority (88%) reported that the intervention motivated them to exercise and that they were satisfied with their experience. No statistically significant difference in change in moderate-to-vigorous physical activity was found from baseline to 12 weeks between arms.

**Conclusion:** A 12-week physical activity intervention with a Fitbit and text messages was feasible and acceptable among colorectal cancer patients after curative treatment. Larger studies are needed to determine whether the intervention increases physical activity.

**Trial registration:** [Clinicaltrials.gov](https://clinicaltrials.gov) Identifier [NCT02966054](https://clinicaltrials.gov/ct2/show/study/NCT02966054). Registered 17 November 2016, retrospectively registered.

**Keywords:** Fitbit, Digital health, Exercise, Accelerometer, Lifestyle, Intervention, Cancer survivor, Colorectal cancer

\* Correspondence: [erin.vanblarigan@ucsf.edu](mailto:erin.vanblarigan@ucsf.edu)

<sup>1</sup>Department of Epidemiology and Biostatistics, University of California San Francisco, UCSF Box 0560, 550 16th St. 2nd Floor, San Francisco, CA 94158, USA

<sup>2</sup>Department of Urology, University of California San Francisco, San Francisco, CA, USA

Full list of author information is available at the end of the article



## Background

Over 1.3 million individuals currently live with colorectal cancer in the United States (US), representing nearly 10% of all US cancer survivors [1]. Evidence from prospective studies strongly suggests that physical activity after colorectal cancer diagnosis reduces the risk of cancer-specific and overall mortality [2–4]. National guidelines recommend that cancer survivors engage in 150 min per week or more of moderate physical activity [5]. Yet, less than half of colorectal cancer survivors currently meet these recommendations [6]. Interventions that increase physical activity and ultimately improve outcomes among colorectal cancer survivors could provide a great public health benefit.

Digital health tools (e.g., text messaging, physical activity trackers) are a promising approach for increasing physical activity at a lower cost and burden compared to in-person or telephone counseling. A recent systematic review concluded that the majority of randomized controlled trials (RCTs) that included text messaging reported statistically significant improvements in health behaviors (including physical activity) [7]. Most of this research focused on weight loss or management of type II diabetes or cardiovascular disease. A growing number of digital health behavioral interventions have included cancer patients [8], yet colorectal cancer survivors are underrepresented in these studies. In fact, no studies focused on colorectal cancer, and fewer than 30 individuals with colorectal cancer were included in any one study that enrolled multiple cancer types [8].

The benefits of physical activity for colorectal cancer patients is among the most consistent association in the epidemiologic literature of cancer survivorship, yet colorectal cancer survivors are under-represented in lifestyle intervention trials [9, 10]. The lower accrual of colorectal cancer survivors may reflect their lower survival rate (65% 5-y survival vs. 100% for prostate and 89% for breast) as well as the disease's distinct burden on quality-of-life [11, 12]. Colorectal cancer patients experience unique physical and psychosocial challenges due to their disease, such as living with an ostomy and bowel dysfunction [12]. Consequently, the appropriate content and format of a physical activity intervention among colorectal cancer survivors is unknown, and may differ compared to healthy adult populations or populations of other cancer survivors.

Our long-term goal is to conduct a definitive randomized controlled trial to test the effect of a digital health physical activity intervention on clinical outcomes among colorectal cancer survivors. However, the feasibility and acceptability of a digital health physical activity intervention in these patients is still unknown. Therefore, we developed a digital health intervention, using a Fitbit Flex™ and daily text messages, aimed at increasing physical activity after completion of treatment for colorectal cancer.

We conducted a 12-week pilot RCT with 42 colorectal cancer survivors to determine the feasibility and acceptability of the intervention and explore the potential effect of the intervention on accelerometer-measured physical activity.

## Methods


### Study population, consent, and randomization

The University of California, San Francisco (UCSF) Institutional Review Board approved this study. Study data were collected and managed using REDCap electronic data capture tools hosted at UCSF [13].

The target population for this pilot RCT was individuals with non-metastatic colon or rectal cancer who had completed curative therapy. Potentially eligible patients were identified through the Cancer Registry at UCSF or through recruitment in the UCSF Gastrointestinal Oncology Clinic or Gastrointestinal Oncology Survivorship Clinic. Our initial eligibility criteria included: stage II-III colon or rectal cancer, previous completion of cancer-directed therapy  $\geq 3$  months and  $< 2$  years prior to enrollment, disease-free status at enrollment, ability to speak and read English, and ability to reliably access the internet and a mobile phone and navigate websites. We excluded individuals with any of 19 specific contraindications to moderate-to-vigorous physical activity (MVPA; e.g., acute myocardial infarction within six months, ongoing unstable angina, uncontrolled cardiac arrhythmia with hemodynamic compromise, active endocarditis) or who were already very active at baseline (defined as engaging in exercise for  $\geq 30$  min for  $\geq 5$  days a week). Due to initial slow accrual, we expanded the eligibility criteria to include patients with stage I and completely resected stage IV disease and dropped the criteria regarding time after completion of therapy.

We reviewed the medical records of 406 potentially eligible patients (Fig. 1). Of these, 221 (54%) were ineligible, 92 declined to participate (23%), and 51 never responded to a mailed recruitment letter (13%). The primary reasons for ineligibility included: non-English speaking (79; 36%), contraindications to exercise (47; 21%), and baseline physical activity  $\geq 30$  min for  $\geq 5$  days a week (exceeding the intervention goal) (41; 19%). Four patients (2%) were excluded for not having a mobile phone.

After screening, informed consent was obtained electronically from all participants included in the study. We randomized 42 individuals who had completed curative treatment for colon or rectal cancer. Participants were randomized 1:1 to intervention or control using a computer-generated randomization scheme developed by a study biostatistician (LZ) who was blinded to group assignment. This scheme was uploaded to REDCap, and a clinical research coordinator utilized the randomization tool in REDCap to obtain the group assignment for each participant.



**Intervention development**

We conducted a semi-structured, in-person, approximately 90 min, group interview with three colorectal cancer survivors (two women and one man) who met our study eligibility criteria to develop our text message content. The participants were 37, 49, and 52 years of age. During the session, we asked participants about their knowledge of physical activity recommendations for cancer survivors, current physical activity behaviors, barriers to physical activity, and gathered feedback on example text messages. While we may not have obtained a diversity of views or saturation of ideas from three individuals, key suggestions from the session included only asking for one-character responses to text messages (e.g., Y for yes) and sending messages either in the morning or evening so they did not arrive during work or family time.

**Intervention description**

After completing the baseline assessment and randomization, patients in the intervention arm received print education materials on physical activity after cancer, a Fitbit Flex™, and daily text messages [14–20]. Patients were counseled through the print materials and text messages to build up to 150 min per week of moderate activities, such as walking at a brisk pace, or 75 min per week of vigorous activities, such as jogging or biking, and resistance (strength training) exercise 2–3 times per week.

The Fitbit Flex is a popular, relatively inexpensive, and widely available small wristband that tracks physical activity, including steps, distance, active minutes, and calories burned. The Fitbit Flex provides a valid measure of physical activity [21]. The Fitbit Flex wirelessly synchronizes data to a computer, tablet, or phone and provides participants with feedback through a user-friendly website. With

participants' consent, we accessed their Fitbit data at Weeks 4, 8, and 12 of the intervention period by logging onto their accounts and downloading their data. Participants kept their Fitbit after completion of the study.

Participants received daily text messages to their cellular phone. The text messages were based on the Theory of Planned Behavior and results of the group interview [14]. The message content included information about the benefits of physical activity for colorectal cancer survivors, prompts for goal setting/planning, advice and tips for incorporating activity into daily life, and challenges and quizzes to increase engagement (see Additional file 1 for sample text messages). Text messages were sent once per day in the morning (8a or 9a) or evening (6p or 8p).

Overall, the intervention took approximately 120 min per participant for the research staff to set up and manage over the 12-week intervention period.

#### Control arm

After completing the baseline assessment and randomization, control participants received print educational materials about physical activity after cancer. We mailed control participants a Fitbit Flex after completion of the 12-week follow-up accelerometer assessment.

#### Measures

**Feasibility and acceptability assessment** We assessed the feasibility of the intervention through an evaluation of adherence (e.g., Fitbit wear time, response rates to interactive text messages) and attrition (proportion of participants who completed the 12-week follow-up accelerometer assessment). A priori, we stated that the intervention was feasible if we were able to achieve  $\geq 70\%$  adherence and  $\leq 20\%$  attrition in the intervention arm. These values were based on the reported adherence for the CanChange trial, a telephone-delivered health behavior intervention in colorectal cancer survivors [22].

We assessed the acceptability of the intervention with an investigator-developed 14-item questionnaire (see Additional file 2) that participants in the intervention arm completed online using REDCap [13]. Participants were asked to what degree they agreed (strongly agree, agree, undecided, disagree, strongly disagree) with four statements about the text messages and one statement about the Fitbit (e.g., "The [text messages, Fitbit] motivated me to exercise."). For the Fitbit, participants were asked whether they accessed the Fitbit website on a computer and/or via the Fitbit app on their phone; how often they accessed the website and/or app; and whether they would continue to wear the Fitbit after the study ended. Participants were asked how satisfied they were overall with the text messages and Fitbit, separately (i.e., very satisfied, satisfied, neutral, dissatisfied, or very dissatisfied) and

provided with an open text box where they could provide any additional feedback.

**Physical activity assessment** Moderate, moderate-to-vigorous, and vigorous physical activity and daily steps were measured at baseline and 12 weeks in both arms using Actigraph GTX3+ accelerometers [23]. Participants wore the accelerometers around their waist for seven consecutive days. We required at least 3 days of valid wear time, defined as at least 10 h of wear per day [24, 25]. We identified non-wear time using the Troiano 2007 default settings in the ActiLife v6.13.3 software. We used the Troiano 2008 cut-points to identify average min/d engaged in sedentary (0–99 counts per minute), light (100–2019 counts per minute), moderate (2020–5998 counts per minute), and vigorous physical activity (5999 or more counts per minute) [25]. These cut-points were chosen to facilitate comparison with NHANES data, a nationally representative sample. For each patient, we calculated MVPA as the sum of time spent in moderate and vigorous physical activity.

**Adverse events assessment** At 4, 8, and 12 weeks, participants in both arms were asked to complete a brief online health check-in survey to assess potential adverse events. Participants who did not complete the online survey were first sent a reminder email and were subsequently called by research staff to obtain the information.

#### Statistical analysis

To determine if the intervention was feasible, we calculated the proportion of the intervention group who wore their Fitbit Flex  $N$  days out of the total 84 study days and the proportion who responded to  $N$  text messages out of a total of 46 text messages that asked for a reply. We compared these proportions to the nominal level 70% that we set a priori. For attrition, we calculated the proportion of participants who completed the 12-week accelerometer assessment, and compared that to the nominal level 80%, which was set a priori. To describe physical activity and estimate the effect of the intervention on physical activity, we calculated each patients' average minutes per day or steps per day based on the 7-day accelerometer at enrollment and 12-weeks. We next calculated each patients' change in activity levels as the follow-up average minus baseline average, and then for all patients in the intervention arm calculated the mean change and for all patients in the control arm calculated the mean change. We compared the mean change in activity between the two groups using the difference calculated as intervention mean change minus control mean change and 95% confidence interval. Analyses were conducted using SAS v. 9.4.

## Results

We randomized 42 patients with colon or rectal cancer between April 2015 – March 2017 (Fig. 1). Of these 42 randomized patients, 41 participants received their assigned intervention. One patient assigned to the intervention arm was diagnosed with a new primary lung cancer prior to receipt of the intervention and was withdrawn from the study. Follow-up (based on completion and return of the accelerometer at 12-weeks) was 95% complete in the intervention arm and 90% complete in the control arm. The two patients who did not complete the 12-week accelerometer assessment in the control arm withdrew from the study (one due to cancer recurrence and the other due to obtaining her own Fitbit during the study period).

Characteristics of the intervention and control arms at enrollment are presented in Table 1. Both groups were more active according to the Actigraph GT3X+ accelerometer data then compared to their self-reported activity during screening. Overall, the study sample participated in 42 min per day of MVPA at enrollment. In addition, the control arm was more active ( $51 \pm 21$  min per day of MVPA and  $11,830 \pm 4052$  steps per day) than the intervention arm

( $33 \pm 18$  min per day of MVPA and  $9008 \pm 3639$  steps per day) at enrollment.

The intervention was feasible and acceptable to participants. Participants randomized to the intervention arm wore their Fitbits a median of 74 out of 84 days [88% of study days; interquartile range (IQR): 23–83 days] (Fig. 2a). Four participants did not wear their Fitbits (2 had 0 days and 2 had only 1 day of wear), while five participants wore their Fitbits every day.


While participants were overall responsive to the text messages, the response rate decreased over time (Fig. 2b). For example, 15/20 intervention arm participants responded to the first text message, while only 6 responded to the last message that asked for a response. Overall, intervention arm participants responded to a median of 34 out of the 46 text messages that asked for a reply (74%; IQR: 13–38 texts). One participant did not respond to any text messages and one participant responded to every message that asked for a reply.

Consistent with the Fitbit wear time and text message data, participants reported that the intervention was highly acceptable (Fig. 3). Sixteen of the 20 participants in the

**Table 1** Demographic characteristics, clinical factors, and physical activity at enrollment of 41 colorectal cancer survivors in a pilot randomized controlled trial of a physical activity intervention

Characteristic, mean $\pm$ SD or N (%)	Total	Intervention	Control
No. of participants	41	20	21
Age, years	$54 \pm 11$	$56 \pm 12$	$54 \pm 11$
Body mass index, kg/m <sup>2</sup>	$28.4 \pm 5.9$	$29.7 \pm 7.2$	$27.1 \pm 4.3$
Male	17 (41)	8 (40)	9 (43)
Race			
White	30 (73)	14 (70)	16 (76)
African American/Black	1 (2)	1 (5)	0 (0)
Asian	5 (12)	2 (10)	3 (14)
American Indian/Alaskan Native, Native Hawaiian or Pacific Islander, or Other	5 (12)	3 (15)	2 (10)
College degree	38 (93)	17 (85)	21 (100)
Works full-time	26 (63)	14 (70)	12 (57)
Married	20 (49)	9 (45)	11 (52)
Cancer site			
Colon	23 (56)	11 (55)	12 (57)
Rectum	18 (44)	9 (45)	9 (43)
Years since diagnosis	$1.5 \pm 1.5$	$1.8 \pm 1.9$	$1.4 \pm 1.3$
Tumor Stage			
I	8 (20)	4 (20)	4 (19)
II	8 (20)	2 (10)	6 (29)
III	24 (59)	13 (65)	11 (52)
IV	1 (2)	1 (5)	0 (0)
Moderate-to-vigorous physical activity, minutes/day	$42 \pm 21$	$33 \pm 18$	$51 \pm 21$
Steps per day	$10,453 \pm 4066$	$9008 \pm 3639$	$11,830 \pm 4052$





intervention arm completed the feedback questionnaire. Of these, 14 (88%) agreed that the text messages motivated them to exercise, the content of the text messages was interesting to them, and the frequency of the messages (1 per day) was ideal for them. Thirteen (81%) agreed that the timing of the text messages (morning and evening) was ideal for them. Overall, 88% of participants were satisfied or very satisfied with their experience with the text messages and the Fitbit. Fourteen (88%) of the participants said that they would continue to wear the Fitbit after the study ended.

The mean and standard deviation MVPA, moderate physical activity, vigorous activity, and steps per day measured by accelerometer at enrollment and 12 weeks for participants in the intervention and control groups are in Table 2. The intervention group engaged in a mean of 32.9 min/d of MVPA at enrollment and 46.6 min/d of MVPA at 12-weeks. In contrast, the control group engaged in a mean of 50.8 min/d of MVPA at enrollment and 54.5 min/d of MVPA at 12-weeks. On average, the intervention arm increased their MVPA by 13 min per day more than the control arm (mean difference: 13.1 min per day; 95% CI: -13.5, 39.7) (Table 2).

No serious adverse events occurred during the study. The proportions of participants in the intervention and controls arms who reported a priori-identified non-serious

adverse events potentially related to physical activity during the study period are in Table 3.


**Discussion**

In this novel pilot RCT, we developed a feasible and acceptable physical activity intervention for colorectal cancer survivors utilizing a physical activity tracker and daily text messages. On average, patients randomized to the intervention arm wore their Fitbit nearly 90% of the time and responded to 74% of text messages that asked for a reply. The vast majority of patients (88%) reported that they were satisfied with their overall experience with the intervention.

Our study had a number of strengths. We developed an interactive text message program to promote physical activity in colorectal cancer survivors. In addition, we enrolled more colorectal cancer survivors than any prior digital health randomized controlled trial, included a racially diverse study sample (27% minorities), and used an accelerometer to measure physical activity at enrollment and 12-weeks.

Limitations of our study included the small sample size; the high level of physical activity, on average, that participants performed prior to enrollment; and a well-educated sample. We used a high cutoff for baseline activity (i.e., 30 min or more on 5 or more days per week based on self-report) when enrolling patients for this pilot study in order to be more inclusive. However, this meant that some patients did not have to change their activity levels very much to meet the intervention goals. Moreover, in contrast to what has been reported by validation studies, participants’ enrollment accelerometer-measured physical activity was even higher than their self-reported activity during screening [26]. Given that our participants had all volunteered for a physical activity intervention study, we postulate that the act of wearing the monitor motivated these individuals to exercise more compared to their usual activity levels [27]. Future studies should focus on less active patients; screen patients activity levels using accelerometers rather than self-report; and consider stratifying on enrollment activity levels to ensure balanced groups. Lastly, 93% of the participants in our study were college educated. Future studies are needed to determine the feasibility and acceptability of this intervention in individuals with different levels of education and socioeconomic status.

While the intervention was determined to be feasible and acceptable, there are aspects that could be improved in future studies. First, we applied pre-determined cut-points for Fitbit wear and text message response rates to assess feasibility. However, the optimal level of physical activity tracker use (e.g., daily, several times a week) to achieve behavior change is not known [28]. Second, we sent daily text messages in the morning or evening based on input from cancer survivors. However, it is not known



**Fig. 3** Acceptability of 12-weeks of text messages and a Fitbit Flex to individuals with colorectal cancer. 16 intervention arm participants completed the feedback survey. All values reported are N, %

**Table 2** Accelerometer-measured physical activity at enrollment and 12 weeks among 41 colorectal cancer survivors participating in a pilot randomized controlled trial of a Fitbit Flex and daily text messages

Type of Physical Activity	Intervention (n = 20)		Control (n = 19) <sup>a</sup>		Mean difference in change in activity between groups (95% CI)
	Baseline Mean ± SD	12 Weeks Mean ± SD	Baseline Mean ± SD	12 Weeks Mean ± SD	
Moderate-to-vigorous, min/day	32.9 ± 17.9	46.6 ± 48.4	50.8 ± 20.7	54.5 ± 24.9	13.1 (– 13.5, 39.7)
Moderate, min/day	32.1 ± 17.6	43.5 ± 39.0	46.9 ± 18.9	51.7 ± 23.2	9.3 (–12.8, 31.3)
Vigorous, min/day	0.87 ± 0.96	3.1 ± 10.3	3.9 ± 5.2	2.8 ± 3.3	3.8 (– 1.5, 9.1)
Steps per day	9007 ± 3638	10,047 ± 4461	11,829 ± 4052	12,541 ± 5535	649 (– 1581, 2878)

<sup>a</sup>12 week accelerometer data was not available from two participants in the control arm



**Table 3** Non-serious adverse events potentially related to physical activity reported during a 12-week Fitbit and text messaging physical activity intervention among 41 colorectal cancer survivors<sup>a</sup>

	Intervention (n = 20)	Control (n = 21)
Low back pain, N (%)	6 (30)	8 (38)
Knee pain, N (%)	7 (35)	5 (24)
Inflammation of the joints, arthritis, N (%)	3 (15)	5 (24)
Joint pain, arthralgia, N (%)	3 (15)	1 (5)
Chest pain, N (%)	2 (10)	1 (5)
Shortness of breath, N (%)	2 (10)	2 (10)
Fatigue, N (%)	7 (35)	9 (43)
Leg cramping, N (%)	5 (25)	1 (5)
Muscle pain, N (%)	4 (20)	4 (19)

<sup>a</sup>Participants in both arms were asked to report if they had experienced any of the issues listed in Table 3 at 4, 8, and 12 weeks via an online questionnaire

if this frequency and/or timing of messages optimizes participants' adherence with the intervention. In addition, participants provided constructive feedback on ways to improve the intervention. One participant suggested that it would be helpful to individualize the text messages based on what participants were recording with their Fitbits: "someone needed to monitor this [the Fitbit data] so other questions or prompts [via text message] could have been given to me." Future larger studies may be strengthened by having real-time access to participants' Fitbit data (versus downloading the data periodically) and research staff available to customize messages. Lastly, participants agreed that the Fitbit was a "great motivator," but one participant had trouble with the band falling off and one participant did not like wearing it at work due to hygiene concerns. Future studies with a larger sample size focused on optimization of the intervention components, including determining whether both components are necessary for behavior change, would be of interest.

## Conclusions

Overall, our pilot study demonstrated that a digital health physical activity intervention is feasible and acceptable to colorectal cancer survivors. A full-scale randomized controlled trial to determine the effect of a digital health physical activity intervention on physical activity in colorectal cancer survivors is warranted.

## Additional files

**Additional file 1:** Sample text messages in the Smart Pace pilot trial. The first two weeks of text messages sent to participants in the Smart Pace randomized controlled trial. (DOCX 15 kb)

**Additional file 2:** Smart Pace Feedback Survey. Investigator-developed survey to solicit feedback regarding the Fitbit and text messaging intervention. The survey was self-administered online. (PDF 73 kb)

## Abbreviations

BMI: Body mass index; IQR: Interquartile range; MVPA: Moderate-to-vigorous physical activity; RCT: Randomized controlled trial; UCSF: University of California, San Francisco; US: United States

## Acknowledgments

We thank the patients at the UCSF Helen Diller Family Comprehensive Cancer Center for their participation in this research study.

## Funding

This research was supported by the NIH (K07CA197077, KL2TR000143, K08CA175143), Mount Zion Health Fund, and UCSF Helen Diller Family Comprehensive Cancer Center. The funders had no role in the design of the study; collection, analysis, or interpretation of data; or writing the manuscript.

## Availability of data and materials

The dataset analyzed during the current study is available from the corresponding author on reasonable request.

## Authors' contributions

Authors contributed to the manuscript in the following ways: study concept and design (EVB, SAK, JMC, KVL, APV); acquisition, analysis, and interpretation of the data (EVB, HC, KVL, EM, LZ, AP, GJ, AL, CEA); drafting the manuscript (EVB); critical revision of the manuscript for important intellectual content (all authors); statistical analysis (EVB, LZ, AP); obtained funding (EVB); administrative, technical, or material support (APV); and study supervision (YF, CM, JAM, APV). All authors read and approved the final manuscript.

## Ethics approval and consent to participate

The University of California, San Francisco (UCSF) Institutional Review Board approved this study, and all participants provided informed consent.

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Author details

<sup>1</sup>Department of Epidemiology and Biostatistics, University of California San Francisco, UCSF Box 0560, 550 16th St. 2nd Floor, San Francisco, CA 94158, USA. <sup>2</sup>Department of Urology, University of California San Francisco, San Francisco, CA, USA. <sup>3</sup>School of Medicine, University of California San Francisco, San Francisco, CA, USA. <sup>4</sup>Division of Hematology/Oncology, Department of Medicine, University of California San Francisco, San Francisco, CA, USA. <sup>5</sup>Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San Francisco, CA, USA. <sup>6</sup>Department of Anthropology, History, and Social Medicine, University of California San Francisco, San Francisco, CA, USA. <sup>7</sup>Department of Physiological Nursing, University of California San Francisco, San Francisco, CA, USA. <sup>8</sup>Dana-Farber/Partners Cancer Care, Boston, MA, USA.

Received: 22 February 2018 Accepted: 1 March 2019

Published online: 11 March 2019

## References

1. Cancer stat fact sheets. [seer.cancer.gov/statfacts/](http://seer.cancer.gov/statfacts/). Accessed Jan 2018.
2. Meyerhardt JA, Giovannucci EL, Holmes MD, Chan AT, Chan JA, Colditz GA, Fuchs CS. Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol.* 2006;24(22):3527–34.
3. Meyerhardt JA, Heseltine D, Niedzwiecki D, Hollis D, Saltz LB, Mayer RJ, Thomas J, Nelson H, Whittom R, Hantel A, et al. Impact of physical activity

- on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin Oncol*. 2006;24(22):3535–41.
4. Je Y, Jeon JY, Giovannucci EL, Meyerhardt JA. Association between physical activity and mortality in colorectal cancer: a meta-analysis of prospective cohort studies. *Int J Cancer*. 2013;133(8):1905–13.
  5. Rock CL, Doyle C, Demark-Wahnefried W, Meyerhardt J, Courneya KS, Schwartz AL, Bandera EV, Hamilton KK, Grant B, McCullough M, et al. Nutrition and physical activity guidelines for cancer survivors. *CA Cancer J Clin*. 2012;62(4):243–74.
  6. Vallance JK, Boyle T, Courneya KS, Lynch BM. Accelerometer-assessed physical activity and sedentary time among colon cancer survivors: associations with psychological health outcomes. *J Cancer Surviv*. 2015;9(3):404–11.
  7. Allen JK, Stephens J, Patel A. Technology-assisted weight management interventions: systematic review of clinical trials. *Telemed J E Health*. 2014;20(12):1103–20.
  8. Roberts AL, Fisher A, Smith L, Heinrich M, Potts HWW. Digital health behaviour change interventions targeting physical activity and diet in cancer survivors: a systematic review and meta-analysis. *J Cancer Surviv*. 2017.
  9. Ligibel JA, Meyerhardt J, Pierce JP, Najita J, Shockro L, Campbell N, Newman VA, Barbier L, Hacker E, Wood M, et al. Impact of a telephone-based physical activity intervention upon exercise behaviors and fitness in cancer survivors enrolled in a cooperative group setting. *Breast Cancer Res Treat*. 2012; 132(1):205–13.
  10. Demark-Wahnefried W, Morey MC, Sloane R, Snyder DC, Miller PE, Hartman TJ, Cohen HJ. Reach out to enhance wellness home-based diet-exercise intervention promotes reproducible and sustainable long-term improvements in health behaviors, body weight, and physical functioning in older, overweight/obese cancer survivors. *J Clin Oncol*. 2012;30(19):2354–61.
  11. SEER stat fact sheets: Colon and Rectum Cancer. [seer.cancer.gov/statfacts/html/colorect.html](http://seer.cancer.gov/statfacts/html/colorect.html). Accessed Jan 2018.
  12. Drury A, Payne S, Brady AM. Cancer survivorship: advancing the concept in the context of colorectal cancer. *Eur J Oncol Nurs*. 2017;29:135–47.
  13. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377–81.
  14. Ajzen I. From intentions to actions: a theory of planned behavior. In: Action-control: From cognition to behavior. In: Kuhl J, Beckmann J, editors. Berlin, Heidelberg: Springer; 1985. p. 11–39.
  15. Kim BH, Glanz K. Text messaging to motivate walking in older African Americans: a randomized controlled trial. *Am J Prev Med*. 2013;44(1):71–5.
  16. Donaldson EL, Fallows S, Morris M. A text message based weight management intervention for overweight adults. *J Hum Nutr Diet*. 2014; 27(Suppl 2):76–90.
  17. Bouhaidar CM, DeShazo JP, Puri P, Gray P, Robins JL, Salyer J. Text messaging as adjunct to community-based weight management program. *Comput Inform Nurs*. 2013;31(10):469–76.
  18. Fjeldsoe BS, Miller YD, Marshall AL. MobileMums: a randomized controlled trial of an SMS-based physical activity intervention. *Ann Behav Med*. 2010; 39(2):101–11.
  19. Nguyen HQ, Gill DP, Wolpin S, Steele BG, Benditt JO. Pilot study of a cell phone-based exercise persistence intervention post-rehabilitation for COPD. *Int J Chron Obstruct Pulmon Dis*. 2009;4:301–13.
  20. Hurling R, Catt M, Boni MD, Fairley BW, Hurst T, Murray P, Richardson A, Sodhi JS. Using internet and mobile phone technology to deliver an automated physical activity program: randomized controlled trial. *J Med Internet Res*. 2007;9(2):e7.
  21. Alharbi M, Bauman A, Neubeck L, Gallagher R. Validation of Fitbit-flex as a measure of free-living physical activity in a community-based phase III cardiac rehabilitation population. *Eur J Prev Cardiol*. 2016;23(14):1476–85.
  22. Hawkes AL, Chambers SK, Pakenham KI, Patrao TA, Baade PD, Lynch BM, Aitken JF, Meng X, Courneya KS. Effects of a telephone-delivered multiple health behavior change intervention (CanChange) on health and behavioral outcomes in survivors of colorectal cancer: a randomized controlled trial. *J Clin Oncol*. 2013;31(18):2313–21.
  23. Vanhelst J, Mikulovic J, Bui-Xuan G, Dieu O, Blondeau T, Fardy P, Beghin L. Comparison of two ActiGraph accelerometer generations in the assessment of physical activity in free living conditions. *BMC Res Notes*. 2012;5:187.
  24. Skender S, Schrotz-King P, Bohm J, Abbenhardt C, Gigic B, Chang-Claude J, Siegel EM, Steindorf K, Ulrich CM. Repeat physical activity measurement by accelerometry among colorectal cancer patients—feasibility and minimal number of days of monitoring. *BMC Res Notes*. 2015;8:222.
  25. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc*. 2008;40(1):181–8.
  26. Ruiz-Casado A, Alejo LB, Santos-Lozano A, Soria A, Ortega MJ, Pagola I, Fiuzza-Luces C, Palomo I, Garatachea N, Cebolla H, et al. Validity of the physical activity questionnaires IPAQ-SF and GPAQ for Cancer survivors: insights from a Spanish cohort. *Int J Sports Med*. 2016;37(12):979–85.
  27. Lee I, Blair S, Manson J. In: RSJ P, editor. epidemiologic methods in physical activity studies. New York: Oxford University Press; 2009.
  28. Phillips SM, Cadmus-Bertram L, Rosenberg D, Buman MP, Lynch BM. Wearable technology and physical activity in chronic disease: opportunities and challenges. *Am J Prev Med*. 2018;54(1):144–50.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

