

UCLA

UCLA Previously Published Works

Title

Greater VO₂peak is correlated with greater skeletal muscle deoxygenation amplitude and hemoglobin concentration within individual muscles during ramp-incremental cycle exercise

Permalink

<https://escholarship.org/uc/item/4m22355j>

Journal

Physiological Reports, 4(23)

ISSN

2051-817X

Authors

Okushima, Dai
Poole, David C
Barstow, Thomas J
[et al.](#)

Publication Date

2016-12-01

DOI

10.14814/phy2.13065

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

ORIGINAL RESEARCH

Greater $\dot{V}O_{2\text{peak}}$ is correlated with greater skeletal muscle deoxygenation amplitude and hemoglobin concentration within individual muscles during ramp-incremental cycle exercise

Dai Okushima¹, David C. Poole², Thomas J. Barstow², Harry B. Rossiter³, Narihiko Kondo⁴, T. Scott Bowen⁵, Tatsuro Amano⁶ & Shunsaku Koga¹

1 Applied Physiology Laboratory, Kobe Design University, Kobe, Japan

2 Departments of Kinesiology and Anatomy and Physiology, Kansas State University, Manhattan, Kansas

3 Rehabilitation Clinical Trials Center, Division of Respiratory & Critical Care Physiology & Medicine, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, California

4 Laboratory for Applied Human Physiology, Graduate School of Human Development and Environment, Kobe University, Kobe, Japan

5 Department of Internal Medicine & Cardiology, Heart Center, Leipzig University, Leipzig, Germany

6 Faculty of Education, Niigata University, Niigata, Japan

Keywords

Diffusive oxygen potential, fractional oxygen extraction, peak oxygen uptake, time-resolved NIRS.

Correspondence

Shunsaku Koga, Applied Physiology Laboratory, Kobe Design University, 8-1-1 Gakunennishi-machi, Nishi-ku, Kobe, 651-2196, Japan.

Tel: +81 78 795 7447

Fax: +81 78 796 2559

Email: s-koga@kobe-du.ac.jp

Funding Information

D. Okushima was supported by grants from Japan Society for the Promotion of Science (KAKENHI- 15K16476). S. Koga was supported by grants from Japan Society for the Promotion of Science (KAKENHI- 24247046, 26560362).

Received: 25 October 2016; Accepted: 3 November 2016

doi: 10.14814/phy2.13065

Physiol Rep, 4 (23), 2016, e13065,
doi: 10.14814/phy2.13065

Abstract

It is axiomatic that greater aerobic fitness ($\dot{V}O_{2\text{peak}}$) derives from enhanced perfusive and diffusive O_2 conductances across active muscles. However, it remains unknown how these conductances might be reflected by regional differences in fractional O_2 extraction (i.e., deoxy [Hb+Mb] and tissue O_2 saturation [S_tO_2]) and diffusive O_2 potential (i.e., total[Hb+Mb]) among muscles spatially heterogeneous in blood flow, fiber type, and recruitment (*vastus lateralis*, VL; *rectus femoris*, RF). Using quantitative time-resolved near-infrared spectroscopy during ramp cycling in 24 young participants ($\dot{V}O_{2\text{peak}}$ range: ~ 37.4 – 66.4 mL kg⁻¹ min⁻¹), we tested the hypotheses that (1) deoxy [Hb+Mb] and total[Hb+Mb] at $\dot{V}O_{2\text{peak}}$ would be positively correlated with $\dot{V}O_{2\text{peak}}$ in both VL and RF muscles; (2) the pattern of deoxygenation (the deoxy[Hb+Mb] slopes) during submaximal exercise would not differ among subjects differing in $\dot{V}O_{2\text{peak}}$. Peak deoxy [Hb+Mb] and S_tO_2 correlated with $\dot{V}O_{2\text{peak}}$ for both VL ($r = 0.44$ and -0.51) and RF ($r = 0.49$ and -0.49), whereas for total[Hb+Mb] this was true only for RF ($r = 0.45$). Baseline deoxy[Hb+Mb] and S_tO_2 correlated with $\dot{V}O_{2\text{peak}}$ only for RF ($r = -0.50$ and 0.54). In addition, the deoxy[Hb+Mb] slopes were not affected by aerobic fitness. In conclusion, while the pattern of deoxygenation (the deoxy[Hb+Mb] slopes) did not differ between fitness groups the capacity to deoxygenate [Hb+Mb] (index of maximal fractional O_2 extraction) correlated significantly with $\dot{V}O_{2\text{peak}}$ in both RF and VL muscles. However, only in the RF did total [Hb+Mb] (index of diffusive O_2 potential) relate to fitness.

Introduction

For large muscle mass exercise such as cycling or running the maximum pulmonary O_2 flux ($\dot{V}O_{2\text{max}}$) is the product

of whole-body perfusive ($\dot{Q}O_2 = \text{cardiac output } [\dot{Q}] \times \text{arterial } O_2 \text{ concentration } [CaO_2]$) and diffusive (muscle transcapillary) O_2 conductances. The Fick relationship ($\dot{V}O_2 =$

$\dot{Q} \times [\text{CaO}_2 - \text{CvO}_2]$ where \bar{v} is mixed venous), as measured across the lungs, provides a volume-weighted average of fractional O_2 extractions across all body compartments (Benson et al. 2013; Fick 1870); each with its own unique $\dot{Q}\text{O}_2/\dot{V}\text{O}_2$ ratio, and thus effluent venous blood O_2 concentration. Thus, without discrete measurements within each tissue, assessments across the lungs provide little insight into $\dot{Q}\text{O}_2$ -to- $\dot{V}\text{O}_2$ matching and the roles of perfusional and diffusional O_2 transport within each tissue.

During progressive exercise from rest to $\dot{V}\text{O}_{2\text{max}}$ whole-body O_2 extraction increases approximately hyperbolically from 25% to 80% or more according to the approximation (Stringer et al. 2005): $(\text{CaO}_2 - \text{CvO}_2) = (20 \times \dot{V}\text{O}_2) / (1 + \dot{V}\text{O}_2)$ (Whipp and Ward 1982). This relationship supports the observations that: individuals with greater $\dot{V}\text{O}_{2\text{max}}$ demonstrate a greater fractional O_2 extraction than their less aerobically-fit counterparts in both health and pathophysiological conditions (Grassi et al. 2007; Porcelli et al. 2010; Salvadego et al. 2016) and endurance training increases both fractional O_2 extraction and $\dot{V}\text{O}_{2\text{max}}$, although not necessarily in linear proportion (Klausen et al. 1982; Roca et al. 1992; Beere et al. 1999). To date, in human muscles, our understanding of these relationships is based overwhelmingly upon measurements made across the whole exercising limb. Within small regions of active muscle, data from both positron emission tomography imaging (Koga et al. 2014; Heinonen et al. 2015) and near-infrared spectroscopy (NIRS) investigations (Koga et al. 2007, 2011, 2014; Okushima et al. 2015) indicated a substantial heterogeneity of the $\dot{Q}\text{O}_2/\dot{V}\text{O}_2$ ratio. This heterogeneity likely arises, in part, from differential motor unit recruitment (Chin et al. 2011) and fiber-type expression (Johnson et al. 1973). In addition, the degree to which $\dot{Q}\text{O}_2$ and $\dot{V}\text{O}_2$ are well matched within muscles, appears to be greater in trained compared with untrained men (Kalliokoski et al. 2005). However, during cycling exercise from rest to $\dot{V}\text{O}_{2\text{peak}}$, whether or not regional $\dot{Q}\text{O}_2/\dot{V}\text{O}_2$ ratio differs among individuals differing in $\dot{V}\text{O}_{2\text{peak}}$ remains unclear. In whole body and thigh muscle $\dot{Q}\text{O}_2$ and $\dot{V}\text{O}_2$ increase linearly with work rate (Knight et al. 1993; Gifford et al. 2016) and therefore the muscle deoxygenation/fractional O_2 extraction will reach a greater peak in fitter individuals (Ferreira et al. 2006; Gifford et al. 2016; Roca et al. 1992). Based upon this logic, we hypothesize that the different regions of active muscle assessed using NIRS will have a similar overall profile of deoxygenation but the regional peak $\dot{Q}\text{O}_2/\dot{V}\text{O}_2$ values (i.e., deoxy[Hb+Mb]) will correlate with whole-body $\dot{V}\text{O}_{2\text{peak}}$.

Recently we employed a time-resolved (TRS) NIRS system, with adipose tissue thickness correction, to determine muscle optical coefficients and therefore measure absolute deoxy- and total-[Hb+Mb] and tissue O_2 saturation ($S_t\text{O}_2$) (Koga et al. 2011, 2015a; Okushima et al. 2015), which are

not impacted significantly by skin blood flow changes (Koga et al. 2015b). Although TRS-NIRS variables do not directly measure muscle $\dot{V}\text{O}_2$ and $\dot{Q}\text{O}_2$, evidence from direct measurements during muscle contractions suggest that deoxy[Hb+Mb] is an index of regional fractional O_2 extraction in the face of presiding blood flow (i.e., microvascular O_2 partial pressure (Koga et al. 2012)) and femoral venous O_2 saturation (Vogiatzis et al. 2015; Sun et al. 2016). Moreover, the total[Hb+Mb] is an index of microvascular blood volume (Ijichi et al. 2005), which reflects local O_2 diffusing capacity (Groebe and Thews 1990). The purpose of this study was to use the TRS-NIRS technology to investigate O_2 perfusive ($\dot{Q}\text{O}_2/\dot{V}\text{O}_2$) and diffusive profiles across the range of achievable power output in individuals ranging broadly in aerobic capacity. We addressed the following specific hypothesis during maximal ramp cycle ergometry: (1) deoxy[Hb+Mb] and total[Hb+Mb] in both VL and RF muscles at peak power output would be positively correlated with $\dot{V}\text{O}_{2\text{peak}}$, and (2) the pattern of deoxygenation (the deoxy[Hb+Mb] slopes) during submaximal exercise would not differ among subjects differing in $\dot{V}\text{O}_{2\text{peak}}$.

Methods

Subjects

Twenty-four young males participated in this study (age, 22 ± 2 years; height, 174 ± 5 cm; weight, 62 ± 5 kg). All participants were nonsmokers and free of known cardiovascular, respiratory, and metabolic disease. This study was approved by the Human Subjects Committee of Kobe Design University, in compliance with the Declaration of Helsinki. Each participant signed an informed consent form after explanation of experimental procedures and the potential risks and benefits of taking part.

Experimental procedure

Participants started the experimental procedure at least 2 h postprandial and refrained from caffeine, alcohol, and intense exercise for 24 h before testing. All experiments were conducted in environmental chambers (SR-3000, Nagano science, Osaka, Japan) maintained at the ambient temperature of 21–24°C and relative humidity of 40–60%, in order to maintain ambient conditions for the studies across Japanese seasons.

Exercise testing was completed on an electromagnetically braked cycling ergometer (XL-75III, Combi, Tokyo, Japan) in the upright position. The exercise protocol began with 2 min of rest, followed by 4 min of 20 W baseline exercise and then a ramp-incremental exercise test. The ramp increase was 20 W min^{-1} . Participants were asked to maintain pedaling frequency at 60 rpm throughout the exercise

test, and the test was terminated when participants could no longer maintain 60 rpm despite strong verbal encouragement. At the limit of tolerance the power output was reduced to 20 W and the participants were monitored for at least 6 min of active recovery.

Measurement

Pulmonary oxygen uptake ($\dot{V}O_2$)

Gas exchange measurement methods were the same as in previous studies (Koga et al. 2011; Bowen et al. 2013; Spencer et al. 2014; Okushima et al. 2015). The breath-by-breath gas exchange system (AE-300S, Minato-Medical, Osaka, Japan) was calibrated according to the manufacturer's recommendation before each exercise test. Participants breathed through a low resistance mouthpiece containing a hot-wire flowmeter for measurement of inspiratory and expiratory flows and volumes. Inspired and expired gases were continuously sampled from the mouth and O_2 and CO_2 fractional concentration was measured by fast-responding paramagnetic and infrared analyzers, respectively. Gas volume and concentration signals were time-aligned to account for the time lag between the signals to calculate $\dot{V}O_2$ and $\dot{V}CO_2$ on a breath-by-breath basis.

Near-infrared spectroscopy

Absolute values of oxygenated (oxy[Hb+Mb]), deoxygenated (deoxy[Hb+Mb]), and total hemoglobin and myoglobin concentration (total[Hb+Mb]) were sampled from the mid-belly of the *vastus lateralis* (VL) and *rectus femoris* (RF) on the dominant leg by two TRS-NIRS instruments (TRS-20, Hamamatsu photonics KK, Hamamatsu, Japan). These muscles were selected because they represent major contributors to power generation in cycle ergometry, have a somewhat different fiber-type composition (i.e., superficial-deep Type I %, VL 38–47 %, RF 30–42 %, Johnson et al. 1973) and are differentially recruited during maximal ramp exercise (Chin et al. 2011).

Before applying the probes, the skin under the probes was shaved. The NIRS optodes were housed in black rubber holders and fixed to the skin with adhesive tape to minimize incidental movement and intrusion of ambient light on the NIRS detectors. The interoptode spacing between irradiation and detection probes was 3 cm for all measurement sites. The sample rate was set to 1 Hz.

Ultrasonographic imaging

Adipose tissue thickness (ATT) at each NIRS site was measured using B-mode Doppler ultrasound (Logiq 400, GE-Yokogawa Medical Systems, Tokyo, Japan).

Ultrasound images were collected with care to prevent pressure and distortion of the thickness of the skin and adipose tissues under the probe.

Data analysis

Peak pulmonary $\dot{V}O_2$ ($\dot{V}O_{2\text{peak}}$) was determined as the mean value of the last 30 sec during the ramp exercise test. Participants were grouped into those with low aerobic capacity (LO: $n = 8$, $\leq 45 \text{ mL kg}^{-1} \text{ min}^{-1}$), mid-range (MID, $n = 9$, $46\text{--}55 \text{ mL kg}^{-1} \text{ min}^{-1}$) and those with high aerobic capacity (HI, $n = 7$, $\geq 56 \text{ mL kg}^{-1} \text{ min}^{-1}$) using similar reference ranges as in a previous study (Gifford et al. 2016). Analysis of deoxy- and total[Hb+Mb] was performed after correction for ATT to a thickness of 0 mm using linear regression of the relationship between total[Hb+Mb] and ATT at rest (Bowen et al. 2013; Okushima et al. 2015) ($\text{total [Hb+Mb]} = 15.7 \times [\text{ATT}] + 201$, $r^2 = 0.801$, $P < 0.001$). For NIRS measurements (deoxy- and total[Hb+Mb] and S_tO_2), 1 Hz optical measurements were averaged over 6 data points, resulting in one NIRS datum calculation every 6 sec. The baseline of each NIRS measurement was calculated as the mean value of 60 sec prior to the start of ramp exercise. The absolute value of each NIRS measurement during ramp exercise was calculated at 20 W, 50 W and each 50 W increment thereafter, as well as for each 10 % of peak power (W_{peak}) increment between 10 and 100 % W_{peak} . NIRS variable amplitudes were derived from the difference between baseline and the mean value of the final 18 sec of ramp exercise. The slope of NIRS variables across absolute power (i.e., watt) was determined from linear regression in two regions of the response: below 50 % W_{peak} and above 70 % W_{peak} (Okushima et al. 2015).

Statistical analysis

All values were expressed as mean \pm standard deviation. Pearson product moment correlation was calculated to quantify the relationship between NIRS variables and $\dot{V}O_{2\text{peak}}$. Comparisons of $\dot{V}O_{2\text{peak}}$ and W_{peak} were analyzed by one-way ANOVA with the main effect of fitness (LO, MID, HI). Comparisons of baseline and amplitude in each NIRS variable and ATT were analyzed by two-way repeated-measures ANOVA, with main effects of aerobic fitness and muscle site (VL, RF). Comparison of the temporal profile for NIRS variables was analyzed by two-way repeated-measures ANOVA, with main effects of aerobic fitness and exercise intensity (normalized power output: every 10 % W_{peak} from 10 to 100 % W_{peak} ; absolute power output: 20 W, peak power and every 50 W from 50 to 200 W). Comparison of the slopes of each NIRS variable was made by two-way repeated-measures

ANOVA, with main effects of aerobic fitness and exercise intensity ($<50\%W_{peak}$, $>70\%W_{peak}$). A significant F ratio was analyzed using Bonferroni's post-hoc test. Effect size (ES: using Cohen's *d*) and statistical power ($1-\beta$) were also calculated for the comparison of each NIRS variable. Significance was accepted at $P < 0.05$.

Results

By design, participants achieved a wide range of peak power (W_{peak}), ranging from 204 to 385 W, and a $\dot{V}O_{2peak}$ ranging from 37.4 to 66.4 mL kg⁻¹ min⁻¹ (Table 1). ATT averaged 4.0 ± 1.2 mm and 5.5 ± 1.8 mm for the VL and RF, respectively, and was significantly less for the VL than the RF muscle (Table 1). However, ATT was not significantly different across participants differing in $\dot{V}O_{2peak}$.

Across absolute submaximal power outputs up to 200 W, deoxy[Hb+Mb] was greater in LO than MID at 150 and 200 W ($P < 0.05$) for RF muscle, whereas total[Hb+Mb] and S_tO_2 were greater in HI than MID ($P < 0.05$, except 200 W) and LO ($P < 0.05$, except 100 W) for only the RF muscle (Fig. 1). At peak power output, deoxy[Hb+Mb] in VL and total[Hb+Mb] in RF were greater in HI than LO and MID ($P < 0.05$), while S_tO_2 in VL tended to be less in HI compared to LO ($P = 0.05$). When exercise intensity was normalized (Fig. 2), deoxy[Hb+Mb] was greater in HI than LO at 100 $\%W_{peak}$ and MID at 80–100 $\%W_{peak}$ in the VL ($P < 0.05$, Fig. 2). In addition, S_tO_2 was less in HI than LO at 100 $\%W_{peak}$ and MID at 90 $\%W_{peak}$ in the VL ($P < 0.05$). In the RF muscle, total[Hb+Mb] was greater in HI than LO at 100 $\%W_{peak}$ and greater than MID at all intensities ($P < 0.05$).

The VL muscle evinced a lower baseline deoxy[Hb+Mb] ($P < 0.001$, ES = 0.89, $1-\beta = 0.99$, Table 2) and greater baseline S_tO_2 ($P < 0.001$, ES = 1.05, $1-\beta = 0.97$) compared to the RF muscle, in all $\dot{V}O_{2peak}$ groups. Baseline total[Hb+Mb] was greater in HI than MID for both RF and VL muscles ($P = 0.019$, ES = 1.04, $1-\beta = 0.81$). The

exercise-induced increase in deoxy[Hb+Mb] ($P = 0.041$, ES = 1.08, $1-\beta = 0.81$) and decrease in S_tO_2 ($P = 0.035$, ES = 1.14, $1-\beta = 0.92$) were greater in HI than LO in both VL and RF muscles. The deoxy[Hb+Mb] and S_tO_2 amplitudes were significantly correlated with $\dot{V}O_{2peak}$ (Fig. 3) in both VL and RF muscles. At each muscle site, the deoxy[Hb+Mb] and S_tO_2 baselines (negative correlation) and the amplitude of total[Hb+Mb] (positive correlation) were significantly correlated with $\dot{V}O_{2peak}$ in RF (Figs. 3 and 4) but not VL.

In each aerobic fitness group, deoxy[Hb+Mb] increased systematically during exercise up to $\dot{V}O_{2peak}$ in the RF muscle, but in the VL muscle deoxy[Hb+Mb] plateaued above 70 $\%W_{peak}$ (Figs. 1 and 2, Table 3). Thus, in the VL muscle, deoxy[Hb+Mb] increased far more slowly above 70 $\%W_{peak}$ than below 50 $\%W_{peak}$ irrespective of $\dot{V}O_{2peak}$ ($P = 0.033$, ES = 0.54, $1-\beta = 0.72$). In contrast, in the RF muscle deoxy[Hb+Mb] increased more rapidly above 70 $\%W_{peak}$ than below 50 $\%W_{peak}$ ($P < 0.001$, ES = 1.26, $1-\beta = 0.99$). The total[Hb+Mb] increased systematically to $\dot{V}O_{2peak}$ in the RF muscle but decreased above 70 $\%W_{peak}$ in the VL muscle independent of aerobic capacity group ($P < 0.001$, ES = 1.55, $1-\beta = 1.00$). The S_tO_2 in the VL decreased systematically up to $\dot{V}O_{2peak}$ whereas the S_tO_2 in RF muscle decreased more rapidly above 70 $\%W_{peak}$ than below 50 $\%W_{peak}$ irrespective of aerobic capacity group ($P < 0.001$, ES = 1.21, $1-\beta = 0.99$).

Discussion

This investigation resolved the magnitude and pattern of $\dot{Q}O_2/\dot{V}O_2$ matching and O_2 diffusive potential in the VL and RF muscles across groups of individuals ranging broadly in aerobic capacity using quantitative TRS-NIRS during ramp-incremental cycling exercise. The principal original findings are: (1) the magnitude of absolute deoxy[Hb+Mb] increase (i.e., the amplitude of the exercise-induced reduction in $\dot{Q}O_2/\dot{V}O_2$) and S_tO_2 decrease

Table 1. Participant's characteristics.

	W_{peak} (W)	$\dot{V}O_{2peak}$ (mL kg ⁻¹ min ⁻¹)	Adipose tissue thickness (mm)	
			VL	RF
LO ($n = 8$)	240 ± 30	40.1 ± 2.7	4.8 ± 1.6	6.7 ± 2.0 ³
MID ($n = 9$)	285 ± 30 ¹	50.3 ± 2.8 ¹	3.6 ± 1.0	4.8 ± 1.5 ³
HI ($n = 7$)	325 ± 34 ^{1,2}	61.0 ± 3.6 ^{1,2}	3.7 ± 0.5	5.0 ± 1.4 ³
All ($n = 24$)	282 ± 45	50.0 ± 8.9	4.0 ± 1.2	5.5 ± 1.8

Data are grouped by aerobic capacity ($\dot{V}O_{2peak}$): low (LO, ≤ 45 mL kg⁻¹ min⁻¹); middle (MID, 46–55 mL kg⁻¹ min⁻¹); and high (HI, ≥ 56 mL kg⁻¹ min⁻¹).

¹Indicates a significant difference from LO ($P < 0.05$).

²Indicates a significant difference from MID ($P < 0.05$).

³Indicates a significant difference between VL and RF ($P < 0.05$).




Figure 1. The mean responses of NIRS variables for low (LO, $\dot{V}O_{2peak} \leq 45 \text{ mL kg}^{-1} \text{ min}^{-1}$, open), middle (MID, $46\text{--}55 \text{ mL kg}^{-1} \text{ min}^{-1}$, gray), and high aerobic capacity (HI, $\geq 56 \text{ mL kg}^{-1} \text{ min}^{-1}$, filled) across absolute power. Left and right panels show the NIRS variables in VL and RF muscle, respectively. " α ", " β ", and " γ " indicate the significant difference ($P < 0.05$) between HI and LO, HI and MID, and LO and MID, respectively. " α' " indicates the tendency of significant difference ($P = 0.05$) between HI and LO.

were correlated with $\dot{V}O_{2peak}$ in both VL and RF muscles; (2) a high baseline deoxygenation and S_tO_2 in the RF muscle (i.e., low preexercise $\dot{Q}O_2/\dot{V}O_2$) was negatively associated with $\dot{V}O_{2peak}$; (3) the magnitude of increase in the total[Hb+Mb] (i.e., the exercise-induced increase in O_2 diffusive potential) was positively correlated with $\dot{V}O_{2peak}$ in the RF muscle only; (4) the rate of change in each NIRS variable across absolute work rates was not related to aerobic fitness (i.e., $\dot{V}O_{2peak}$).

The absolute peak amplitude and baseline of muscle deoxygenation

The peak absolute amplitude of deoxy[Hb+Mb] increase (positively) and S_tO_2 decrease (negatively) were correlated

with $\dot{V}O_{2peak}$ for both muscle sites (Table 2, Figs. 1–3) and this represents the first demonstration of this relationship with technology capable of measuring absolute deoxy[Hb+Mb] together with ATT correction. This is consistent with the notion that the capacity for O_2 extraction between the capillary and myocyte relates to the aerobic capacity of active muscle. Previous studies have reported that the systemic arteriovenous O_2 difference and O_2 extraction capacity (Murias et al. 2010) and leg O_2 extraction capacity (Roca et al. 1992; Sala et al. 1999; Proctor et al. 2001) are associated with $\dot{V}O_{2max}$. These data extend that notion to the level of individual muscles. Blomstrand et al. (1997) and Robinson et al. (1994) reported that a greater $\dot{V}O_{2max}$ in human and rat muscle is attributed to a greater O_2 flux capacity, consequent to




Figure 2. The mean responses of NIRS variables for low (LO, $\dot{V}O_{2peak} \leq 45$ mL kg⁻¹ min⁻¹, open), middle (MID, 46–55 mL kg⁻¹ min⁻¹, gray), and high aerobic capacity (HI, ≥ 56 mL kg⁻¹ min⁻¹, filled) across normalized power. Left and right panels show the NIRS variables in VL and RF muscle, respectively. “α” and “β” indicate the significant differences ($P < 0.05$) between HI and LO, and HI and MID, respectively.

a greater muscle mitochondrial enzyme activity. The present results suggest that the increase in the peak amplitude of muscle deoxygenation reflects the peripheral adaptation of fractional O₂ extraction to exercise at each muscle site. In addition, this was not different between VL and RF muscles, irrespective of $\dot{V}O_{2peak}$ (Table 2). Thus, the amplitude of muscle deoxygenation at $\dot{V}O_{2peak}$ may be limited to a similar magnitude among active muscle sites.

Baseline deoxy[Hb+Mb] were greater in RF compared with VL independent of $\dot{V}O_{2peak}$ (Table 2), and was associated with $\dot{V}O_{2peak}$ in RF but not in VL (Fig. 4), extending our previous findings (Chin et al. 2011;

Okushima et al. 2015). This suggests that less aerobically-fit individuals may have a lower $\dot{Q}O_2/\dot{V}O_2$ of leg muscles at rest or during very low exercise intensities. A low $\dot{Q}O_2/\dot{V}O_2$ at baseline is consistent with a small intercept for the $\dot{Q}O_2/\dot{V}O_2$ relationship, which predisposes capillary-myocyte O₂ flux to be highly sensitive to the adjustment in convective O₂ delivery: only a small “underperfusion” would rapidly limit capillary PO₂ (Ferreira et al. 2006; Benson et al. 2013). However, the mechanistic bases for this effect, be they structural (e.g., muscle fiber composition and vascular bed) and/or functional (e.g., regulation of O₂ distribution and muscle recruitment), remain unclear at present.

Table 2. Comparison of baseline and peak amplitude in deoxy- and total[Hb+Mb] and tissue O_2 saturation (S_tO_2) in the *vastus lateralis* (VL) and *rectus femoris* (RF) muscle during incremental cycle ergometry.

	VL	RF
Baseline		
Deoxy[Hb+Mb] ($\mu\text{mol/L}$)		
LO ($n = 8$)	50.8 \pm 4.8	66.0 \pm 8.4 ³
MID ($n = 9$)	48.6 \pm 4.6	57.0 \pm 7.3 ³
HI ($n = 7$)	51.2 \pm 10.7	54.8 \pm 12.6 ³
Total[Hb+Mb] ($\mu\text{mol/L}$)		
LO ($n = 8$)	190.3 \pm 6.8	195.1 \pm 15.0
MID ($n = 9$)	188.3 \pm 15.4	183.9 \pm 13.9
HI ($n = 7$)	196.9 \pm 15.9 ²	204.9 \pm 11.6 ²
S_tO_2 (%)		
LO ($n = 8$)	73.3 \pm 2.2	66.1 \pm 4.8 ³
MID ($n = 9$)	74.1 \pm 2.7	69.1 \pm 2.6 ³
HI ($n = 7$)	75.2 \pm 4.5	73.6 \pm 7.0 ³
Peak amplitude		
Deoxy[Hb+Mb] ($\mu\text{mol/L}$)		
LO ($n = 8$)	25.6 \pm 13.2	23.4 \pm 13.2
MID ($n = 9$)	26.1 \pm 6.8	29.5 \pm 11.3
HI ($n = 7$)	41.4 \pm 17.1 ¹	43.0 \pm 22.7 ¹
Total [Hb+Mb] ($\mu\text{mol/L}$)		
LO ($n = 8$)	24.4 \pm 6.5	14.8 \pm 3.5
MID ($n = 9$)	19.8 \pm 11.0	19.5 \pm 9.3
HI ($n = 7$)	24.9 \pm 20.3	26.4 \pm 11.3
S_tO_2 (%)		
LO ($n = 8$)	-8.9 \pm 6.3	-8.6 \pm 5.8
MID ($n = 9$)	-10.1 \pm 3.2	-11.3 \pm 4.3
HI ($n = 7$)	-16.5 \pm 5.5 ¹	-15.7 \pm 8.4 ¹

Data are grouped by aerobic capacity ($\dot{V}O_{2peak}$): low (LO, $\leq 45 \text{ mL kg}^{-1} \text{ min}^{-1}$); middle (MID, $46\text{--}55 \text{ mL kg}^{-1} \text{ min}^{-1}$); and high (HI, $\geq 56 \text{ mL kg}^{-1} \text{ min}^{-1}$).

¹Indicates a significant difference from LO ($P < 0.05$).

²Indicates a significant difference from MID ($P < 0.05$).

³Indicates a significant difference between VL and RF ($P < 0.01$).

The absolute peak amplitude and baseline of total[Hb+Mb]

The peak absolute amplitude of total[Hb+Mb] for RF, but not for VL, was significantly related to $\dot{V}O_{2peak}$ (Fig. 3). Our previous study reported that RF activation increased slower than that of the VL and was subsequently speeded further from high submaximal exercise intensities up to $\dot{V}O_{2peak}$ (Chin et al. 2011). Another study reported that in the VL muscle there was a relationship between the total[Hb+Mb] plateau onset and the steep reduction in the EMG mean power frequency (MPF) (Boone et al. 2016), where the MPF reduction is thought to reflect a reduction in fast twitch fiber activity related to muscle fatigue (Komi and Tesch 1979)

(although this interpretation of MPF is controversial). These findings support that the RF muscle increases capillary O_2 diffusional capacity by longitudinal capillary recruitment (Poole et al. 2013), whereas the VL muscle is limited in this regard. Thus in the RF muscle the relationship of the absolute total[Hb+Mb] amplitude with $\dot{V}O_{2peak}$ may reflect a higher capacity of hematocrit regulation than for VL. Accordingly, the capacity for diffusive O_2 conductance may be more important for aerobic exercise performance in some muscles (i.e., RF) compared to others (i.e., VL).

The total[Hb+Mb] concentration at submaximal exercise intensities is greater for HI compared with MID in RF muscle (Figs. 1, 2). It is quite possible that this may also reflect differences in muscle capillarity among individual participants.

Kinetics of regional muscle deoxygenation pattern

The slope of NIRS variables measured as a function of absolute work (watts) did not differ among participants varying in $\dot{V}O_{2peak}$ per se (Table 3). Although in the VL muscle each NIRS variable, except S_tO_2 , plateaued above $70 \%W_{peak}$ in the RF muscle these NIRS variables increased up to $\dot{V}O_{2peak}$ systematically or at least more rapidly above $70 \%W_{peak}$ than below $50 \%W_{peak}$, except total[Hb+Mb] (Table 3). This suggests that the balance of $\dot{Q}O_2/\dot{V}O_2$ in active muscle is adjusted according to the metabolic requirements of the myocytes as the exercise intensity increases irrespective of the differential $\dot{V}O_{2peak}$. In previous studies, a controversy has arisen over whether or not the increase in the deoxy[Hb+Mb] slope across normalized power is actually delayed (Boone et al. 2009; Gravelle et al. 2012). In this respect, it is true that, for the whole body, the regression line between $\dot{V}O_2$ and $\dot{Q}O_2$ is largely independent of training status, age and, perhaps, sex differences (Ogawa et al. 1992; Roca et al. 1992; Proctor et al. 1998). Moreover, in rat muscle the slope of the blood flow to $\dot{V}O_2$ is similar between slow and fast twitch muscle fibers (Ferreira et al. 2006). Thus, it may not be requisite to increase the $\dot{Q}O_2/\dot{V}O_2$ ratio within active muscles to elevate $\dot{V}O_{2peak}$.

The marked increase in deoxy[Hb+Mb] in RF muscle close to $\dot{V}O_{2peak}$ suggests that the ability to activate the RF muscle during severe intensity exercise is related to the capacity for high $\dot{V}O_2$'s during cycling exercise (Table 3). Using magnetic resonance T_2 imaging, Reid et al. (2001) reported that trained subjects had a greater activation of their RF muscle during severe compared with moderate intensity exercise. Moreover, Cannon et al. (2013) showed during incremental knee-extension exercise that rates of PCr and pH decline varied by muscle region,




Figure 3. Relationship of the peak amplitude of muscle deoxygenation (deoxy[Hb+Mb]), total tissue hemoglobin and myoglobin concentration (total[Hb+Mb]), and tissue O₂ saturation (S_tO₂) with $\dot{V}O_{2peak}$ in the *vastus lateralis* (left panel) and *rectus femoris* muscles (right panel) during ramp-incremental cycle ergometry. * $P < 0.05$.

with one quadriceps muscle often “leading” others. Thus, supplemental activation of other skeletal muscles (e.g., deeper RF and possibly VM, Okushima et al. 2015) may be requisite if the severe intensity exercise is to be continued.

Methodological consideration

In our preliminary analyses, we fitted NIRS variables with both sigmoidal (Boone et al. 2009; Chin et al. 2011; Spencer et al. 2014) and double linear models (Spencer et al. 2012, 2014; Boone et al. 2016). We found several patterns in the deoxy- and total-[Hb+Mb] responses, for example, a systematic increase, a plateau after reaching the apex, and a decrease after reaching that apex. Therefore we

concluded that it was inappropriate to fit a single model to all deoxy- and total[Hb+Mb] data. This suggests that other factors besides aerobic capacity also exert an influence over muscle deoxygenation and capillary hematocrit responses, for example, muscle fiber distribution, capillarity, and muscle recruitment patterns. Thus, rather than presuming that one (i.e., sigmoidal) model fits all, future investigations should clarify the disparate patterns of muscle deoxygenation extant during ramp exercise.

Limitations

The VL and RF muscles measured in this study are two of the dominant locomotor muscles during cycling exercise.




Figure 4. Relationship of the baseline in muscle deoxygenation (deoxy[Hb+Mb]), total tissue hemoglobin and myoglobin concentration (total [Hb+Mb]), and tissue O_2 saturation (S_tO_2) with $\dot{V}O_{2peak}$ in the vastus lateralis (left panel) and rectus femoris muscles (right panel) during ramp-incremental cycle ergometry. * $P < 0.05$.

These muscles have different EMG activation patterns during incremental cycling (Chin et al. 2011) and their timing of activation is different during a single crank revolution (Heinonen et al. 2015). In addition, there are individual differences in these characteristics due to handle and saddle position, foot position on the pedal, and experience of cycling (Ericson et al. 1985; Takaishi et al. 1998). In our study, we did not measure muscle EMG activity during cycling exercise, and therefore it is not known whether or not the temporal profile of VL and RF activation across the incremental exercise is different among aerobic fitness groups. In addition, we did not assess arterial O_2 saturation, so we do not know whether exercise-induced arterial hypoxemia was greater in the HI group compared with MID and LO (Powers et al. 1989; Harms et al. 2000). Should

arterial desaturation occur, it could impact the muscle deoxy[Hb+Mb] and S_tO_2 responses, potentially contributing to the observed greater muscle deoxygenation in the HI group. The sample size was small for the analysis by two-way repeated-measures ANOVA of some (but not all) variables presented in Figures 1, 2. This increases the potential for type II error; therefore the main effect and interaction in each test of Figures 1, 2 may be an underestimate.

Conclusions

During ramp-incremental cycling exercise the amplitude of absolute deoxy[Hb+Mb] and S_tO_2 for both VL and RF muscles was significantly related to $\dot{V}O_{2peak}$ (i.e., higher and lower, respectively, in individuals with a greater

Table 3. Comparison of the slope per watt of deoxy- and total[Hb+Mb] and S_tO_2 in the *vastus lateralis* (VL) and *rectus femoris* (RF) muscle during incremental cycle ergometry.

	VL		RF	
	<50 % W_{peak}	>70 % W_{peak}	<50 % W_{peak}	>70 % W_{peak}
Deoxy[Hb+Mb] ($\mu\text{mol/L/W}$)				
LO ($n = 8$)	0.15 \pm 0.06	0.05 \pm 0.13 ¹	0.09 \pm 0.03	0.16 \pm 0.10 ¹
MID ($n = 9$)	0.11 \pm 0.05	0.09 \pm 0.05 ¹	0.06 \pm 0.03	0.21 \pm 0.11 ¹
HI ($n = 7$)	0.14 \pm 0.08	0.13 \pm 0.12 ¹	0.09 \pm 0.05	0.25 \pm 0.10 ¹
Total[Hb+Mb] ($\mu\text{mol/L/W}$)				
LO ($n = 8$)	0.24 \pm 0.06	-0.04 \pm 0.11 ¹	0.10 \pm 0.06	0.06 \pm 0.07
MID ($n = 9$)	0.15 \pm 0.05	-0.04 \pm 0.09 ¹	0.04 \pm 0.05	0.08 \pm 0.10
HI ($n = 7$)	0.17 \pm 0.11	-0.09 \pm 0.16 ¹	0.05 \pm 0.06	0.17 \pm 0.12
S_tO_2 (% W)				
LO ($n = 8$)	-0.04 \pm 0.03	-0.03 \pm 0.06	-0.03 \pm 0.01	-0.06 \pm 0.04 ¹
MID ($n = 9$)	-0.03 \pm 0.02	-0.05 \pm 0.02	-0.02 \pm 0.02	-0.09 \pm 0.05 ¹
HI ($n = 7$)	-0.04 \pm 0.02	-0.07 \pm 0.03	-0.04 \pm 0.02	-0.08 \pm 0.02 ¹

Data are grouped by aerobic capacity ($\dot{V}O_{2peak}$): low (LO, $\leq 45 \text{ mL kg}^{-1} \text{ min}^{-1}$); middle (MID, $46\text{--}55 \text{ mL kg}^{-1} \text{ min}^{-1}$); and high (HI, $\geq 56 \text{ mL kg}^{-1} \text{ min}^{-1}$).

¹Indicates a significant difference from <50 % W_{peak} ($P < 0.05$).

$\dot{V}O_{2peak}$). This implies that the capacity for muscle O_2 extraction, down to the individual active muscles, is correlated with whole-body aerobic capacity. A high baseline deoxygenation in the RF muscle was negatively associated with $\dot{V}O_{2peak}$, which implies a greater reliance on increased regional muscle O_2 delivery during exercise rather than enhanced O_2 extraction in low- $\dot{V}O_{2peak}$ individuals (see Ferreira et al. 2006). In addition, the increase in absolute total[Hb+Mb] (an estimate of diffusive O_2 conductance) within the RF muscle was significantly correlated with $\dot{V}O_{2peak}$. These data suggest that the capacity for diffusive O_2 conductance may be more important for aerobic exercise performance in some muscles (RF) than others (VL). Overall, however, the pattern of muscle $\dot{Q}O_2$ -to- $\dot{V}O_2$ balance (i.e., the relative slopes of deoxygenation) and diffusive O_2 potential (total[Hb+Mb]) were less strongly associated with aerobic capacity, while peak deoxy[Hb+Mb] and S_tO_2 (reflecting maximal O_2 extraction) were more strongly related to $\dot{V}O_{2peak}$.

Acknowledgments

The authors express our gratitude to the subjects in this study. We also express our gratitude to Drs. M. D. Spencer and N. Gerrett for very helpful insights during the preparation of the manuscript.

Conflict of Interest

None declared.

References

- Beere, P. A., S. D. Russell, M. C. Morey, D. W. Kitzman, and M. B. Higginbotham. 1999. Aerobic exercise training can reverse age-related peripheral circulatory changes in healthy older men. *Circulation* 100:1085–1094.
- Benson, A. P., B. Grassi, and H. B. Rossiter. 2013. A validated model of oxygen uptake and circulatory dynamic interactions at exercise onset in humans. *J. Appl. Physiol.* 115:743–755.
- Blomstrand, E., G. Radegran, and B. Saltin. 1997. Maximum rate of oxygen uptake by human skeletal muscle in relation to maximal activities of enzymes in the Krebs cycle. *J. Physiol.* 501:455–460.
- Boone, J., K. Koppo, T. J. Barstow, and J. Bouckaert. 2009. Pattern of deoxy[Hb+Mb] during ramp cycle exercise: influence of aerobic fitness status. *Eur. J. Appl. Physiol.* 105:851–859.
- Boone, J., T. J. Barstow, B. Celie, F. Prieur, and J. Bourgeois. 2016. The interrelationship between muscle oxygenation, muscle activation, and pulmonary oxygen uptake to incremental ramp exercise: influence of aerobic fitness. *Appl. Physiol. Nutr. Metab.* 41:55–62.
- Bowen, T. S., H. B. Rossiter, A. P. Benson, T. Amano, N. Kondo, J. M. Kowalchuk, et al. 2013. Slowed oxygen uptake kinetics in hypoxia correlate with the transient peak and reduced spatial distribution of absolute skeletal muscle deoxygenation. *Exp. Physiol.* 98:1585–1596.
- Cannon, D. T., F. A. Howe, B. J. Whipp, S. A. Ward, D. J. McIntyre, C. Ladroue, et al. 2013. Muscle metabolism and activation heterogeneity by combined ³¹P chemical shift and

- T₂ imaging, and pulmonary O₂ uptake during incremental knee-extensor exercise. *J. Appl. Physiol.* 115:839–849.
- Chin, L. M., J. M. Kowalchuk, T. J. Barstow, N. Kondo, T. Amano, T. Shiojiri, et al. 2011. The relationship between muscle deoxygenation and activation in different muscles of the quadriceps during cycle ramp exercise. *J. Appl. Physiol.* 111:1259–1265.
- Ericson, M. O., R. Nisell, U. P. Arborelius, and J. Ekholm. 1985. Muscular activity during ergometer cycling. *Scand. J. Rehabil. Med.* 17:53–61.
- Ferreira, L. F., P. McDonough, B. J. Behnke, T. I. Musch, and D. C. Poole. 2006. Blood flow and O₂ extraction as a function of O₂ uptake in muscles composed of different fiber types. *Respir. Physiol. Neurobiol.* 153:237–249.
- Fick, A. 1870. Über die Messung des Blutquantums in den Herzventrikeln. *Sitz Physik Med. Ges. Würzburg.* 2:16–27.
- Gifford, J. R., R. S. Garten, A. D. Nelson, J. D. Trinity, G. Layec, M. A. Witman, et al. 2016. Symmorphosis and skeletal muscle $\dot{V}O_{2max}$: in vivo and in vitro measures reveal differing constraints in the exercise-trained and untrained human. *J. Physiol.* 594:1741–1751.
- Grassi, B., M. Marzorati, F. Lanfranconi, A. Ferri, M. Longaretti, A. Stucchi, et al. 2007. Impaired oxygen extraction in metabolic myopathies: detection and quantification by near-infrared spectroscopy. *Muscle Nerve* 35:510–520.
- Gravelle, B. M., J. M. Murias, M. D. Spencer, D. H. Paterson, and J. M. Kowalchuk. 2012. Adjustments of pulmonary O₂ uptake and muscle deoxygenation during ramp incremental exercise and constant-load moderate-intensity exercise in young and older adults. *J. Appl. Physiol.* 113:1466–1475.
- Groebe, K., and G. Thews. 1990. Calculated intra- and extracellular PO₂ gradients in heavily working red muscle. *Am. J. Physiol.* 259:H84–H92.
- Harms, C. A., S. R. McClaran, G. A. Nিকেle, D. F. Pegelow, W. B. Nelson, and J. A. Dempsey. 2000. Effect of exercise-induced arterial O₂ desaturation on $\dot{V}O_{2max}$ in women. *Med. Sci. Sports Exerc.* 32:1101–1108.
- Heinonen, I., S. Koga, K. K. Kalliokoski, T. I. Musch, and D. C. Poole. 2015. Heterogeneity of Muscle Blood Flow and Metabolism: Influence of Exercise, Aging, and Disease States. *Exerc. Sport Sci. Rev.* 43:117–124.
- Ijichi, S., T. Kusaka, K. Isobe, K. Okubo, K. Kawada, M. Namba, et al. 2005. Developmental changes of optical properties in neonates determined by near-infrared time-resolved spectroscopy. *Pediatr. Res.* 58:568–573.
- Johnson, M. A., J. Polgar, D. Weightman, and D. Appleton. 1973. Data on the distribution of fibre types in thirty-six human muscles. An autopsy study. *J. Neurol. Sci.* 18:111–129.
- Kalliokoski, K. K., J. Knuuti, and P. Nuutila. 2005. Relationship between muscle blood flow and oxygen uptake during exercise in endurance-trained and untrained men. *J. Appl. Physiol.* 98:380–383.
- Klausen, K., N. H. Secher, J. P. Clausen, O. Hartling, and J. Trap-Jensen. 1982. Central and regional circulatory adaptations to one-leg training. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* 52:976–983.
- Knight, D. R., W. Schaffartzik, D. C. Poole, M. C. Hogan, D. E. Bebout, and P. D. Wagner. 1993. Effects of hyperoxia on maximal leg O₂ supply and utilization in men. *J. Appl. Physiol.* 75:2586–2594.
- Koga, S., D. C. Poole, L. F. Ferreira, B. J. Whipp, N. Kondo, T. Saitoh, et al. 2007. Spatial heterogeneity of quadriceps muscle deoxygenation kinetics during cycle exercise. *J. Appl. Physiol.* 103:2049–2056.
- Koga, S., D. C. Poole, Y. Fukuoka, L. F. Ferreira, N. Kondo, E. Ohmae, et al. 2011. Methodological validation of the dynamic heterogeneity of muscle deoxygenation within the quadriceps during cycle exercise. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 301:R534–R541.
- Koga, S., Y. Kano, T. J. Barstow, L. F. Ferreira, E. Ohmae, M. Sudo, et al. 2012. Kinetics of muscle deoxygenation and microvascular PO₂ during contractions in rat: comparison of optical spectroscopy and phosphorescence-quenching techniques. *J. Appl. Physiol.* 112:26–32.
- Koga, S., H. B. Rossiter, I. Heinonen, T. I. Musch, and D. C. Poole. 2014. Dynamic heterogeneity of exercising muscle blood flow and O₂ utilization. *Med. Sci. Sports Exerc.* 46:860–876.
- Koga, S., T. J. Barstow, D. Okushima, H. B. Rossiter, N. Kondo, E. Ohmae, et al. 2015a. Validation of a high-power, time-resolved, near-infrared spectroscopy system for measurement of superficial and deep muscle deoxygenation during exercise. *J. Appl. Physiol.* 118:1435–1442.
- Koga, S., D. C. Poole, N. Kondo, A. Oue, E. Ohmae, and T. J. Barstow. 2015b. Effects of increased skin blood flow on muscle oxygenation/deoxygenation: comparison of time-resolved and continuous-wave near-infrared spectroscopy signals. *Eur. J. Appl. Physiol.* 115:335–343.
- Komi, P. V., and P. Tesch. 1979. EMG frequency spectrum, muscle structure, and fatigue during dynamic contractions in man. *Eur. J. Appl. Physiol. Occup. Physiol.* 42:41–50.
- Murias, J. M., J. M. Kowalchuk, and D. H. Paterson. 2010. Time course and mechanisms of adaptations in cardiorespiratory fitness with endurance training in older and young men. *J. Appl. Physiol.* 108:621–627.
- Ogawa, T., R. J. Spina, W. H. 3rd Martin, W. M. Kohrt, K. B. Schechtman, J. O. Holloszy, et al. 1992. Effects of aging, sex, and physical training on cardiovascular responses to exercise. *Circulation* 86:494–503.
- Okushima, D., D. C. Poole, H. B. Rossiter, T. J. Barstow, N. Kondo, E. Ohmae, et al. 2015. Muscle deoxygenation in the quadriceps during ramp incremental cycling: Deep vs. superficial heterogeneity. *J. Appl. Physiol.* 119:1313–1319.
- Poole, D. C., S. W. Copp, S. K. Ferguson, and T. I. Musch. 2013. Skeletal muscle capillary function: contemporary

- observations and novel hypotheses. *Exp. Physiol.* 98:1645–1658.
- Porcelli, S., M. Marzorati, F. Lanfrancioni, P. Vago, R. Pisot, and B. Grassi. 2010. Role of skeletal muscles impairment and brain oxygenation in limiting oxidative metabolism during exercise after bed rest. *J. Appl. Physiol.* 109:101–111.
- Powers, S. K., J. Lawler, J. A. Dempsey, S. Dodd, and G. Landry. 1989. Effects of incomplete pulmonary gas exchange on $\dot{V}O_{2\text{max}}$. *J. Appl. Physiol.* 66:2491–2495.
- Proctor, D. N., K. C. Beck, P. H. Shen, T. J. Eickhoff, J. R. Halliwill, and M. J. Joyner. 1998. Influence of age and gender on cardiac output. $\dot{V}O_2$ relationships during submaximal cycle ergometry. *J. Appl. Physiol.* 84:599–605.
- Proctor, D. N., J. D. Miller, N. M. Dietz, C. T. Minson, and M. J. Joyner. 2001. Reduced submaximal leg blood flow after high-intensity aerobic training. *J. Appl. Physiol.* 91:2619–2627.
- Reid, R. W., J. M. Foley, R. C. Jayaraman, B. M. Prior, and R. A. Meyer. 2001. Effect of aerobic capacity on the T_2 increase in exercised skeletal muscle. *J. Appl. Physiol.* 90:897–902.
- Robinson, D. M., R. W. Ogilvie, P. C. Tullson, and R. L. Terjung. 1994. Increased peak oxygen consumption of trained muscle requires increased electron flux capacity. *J. Appl. Physiol.* 77:1941–1952.
- Roca, J., A. G. Agusti, A. Alonso, D. C. Poole, C. Viegas, J. A. Barbera, et al. 1992. Effects of training on muscle O_2 transport at $\dot{V}O_{2\text{max}}$. *J. Appl. Physiol.* 73:1067–1076.
- Sala, E., J. Roca, R. M. Marrades, and J. Alonso. 1999. Gonzalez De Suso JM, Moreno A, Barbera JA, Nadal J, de Jover L, Rodriguez-Roisin R, and Wagner PD. Effects of endurance training on skeletal muscle bioenergetics in chronic obstructive pulmonary disease. *Am. J. Respir. Crit. Care Med.* 159:1726–1734.
- Salvadego, D., M. E. Keramidas, L. Brocca, R. Domenis, I. Mavelli, J. Rittweger, et al. 2016. Separate and combined effects of a 10-d exposure to hypoxia and inactivity on oxidative function in vivo and mitochondrial respiration ex vivo in humans. *J. Appl. Physiol.* 121:154–163.
- Spencer, M. D., J. M. Murias, and D. H. Paterson. 2012. Characterizing the profile of muscle deoxygenation during ramp incremental exercise in young men. *Eur. J. Appl. Physiol.* 112:3349–3360.
- Spencer, M. D., T. Amano, N. Kondo, J. M. Kowalchuk, and S. Koga. 2014. Muscle O_2 extraction reserve during intense cycling is site-specific. *J. Appl. Physiol.* 117:1199–1206.
- Stringer, W. W., B. J. Whipp, K. Wasserman, J. Porszasz, P. Christenson, and W. J. French. 2005. Non-linear cardiac output dynamics during ramp-incremental cycle ergometry. *Eur. J. Appl. Physiol.* 93:634–639.
- Sun, Y., B. S. Ferguson, M. J. Rogatzki, J. R. McDonald, and L. B. Gladden. 2016. Muscle NIRS Signals vs. Venous Blood Hemoglobin Oxygen Saturation in Skeletal Muscle. *Med. Sci. Sports Exerc.* 48:2013–2020.
- Takaishi, T., T. Yamamoto, T. Ono, T. Ito, and T. Moritani. 1998. Neuromuscular, metabolic, and kinetic adaptations for skilled pedaling performance in cyclists. *Med. Sci. Sports Exerc.* 30:442–449.
- Vogiatzis, I., H. Habazettl, Z. Louvaris, V. Andrianopoulos, H. Wagner, S. Zakynthinos, et al. 2015. A method for assessing heterogeneity of blood flow and metabolism in exercising normal human muscle by near-infrared spectroscopy. *J. Appl. Physiol.* 118:783–793.
- Whipp, B. J., and S. A. Ward. 1982. Cardiopulmonary coupling during exercise. *J. Exp. Biol.* 100:175–193.