

UCSF

UC San Francisco Previously Published Works

Title

Mutations in ANKLE2, a ZIKA Virus Target, Disrupt an Asymmetric Cell Division Pathway in *Drosophila* Neuroblasts to Cause Microcephaly

Permalink

<https://escholarship.org/uc/item/4np6183x>

Journal

Developmental Cell, 51(6)

ISSN

1534-5807

Authors

Link, Nichole
Chung, Hyunglok
Jolly, Angad
[et al.](#)

Publication Date

2019-12-01

DOI

10.1016/j.devcel.2019.10.009

Peer reviewed



Published in final edited form as:

Dev Cell. 2019 December 16; 51(6): 713–729.e6. doi:10.1016/j.devcel.2019.10.009.

Mutations in *ANKLE2*, a ZIKA virus target, disrupt an asymmetric cell division pathway in *Drosophila* neuroblasts to cause microcephaly.

Nichole Link^{1,2,3}, Hyunglok Chung^{1,2,3}, Angad Jolly^{2,4}, Marjorie Withers², Burak Tepe^{2,3,5}, Benjamin R. Arenkiel^{2,3,5}, Priya S. Shah⁶, Nevan J. Krogan^{7,8}, Hatip Aydin⁹, Bilgen B. Geckinli¹⁰, Tulay Tos¹¹, Sedat Isikay¹², Beyhan Tuysuz¹³, Ganesh H. Mochida^{14,15,16}, Ajay X. Thomas^{17,18}, Robin D. Clark¹⁹, Ghayda M. Mirzaa^{20,21}, James R. Lupski^{2,22,23}, Hugo J. Bellen^{1,2,3,4,5,24}

¹Howard Hughes Medical Institute, BCM, Houston, TX 77030, USA

²Department of Molecular and Human Genetics (MHG), BCM, Houston, TX 77030, USA

³Neurological Research Institute, Texas Children's Hospital, Houston, TX 77030, USA

⁴MD/PhD Medical Scientist Training Program and MHG Graduate program, BCM, Houston, TX 77030, USA

⁵Program in Developmental Biology. BCM, Houston, TX 77030, USA

⁶Department of Chemical Engineering, Department of Microbiology and Molecular Genetics, University of California, Davis, CA 95616, USA

⁷Department of Cellular and Molecular Pharmacology, University of California, San Francisco, San Francisco, CA 94158, USA

⁸California Institute for Quantitative Biosciences, QB3, University of California, San Francisco, San Francisco, CA 94158, USA

⁹Center of Genetics Diagnosis, Zeynep Kamil Maternity and Children's Training and Research Hospital, Istanbul, Turkey

¹⁰Department of Medical Genetics, Marmara University School of Medicine, Istanbul, Turkey

¹¹Department of Medical Genetics, Dr. Sami Ulus Research and Training Hospital of Women's and Children's Health and Diseases, Ankara, Turkey

Author of correspondence: Hugo J. Bellen, hbellen@bcm.edu.

AUTHOR CONTRIBUTIONS

N.L. and H.J.B. conceived the project and designed experiments, and wrote and revised the manuscript with J.R.L. H.C. performed *in vivo* Ankle2 immunoprecipitations and assessed phosphorylation/total protein levels. A.J. assisted with brain volume measurements. A.J., M.W. and J.R.L. performed primary fibroblast experiments, and human mutation studies. H.A., B.B.G., T.T., S.I., B.T., G.M.M., G.H.M., A.X.J., and R.D.C. ascertained clinical and molecular data of children with variants. B.T., B.A., P.S., and N.K. assisted with Zika virus experiments. N.L. performed all other experiments.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

DECLARATION OF INTERESTS

The authors declare no competing interests.

¹²Department of Physiotherapy and Rehabilitation, Hasan Kalyoncu University, School of Health Sciences, Gaziantep, Turkey

¹³Department of Pediatrics, Istanbul University-Cerrahpasa, Medical Faculty, Istanbul, Turkey

¹⁴Division of Genetics and Genomics, Department of Pediatrics, and Manton Center for Orphan Disease Research, Boston Children's Hospital, Boston, MA 02115, USA

¹⁵Department of Pediatrics, Harvard Medical School, Boston, MA 02115, USA

¹⁶Pediatric Neurology Unit, Department of Neurology, Massachusetts General Hospital, Boston, MA 02114, USA

¹⁷Department of Pediatrics, Section of Neurology and Developmental Neuroscience, BCM, Houston, TX 77030, USA

¹⁸Section of Child Neurology, Texas Children's Hospital, Houston, TX 77030, USA

¹⁹Division of Medical Genetics, Department of Pediatrics, Loma Linda University Medical Center, Loma Linda, CA 92354, USA

²⁰Department of Pediatrics, University of Washington, Seattle, WA 98195, USA

²¹Center for Integrative Brain Research, Seattle Children's Research Institute, Seattle, WA 98105, USA

²²Department of Pediatrics, BCM, Houston, TX; Texas Children's Hospital, Houston, TX 77030, USA


²³Human Genome Sequencing Center, Baylor College of Medicine, Houston, TX 77030, USA

²⁴Lead Contact

SUMMARY

The apical Par complex, which contains aPKC, Bazooka (Par-3), and Par-6, is required for establishing polarity during asymmetric division of neuroblasts in *Drosophila*, and its activity depends on L(2)gl. We show that loss of Ankle2, a protein associated with microcephaly in humans and known to interact with Zika protein NS4A, reduces brain volume in flies and impacts the function of the Par complex. Reducing Ankle2 levels disrupts ER and nuclear envelope morphology, releasing the kinase Ballchen/VRK1 into the cytosol. These defects are associated with reduced phosphorylation of aPKC, disruption of Par complex localization, and spindle alignment defects. Importantly, removal of one copy of *ballchen/VRK1* or *l(2)gl* suppresses *Ankle2* mutant phenotypes and restores viability and brain size. Human mutational studies implicate the above mentioned genes in microcephaly and motor neuron disease. We suggest that NS4A, ANKLE2, VRK1 and LLGL1 define a pathway impinging on asymmetric determinants of neural stem cell division.

Graphical Abstract



eTOC Blurp

The Zika virus protein NS4A interacts with ANKLE2, a protein linked to hereditary microcephaly. Mutations in ANKLE2 also cause microcephaly-like phenotypes in flies. Link et al. now connect these phenotypes with disruption of an asymmetric cell division pathway in fly neuroblasts via an interaction between ANKLE2 and the kinase Bällchen/VRK1.

Keywords

VRK1; aPKC; L(2)gl; Miranda; Bazooka; Ballchen; brain development; congenital infection; MCPH16; NS4A

INTRODUCTION

Proper development of the human brain requires an exquisitely coordinated series of steps and is disrupted in disorders associated with congenital microcephaly. Congenital microcephaly in humans is characterized by reduced brain size (using occipital frontal circumference, OFC, as a surrogate measure) more than two standard deviations below the mean (Z -score < -2) at birth. It is associated with neurodevelopmental disorders, such as developmental delay and intellectual disability and can be caused by external exposures to toxins, *in utero* infections, or gene mutations. Pathogenic gene variants for microcephaly have been identified through targeted genetic testing, genomic copy number studies, and

exome sequencing (ES) (Brunetti-Pierrri *et al.*, 2008; Dumas *et al.*, 2012; Lupski, 2015; Shinawi *et al.*, 2010; Shaheen *et al.*, 2018), identifying 18 primary microcephaly loci. Many syndromes significantly overlap with classic microcephaly phenotypes, and together, these disorders can be caused by defects in a wide variety of biological processes, including centriole biogenesis, DNA replication, DNA repair, cell cycle and cytokinesis, genome stability, as well as multiple cell signaling pathways (Jayaraman *et al.*, 2018). In flies, we refer to microcephalic phenotypes when the third instar brain lobes are reduced in size (Yamamoto *et al.*, 2014) or when adult flies exhibit small heads relative to their body size (Oláhová *et al.*, 2018). As in humans, microcephaly in flies can be a result of mutations that affect cell division and centrosome biology as demonstrated with mutations in WDR62 (Ramdas Nair *et al.*, 2016; Lim *et al.*, 2017) and ASPM/ASP (Rujano *et al.*, 2013), but also those that affect the spindle assembly checkpoint (Poulton *et al.*, 2017) and neuroblast proliferation (Kanai *et al.*, 2018).

A forward, mosaic screen for neurodevelopmental and neurodegenerative phenotypes associated with lethal mutations on the X-chromosome in *Drosophila* identified 165 loci, many with corresponding human genetic disease trait phenotypes (Yamamoto *et al.*, 2014). Among them, a mutation in *Ankle2* (*Ankryin repeat and LEM domain containing 2*) causes loss of Peripheral Nervous System (PNS) organs in adult mutant clones and severely reduced brain size in hemizygous third instar larvae. To identify patients with pathogenic variants in *ANKLE2*, we surveyed the exome database of the Baylor-Hopkins Center for Mendelian Genomics (BHCMG) (Bamshad *et al.*, 2012; Posey *et al.*, 2019) and identified compound heterozygous mutations in *ANKLE2* in two siblings. Both infants exhibited severe microcephaly (Z -score = -9), and the surviving patient displayed cognitive and neurological deficits alongside extensive intellectual and developmental disabilities. We showed that mutations in *Ankle2* led to cell loss of neuroblasts and affected neuroblast division in the developing third instar larval brain. Remarkably, expression of the wild type human *ANKLE2* in flies rescued the observed mutant phenotypes (Yamamoto *et al.*, 2014). Here we explore the molecular pathways and proteins that are affected by *Ankle2* loss.

ANKLE2 belongs to a family of proteins containing LEM (LAP2, Emerin, MAN1) domains that typically associate with the inner nuclear membrane (Lin *et al.*, 2000; Barton *et al.*, 2015). Conventional LEM proteins have been shown to interact with BAF (Barrier to Autointegration Factor), which binds to both DNA and the nuclear lamina (Segura-Totten *et al.*, 2002) to organize nuclear and chromatin structure. However, the LEM domain in *Drosophila* and *C. elegans* *Ankle2* is not obviously conserved (Marchler-Bauer *et al.*, 2017). Studies in *C. elegans* indicate that a homolog of *ANKLE2* regulates nuclear envelope morphology and functions in mitosis to promote reassembly of the nuclear envelope upon mitotic exit (Asencio *et al.*, 2012; Snyers *et al.*, 2018). During this process, *ANKLE2* modulates the activities of VRK1 (Vaccinia Related Kinase 1) and PP2A (Protein Phosphatase 2A) (Asencio *et al.*, 2012). However, all experiments in worms were performed at the embryonic two-cell stage, and no other phenotypes were reported except early lethality. Whilst mutations in *ANKLE2* have been associated with severe microcephaly (OFC z -score = -2.5 to -9), human *VRK1* pathogenic variant alleles can cause a neurological disease trait consisting of complex motor and sensory axonal neuropathy and microcephaly (Gonzaga-Jauregui *et al.*, 2013).

Mutations in both *Ankle2* and the fly homologue of *VRK1*, *ballchen*, cause a loss of neuroblasts in 3rd instar larval brains in *Drosophila* (Yamamoto *et al.*, 2014; Yakulov *et al.*, 2014). Neuroblasts (NBs) divide asymmetrically and are often used as a model to investigate stem cell biology (Homem and Knoblich, 2012) and asymmetric cell division (Gallaud *et al.*, 2017). Most NBs in the larval central brain give rise to another NB and a smaller ganglion mother cell (GMC), which then divides once again to produce neurons or glia. Proper NB maintenance and regulation is essential for precise development of the adult nervous system, and misregulation of NB number or function can lead to defects in brain size (Wang *et al.*, 2009; Gateff and Schneiderman, 1974).

Congenital Zika virus infection in humans during pregnancy has been associated with severe microcephaly that can be as dramatic as certain genetic forms of microcephaly including phenotypes associated with bi-allelic mutations in *MCPH16/ANKLE2* (Moore *et al.*, 2017; Yamamoto *et al.*, 2014). Recently, we showed that a Zika virus protein, NS4A, physically interacts with ANKLE2 in human cells. Expression of NS4A in larval brains causes microcephaly, induces apoptosis, and reduces proliferation. Importantly, expression of human ANKLE2 in flies that express NS4A suppresses the associated phenotypes, demonstrating that NS4A interacts with the ANKLE2 protein and inhibits its function (Shah *et al.*, 2018). Interestingly, Zika virus crosses the blood brain barrier and targets radial glial cells, the neural progenitors in the vertebrate cortex (Devhare *et al.*, 2017; Tang *et al.*, 2016).

Here, we show that Ankle2 is localized to the endoplasmic reticulum and nuclear envelope, similar to NS4A, and genetically interacts with *ball/VRK1* to regulate brain size in flies. An allelic series at the *ANKLE2* and *VRK1* loci shows that perturbation of this pathway results in neurological disease including microcephaly. Our data indicate that the Ankle2-Ball/VRK1 pathway is required for proper localization of asymmetric proteins and spindle alignment during NB cell division by affecting two proteins, aPKC and L(2)gl, that play critical roles in the asymmetric segregation of cell fate determinants. In addition, NS4A expression in neuroblasts mimics phenotypes seen in *Ankle2* mutants, and NS4A induced microcephaly is suppressed by removing a single copy of *ball/VRK1* or *l(2)gl*. Human genomics variant data and disease trait correlations extend this asymmetric cell division pathway from proteins identified in flies and reveal insights into neurological disease. In summary, NS4A hijacks the Ankle2-Ball/VRK1 pathway, which regulates progenitor stem cell asymmetric division during brain development and defines a human microcephaly pathway.

RESULTS

Human ANKLE2 variants cause microcephaly

We previously reported that compound heterozygous variants in *ANKLE2* are associated with microcephaly (Z-score = -9) (*MCPH16*, MIM#616681) in two affected siblings (Yamamoto *et al.*, 2014). Here, we report two additional probands carrying unique variants in *ANKLE2* identified in Seattle (LR17-511 and LR18-033; Figure S1 and S2, Table S1). Brain MRIs of an age matched control (Figure S1A) and a proband with microcephaly from family LR17-511 document one of the more severe cases of microcephaly (Z-score = -8) (Figure S1B). To investigate potential genotype-phenotype correlations, we explored the

spectrum of reported neurological disease trait manifestations associated with *ANKLE2* present in the Baylor Genetics (BG) Laboratories databases. These contain results of clinical exome sequencing (ES) of patients with presumed genetic disorders that remain unsolved for a molecular diagnosis but patients/families have consented to research analyses. We screened for rare biallelic variants, predicted damaging, in *ANKLE2* that fulfill Mendelian expectations for a recessive disease trait. Three families were found to fulfill these criteria in probands with neurologically associated phenotypes (Figure S1D and S2, Table S2). These cases suggest that a diverse set of variants in *ANKLE2* may be associated with a spectrum of neurologic disease and reveal either sporadic disease, apparent vertical transmission, and in some cases, consanguineous parentage (Yamamoto *et al.*, 2014; Shaheen *et al.*, 2018). The identified mutations are missense, nonsense, or splicing variants that lead to premature stop codons; all subjects have biallelic variants, either compound heterozygous or homozygous alleles (Figure S1 and S2). Probands exhibit congenital microcephaly, but some also present with severe brain MRI abnormalities and skin pigmentation abnormalities (Figure S1D). These aggregate data demonstrate that mutations in *ANKLE2* cause autosomal recessive microcephaly.

Null alleles of *Ankle2* are associated with reduced brain size in flies

Given the human genetic implications noted above, we used *Drosophila* to elucidate molecular mechanisms underlying *ANKLE2* associated microcephaly. The mutation originally identified in flies, *Ankle2^A* (L326H), causes reduced brain size in third instar larvae and leads to pupal lethality at temperatures $\geq 22^{\circ}\text{C}$ (Figure S3A). It results in decreased neuroblast number, reduced cell divisions when assessed in MARCM clones, and a high incidence of apoptotic cell death throughout the brain, including neuroblasts and neurons (Yamamoto *et al.*, 2014). However, first instar larvae are born with a slight, but not significant, reduction in the number of neuroblasts or Dpn positive cells (Figure S3B and C) indicating that *Ankle2* loss causes a premature differentiation, loss of stem cell markers, or death of neuroblasts.

To create a severe loss of function allele for *Ankle2*, we integrated a CRIMIC construct containing *attP-FRT-SA-3XSTOP-polyA-3xP3-EGFP-FRT-attP* sequences using CRISPR-Cas9 in the fifth intron shared by all isoforms (Figure 1A; pM14; (Lee *et al.*, 2018). The construct leads to a truncated transcript that likely corresponds to a null allele (*Ankle2^{CRIMIC}*, Figure 1A). These animals die as 3rd instar larvae (Figure 1B), are smaller than wild type and *Ankle2^A* animals, and show a very severely reduced brain volume (Figure 1E versus I, and M) with complete disruption of brain morphology, especially the optic lobe (Figure 1I). Since the severe phenotypes make it difficult to assess the biological questions presented below, we assayed the majority of phenotypes with the hypomorphic *Ankle2^A* missense allele.

To determine whether *Ankle2* is expressed in the brain, we used the CRIMIC allele to introduce an artificial exon that contains SA-GFP-SD in frame which produces a tagged fusion protein (*Ankle2^{GFP}*, Figure 1A–B) (Lee *et al.*, 2018). We readily detect *Ankle2^{GFP}* protein in brains of heterozygous animals (Figure S4A). However, homozygous animals are lethal and exhibit very small brains, indicating that integration of this exon disrupts protein

function. Based on complementation tests, the strength of the allelic series is $Ankle2^A < Ankle2^{CRIMIC} = Ankle2^{IGFP}$ (Figure 1B and C). We therefore used recombineering (Venken *et al.*, 2006) to add a C-terminal GFP tag to Ankle2 in a bacterial artificial chromosome (BAC CH321–85N12; referred to as *Ankle2-GFP*, Figure 1A and D) (Venken *et al.*, 2008; Venken *et al.*, 2009). When this P[acman] clone was introduced in all three *Ankle2* mutant backgrounds, *Ankle2-GFP* rescued brain phenotypes and lethality of these alleles (Figure 1B, G, and J). Hence, the chromosomes carrying the three *Ankle2* alleles do not carry second site mutations that affect brain size or viability and the tagged protein is likely to reflect the endogenous Ankle2 protein distribution.

The human reference *ANKLE2* gene rescues lethality and small brain phenotypes of *Ankle2^A* animals when expressed ubiquitously (*da-GAL4 > UAS-hANKLE2*, Figure 1H, M) or in neuroblasts only (*insc-GAL4 > UAS-hANKLE2*, Figure 1N). To determine whether the microcephaly associated mutations in human ANKLE2 are loss of function alleles, we next expressed *ANKLE2 p.L573V*, *ANKLE2 p.Q782**, *ANKLE2 p.A109P*, *ANKLE2 p.G201W*, in *Ankle2^A* mutant animals. The p.Q782*, p.A109P, and p.G201W variants failed to rescue lethality or reduced brain sizes (Figure 1L–M) consistent with them being severe loss-of-function variant alleles. However, p.L573V restored both viability and brain size (Figure 1K, M) in some *Ankle2^A* animals (Figure 1M), suggesting that this variant is a mild hypomorphic allele.

Ankle2 localizes to the ER and nuclear envelope and is required for their integrity

The endogenously tagged Ankle2^{IGFP} (Figure S4A) as well as the tagged genomic rescue construct, *Ankle2-GFP* (Figure 2A–B, Figure S4B–G) show that *Ankle2* is expressed in most tissues including larval discs (Figure S4B–C), embryos (Figure S4D–F), and cells of the third instar larval brain. Neuroblasts, ganglion mother cells, and neurons (Figure S4G) all show high expression of Ankle2. The protein appears to be localized to the cytoplasm of all cells including neuroblasts (arrows in Figure 2A–B). However, in a subset of cells, the protein is clearly enriched at the nuclear envelope (arrowhead). To determine the dynamics of Ankle2 protein localization, we performed live imaging. As shown in Movie S1, the protein is recruited to the nuclear envelope at the initiation of mitosis and remains associated with the nuclear envelope until briefly after cytokinesis. Indeed, Ankle2 is enhanced at the nuclear envelope in cells positive for phospho-histone H3 (pHH3), a mitotic marker (Figure S4H–I, arrowhead). Hence, Ankle2 is localized to the nuclear envelope in neuroblasts undergoing mitosis.

To determine precisely where Ankle2 is localized, we performed live imaging of brains from animals carrying *Ankle2-GFP* and a transgene that labels the ER: *da-Gal4 > UAS-Sec61β-tdTomato* (Summerville *et al.*, 2016). In neuroblasts (large cell in Figure 2C–E), the Ankle2 protein fully colocalizes with Sec61β at the nuclear envelope as well as the ER. In the surrounding neurons (small cells), much of the cytoplasm is co-labeled. We also counter-stained fixed samples with Calnexin 99a, another ER marker (Riedel *et al.*, 2016). Again, Ankle2 localizes to the nuclear envelope and the ER, but in fixed samples, the ER structure is less obvious than in live imaging (Figure 2G–I).

To determine if Ankle2 is required for proper ER structure, we performed live imaging of *Ankle2^A* mutant neuroblasts expressing Sec61 β -tdTomato (Figure 2F). When compared to wild type (Figure 2E), *Ankle2^A* mutants display highly aberrant Sec61 β localization in many NBs (25°C). The neuroblast in Figure 2F appears larger than normal and displays Sec61 β folds within the nucleus. In addition, we stained fixed *Ankle2^A* mutant neuroblasts with Calnexin 99a and found that *Ankle2^A* mutants also display irregular Calnexin 99a localization (Figure 2J versus 2I), similar to defects shown in Figure 2F, suggesting that even a partial loss of Ankle2 disrupts ER and possibly nuclear envelope structure. Indeed, the morphology of the nuclear envelope is aberrant and convoluted in some *Ankle2^A* mutant cells when stained with Lamin Dm0 (Riemer *et al.*, 1995), a nuclear envelope marker (compare Figure 2N with Figure 2K–M). Hence, Ankle2 is required for proper ER and nuclear envelope morphology.

Ankle2 mutations affect the asymmetric localization of neuroblast determinants

Due to the reduced cell proliferation and reduced neuroblast number in *Ankle2^A* third instar brains (Yamamoto *et al.*, 2014), we sought to explore neuroblast division in more detail. Neuroblast polarity during division relies on the function of the highly conserved apically localized Par complex, which consists of Bazooka (Baz, Par-3) (Schober *et al.*, 1999), Par-6 (Petronczki and Knoblich, 2001), and atypical Protein Kinase C (aPKC) (Rolls *et al.*, 2003). Once activated, the Par complex is responsible for restricting Miranda (Mira) and other cell fate determinants to the basal domain of neuroblasts. After division in most neuroblasts, the basal domain will become the ganglion mother cell, which divides again to produce neurons or glia (Betschinger *et al.*, 2003; Atwood and Prehoda, 2009). Several proteins have been implicated in regulating the Par complex (Chabu and Doe, 2009; Andersen *et al.*, 2012; Bonaccorsi *et al.*, 2007; Atwood *et al.*, 2007), including those associated with cell cycle regulation (Chabu and Doe, 2008; Lee *et al.*, 2006; Wang *et al.*, 2007; Wang *et al.*, 2006).

A previous inspection of dividing neuroblasts revealed no obvious defects in Mira staining (Yamamoto *et al.*, 2014). However, a quantitative analysis of dividing *Ankle2^A* neuroblasts stained with anti-Baz, Par-6, aPKC, and Mira revealed severe localization defects of these proteins in greater than 40% of metaphase neuroblasts during asymmetric division (Figure 3A–L, quantified in Figure 3M–P) in both *Ankle2^A* and trans-heterozygous animals (*Ankle2^A/Ankle2^{CRIMIC}*). These defects are rescued by the genomic construct (Figure 1D), *Ankle2-GFPR* (Figure 3D, H, L, M–P). Finally, we performed live imaging of 3rd instar larval brains of wild type (Movie S2) and *Ankle2^A* mutants labeled with Mira-RFP and Histone-GFP (Movies S3–S5). As shown in Movies S3–S5, neuroblasts exhibit abnormal Mira localization as well as instances of failed division including DNA segregation defects, chromatin bridges, and cytokinesis defects (Movies S3–S4).

For proper neuroblast division to occur, cells must not only asymmetrically localize Par complex members and cell fate determinants, they must also align the mitotic spindle so that divisions segregate cell fate determinants to the proper daughter cell (Cabernard and Doe, 2009). In wild type neuroblasts, the mitotic spindle is aligned parallel to the polarity axis (Figure 3Q, T). Our initial observations in *Ankle2^A* mutants suggested no defects in mitotic spindle alignment in many cells (Yamamoto *et al.*, 2014), but after a quantitative analysis of

metaphase neuroblasts, we noted that spindle alignment appeared disrupted in some *Ankle2^A* mutant cells. To quantify these defects, we measured the axis of division using DNA and Centrosomin (CNN) (Lucas and Raff, 2007) to highlight centrosome placement relative to the localization of cell polarity proteins aPKC and Mira (Figure 3Q–U). Surprisingly, we found that nearly 40% of *Ankle2^A* mutant neuroblasts contained supernumerary centrosomes (Figure 3S). In the remaining 60% of *Ankle2^A* mutant metaphase neuroblasts with obvious aPKC/Mira localization, we also found varying degrees of mitotic spindle alignment defects (compare Figure 3Q–R and 3T–U), showing that Ankle2 is also required for proper spindle alignment in neuroblast division. Together, these results show that Ankle2 plays a prominent role in asymmetric protein localization, spindle alignment, and cell division of neuroblasts.

Ankle2 interacts with VRK1/Ballchen

A *C. elegans* homologue of Ankle2, Lem4L, was previously shown to physically and genetically interact with VRK1, the homologue of Ballchen (Ball) in flies (Asencio *et al.*, 2012). Lem4L and VRK1 in worms localize to the nuclear envelope of the 2-cell stage embryo (Asencio *et al.*, 2012). In contrast, Ball appears to be nuclear during interphase/prophase in all *Drosophila* cells (Figure S3D) (Yakulov *et al.*, 2014) as well as in mammalian cells (Figure 4G–H), and both fly and human proteins have nuclear localization signals. Interestingly, human *VRK1* pathogenic variants cause reduced brain size and microcephaly as well as axonal neuropathy in affected patients (Gonzaga-Jauregui *et al.*, 2013; Renbaum *et al.*, 2009). Hence, to characterize the relationship between Ankle2 and Ball/VRK1, we analyzed the expression and localization of Ball and Ankle2 during neuroblast cell division (Figure 4A–D). During interphase, Ankle2 and Ball do not co-localize as Ankle2 is in the cytoplasm and ER whereas Ball is in the nucleus (Figure 4A). During the mitotic prophase, Ankle2 accumulates at the nuclear envelope but the proteins do not seem to co-localize (Figure 4B). However, at metaphase, the nuclear envelope becomes fragmented but it does not completely dissociate in flies (Katsani *et al.*, 2008). Ankle2 can still be seen localized to the fragmented nuclear lamina, and Ball is briefly present throughout the cytoplasm (Figure 4C). Yet, after telophase, Ball is quickly recruited back to the nucleus and briefly enriched at the nuclear envelope (Figure 4D; Movie S6). After mitosis and once the chromatin is no longer condensed, Ball is nuclear. Interestingly, the spatial restriction of Ball in *Ankle2^A* mutants during interphase and prophase is abolished in many neuroblasts as Ball localizes throughout the cell, a phenotype that is not observed in wild type brains (Figure 4E–F, quantified in Figure S3E). In summary, Ankle2 is required for proper nuclear localization of Ball in *Drosophila*.

To determine whether ANKLE2 regulates Ball/VRK1 subcellular localization in human cells, we assayed VRK1 localization in human fibroblasts. In reference human primary fibroblasts (parental variant p.L573V/+), VRK1 is localized to the nucleus (Figure 4G) consistent with published literature indicating that VRK1 is nuclear localized (Nichols and Traktman, 2004). However, fibroblasts from microcephaly patients carrying compound heterozygous variants in *ANKLE2* (p.L573V/p.Q782* and p.V229G/p.V229G) display significantly reduced VRK1 intensity in the nucleus (Figure 4H, quantified in Figure 4I) and increased cytoplasmic staining in nondividing cells (arrows in Figure 4H) with no significant

change in overall VRK1 intensity (Figure S3F). These data argue for a conserved role between fruit flies and human for ANKLE2 in restricting VRK1 to the nucleus.

Given that Ankle2 is required to maintain Ball/VRK1 in the nucleus during interphase, it is possible that Ball/VRK1 is ectopically active in the cytoplasm of *Ankle2^A* mutants and inhibits or promotes phosphorylation of proteins not normally encountered in the biological homeostatic state. Reducing the level of Ball/VRK1 may therefore alleviate the phenotype associated with the reduction in Ankle2 protein. Indeed, we observe evidence for strong dominant interactions between *Ankle2^A* and *ball* (multiple alleles). *Ankle2^A* animals are pupal lethal and have reduced brain volumes (compare Figure 4J to 4K). However, removal of one copy of *ball*, akin to a heterozygous deletion CNV resulting in haploinsufficiency in human, restores brain development (Figure 4L–M) and suppresses the lethality of *Ankle2^A* mutants (Figure 4N). Importantly, loss of one copy of *ball* (*ball^{e107}*) in *Ankle2^A* mutants also restores the asymmetric protein localization of aPKC and Mira crescents in metaphase neuroblasts (Figure 4O–P). Hence, a partial reduction of Ball/VRK1 activity rescues *Ankle2^A* mutants, providing strong evidence for a gene dosage sensitive locus. However, removing both copies of *ball* in wild type animals leads to pupal lethality (Cullen *et al.*, 2005), causes severely reduced brain volumes in 3rd instar larvae (Herzig *et al.*, 2014), and does not rescue *Ankle2^A* animals, emphasizing that the gene dosage and balance of the protein levels is critical. Indeed, a severe loss of function allele, *Ankle2^{CRIMIC}*, cannot be suppressed by reducing Ball/VRK1 activity (Figure 4N). In summary, these data demonstrate that both Ankle2 and Ball/VRK1 control the distribution of asymmetric determinants, and experimental evidence reveals an antagonistic relationship between both proteins.

The Ankle2-Ball pathway modulates aPKC and L(2)gl

Due to the similarities in defects observed with loss of *Ankle2* or *aPKC*, including mislocalization of Par-6 and Mira (Kim *et al.*, 2009), decreased cell divisions, and reduced neuroblast clone volume (Rolls *et al.*, 2003), we hypothesized that the activity of aPKC, an important mediator of neuroblast asymmetric division (Figure 5A), might be affected. aPKC phosphorylation (Kim *et al.*, 2009) or abundance could be modulated by Ankle2. We therefore assessed both total and phosphorylated aPKC levels in third instar larval brains using an antibody specific for human p-aPKC T410 (T422 in flies). This phosphorylation site is located in its activation loop and was shown to be important for its kinase activity (Kim *et al.*, 2009). Phosphorylation of aPKC (T422) relative to total aPKC is decreased in *Ankle2* mutants (Figure 5B) and is restored with either addition of Ankle2-GFP or reduction of *ball* (Figure 5B), consistent with the data presented in Figure 4. However, overexpression of aPKC or constitutively active aPKC (aPKC^N) (Betschinger *et al.*, 2003) in *Ankle2^A* mutants did not rescue brain size or viability (data not shown).

aPKC has been shown to physically interact with L(2)gl (Betschinger *et al.*, 2003), a regulator of apico-basal polarity that inhibits the function of aPKC (Atwood and Prehoda, 2009; Wirtz-Peitz *et al.*, 2008). *aPKC* and *l(2)gl* genetically interact as removal of one copy of *aPKC* suppresses *l(2)gl* loss of function phenotypes (Rolls *et al.*, 2003), and aPKC has been shown to phosphorylate L(2)gl to control its plasma membrane or cortical release

(Betschinger *et al.*, 2003). When aPKC is active, L(2)gl is phosphorylated and released from the cortex; once released, it no longer binds to aPKC or inhibits its function. Because aPKC and L(2)gl interact, the Ankle2-Ball pathway may affect L(2)gl. We therefore assessed whether L(2)gl physically interacts with the Ankle2-Ball pathway using immunoprecipitation of a GFP-tagged L(2)gl from third instar larval brains and found that Ball indeed interacts with L(2)gl (Figure 5C).

The reduced aPKC activity that we observe may be associated with a gain of function of L(2)gl. Therefore, to determine whether removal of one copy of *l(2)gl* suppresses *Ankle2* associated phenotypes, we introduced a temperature sensitive mutation of *l(2)gl* (*l(2)gl^{ts3}*) into the *Ankle2^A* mutant background and found that reducing *l(2)gl* in *Ankle2^A* mutants at 22°C (Figure 5D–F) and 25°C indeed partially restored brain size. *Ankle2^A* is pupal lethal at 22°C, but when combined with a heterozygous *l(2)gl* mutant allele, some *Ankle2^A* animals survive to adulthood. However, unlike the removal of one copy of *ball*, these animals die a few days after eclosion. In summary, Ankle2 and Ball interact with the apical-basal polarity regulators aPKC and L(2)gl (Figure 5A) and affect aPKC and L(2)gl activity by disturbing the asymmetric segregation of apical-basal polarity factors in neuroblasts. Our data suggest that in *Ankle2* mutants, L(2)gl acts as a gain of function (increased activity), and aPKC activity is reduced. Therefore, by reducing L(2)gl activity, aPKC function is restored.

Disease associated variants in *VRK1* and its paralogs

Ten families have been described with biallelic variants in *VRK1* that cause a spectrum of neurologic diseases including 6 individuals with microcephaly (Feng *et al.*, 2018; Gonzaga-Jauregui *et al.*, 2013; Najmabadi *et al.*, 2011; Nguyen *et al.*, 2015; Renbaum *et al.*, 2009; Shaheen *et al.*, 2018; Stoll *et al.*, 2016) (Table S1; Figure S5). The family structures suggest either a sporadic or recessive neurological disease trait; historical consanguinity in 3/10 pedigrees implicate an autosomal recessive locus. Screening the BHCMG and BG databases identified two additional families with potentially biallelic variants in *VRK1*. (Table S2, Figure S5). These cases suggest that like *ANKLE2* (Figure S1), a heterogenous set of variant alleles in *VRK1* are associated with neurologic disease and microcephaly.

It was previously shown that fly genes with more than one human homolog, especially those that are evolutionarily conserved, have an enriched association with Online Mendelian Inheritance in Man (OMIM) disease phenotypes (Yamamoto *et al.*, 2014), which is a comprehensive and authoritative catalogue of human genes and genetic phenotypes associated with mendelian disorders. We searched the BHCMG database to establish if damaging variants in paralogs of *VRK1* are associated with disease. Predicted deleterious, biallelic variants were found in two paralogs of *VRK1*: *VRK2* is associated with very small eyes and *VRK3* with severe microcephaly (Table S2, Figure S6).

NS4A targets the Ankle2 pathway

Drosophila has been developed as a model of viral infection (Harsh *et al.*, 2018; Liu *et al.*, 2018), and we recently showed that expression of the Zika virus protein NS4A results in reduced brain size in *Drosophila* (Shah *et al.*, 2018). Strikingly, NS4A expression in *Ankle2^A/+* heterozygous animals leads to a more severe phenotype than NS4A expression in

a wild type background, and these animals display brain phenotypes that mimic *Ankle2^{CRIMIC}* null mutants (Shah *et al.*, 2018). These data again suggest that levels of Ankle2 protein are critical. In our previous experiments, we were unable to detect NS4A using immunohistochemistry. We therefore generated an additional NS4A construct with a C-terminal HA tag (*UAS-NS4A-HA*) and drove expression with *insc-GAL4>UAS-NS4A-HA*. As shown in Figure 6A, NS4A is localized to the nuclear envelope and ER (Figure 6A), similar to the subcellular localization of Ankle2 (Figure 2). Expression of NS4A may cause brain defects by affecting aPKC and Miranda localization. Indeed, expression of NS4A in neuroblasts (*insc-GAL4>UAS-NS4A*) affects the apical aPKC localization and leads to an expansion of the Mira domain (Figure 6, compare B to C-D and quantified in Figure 6F-G). In the metaphase neuroblasts that express NS4A, we also note spindle orientation defects in some cells (Figure 6J-K), similar to *Ankle2^A* animals shown in Figure 3. These data indicate that NS4A targets the Ankle2 pathway; this is further strengthened with the observation that when NS4A is expressed in neuroblasts of *ball* heterozygous animals, aPKC and Mira crescents are restored to their wild type patterns (Figure 6E-G) and spindle orientation defects are rescued (compare Figure 6L with K). Finally, ubiquitous expression of NS4A-HA (Figure 6H-I) using two different *Act-GAL4* insertions causes reduced brain volume (Figure 6H) that is rescued by removal of one copy of *ball*, (Figure 6I) co-expression of human ANKLE2, or (Figure 6I) removal of one copy of *l(2)gl*. In summary, the Zika virus protein NS4A targets the Ankle2 pathway and affects asymmetric distribution of cell fate determinants, leading to defects in neuroblast division and brain development.

DISCUSSION

We investigated the biological basis for *ANKLE2* associated microcephaly. We report six additional patients with microcephaly that carry mutations in *ANKLE2* and show that three variants identified in probands cause a loss of *ANKLE2* function when tested in flies (Figure 1 and S1), providing compelling evidence that its loss causes reduced brain size in flies and severe microcephaly (Z-score < -2.5) in humans. *Ankle2* is a dosage sensitive locus whose product is inhibited by the Zika virus protein NS4A. We show that Ankle2, like NS4A, is localized to the ER, and that it targets the nuclear envelope during mitosis. Loss of *Ankle2* affects the nuclear envelope and ER distribution and results in a redistribution of Ball/VRK1, a kinase that is normally localized to the nucleus except when the nuclear envelope breaks down during mitosis (Figure 4). Loss of *Ankle2* disrupts the localization of neuroblast apical-basal polarity determinants such as aPKC, Par-6, Baz, and Miranda, and aPKC phosphorylation is reduced by *Ankle2* mutations. Importantly, loss of one copy of *ball* or *l(2)gl* suppresses the reduced brain volume associated with a partial loss of *Ankle2*, suggesting that much of the biological function of Ankle2 is modulated by aPKC and *l(2)gl*. Finally, the negative influence of NS4A on the activity of ANKLE2 can also be suppressed by removal of one copy of *ball* or *l(2)gl*, suggesting the following pathway: NS4A ⊣ ANKLE2 ⊣ Ball/VRK1 → *l(2)gl*/LLGL1 ⊣ aPKC. This pathway regulated by ANKLE2 plays an important role in neuroblast stem cell divisions in flies and microcephaly and potentially other neurological disease phenotypes in humans.

Interestingly, the above pathway links environmental cues with several genetic causes of sporadic and autosomal recessive microcephaly in human; moreover, it implicates this

pathway in microcephaly accompanying congenital infection. As one example of the latter, the Zika virus has been shown to cross the infant Blood Brain Barrier (Mlakar *et al.*, 2016) and has been identified in radial glial cells (Li *et al.*, 2016), as well as intermediate progenitor cells and neurons (Lin *et al.*, 2017). We propose that NS4A affects the function of Ankle2 leading to the release of Ball/VRK1 from the nucleus. We speculate that this in turn affects the phosphorylation of aPKC and L(2)gl directly by masking phosphorylation sites or indirectly by promoting the activity of one or more phosphatases. Loss of VRK1 has been shown to cause microcephaly and some variant alleles are also associated with pontocerebellar hypoplasia (PCH) in humans (Gonzaga-Jauregui *et al.*, 2013; Renbaum *et al.*, 2009), consistent with the loss of *ball* in flies that causes a severe reduction in brain size (Yakulov *et al.*, 2014). Note that ANKLE2, VRK1, LLGL1, and aPKC, as well as other components of the apical complex like PARD3 are all present in radial glial cells during cortical development (Ayoub *et al.*, 2011). These data suggest that ANKLE2 and its partners such as LLGL1 and asymmetric determinants are important proteins during neural cell proliferation and that the proper levels and relative amounts of these proteins determine how many neurons will eventually be formed in vertebrates. Our data also indicate that variant alleles at either *ANKLE2* or *VRK1* are responsible for some causes of embryonic lethality and severe congenital microcephaly.

LLGL1 has recently been shown to play an important role in radial glia in mice during neurogenesis, and its loss in clones increases the number of divisions (Beattie *et al.*, 2017). In addition, aPKC ζ/λ localizes at the apical membrane of proliferating neural stem cells in chicken embryos during division and has been shown to provide an instructive signal for apical assembly of adherens junctions (Ghosh *et al.*, 2008). Mouse knockouts of *aPKC λ* (Soloff *et al.*, 2004) and *aPKC ζ* (Seidl *et al.*, 2013) are embryonic lethal; however *aPKC ζ* knockouts are viable (Leitges *et al.*, 2001), perhaps suggesting redundant functions within the atypical PKC family. These proteins have not been linked to microcephaly in mice, but conditional removal of an apical complex protein Pals1 in cortical progenitors resulted in complete cortex loss (Kim *et al.*, 2010). Finally, Numb is asymmetrically localized by the Par complex protein in *Drosophila*, segregated to the daughter cell during asymmetric cell division (Wirtz-Peitz *et al.*, 2008), and essential for daughter cells to adopt distinct fates (Bhalerao *et al.*, 2005). In mice, Numb localization is also asymmetric and null mutations exhibit embryonic lethality, neural tube closure defects, and premature neuron development (Zhong *et al.*, 2000). These data indicate that asymmetric division may be important for vertebrate neuronal development, but microcephaly is not a phenotype that typically associates with loss of the mice homologues of asymmetric localized determinants identified in *Drosophila*. However, the observations reported here indicate that the ANKLE2/PAR complex pathway is evolutionarily conserved from flies to human, although the precise mechanisms remain to be determined as different cells may use this pathway in different contexts (Suzuki and Ohno, 2006).

In order to determine whether predicted deleterious biallelic variants in PAR complex encoding genes or their paralogs associated with a neurologic disease trait, we searched the BHCMG database for mutations associated with neurological disease. We found homozygous predicted deleterious missense variants in *PARD3B* (c.1222G>A, p.G408S) in a patient that has microcephaly (Table S2, Figure S6) and compound heterozygous

mutations in *PAR3B* (c.1654G>A;p.A552T) that are associated with other neurological defects (Table S2, Figure S6). The human orthologue of *L(2)gl*, *LLGL1*, is deleted in Smith-Magenis syndrome (SMS) (Smith *et al.*, 1986) and 86–89% of the SMS patients have brachycephaly (Greenberg *et al.*, 1996). These observations extend the mutational load beyond *ANKLE2* and *VRK1* and suggest an association between congenital disease and variants within the PAR complex (Table S2, Figure S6) potentially by a compound inheritance gene dosage model (Wu *et al.*, 2015).

Aurora A (AurA) kinase has been shown to phosphorylate the Par complex (Wirtz-Peitz, Nishimura and Knoblich, 2008) as well as *L(2)gl* (Carvalho *et al.*, 2015) and regulates cortical polarity and spindle orientation in neuroblasts (Lee *et al.*, 2006; Wang *et al.*, 2006). The aberrant localization of Ball/VRK1 in *Ankle2* mutants may lead to gain of function phenotypes that are highly dosage sensitive, as they can be repressed by removing a single copy of Ball/VRK1 in *Ankle2^A*. Mislocalized Ball/VRK1 may mask or interfere with the function of AurA in neuroblast asymmetric division as they share similar kinase substrate consensus sequences (Sanz-García *et al.*, 2011; Ferrari *et al.*, 2005). Future studies are needed to assess Ball/VRK1 redundancy or interference with AurA function.

Another possible evolutionarily parallel with implications in multicellular organismal development is the genetic interaction between the *C. elegans* homologue of VRK1 and an ANKLE2-like protein at the two cell stage (Asencio *et al.*, 2012). Whereas VRK1 in both *Drosophila* and humans (Nichols and Traktman, 2004) is localized to the nucleus, except during mitosis when the nuclear envelope is broken down (Figure 4), the worm VRK1 protein is localized to the nuclear envelope. The worm ANKLE2-like protein, Lem-4L, also interacts with the phosphatase PP2A (Asencio *et al.*, 2012), and the fly PP2A regulates neuroblast asymmetric division by interacting with aPKC and excluding it from the basal cortex (Chabu and Doe, 2009; Ogawa *et al.*, 2009; Wang *et al.*, 2009). PP2A also antagonizes the phosphorylation of Baz by PAR-1 to control apical-basal polarity in dividing embryonic neuroblasts (Krahn *et al.*, 2009) and regulates Baz localization in other cells such as neurons (Nam *et al.*, 2007). This raises the possibility that the Ankle2 pathway also acts with PP2A in neuroblast asymmetric division.

Here, we identified a pathway that plays a significant role in neuroblast asymmetric division. By combining functional studies in *Drosophila* together with human subject data, we have linked several microcephaly-associated genes and congenital infection to a single genetic pathway. These studies allowed us to highlight conserved functions of the ANKLE2 pathway, and provide mechanistic insight into how a Zika infection might affect asymmetric division. This *ANKLE2-VRK1* gene dosage sensitive pathway can be perturbed by genetic variants that disturb biological homeostasis resulting in neurological disease traits or by environmental insults such as Zika virus impinging on neurodevelopment. Hence, lessons learned from the study of rare diseases such as MCPH16/*ANKLE2* can provide insights into more common disease and potential gene by environmental interactions.

STAR Methods

Lead Contact and Materials Availability

All unique/stable reagents generated in this study are available from the Lead Contact without restriction. Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Hugo J. Bellen (hbellen@bcm.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Drosophila melanogaster

The following fly lines were used: *FRT19a* (Yamamoto *et al.*, 2014), *Ankle2^A* (Yamamoto *et al.*, 2014), *Ankle2^{CRIMIC}* (this study), *Ankle2^{GFP}* (this study), *Ankle2-GFPR* (this study), *P{UAS-hANKLE2}VK37* (Yamamoto *et al.*, 2014), *P{UAS-hANKLE2 p.L537V}VK37* (this study), *P{UAS-hANKLE2 p.Q782*}VK37* (this study), *P{UAS-hANKLE2 p.A109P}VK37* (this study), *P{UAS-hANKLE2 p.G201W}VK37* (this study), *P{20XUAS-tdTomato-Sec61beta}attP2* (Summerville *et al.*, 2016), *ball^{e107}* (Cullen *et al.*, 2005), *ball²* (Herzig *et al.*, 2014), *l(2)gIM107575-GFSTF.0* (Nagarkar-Jaiswal *et al.*, 2015a), *l(2)g^{ts3}cn¹sp¹³* (Manfrulli *et al.*, 1996), *P{UAS-NS4Aug}* (Shah *et al.*, 2018), *P{GW UAS-NS4Aug-HA}* (this study), *P{UAS-CD8-GFP}* (Lee and Luo, 2001), *Actin-GAL4 (P{Act5C-GAL4}17bFOI)* (Ito *et al.*, 1997), *inscuteable-GAL4 (P{w[+mW.hs]=GawB}insc[Mz1407])* (Luo *et al.*, 1994), *daughterless-GAL4 (P{w[+mW.hs]=GAL4-da.G32}UHI)* (Wodarz *et al.*, 1995), *ball-GFP (fTRG-823)* (Sarov *et al.*, 2016), *wor-mira-cherry P{w[+mC]=wor.GAL4.A}2,P{w[+mC]=UAS-mira.cherry}2/CyO* (Cabernard and Doe, 2009), *P{His2Av[T:Avic|GFP-S65T]}62A* (Clarkson and Saint, 1999), *P{w[+mC]=UAS-aurA.Exel}2, M{UAS-aPKC.ORF.3xHA}ZH-86Fb* (Bischof *et al.*, 2013), *P{w[+mC]=UAS-aPKC.DeltaN}3* (Drier *et al.*, 2002). All flies were maintained at 22°C and grown on standard cornmeal and molasses medium in plastic vials. Crosses were performed at temperature indicated (18°C, 22°C, 25°C, or 29°C). Hemizygous males were analyzed as *Ankle2* mutants (which is on the X chromosome) and females were used for *Ankle2* heterozygous studies. All other studies contained males and females. Brain volume measurements were conducted in late 3rd instar larvae (gut clearance, extruding spiracles). *Act-GAL4* was used to ubiquitously express Zika virus NS4A and Sec61β, *da-GAL4* was used to express human ANKLE2 constructs and aPKC, *insc-GAL4* was used to express NS4A in neuroblasts.

Human studies

All study subjects enrolled into the Baylor-Hopkins Center for Mendelian Genomics (BHCMG) provided informed consent for exome sequencing and study participation under the Baylor College of Medicine Institutional Review Board-approved protocol H-29697. BAB701 provided informed consent for molecular and genomic analysis under the Baylor College of Medicine Institutional Review Board-approved protocol H-9170. All study subjects enrolled through Baylor Genetics Laboratory (BGL) were analyzed on a retrospective basis, and only deidentified information is provided under the Baylor College of Medicine Institutional Board approved protocol H-41191. Patients were ascertained from the 7148 sequenced individuals in BCHMG or the ~12500 sequenced individuals in the BGL

by searching for biallelic variants with CADD scores >15 in conjunction with phenotypes of interest. Six male and two female patients were ascertained from the BHCMG database. The age of the patient is known for 3 individuals (2y, 7y, 32y). Two female and three male patients were ascertained from the BGL database. Ages at referral were 2 months, 7y, 12y, 20y, and 41y.

Human Clinical Information

BAB4821—A homozygous missense variant in *VRK2* (c.1234G>A;p.D412N; Table S2, Figure S6) was observed in a proband, BAB4821 born to consanguineous Turkish parents, with nanophthalmos. The variant was in a region of absence of heterozygosity (AOH) ~7.7Mb in length gleaned from unphased ES data (Karaca *et al.*, 2018) and consistent with identity-by-descent.

BAB7812—In BAB7812, a patient with severe microcephaly ($Z = -6.7$) and additional brain malformations including cortical dysplasia and agenesis of the corpus callosum, a homozygous variant with a predicted detrimental effect on splicing in a canonical splice site within *VRK3* (c.139+2 T>G) was found (Table S2, Figure S6). Additional homozygous variants were found in the same (*PNKP*) and an adjacent (*RNT2*) AOH block as *VRK3* (Table S2, Figure S6). All three homozygous variants are predicted deleterious. *PNKP* has a known disease association with microcephaly, seizures, and developmental delay (MIM# 613402) while *RNT2* has a known disease association with Spastic paraplegia 12 (MIM# 604805). Multilocus variation may contribute to the phenotype in BAB7812. The proband was born full term to consanguineous Turkish parents. At 5 months of age, he developed seizures. At 32 months, he was unable to walk, and his weight was 9.8kg ($-3.34SD$), height was 85cm ($-2.03SD$), and head circumference was 36.7cm ($-6.7SD$). Physical examination revealed thick eyebrows, protuberant ears, and almond shaped eyes. Metabolic and ocular examinations were within normal limits. He received continued follow-up care for microcephaly and refractory epilepsy, and special education for associated intellectual disability and developmental delay. His occipital frontal circumference was below the 3rd percentile when measured at 8–11 years of age. Parents and an unaffected sibling were heterozygous for the variant, demonstrating segregation of the variant with disease.

BAB10531—A BHCMG search database found a homozygous predicted deleterious missense variant in *PARD3B* (c.1222G>A, p.G408S) in BAB10531 who has microcephaly (Table S2, Figure S6). At 15 months, weight he was 9kg ($-2.1SD$), height was 79cm (0.4SD), and head circumference was 44.5cm ($Z = -2.2$ measured at 15m and 18m). Both unaffected consanguineous parents of Turkish descent were heterozygous for the variant that mapped to a region of AOH ~22Mb in length. BAB10531 has had frequent upper respiratory tract infections and was admitted at 17 months of age to investigate a cause for with no clear etiology found. Follow-up head circumference was performed at 18 months, which demonstrated microcephaly was nonprogressive (45cm, $-2.2SD$). Notably, no follow-up for intellectual disability was described on recruitment to the BHCMG.

BAB8223—In BAB8223, a proband with distal arthrogryposis, a homozygous missense variant in *PARD3B* (c.1654G>A;p.A552T) which segregated with disease was found (Table

S2, Figure S6). The proband was born to consanguineous Turkish parents and the homozygous variant was found within a region of AOH ~37Mb in length. At age 5.5y head circumference was 49cm (13th percentile, ~ -1SD). Cognitive and motor development is within normal limits.

BGL – 1—Referral diagnosis: Microcephaly, intrauterine growth restriction, radial microbrain and immature gyral pattern.

BGL – 2—Referral diagnosis: Global developmental delay with aphasia, microcephaly and seizures, progressive weakness, lethargy with episodes of irregular breathing and respiratory acidosis, MRI showing progressive diffuse central white matter atrophy and gliosis, hypomyelination, atrophy of thalami and pons. (Table S2, Figure S2)

BGL – 3—Referral diagnosis: Intellectual disability, ataxia, spasticity, autism spectrum disorder, speech delay, and anxiety disorder. (Table S2, Figure S2)

BGL – 4—Referral diagnosis: Spinal muscular atrophy complicated by failure to thrive and respiratory insufficiency, microcephaly, short stature, scoliosis, delayed motor milestones, progressive weakness, hypotonia, history of prematurity and intrauterine growth retardation (Table S2, Figure S5).

BGL – 5—Referral diagnosis: Prematurity, spasticity of inferior limbs, seizures, contractures of the Achilles, myopia (Table S2, Figure S5).

Human primary cultures

Fibroblasts were cultured in flasks containing Gibco DMEM (1x) with 4.5g/L D-Glucose, L-Glutamine, 25mM HEPES, HyClone FBS (10%), and Gibco Anti-Anti (1%) at 37 degrees Celsius with 5% CO₂. Cultures of p.L573V/+ and p.L573V/p.Q782* were male; p.V229G/p.V229G was female.

METHOD DETAILS

Generation of Ankle2 mutations and constructs

To generate *Ankle2*^{CRIMIC} by CRISPR-Cas9, two guide RNAs targeting *Ankle2*, 5'-ATAAAGTATTTCTTAACGGTGG-3' and 5'-TAATAATTTTAAATTCTCATTGG-3' with PAM sites underlined, were cloned into pCDF3 (Port *et al.*, 2014). Regions of homology targeting the 4th coding intron were cloned into PM14 (Lee *et al.*, 2018) using 5'-ccatagcatggGCAATTCCTCAATGTCGAATTTACTGCTCA-3' and 5'-ttatgcatATTTTCTTAACGGTGGGAAATTATAC-3' to amplify the left arm for BstXI/NsiI cloning and 5'-tagcatcATACTTTATTATTGCATTTGTTATAAGTATGAGA-3' and 5'-tactcagGCAAAGTTCCAGACCGTTTCTGATTTATC-3' to amplify the right arm for conventional cloning with SphI/XhoI. This donor construct and two guide RNA constructs were injected into *w;attP40(y+){nos-Cas9(v+)}/CyO* (Kondo and Ueda, 2013) embryos, and positive expression of *3XP3-GFP* was used to isolate animals with targeted events. PCR and genomic sequencing of surrounding regions validated the *Ankle2*^{CRIMIC} allele. *Ankle*^{IGFP} was generated by RMCE by injecting a plasmid expressing integrase with pBS-

KS-attB1–2-PT-SA-SD-EGFP-FlAsH-StrepII-TEV-3xFlag (Nagarkar-Jaiswal *et al.*, 2015b) into *Ankle2^{CRIMIC}* animals. Animals with 3XP3-GFP loss were screened using PCR for targeted cassette exchange. Regions flanking the targeted event were sequenced to verify the allele. The *Ankle2-GFP* was created using recombineering (Venken *et al.*, 2008) of BAC CH321–85N12 (Venken *et al.*, 2006). A GFP donor construct was generated by amplifying the GFP coding region and a selection cassette from plasmid PL-452 C-EGFP with primers containing 50bp homology with the C-terminal end of Ankle2 (5'-
GGGATCAACGGTCTATAACGAGGGGGACACGCCGCTGGGCAATCGGAACGCAG
CCCAATTCCGATCATATTC-3' and 5'-
 CATCAATCAGTCGCTGTTTCTGTTTCTGTTTCCGGGCCGATT
 CCGTTTCATTACTTGTACAGCTCGTCCATG-3'. Regions matching PL-452 C-EGFP are underlined. CH321–85N12 was transformed into DY380 cells using electroporation. Stable colonies were grown overnight at 30°C, induced for recombination functions at 42°C for 15 min, and transformed using electroporation (1.8kV, 200Ohm, 25μFD) with the amplified donor construct. Colonies were selected for both the BAC (chloramphenicol) and insertion of GFP (kanamycin). Resulting colonies were verified using PCR, restriction enzyme digestion, and sequencing. The GFP tag selection cassette was removed using Cre mediated excision by transforming the *Ankle2-GFP* BAC into induced EL350 cells. Properly excised events were verified by PCR, absence of growth on kanamycin selection plates, and sequencing.

Generation of human ANKLE2 and NS4A-HA expression constructs

NEB Q5 Site-directed mutagenesis was performed on *P{UAS^t-hANKLE2}*. Each plasmid was sequence verified and injected into VK37 flies with a plasmid expressing integrase for site-specific integration. NS4A, including the 2K peptide, from strain MR-766 was PCR amplified from UAS-NS4A (Shah *et al.*, 2018) and Gibson assembly was used to insert NS4A into pGW-HA.attB linearized with AgeI and KpnI to remove the ccdB region. Resulting vector was injected into VK37 flies as above.

Brain immunostaining

Late 3rd instar (based on gut clearance and extruding spiracles) larval brains were dissected in PBS and fixed with 4% PFA/PBS/0.3% Triton for 20 minutes. For immunostaining, brains were blocked in PBS/0.3% Triton/1% BSA/5% normal goat serum and incubated in primary antibody in PBS/0.3% Triton/1% BSA overnight. Primary antibodies include rat anti-Deadpan (Abcam Cat# ab195172, 1:250 or 1:500), mouse anti-Prospero MR1A (Developmental Studies Hybridoma Bank, 1:1000), rat anti-Miranda (1:500, Abcam Cat# ab197788), rabbit anti-aPKC (1:1000, PKCz (C-20) Santa Cruz, discontinued), rabbit anti-GFP (1:1000, Invitrogen Cat# A11122), mouse anti-Calnexin 99a (Developmental Studies Hybridoma Bank, 1:100), mouse anti-Lamin Dm0 ADL67.10 (Developmental Studies Hybridoma Bank, 1:250), guinea pig anti-Bazooka (1:1000) (Siller *et al.*, 2006), rat anti-Par-6 (1:50) (Rolls *et al.*, 2003), rabbit antiphospho-Histone H3 (1:1000, Millipore Cat# 06–570), rabbit anti-Ball (1:1000) (Yakulov *et al.*, 2014), rabbit anti-VRK1 (1:1000, Abcam Cat# ab151706), rabbit anti-CNN (1:1000) (Lucas and Raff, 2007), and mouse anti-Strep (Qiagen Cat# 34850, 1:500) with goat or donkey secondary antibodies from Jackson

ImmunoResearch used 1:500. Brains were mounted with double sided tape spacers and imaged using a Leica Sp8 with 2 μ m or 3 μ m sections through the entire brain lobe.

Live imaging

3rd instar larvae were dissected in sterile PBS supplemented with 1% FBS and 0.5mM ascorbic acid, fine dissected on an inverted Sarstedt lumox dish 50 in a petroleum jelly well. Samples were imaged on a Leica Sp8 with optimized settings for high quality images without bleaching or a Zeiss 880 with Airy scan (wild type Ankle2-GFP and Sec61 β colocalization).

Protein immunoprecipitation and western analysis

3rd instar larvae or dissected larval brains from *l(2)gIMI07575-GFSTF.0* animals were dissociated in 0.1% CHAPS buffer supplemented with protease and phosphatase inhibitors for at least 30 min on ice, centrifuged for 10 min at 4°C, and supernatant was used for immunoprecipitation or western analysis. For immunoprecipitation, 25ul of Allele Biotechnology GFP nanoantibody agarose (nAb, Cat# ABP-NAB-GFPA100) was equilibrated and incubated with lysate 2hrs - overnight at 4°C with rotation. Agarose was spun down for 1 min at 1000 x *g* at 4°C, supernatant was removed, and pellet was washed 3X (1X binding buffer (10mM Tris-HCl pH 7.5, 150mM NaCl), 2X wash buffer (10mM Tris-HCl pH 7.5, 500mM NaCl)). Remaining agarose pellet was eluted for western analysis in loading buffer. For western analysis, larval brains were dissected and dissociated as stated above, and were lysed in 0.1% CHAPS buffer [[50mM NaCl, 200mM HEPES, 1mM EDTA and protease inhibitor cocktail (Roche)] Loading input was adjusted for brain size and protein concentration. Primary antibodies include rabbit anti-GFP (1:2500, Invitrogen Cat# A11122), rabbit anti-Ball (1:1000) (Herzig *et al.*, 2014), rat anti-L(2)gl (Peng *et al.*, 2000), rabbit anti-aPKC c-20 (1:1000, Santa Cruz, discontinued), rabbit anti-aPKC phosphoT410 (1:1000, Santa Cruz, discontinued), and mouse anti-Actin-c4 (1:5000, Millipore Cat# MAB1501). Secondary antibodies include Rockland DyLight 600 and 800 (1:1000), BioRad Star Bright Blue 700 (1:1000) and Jackson ImmunoResearch HRP conjugated (1:5000). Blots were imaged on a Bio-Rad ChemiDocMP.

Food ingestion assay

Ankle^{2A/Y} or Ankle^{2A/+} third instar larvae from the same vial were washed in PBS and placed on yeast paste containing 0.08% Brilliant Blue R dye in 10mm petri dishes for 20 minutes. Larvae were transferred, rinsed, boiled for 10 sec, and aligned on a glass slide ventral side up. Larvae from both genotypes were imaged at the same time using a Biorad ChemiDoc MP Imaging system using the 715/30 (far red/680) filter. Total intensity of blue food in the gut was measured using Biorad's ImageLab software and plotted as total intensity per larvae.

Human cell immunohistochemistry

Cells were detached using Trypsin-EDTA 0.05% and plated onto 18mm glass cover slips in 6 well plates. Cells were cultured for an additional 3 days under the same conditions before fixing and staining. Cells were rinsed with PBS followed by fixing in 4% paraformaldehyde

in PBS. Cells were rinsed and washed 3x in PBST, washed 2x in PBST + 1% BSA (PBSTB), and then blocked in PBSTB + 5% normal goat serum. They were then incubated in PBSTB and primary antibody overnight at 4 degrees Celsius. Cells were then washed 3x in PBSTB, incubated in anti-rabbit Cy5 secondary antibody (1:500) for 2 hours, and washed 3x in PBST. The cells were given a final wash in PBST with DAPI (1:1000) for 30m before mounting using SlowFade glow on glass slides and sealing with nail polish.

Exome and Sanger Sequencing

Exome sequencing was performed under the Baylor Hopkins Center for Mendelian Genomics (BHCMG) research initiative as previously described (Lupski *et al.*, 2013). Exome capture was performed with Nimblegen reagents and a custom capture reagent, VCRome2.1. Raw data was processed using the Mercury pipeline, available on DNANexus (<http://blog.dnanexus.com/2013-10-22-run-mercury-variant-calling-pipeline/>) (Reid *et al.*, 2014) and the ATLAS2 method was used for variant calling followed by an in-house Cassandra annotation pipeline based on Annotation of Genetic Variants (ANNOVAR). The *LLGL1* variant was orthogonally validated and segregated with disease by dideoxy Sanger sequencing of PCR amplicons (Sanger *et al.*, 1977).

QUANTIFICATION AND STATISTICAL ANALYSIS

Brain volume

Brains from third instar larvae were stained and mounted with tape spacers and imaged using a Leica Sp8 with 2 μm or 3 μm sections through the entire brain lobe. Resulting stacks were analyzed using the Surfaces function in Imaris (Bitplane) to quantify brain lobe volume as total microns cubed. One lobe from each brain was imaged and a total of 5–10 brains were analyzed per genotype or condition. Brain lobe volumes are displayed as box plots with hinges representing the 25th to 75th percentiles, a line represents the median, and whiskers represent min to max. Statistical significance was determined using one-way ANOVA with multiple comparisons post-test calculated using GraphPad Prism. Brain volumes in Figure 1 are normalized to wild type (*FRT19a*). Average volume from wild type is set to 100%, and each mutant or condition is normalized as percentage of wild type volume. Brain volumes from Figure 4–5 are displayed as total brain volume (μm^3).

Asymmetric phenotypes

3rd instar larvae were immunostained for pH3, Baz, Par-6, aPKC, or Mira as described above. Metaphase neuroblasts (pH3 positive, chromosomes aligned at the metaphase plate) were imaged on a Leica Sp8 (63X). Only metaphase neuroblasts in the correct plane for imaging were analyzed. Mild disruption refers to weak or incomplete crescent localization, and strong disruption indicates no crescent localization. To quantify spindle orientation, CNN was used to mark the plane of division, and aPKC, and Mira were used to establish cortex polarity. Only metaphase neuroblasts in the correct plane for imaging were analyzed. The angle between spindle orientation and cortex polarity was measured using the angle function of ImageJ. Phenotypes are portrayed as percentage of total counted metaphase neuroblasts. For all samples, n is noted in the figure.

VRK1 intensity

Human fibroblasts were stained as described above, imaged on a Zeiss 710 as Z-stacks with equivalent laser power and confocal settings in the same imaging session. Resulting images were analyzed in Imaris (Bitplane) using the Surfaces function to mark nuclear volume. Total intensity sum of the VRK1 channel within the nucleus and nuclear volume were recorded. VRK1 intensity is displayed as intensity sum normalized to volume. One-way ANOVA with multi-comparisons post-test from GraphPad Prism was used to assess significance. Each dot represents one nucleus. Three fields from each cell line were assessed.

DATA AND CODE AVAILABILITY

The dbGaP accession number for all exome sequences reported in this paper and for which informed consent for data sharing in controlled-access databases has been provided under the Baylor College of Medicine Institutional Review Board Protocol Number H-29697 is dbGaP: phs000711.v5.p1. BAB701 was consented under the Baylor College of Medicine Institutional Review Board Protocol Number H-9170 and subjects enrolled through Baylor Genetics Laboratories were consented under the Baylor College of Medicine Institutional Review Board Protocol Number H-41191 which do not allow for data sharing through public repository.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

We would like to thank many members of the Bellen lab for suggestions, and Megan Campbell and Karen Schulze for critical reading. We thank Chris Doe, Jurgen Knoblich, Jim Skeath, Kenneth Prehoda, Alf Herzig, Hiroyuki Ohkura, and the Bloomington Drosophila Stock Center for providing stocks and reagents, and the IDDR Microscopy Core (NIH/NICHD U54 HD083092) for valuable input. This work was supported by NIH/NINDS F32NS092270 to N.L., Howard Hughes Medical Institute (HHMI) Medical Research Fellowship to A.J., the NIH/NINDS K08NS092898 and Jordan's Guardian Angels to G.M.M., a jointly funded NHGRI and NHLBI grant to the Baylor-Hopkins Center for Mendelian Genomics (UM1 HG006542) and NIH/NINDS R35NS105078 to J.R.L., NIH U54NS093793, NIH R24OD022005, and the Huffington Foundation to H.J.B.. N.L. and H.C. are supported by HHMI, and H.J.B. is an Investigator of the Howard Hughes Medical Institute.

REFERENCES

- Andersen RO, Turnbull DW, Johnson EA and Doe CQ (2012) 'Sgt1 acts via an LKB1/AMPK pathway to establish cortical polarity in larval neuroblasts', *Dev Biol*, 363(1), pp. 258–65. [PubMed: 22248825]
- Asencio C, Davidson IF, Santarella-Mellwig R, Ly-Hartig TB, Mall M, Wallenfang MR, Mattaj IW and Gorjanacz M (2012) 'Coordination of kinase and phosphatase activities by Lem4 enables nuclear envelope reassembly during mitosis', *Cell*, 150(1), pp. 122–35. [PubMed: 22770216]
- Atwood SX, Chabu C, Penkert RR, Doe CQ and Prehoda KE (2007) 'Cdc42 acts downstream of Bazooka to regulate neuroblast polarity through Par-6 aPKC', *J Cell Sci*, 120(Pt 18), pp. 3200–6. [PubMed: 17726059]
- Atwood SX and Prehoda KE (2009) 'aPKC phosphorylates Miranda to polarize fate determinants during neuroblast asymmetric cell division', *Curr Biol*, 19(9), pp. 723–9. [PubMed: 19375318]

- Ayoub AE, Oh S, Xie Y, Leng J, Cotney J, Dominguez MH, Noonan JP and Rakic P (2011) 'Transcriptional programs in transient embryonic zones of the cerebral cortex defined by high-resolution mRNA sequencing', *Proc Natl Acad Sci U S A*, 108(36), pp. 14950–5. [PubMed: 21873192]
- Bamshad MJ, Shendure JA, Valle D, Hamosh A, Lupski JR, Gibbs RA, Boerwinkle E, Lifton RP, Gerstein M, Gunel M, Mane S, Nickerson DA and Genomics, C. f. M. (2012) 'The Centers for Mendelian Genomics: a new large-scale initiative to identify the genes underlying rare Mendelian conditions', *Am J Med Genet A*, 158A(7), pp. 1523–5. [PubMed: 22628075]
- Barton LJ, Soshnev AA and Geyer PK (2015) 'Networking in the nucleus: a spotlight on LEM-domain proteins', *Curr Opin Cell Biol*, 34, pp. 1–8. [PubMed: 25863918]
- Betschinger J, Mechtler K and Knoblich JA (2003) 'The Par complex directs asymmetric cell division by phosphorylating the cytoskeletal protein Lgl', *Nature*, 422(6929), pp. 326–30. [PubMed: 12629552]
- Bhalerao S, Berdnik D, Török T and Knoblich JA (2005) 'Localization-dependent and - independent roles of numb contribute to cell-fate specification in *Drosophila*', *Curr Biol*, 15(17), pp. 1583–90. [PubMed: 16139215]
- Bier E, Vaessin H, Younger-Shepherd S, Jan LY and Jan YN (1992) 'deadpan, an essential pan-neural gene in *Drosophila*, encodes a helix-loop-helix protein similar to the hairy gene product', *Genes Dev*, 6(11), pp. 2137–51. [PubMed: 1427077]
- Bischof J, Björklund M, Furger E, Schertel C, Taipale J and Basler K (2013) 'A versatile platform for creating a comprehensive UAS-ORFeome library in *Drosophila*', *Development*, 140(11), pp. 2434–42. [PubMed: 23637332]
- Bonaccorsi S, Mottier V, Giansanti MG, Bolkan BJ, Williams B, Goldberg ML and Gatti M (2007) 'The *Drosophila* Lkb1 kinase is required for spindle formation and asymmetric neuroblast division', *Development*, 134(11), pp. 2183–93. [PubMed: 17507418]
- Brunetti-Pierri N, Berg JS, Scaglia F, Belmont J, Bacino CA, Sahoo T, Lalani SR, Graham B, Lee B, Shinawi M, Shen J, Kang SH, Pursley A, Lotze T, Kennedy G, Lansky-Shafer S, Weaver C, Roeder ER, Grebe TA, Arnold GL, Hutchison T, Reimschisel T, Amato S, Geraghty MT, Innis JW, Obersztyrn E, Nowakowska B, Rosengren SS, Bader PI, Grange DK, Naqvi S, Garnica AD, Bernes SM, Fong CT, Summers A, Walters WD, Lupski JR, Stankiewicz P, Cheung SW and Patel A (2008) 'Recurrent reciprocal 1q21.1 deletions and duplications associated with microcephaly or macrocephaly and developmental and behavioral abnormalities', *Nat Genet*, 40(12), pp. 1466–71. [PubMed: 19029900]
- Cabernard C and Doe CQ (2009) 'Apical/basal spindle orientation is required for neuroblast homeostasis and neuronal differentiation in *Drosophila*', *Dev Cell*, 17(1), pp. 134–41. [PubMed: 19619498]
- Campbell G, Göring H, Lin T, Spana E, Andersson S, Doe CQ and Tomlinson A (1994) 'RK2, a glial-specific homeodomain protein required for embryonic nerve cord condensation and viability in *Drosophila*', *Development*, 120(10), pp. 2957–66. [PubMed: 7607085]
- Carvalho CA, Moreira S, Ventura G, Sunkel CE and Morais-de-Sá E (2015) 'Aurora A triggers Lgl cortical release during symmetric division to control planar spindle orientation', *Curr Biol*, 25(1), pp. 53–60. [PubMed: 25484294]
- Chabu C and Doe CQ (2008) 'Dap160/intersectin binds and activates aPKC to regulate cell polarity and cell cycle progression', *Development*, 135(16), pp. 2739–46. [PubMed: 18614576]
- Chabu C and Doe CQ (2009) 'Twins/PP2A regulates aPKC to control neuroblast cell polarity and self-renewal', *Dev Biol*, 330(2), pp. 399–405. [PubMed: 19374896]
- Clarkson M and Saint R (1999) 'A His2AvDGFP fusion gene complements a lethal His2AvD mutant allele and provides an in vivo marker for *Drosophila* chromosome behavior', *DNA Cell Biol*, 18(6), pp. 457–62. [PubMed: 10390154]
- Cullen CF, Brittle AL, Ito T and Ohkura H (2005) 'The conserved kinase NHK-1 is essential for mitotic progression and unifying acentrosomal meiotic spindles in *Drosophila melanogaster*', *J Cell Biol*, 171(4), pp. 593–602. [PubMed: 16301329]

- Devhare P, Meyer K, Steele R, Ray RB and Ray R (2017) 'Zika virus infection dysregulates human neural stem cell growth and inhibits differentiation into neuroprogenitor cells', *Cell Death Dis*, 8(10), pp. e3106. [PubMed: 29022904]
- Drier EA, Tello MK, Cowan M, Wu P, Blace N, Sacktor TC and Yin JC (2002) 'Memory enhancement and formation by atypical PKM activity in *Drosophila melanogaster*', *Nat Neurosci*, 5(4), pp. 316–24. [PubMed: 11914720]
- Dumas LJ, O'Bleness MS, Davis JM, Dickens CM, Anderson N, Keeney JG, Jackson J, Sikela M, Raznahan A, Giedd J, Rapoport J, Nagamani SS, Erez A, Brunetti-Pierri N, Sugalski R, Lupski JR, Fingerlin T, Cheung SW and Sikela JM (2012) 'DUF1220-domain copy number implicated in human brain-size pathology and evolution', *Am J Hum Genet*, 91(3), pp. 444–54. [PubMed: 22901949]
- Ferrari S, Marin O, Pagano MA, Meggio F, Hess D, El-Shemerly M, Krystyniak A and Pinna LA (2005) 'Aurora-A site specificity: a study with synthetic peptide substrates', *Biochem J*, 390(Pt 1), pp. 293–302. [PubMed: 16083426]
- Gallaud E, Pham T and Cabernard C (2017) '*Drosophila melanogaster* Neuroblasts: A Model for Asymmetric Stem Cell Divisions', *Results Probl Cell Differ*, 61, pp. 183–210. [PubMed: 28409305]
- Gateff E and Schneiderman HA (1974) 'Developmental capacities of benign and malignant neoplasms of *Drosophila*', *Wilhelm Roux Arch Entwickl Mech Org*, 176(1), pp. 23–65. [PubMed: 28304815]
- Ghosh S, Marquardt T, Thaler JP, Carter N, Andrews SE, Pfaff SL and Hunter T (2008) 'Instructive role of aPKCzeta subcellular localization in the assembly of adherens junctions in neural progenitors', *Proc Natl Acad Sci U S A*, 105(1), pp. 335–40. [PubMed: 18162555]
- Gonzaga-Jauregui C, Lotze T, Jamal L, Penney S, Campbell IM, Pehlivan D, Hunter JV, Woodbury SL, Raymond G, Adesina AM, Jhangiani SN, Reid JG, Muzny DM, Boerwinkle E, Lupski JR, Gibbs RA and Wiszniewski W (2013) 'Mutations in VRK1 associated with complex motor and sensory axonal neuropathy plus microcephaly', *JAMA Neurol*, 70(12), pp. 1491–8. [PubMed: 24126608]
- Greenberg F, Lewis RA, Potocki L, Glaze D, Parke J, Killian J, Murphy MA, Williamson D, Brown F, Dutton R, McCluggage C, Friedman E, Sulek M and Lupski JR (1996) 'Multi-disciplinary clinical study of Smith-Magenis syndrome (deletion 17p11.2)', *Am J Med Genet*, 62(3), pp. 247–54. [PubMed: 8882782]
- Harsh S, Ozakman Y, Kitchen SM, Paquin-Proulx D, Nixon DF and Eleftherianos I (2018) 'Dicer-2 Regulates Resistance and Maintains Homeostasis against Zika Virus Infection in', *J Immunol*, 201(10), pp. 3058–3072. [PubMed: 30305326]
- Herzig B, Yakulov TA, Klinge K, Günesdogan U, Jäckle H and Herzig A (2014) 'Bällchen is required for self-renewal of germline stem cells in *Drosophila melanogaster*', *Biol Open*, 3(6), pp. 510–21. [PubMed: 24876388]
- Homem CC and Knoblich JA (2012) '*Drosophila* neuroblasts: a model for stem cell biology', *Development*, 139(23), pp. 4297–310. [PubMed: 23132240]
- Ito K, Awano W, Suzuki K, Hiromi Y and Yamamoto D (1997) 'The *Drosophila* mushroom body is a quadruple structure of clonal units each of which contains a virtually identical set of neurones and glial cells', *Development*, 124(4), pp. 761–71. [PubMed: 9043058]
- Jayaraman D, Bae BI and Walsh CA (2018) 'The Genetics of Primary Microcephaly', *Annu Rev Genomics Hum Genet*, 19, pp. 177–200. [PubMed: 29799801]
- Kanai MI, Kim MJ, Akiyama T, Takemura M, Wharton K, O'Connor MB and Nakato H (2018) 'Regulation of neuroblast proliferation by surface glia in the *Drosophila* larval brain', *Sci Rep*, 8(1), pp. 3730. [PubMed: 29487331]
- Katsani KR, Karess RE, Dostatni N and Doye V (2008) 'In vivo dynamics of *Drosophila* nuclear envelope components', *Mol Biol Cell*, 19(9), pp. 3652–66. [PubMed: 18562695]
- Kim S, Gailite I, Moussian B, Luschnig S, Goette M, Fricke K, Honemann-Capito M, Grubmüller H and Wodarz A (2009) 'Kinase-activity-independent functions of atypical protein kinase C in *Drosophila*', *J Cell Sci*, 122(Pt 20), pp. 3759–71. [PubMed: 19789180]
- Kondo S and Ueda R (2013) 'Highly improved gene targeting by germline-specific Cas9 expression in *Drosophila*', *Genetics*, 195(3), pp. 715–21. [PubMed: 24002648]

- Krahn MP, Egger-Adam D and Wodarz A (2009) 'PP2A antagonizes phosphorylation of Bazooka by PAR-1 to control apical-basal polarity in dividing embryonic neuroblasts', *Dev Cell*, 16(6), pp. 901–8. [PubMed: 19531360]
- Lee CY, Andersen RO, Cabernard C, Manning L, Tran KD, Lanskey MJ, Bashirullah A and Doe CQ (2006) 'Drosophila Aurora-A kinase inhibits neuroblast self-renewal by regulating aPKC/Numb cortical polarity and spindle orientation', *Genes Dev*, 20(24), pp. 3464–74. [PubMed: 17182871]
- Lee PT, Zirin J, Kanca O, Lin WW, Schulze KL, Li-Kroeger D, Tao R, Devereaux C, Hu Y, Chung V, Fang Y, He Y, Pan H, Ge M, Zuo Z, Housden BE, Mohr SE, Yamamoto S, Levis RW, Spradling AC, Perrimon N and Bellen HJ (2018) 'A genespecific T2A-GAL4 library for Drosophila', *Elife*, 7.
- Lee T and Luo L (2001) 'Mosaic analysis with a repressible cell marker (MARCM) for Drosophila neural development', *Trends Neurosci*, 24(5), pp. 251–4. [PubMed: 11311363]
- Leitges M, Sanz L, Martin P, Duran A, Braun U, García JF, Camacho F, Diaz-Meco MT, Rennert PD and Moscat J (2001) 'Targeted disruption of the zetaPKC gene results in the impairment of the NF-kappaB pathway', *Mol Cell*, 8(4), pp. 771–80. [PubMed: 11684013]
- Li C, Xu D, Ye Q, Hong S, Jiang Y, Liu X, Zhang N, Shi L, Qin CF and Xu Z (2016) 'Zika Virus Disrupts Neural Progenitor Development and Leads to Microcephaly in Mice', *Cell Stem Cell*, 19(5), pp. 672. [PubMed: 27814481]
- Lim NR, Shohayeb B, Zaytseva O, Mitchell N, Millard SS, Ng DCH and Quinn LM (2017) 'Glial-Specific Functions of Microcephaly Protein WDR62 and Interaction with the Mitotic Kinase AURKA Are Essential for Drosophila Brain Growth', *Stem Cell Reports*, 9(1), pp. 32–41. [PubMed: 28625535]
- Lin F, Blake DL, Callebaut I, Skerjanc IS, Holmer L, McBurney MW, Paulin-Levasseur M and Worman HJ (2000) 'MAN1, an inner nuclear membrane protein that shares the LEM domain with lamina-associated polypeptide 2 and emerlin', *J Biol Chem*, 275(7), pp. 4840–7. [PubMed: 10671519]
- Lin MY, Wang YL, Wu WL, Wolseley V, Tsai MT, Radic V, Thornton ME, Grubbs BH, Chow RH and Huang IC (2017) 'Zika Virus Infects Intermediate Progenitor Cells and Post-mitotic Committed Neurons in Human Fetal Brain Tissues', *Sci Rep*, 7(1), pp. 14883. [PubMed: 29093521]
- Liu Y, Gordesky-Gold B, Leney-Greene M, Weinbren NL, Tudor M and Cherry S (2018) 'Inflammation-Induced, STING-Dependent Autophagy Restricts Zika Virus Infection in the Drosophila Brain', *Cell Host Microbe*, 24(1), pp. 57–68.e3. [PubMed: 29934091]
- Lucas EP and Raff JW (2007) 'Maintaining the proper connection between the centrioles and the pericentriolar matrix requires Drosophila centrosomin', *J Cell Biol*, 178(5), pp. 725–32. [PubMed: 17709428]
- Luo L, Liao YJ, Jan LY and Jan YN (1994) 'Distinct morphogenetic functions of similar small GTPases: Drosophila Drac1 is involved in axonal outgrowth and myoblast fusion', *Genes Dev*, 8(15), pp. 1787–802. [PubMed: 7958857]
- Lupski JR (2015) 'Structural variation mutagenesis of the human genome: Impact on disease and evolution', *Environ Mol Mutagen*, 56(5), pp. 419–36. [PubMed: 25892534]
- Lupski JR, Gonzaga-Jauregui C, Yang Y, Bainbridge MN, Jhangiani S, Buhay CJ, Kovar CL, Wang M, Hawes AC, Reid JG, Eng C, Muzny DM and Gibbs RA (2013) 'Exome sequencing resolves apparent incidental findings and reveals further complexity of SH3TC2 variant alleles causing Charcot-Marie-Tooth neuropathy', *Genome Med*, 5(6), pp. 57. [PubMed: 23806086]
- Manfrulli P, Arquier N, Hanratty WP and Sémériva M (1996) 'The tumor suppressor gene, lethal(2)giant larvae (1(2)g1), is required for cell shape change of epithelial cells during Drosophila development', *Development*, 122(7), pp. 2283–94. [PubMed: 8681808]
- Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, Gwadz M, Hurwitz DI, Lu F, Marchler GH, Song JS, Thanki N, Wang Z, Yamashita RA, Zhang D, Zheng C, Geer LY and Bryant SH (2017) 'CDD/SPARCLE: functional classification of proteins via subfamily domain architectures', *Nucleic Acids Res*, 45(D1), pp. D200–D203. [PubMed: 27899674]
- Mlakar J, Korva M, Tul N, Popovi M, Poljšak-Prijatelj M, Mraz J, Kolenc M, Resman Rus K, Vesnaver Vipotnik T, Fabjan Vodusek V, Vizjak A, Pizem J, Petrovec M and Avšič Županc T

- (2016) 'Zika Virus Associated with Microcephaly', *N Engl J Med*, 374(10), pp. 951–8. [PubMed: 26862926]
- Moore CA, Staples JE, Dobyns WB, Pessoa A, Ventura CV, Fonseca EB, Ribeiro EM, Ventura LO, Neto NN, Arena JF and Rasmussen SA (2017) 'Characterizing the Pattern of Anomalies in Congenital Zika Syndrome for Pediatric Clinicians', *JAMA Pediatr*, 171(3), pp. 288–295. [PubMed: 27812690]
- Nagarkar-Jaiswal S, DeLuca SZ, Lee PT, Lin WW, Pan H, Zuo Z, Lv J, Spradling AC and Bellen HJ (2015a) 'A genetic toolkit for tagging intronic MiMIC containing genes', *Elife*, 4.
- Nagarkar-Jaiswal S, Lee PT, Campbell ME, Chen K, Anguiano-Zarate S, Gutierrez MC, Busby T, Lin WW, He Y, Schulze KL, Booth BW, Evans-Holm M, Venken KJ, Levis RW, Spradling AC, Hoskins RA and Bellen HJ (2015b) 'A library of MiMICs allows tagging of genes and reversible, spatial and temporal knockdown of proteins in *Drosophila*', *Elife*, 4.
- Nam SC, Mukhopadhyay B and Choi KW (2007) 'Antagonistic functions of Par-1 kinase and protein phosphatase 2A are required for localization of Bazooka and photoreceptor morphogenesis in *Drosophila*', *Dev Biol*, 306(2), pp. 624–35. [PubMed: 17475233]
- Nichols RJ and Traktman P (2004) 'Characterization of three paralogous members of the Mammalian vaccinia related kinase family', *J Biol Chem*, 279(9), pp. 7934–46. [PubMed: 14645249]
- Ogawa H, Ohta N, Moon W and Matsuzaki F (2009) 'Protein phosphatase 2A negatively regulates aPKC signaling by modulating phosphorylation of Par-6 in *Drosophila* neuroblast asymmetric divisions', *J Cell Sci*, 122(Pt 18), pp. 3242–9. [PubMed: 19690050]
- Oláhová M, Yoon WH, Thompson K, Jangam S, Fernandez L, Davidson JM, Kyle JE, Grove ME, Fisk DG, Kohler JN, Holmes M, Dries AM, Huang Y, Zhao C, Contrepolis K, Zappala Z, Frésard L, Waggott D, Zink EM, Kim YM, Heyman HM, Stratton KG, Webb-Robertson BM, Snyder M, Merker JD, Montgomery SB, Fisher PG, Feichtinger RG, Mayr JA, Hall J, Barbosa IA, Simpson MA, Deshpande C, Waters KM, Koeller DM, Metz TO, Morris AA, Schelley S, Cowan T, Friederich MW, McFarland R, Van Hove JLK, Enns GM, Yamamoto S, Ashley EA, Wangler MF, Taylor RW, Bellen HJ, Bernstein JA, Wheeler MT and Network UD (2018) 'Biallelic Mutations in ATP5F1D, which Encodes a Subunit of ATP Synthase, Cause a Metabolic Disorder', *Am J Hum Genet*, 102(3), pp. 494–504. [PubMed: 29478781]
- Peng CY, Manning L, Albertson R and Doe CQ (2000) 'The tumour-suppressor genes *lgl* and *dlg* regulate basal protein targeting in *Drosophila* neuroblasts', *Nature*, 408(6812), pp. 596–600. [PubMed: 11117748]
- Petronczki M and Knoblich JA (2001) 'DmPAR-6 directs epithelial polarity and asymmetric cell division of neuroblasts in *Drosophila*', *Nat Cell Biol*, 3(1), pp. 43–9. [PubMed: 11146625]
- Port F, Chen HM, Lee T and Bullock SL (2014) 'Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in *Drosophila*', *Proc Natl Acad Sci U S A*.
- Posey JE, O'Donnell-Luria AH, Chong JX, Harel T, Jhangiani SN, Coban Akdemir ZH, Buyske S, Pehlivan D, Carvalho CMB, Baxter S, Sobreira N, Liu P, Wu N, Rosenfeld JA, Kumar S, Avramopoulos D, White JJ, Doheny KF, Witmer PD, Boehm C, Sutton VR, Muzny DM, Boerwinkle E, Günel M, Nickerson DA, Mane S, MacArthur DG, Gibbs RA, Hamosh A, Lifton RP, Matise TC, Rehm HL, Gerstein M, Bamshad MJ, Valle D, Lupski JR and Genomics, C. f. M. (2019) 'Insights into genetics, human biology and disease gleaned from family based genomic studies', *Genet Med*.
- Poulton JS, Cuningham JC and Peifer M (2017) 'Centrosome and spindle assembly checkpoint loss leads to neural apoptosis and reduced brain size', *J Cell Biol*, 216(5), pp. 1255–1265. [PubMed: 28351851]
- Ramdas Nair A, Singh P, Salvador Garcia D, Rodriguez-Crespo D, Egger B and Cabernard C (2016) 'The Microcephaly-Associated Protein Wdr62/CG7337 Is Required to Maintain Centrosome Asymmetry in *Drosophila* Neuroblasts', *Cell Rep*, 14(5), pp. 1100–1113. [PubMed: 26804909]
- Reid JG, Carroll A, Veeraraghavan N, Dahdouli M, Sundquist A, English A, Bainbridge M, White S, Salerno W, Buhay C, Yu F, Muzny D, Daly R, Duyk G, Gibbs RA and Boerwinkle E (2014) 'Launching genomics into the cloud: deployment of Mercury, a next generation sequence analysis pipeline', *BMC Bioinformatics*, 15, pp. 30. [PubMed: 24475911]

- Renbaum P, Kellerman E, Jaron R, Geiger D, Segel R, Lee M, King MC and Levy-Lahad E (2009) 'Spinal muscular atrophy with pontocerebellar hypoplasia is caused by a mutation in the VRK1 gene', *Am J Hum Genet*, 85(2), pp. 281–9. [PubMed: 19646678]
- Riedel F, Gillingham AK, Rosa-Ferreira C, Galindo A and Munro S (2016) 'An antibody toolkit for the study of membrane traffic in *Drosophila melanogaster*', *Biol Open*, 5(7), pp. 987–92. [PubMed: 27256406]
- Riemer D, Stuurman N, Berrios M, Hunter C, Fisher PA and Weber K (1995) 'Expression of *Drosophila* lamin C is developmentally regulated: analogies with vertebrate A-type lamins', *J Cell Sci*, 108 (Pt 10), pp. 3189–98. [PubMed: 7593280]
- Rolls MM, Albertson R, Shih HP, Lee CY and Doe CQ (2003) '*Drosophila* aPKC regulates cell polarity and cell proliferation in neuroblasts and epithelia', *J Cell Biol*, 163(5), pp. 1089–98. [PubMed: 14657233]
- Rujano MA, Sanchez-Pulido L, Pennetier C, le Dez G and Basto R (2013) 'The microcephaly protein Asp regulates neuroepithelium morphogenesis by controlling the spatial distribution of myosin II', *Nat Cell Biol*, 15(11), pp. 1294–306. [PubMed: 24142104]
- Sanger F, Nicklen S and Coulson AR (1977) 'DNA sequencing with chain-terminating inhibitors', *Proc Natl Acad Sci U S A*, 74(12), pp. 5463–7. [PubMed: 271968]
- Sanz-García M, Vázquez-Cedeira M, Kellerman E, Renbaum P, Levy-Lahad E and Lazo PA (2011) 'Substrate profiling of human vaccinia-related kinases identifies coilin, a Cajal body nuclear protein, as a phosphorylation target with neurological implications', *J Proteomics*, 75(2), pp. 548–60. [PubMed: 21920476]
- Sarov M, Barz C, Jambor H, Hein MY, Schmied C, Suchold D, Stender B, Janosch S, K J VV, Krishnan RT, Krishnamoorthy A, Ferreira IR, Ejsmont RK, Finkl K, Hasse S, Kämpfer P, Plewka N, Vinis E, Schloissnig S, Knust E, Hartenstein V, Mann M, Ramaswami M, VijayRaghavan K, Tomancak P and Schnorrer F (2016) 'A genome-wide resource for the analysis of protein localisation in *Drosophila*', *Elife*, 5, pp. e12068. [PubMed: 26896675]
- Schober M, Schaefer M and Knoblich JA (1999) 'Bazooka recruits Inscuteable to orient asymmetric cell divisions in *Drosophila* neuroblasts', *Nature*, 402(6761), pp. 548–51. [PubMed: 10591217]
- Segura-Totten M, Kowalski AK, Craigie R and Wilson KL (2002) 'Barrier-to-autointegration factor: major roles in chromatin decondensation and nuclear assembly', *J Cell Biol*, 158(3), pp. 475–85. [PubMed: 12163470]
- Seidl S, Braun U, Roos N, Li S, Lüttke TH, Kispert A and Leitges M (2013) 'Phenotypical analysis of atypical PKCs in vivo function display a compensatory system at mouse embryonic day 7.5', *PLoS One*, 8(5), pp. e62756. [PubMed: 23690951]
- Shah PS, Link N, Jang GM, Sharp PP, Zhu T, Swaney DL, Johnson JR, Von Dollen J, Ramage HR, Satkamp L, Newton B, Hüttenhain R, Petit MJ, Baum T, Everitt A, Laufman O, Tassetto M, Shales M, Stevenson E, Iglesias GN, Shokat L, Tripathi S, Balasubramaniam V, Webb LG, Aguirre S, Willsey AJ, Garcia-Sastre A, Pollard KS, Cherry S, Gamarnik AV, Marazzi I, Taunton J, Fernandez-Sesma A, Bellen HJ, Andino R and Krogan NJ (2018) 'Comparative Flavivirus-Host Protein Interaction Mapping Reveals Mechanisms of Dengue and Zika Virus Pathogenesis', *Cell*, 175(7), pp. 1931–1945.e18. [PubMed: 30550790]
- Shaheen R, Maddirevula S, Ewida N, Alsahli S, Abdel-Salam GMH, Zaki MS, Tala SA, Alhashem A, Softah A, Al-Owain M, Alazami AM, Abadel B, Patel N, Al-Sheddi T, Alomar R, Alobeid E, Ibrahim N, Hashem M, Abdulwahab F, Hamad M, Tabarki B, Alwadei AH, Alhazzani F, Bashiri FA, Kentab A, ahintürk S, Sherr E, Fregeau B, Sogati S, Alshahwan SAM, Alkhalifi S, Alhumaidi Z, Temtamy S, Aglan M, Otaify G, Girisha KM, Tulbah M, Seidahmed MZ, Salih MA, Abouelhoda M, Momin AA, Saffar MA, Partlow JN, Arold ST, Faqeih E, Walsh C and Alkuraya FS (2018) 'Genomic and phenotypic delineation of congenital microcephaly', *Genet Med*.
- Shinawi M, Liu P, Kang SH, Shen J, Belmont JW, Scott DA, Probst FJ, Craigen WJ, Graham BH, Pursley A, Clark G, Lee J, Proud M, Stocco A, Rodriguez DL, Kozel BA, Sparagana S, Roeder ER, McGrew SG, Kurczynski TW, Allison LJ, Amato S, Savage S, Patel A, Stankiewicz P, Beaudet AL, Cheung SW and Lupski JR (2010) 'Recurrent reciprocal 16p11.2 rearrangements associated with global developmental delay, behavioural problems, dysmorphism, epilepsy, and abnormal head size', *J Med Genet*, 47(5), pp. 332–41. [PubMed: 19914906]

- Siller KH, Cabernard C and Doe CQ (2006) 'The NuMA-related Mud protein binds Pins and regulates spindle orientation in *Drosophila* neuroblasts', *Nat Cell Biol*, 8(6), pp. 594–600. [PubMed: 16648843]
- Smith AC, McGavran L, Robinson J, Waldstein G, Macfarlane J, Zonona J, Reiss J, Lahr M, Allen L and Magenis E (1986) 'Interstitial deletion of (17)(p11.2p11.2) in nine patients', *Am J Med Genet*, 24(3), pp. 393–414. [PubMed: 2425619]
- Snyers L, Erhart R, Laffer S, Pusch O, Weipoltshammer K and Schöfer C (2018) LEM4/ANKLE-2 deficiency impairs post-mitotic re-localization of BAF, LAP2 α and LaminA to the nucleus, causes nuclear envelope instability in telophase and leads to hyperploidy in HeLa cells *European Journal of Cell Biology*: Elsevier.
- Soloff RS, Katayama C, Lin MY, Feramisco JR and Hedrick SM (2004) 'Targeted deletion of protein kinase C lambda reveals a distribution of functions between the two atypical protein kinase C isoforms', *J Immunol*, 173(5), pp. 3250–60. [PubMed: 15322187]
- Summerville JB, Faust JF, Fan E, Pendin D, Daga A, Formella J, Stern M and McNew JA (2016) 'The effects of ER morphology on synaptic structure and function in *Drosophila melanogaster*', *J Cell Sci*, 129(8), pp. 1635–48. [PubMed: 26906425]
- Suzuki A and Ohno S (2006) 'The PAR-aPKC system: lessons in polarity', *J Cell Sci*, 119(Pt 6), pp. 979–87. [PubMed: 16525119]
- Tang H, Hammack C, Ogden SC, Wen Z, Qian X, Li Y, Yao B, Shin J, Zhang F, Lee EM, Christian KM, Didier RA, Jin P, Song H and Ming GL (2016) 'Zika Virus Infects Human Cortical Neural Progenitors and Attenuates Their Growth', *Cell Stem Cell*, 18(5), pp. 587–90. [PubMed: 26952870]
- Venken KJ, Carlson JW, Schulze KL, Pan H, He Y, Spokony R, Wan KH, Koriabine M, de Jong PJ, White KP, Bellen HJ and Hoskins RA (2009) 'Versatile P[acman] BAC libraries for transgenesis studies in *Drosophila melanogaster*', *Nat Methods*, 6(6), pp. 431–4. [PubMed: 19465919]
- Venken KJ, He Y, Hoskins RA and Bellen HJ (2006) 'P[acman]: a BAC transgenic platform for targeted insertion of large DNA fragments in *D. melanogaster*', *Science*, 314(5806), pp. 1747–51. [PubMed: 17138868]
- Venken KJ, Kasprovicz J, Kuenen S, Yan J, Hassan BA and Verstreken P (2008) 'Recombineering-mediated tagging of *Drosophila* genomic constructs for in vivo localization and acute protein inactivation', *Nucleic Acids Res*, 36(18), pp. e114. [PubMed: 18676454]
- Venken KJ, Schulze KL, Haelterman NA, Pan H, He Y, Evans-Holm M, Carlson JW, Levis RW, Spradling AC, Hoskins RA and Bellen HJ (2011) 'MiMIC: a highly versatile transposon insertion resource for engineering *Drosophila melanogaster* genes', *Nat Methods*, 8(9), pp. 737–43. [PubMed: 21985007]
- Wang C, Chang KC, Somers G, Virshup D, Ang BT, Tang C, Yu F and Wang H (2009) 'Protein phosphatase 2A regulates self-renewal of *Drosophila* neural stem cells', *Development*, 136(13), pp. 2287–96. [PubMed: 19502489]
- Wang H, Ouyang Y, Somers WG, Chia W and Lu B (2007) 'Polo inhibits progenitor self-renewal and regulates Numb asymmetry by phosphorylating Pon', *Nature*, 449(7158), pp. 96–100. [PubMed: 17805297]
- Wang H, Somers GW, Bashirullah A, Heberlein U, Yu F and Chia W (2006) 'Aurora-A acts as a tumor suppressor and regulates self-renewal of *Drosophila* neuroblasts', *Genes Dev*, 20(24), pp. 3453–63. [PubMed: 17182870]
- Wirtz-Peitz F, Nishimura T and Knoblich JA (2008) 'Linking cell cycle to asymmetric division: Aurora-A phosphorylates the Par complex to regulate Numb localization', *Cell*, 135(1), pp. 161–73. [PubMed: 18854163]
- Wodarz A, Hinz U, Engelbert M and Knust E (1995) 'Expression of crumbs confers apical character on plasma membrane domains of ectodermal epithelia of *Drosophila*', *Cell*, 82(1), pp. 67–76. [PubMed: 7606787]
- Wu N, Ming X, Xiao J, Wu Z, Chen X, Shinawi M, Shen Y, Yu G, Liu J, Xie H, Gucev ZS, Liu S, Yang N, Al-Kateb H, Chen J, Zhang J, Hauser N, Zhang T, Tasic V, Liu P, Su X, Pan X, Liu C, Wang L, Shen J, Chen Y, Choy KW, Wang J, Wang Q, Li S, Zhou W, Guo J, Wang Y, Zhang C, Zhao H, An Y, Zhao Y, Liu Z, Zuo Y, Tian Y, Weng X, Sutton VR, Wang H, Ming Y, Kulkarni S,

- Zhong TP, Giampietro PF, Dunwoodie SL, Cheung SW, Zhang X, Jin L, Lupski JR, Qiu G and Zhang F (2015) 'TBX6 null variants and a common hypomorphic allele in congenital scoliosis', *N Engl J Med*, 372(4), pp. 341–50. [PubMed: 25564734]
- Yakulov T, Günesdogan U, Jäckle H and Herzig A (2014) 'Bällchen participates in proliferation control and prevents the differentiation of *Drosophila melanogaster* neuronal stem cells', *Biol Open*, 3(10), pp. 881–6. [PubMed: 25190057]
- Yamamoto S, Jaiswal M, Charng WL, Gambin T, Karaca E, Mirzaa G, Wiszniewski W, Sandoval H, Haelterman NA, Xiong B, Zhang K, Bayat V, David G, Li T, Chen K, Gala U, Harel T, Pehlivan D, Penney S, Vissers LE, de Ligt J, Jhangiani SN, Xie Y, Tsang SH, Parman Y, Sivaci M, Battaloglu E, Muzny D, Wan YW, Liu Z, Lin-Moore AT, Clark RD, Curry CJ, Link N, Schulze KL, Boerwinkle E, Dobyns WB, Allikmets R, Gibbs RA, Chen R, Lupski JR, Wangler MF and Bellen HJ (2014) 'A drosophila genetic resource of mutants to study mechanisms underlying human genetic diseases', *Cell*, 159(1), pp. 200–14. [PubMed: 25259927]
- Zhong W, Jiang MM, Schonemann MD, Meneses JJ, Pedersen RA, Jan LY and Jan YN (2000) 'Mouse numb is an essential gene involved in cortical neurogenesis', *Proc Natl Acad Sci U S A*, 97(12), pp. 6844–9. [PubMed: 10841580]

Highlights

Ankle2 interacts with Ballchen/VRK1 to regulate brain development in *Drosophila*.

Ankle2/VRK1 pathway affects asymmetric protein localization during neuroblast division.

ANKLE2 and VRK1 are associated with human microcephaly.

The Zika virus protein NS4A interacts with and inhibits the ANKLE2 pathway.






Figure 1: Mutations in ANKLE2 cause microcephaly in humans and *Drosophila*.

The structure and transcripts of the *Ankle2* gene with insertional mutations or tags are shown in (A). *Ankle2*^A, noted by *, is an EMS generated L326H mutation, while *Ankle2*^{CRIMIC} represents a Crispr-Cas9 mediated MIMiC-like (CRIMIC) insertion in the 4th coding intron (Lee *et al.*, 2018). ΦC31 mediated cassette exchange replaced the stop-polyA in *Ankle2*^{CRIMIC} with an artificial exon cassette consisting of SA-GFP-SD (Venken *et al.*, 2011) to produce *Ankle2*^{IGFP}. A C-terminal GFP tagged *Ankle2* genomic rescue construct (*Ankle2*-GFP) was generated using recombineering (Venken *et al.*, 2008). (B) Lethal stages and rescue by *Ankle2*-GFP or (C) complementation tests of *Ankle2*^A, *Ankle2*^{CRIMIC}, and *Ankle2*^{IGFP}. (D) *Ankle2* gene structure and genomic rescue construct (CH321–85N12; Venken *et al.*, 2009). (E–L) Partial projections of 3rd instar larval brains stained with Deadpan (Dpn) (purple, neuroblasts) a neuroblast marker (Bier *et al.*, 1992) and Pros (green, GMC, neurons), a neuronal lineage marker (Campbell *et al.*, 1994), to

document overall brain structure is shown in (E) wild type ($y w FRT19a$), (F) $y w Ankle2^A FRT19a$ (G) $y w Ankle2^A FRT19a; Ankle2-GFP$, (H) $y w Ankle2^A FRT19a; da::hANKLE2^{wt}$, (I) $Ankle2^{CRIMIC}$, (J) $Ankle2^{CRIMIC}; Ankle2-GFP$ (K) $y w Ankle2^A FRT19a; da::hANKLE2 p.L573V$ and (L) $y w Ankle2^A FRT19a; da::hANKLE2 p.Q782^*$ animals. The C-terminal GFP-tagged rescue P[acman] clone CH321–85N12, ($Ankle2-GFP$, diagramed in A and B) rescues brain morphology, brain size, and lethality in both $y w Ankle2^A FRT19a$ (G) and $Ankle2^{CRIMIC}$ (J) animals. Quantification of brain size is shown in (M). Box plots hinges represent the 25th to 75th percentiles, the central line is the median, and whiskers represent min to max. Note that hANKLE2^{wt} and hANKLE2 p.L573V rescue brain size and lethality of $Ankle2^A$ mutants, whereas hANKLE2 p.Q782*, hANKLE2 p.A109P, and hANKLE2 p.G201W do not. Here, $Ankle2-GFP$ is reported as GFP. (N) Neuroblast specific expression of wild type human ANKLE2 ($y w Ankle2^A FRT19a; insc::hANKLE2^{wt}$) partially rescues brain size of $Ankle2^A$ mutants. For quantifications, the total number counted (n) is noted below each graph. See also Figure S1–S3.

Ankle2 localizes with ER markers and is required for proper ER morphology



Ankle2 localizes with the nuclear envelope and is required for its morphology




Figure 2: Ankle2 localizes to the ER and is dynamically expressed in the brain.

(A-B) 3rd instar larval brains from *Ankle2-GFP* animals stained for GFP to document Ankle2 expression and localization in (A-B) single slices (arrows point to neuroblasts). (C-E) Live imaging of Ankle2 (D, green) and the ER labeled with Sec61B (E, *da-GAL4, UAS-Sec61β-tdTomato*, purple) (Summerville *et al.*, 2016) shows strong colocalization. (F) Live *Ankle2^A* mutant neuroblast display aberrant Sec61β-tdTomato expression. (G-I) Fixed *Ankle2-GFP* animals highlighting Ankle2-GFP (green) and another ER marker (Calnexin 99a, purple) (Riedel *et al.*, 2016). (J) Fixed *Ankle2* mutant animals (*Ankle2^A*) display aberrant ER structures (Calnexin 99a, purple). (K-M) Ankle2 (L, green) colocalizes with some portions of the nuclear envelope (Lamin Dm0, purple). (N) *Ankle2* mutant animals (*Ankle2^A*) display disrupted nuclear envelope structure (Lamin Dm0, purple). See also Figure S4 and Movie S1.




Figure 3: *Ankle2* mutations affect asymmetric division, spindle alignment and centrosomes. Metaphase neuroblasts stained with Baz (green, A-D), Par-6 (green, E-H), aPKC (green, I-L) and Mira (red, I-L) are shown in wild type (A, E, I), *Ankle2^A/Y* hemizygous (B, F, J), *Ankle2^A/Ankle2^{CRIMIC}* transheterozygous (C, G, K) and rescued (D, H, L) animals. Phospho-Histone H3 (pHH3, green, A-C, E-K) was used to identify cell cycle stage. * notes samples where pHH3 was not used. (M-P) quantification of phenotype severity demonstrate that *Ankle2* is required for protein localization during asymmetric division in numerous cells. Below each graph, the # of neuroblasts counted for each genotype (n) is noted. (Q-U) Metaphase neuroblasts stained with aPKC (green) and Mira (red) to mark the polarity axis and DNA (white) and CNN (green puncta) to highlight the spindle axis from (Q) wild type (*FRT19a*) and (R-S) *Ankle2* mutant neuroblasts. The angle between the spindle axis and polarity axis is measured and % of metaphase neuroblasts is plotted in 15° intervals and is

shown in (T-U). Below each group, the # of neuroblasts counted for each genotype (n) is noted. See also Movie S2–S5.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 4: Ankle2 controls Ballchen/VRK1 localization and function.

(A-D) Immunostaining of Ankle2-GFP (green) and Ball (red) show dynamic localizations during the cell cycle. (B) Ankle2 is localized at the nuclear envelope at prophase. Ball is nuclear (A-B) until nuclear envelope breakdown (C) and then localizes to the cytoplasm through the end of mitosis (D). (E-F) Immunostaining of Ball (red) and Lamin (white) in *Ankle2^{A/+}* heterozygotes and *Ankle2^{A/Y}* hemizygous mutants. Note that Lamin is disrupted and Ball becomes mislocalized throughout the cytoplasm during interphase in *Ankle2^{A/Y}* hemizygous mutants. Quantification is shown in Figure S3E. (G-H) Confocal projections of immunostaining of VRK1 (green) and DNA (purple) in primary human fibroblasts from (G) parental unaffected (p.L573V/+) and (H) an ANKLE2 compound heterozygous patient (p.L573V/p.Q782*). VRK1 is mislocalized in fibroblasts carrying microcephaly associated ANKLE2 variants (p.L573V/p.Q782* and p.V229G/p.V229G), quantified as nuclear intensity in (I). Arrows in (H) indicate cytoplasmic VRK1 staining, which is minimal in (G)

control fibroblasts. (J-L) Partial projections of 3rd instar larval brains stained with apical marker aPKC (green) and basal marker Miranda (red) in (J) wild type (*y,w, FRT19a*), (K) *Ankle2^A*, and (L) *Ankle2^A;ball^{107/+}* animals. Note that removal of a single copy of *ball* rescues the phenotypes of *Ankle2^A*. (M) Quantification of 3rd instar larval brain size (as shown in (J-L)). N = 6. One-way ANOVA with multi-comparison post-test. ****p<0.0001, ***p<0.001. Box plots hinges represent the 25th to 75th percentiles, a line is at the median, and whiskers represent min to max. (N) *Ankle2^A*, but not *Ankle2^{CRIMIC}*, lethality is rescued with introduction of multiple *ball* heterozygous mutations. (O-P) Quantification of (O) aPKC or (P) Mira crescent intensity in 3rd instar metaphase neuroblasts in wild type (*y w FRT19a*), *Ankle2^A* and rescued *Ankle2^A; ball^{107/+}* animals, demonstrating that *Ankle2* asymmetric division phenotypes are rescued with *ball* heterozygosity. Note that wild type and *Ankle2^A* quantifications were shown in Fig 3. Below each graph, the # of neuroblasts or brains counted for each genotype (n) is noted. See also Figure S3 and S5 and Movie S6.




Figure 5: Ankle2 affects asymmetric division through aPKC and L(2)gl

(A) Ankle2 and Ball regulate asymmetric division. (B) Western analysis of phosphorylated aPKC in larval brains (mammalian T410 corresponds to T422 in *Drosophila*) from genomic rescue (*Ankle2^A; Ankle2-GFP*), *Ankle2^A*, and *Ankle2^A; ball^{e107/+}* animals or total aPKC levels in genomic rescue (*Ankle2^A; Ankle2-GFP*) and *Ankle2^A* mutants. Note that p-aPKC is reduced in *Ankle2* mutants but restored with introduction of genomic rescue (GR) or reduction of Ball and is quantified as the ratio of p-aPKC to total aPKC. N=3 replicates. (C) *In vivo* immunoprecipitation of L(2)gl-GFP using GFP-nAb in *L(2)gl^{M107575-GFSTF}* larvae demonstrates L(2)gl can interact with Ball *in vivo*. (D,F) Partial projections of third instar larval brains stained for Dpn (purple, neuroblasts) and Pros (green, daughter cells and neurons) of (D) *Ankle2^A* and (E) *Ankle2^A; l(2)gl^{ts3}* mutant animals raised at 22°C with brain volume quantified in (F). Note that reduction of L(2)gl in an *Ankle2^A* hemizygous animals rescues brain size defects and lethality at 22°C. Box plots hinges represent the 25th

to 75th percentiles, a line is at the median, and whiskers represent min to max. Below the graph, the # of brains counted for each genotype (n) is noted. See also Figure S6.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 6: Zika virus NS4A targets the Ankle2 pathway.

(A) 3rd instar brains with neuroblast specific (*insc-GAL4*) NS4A-HA expression stained for HA (green) and DNA (white) shows that NS4A localizes in a pattern similar to Ankle2 (Figure 2). (B-E) Metaphase neuroblasts stained for aPKC (green) and Mira (purple) in brains with neuroblast specific (*insc-GAL4*) expression of (B) *CD8-GFP*, (C-D) *NS4A*, or (E) *NS4A* in *ball¹⁰⁷* heterozygous animals. aPKC and Mira crescent intensities are quantified in (F-G). (H) Brain volume quantification from animals with ubiquitous expression of *CD8-GFP* (control), *NS4A-HA*, or *NS4A-HA* in *ball¹⁰⁷* heterozygous animals using *Act-GAL4* on the 2nd chromosome. (I) *Act-GAL4* (3rd chromosome) ubiquitous expression of *CD8-GFP* (control), *NS4A-HA*, *NS4A-HA* and human *ANKLE2*, or *NS4A-HA* in *I(2)g^{ts3}* heterozygous animals. Note that human *ANKLE2* expression or reduction of Ball or L(2)gl activity rescues NS4A induced brain defects. One-way ANOVA with multi-comparison post-test. ****p<0.0001, **p<0.01, *p<0.05. Box plots hinges

represent the 25th to 75th percentiles, a line is at the median, and whiskers represent min to max. (J-L) aPKC (green), CNN (green), and Mira (purple) staining of metaphase neuroblasts with *insc-GAL4* expression of (J) *CD8-GFP*, (K) *NS4A*, and (M) *NS4A* in *balF¹⁰⁷* heterozygous animals. The angle between the spindle axis and polarity axis is measured and % of total metaphase neuroblasts is noted at 15° intervals. Expression of NS4A causes localization defects of aPKC and Mira and spindle orientation defects in metaphase neuroblasts. (M) Zika virus NS4A inhibits the ANKLE2/VRK1 pathway, which regulates asymmetric determinant localization as well as the division axes.