

Lawrence Berkeley National Laboratory

Recent Work

Title

COMPARISON QUARK MODEL PREDICTIONS WITH EXPERIMENT FOR pp ANNIHILATION

Permalink

<https://escholarship.org/uc/item/4pt2d69d>

Author

Kirz, Janos.

Publication Date

1966-06-17

University of California Ernest O. Lawrence Radiation Laboratory

COMPARISON QUARK-MODEL PREDICTIONS WITH EXPERIMENT
FOR $p\bar{p}$ ANNIHILATION

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

To be submitted to Physics Letters

UCRL-16934

Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

COMPARISON OF QUARK MODEL PREDICTIONS WITH
EXPERIMENT FOR $\bar{p}p$ ANNIHILATION

Janos Kirz

June 17, 1966

COMPARISON OF QUARK MODEL PREDICTIONS WITH
EXPERIMENT FOR $\bar{p}p$ ANNIHILATION

Janos Kirz

Lawrence Radiation Laboratory
University of California
Berkeley, California

June 17, 1966

ABSTRACT

Predictions for $\bar{p}p$ annihilation, based on a quark model, are compared with experimental data. The model fails to give a quantitative description of the reaction.

Recently Rubinstein and Stern¹ derived predictions for nucleon-antinucleon annihilations based on a simple and interesting quark model. They assumed that in the annihilation process the three quarks and the three antiquarks that make up the initial state rearrange into three quark-antiquark pairs, i. e. three mesons.² They further assumed a one-to-one correspondence between the quarks in the initial and final states, including their charge, strangeness, and spin state.³ In this note their predictions are compared with the experimental data on $\bar{p}p$ annihilations at rest.⁴⁻¹⁰

Prediction 1: Annihilation results in three and only three mesons.

Unfortunately only 38% of the annihilation events can be completely analyzed, and distributed among the various final states in hydrogen bubble chamber experiments.^{4, 9} (The remainder result in more than one unobserved neutral particle.) Of this 38%, $5 \pm 1\%$ correspond to two-meson,⁵ $13 \pm 1\%$ to three-meson,^{4, 9, 10} and $20 \pm 2\%$ to four-or-more-meson final states.^{4, 9} The findings are in serious disagreement with the prediction.

Prediction 2: No strange particles are produced.

Experimentally the fraction of all annihilations involving $\bar{K}K$ pairs is $6.82 \pm 0.25\%$.⁸ The small rate for the associated production of strange-particle pairs, however, is a common feature of all elementary particle interactions.

Prediction 3: Because of its strange quark content, ϕ -meson production is severely limited.

Results of the Columbia experiment, as quoted in Table V of Ref. (11), give

$$\frac{\overline{pp} \rightarrow \pi^+ \pi^- \phi}{\overline{pp} \rightarrow \pi^+ \pi^- \omega} = 0.006 \pm 0.003,$$

in good agreement with the prediction. The same table shows, however, that a similar suppression of ϕ production exists in $\pi^\pm p$ interactions.¹²

Prediction 4: In Ref. (1) the annihilation products are predicted to be distributed among the various three-meson states as shown on line 2 of Table I.¹³

If we make the additional assumption that the singlet and triplet isotopic spin states contribute equally to the annihilation process, we must modify these predictions. The relative weights of the annihilation process in the available eigenstates of $I^{G_J P}$ become

$$0^+ 0^- : 1^- 0^- : 0^- 1^- : 1^+ 1^- : : 1/8 : 1/8 : 3/8 : 3/8 .$$

With this plausible modification of the model we get the predictions shown on line 3 of Table I.¹⁴ The experimental results shown on line 4 clearly disagree with either set of predictions.

To summarize, predictions 1 and 4 are not fulfilled; the agreement of the experiment with predictions 2 and 3 does not provide a strong argument for the model, since these predictions also hold true for reactions in which the assumptions of the model are inapplicable.

The overall picture indicates that the quark model in its simple intuitively appealing form used in Ref. (1) does not lead to a quantitative understanding of antiproton annihilation. A more sophisticated form will be necessary to obtain agreement with experiment.

It is a pleasure to thank Professor Edward Teller for numerous discussions concerning the model, and Professor Luis W. Alvarez for encouragement and support.

Work done under the auspices of the U. S. Atomic Energy Commission.

Table I. Distribution of final states in the reaction
 $\bar{p}p \rightarrow 3 \text{ mesons.}$

Final state	$\pi^+ \pi^- \pi^0$	$\pi^0 \pi^0 \pi^0$	$\rho^0 \pi^+ \pi^-$	$\omega \pi^+ \pi^-$
Prediction of Ref. (1): % of all annihilations	8	12	25	28
Modified pre- diction (see text): % of all annihilations	4	6	30	24
Experiment: % of all annihilations	2.7 ± 0.7 3.5 ± 0.6 (a)	$< 3.2 \pm 0.5$ $< 3.5 \pm 0.4$ (b)	5.4 ± 0.3 (c)	3.1 ± 0.5 (c)

(a) Reference (6), (7)

(b) These values are given in References (9) and (15) respectively for annihilations with no charged prongs. They therefore constitute an upper limit for $\bar{p}p \rightarrow 3 \pi^0$.

(c) References (4), (9)

REFERENCES

1. H. R. Rubenstein and H. Stern, *Phys. Letters* 21, 447 (1966).
2. Within the framework of the model only members of the pseudo-scalar and vector meson nonets are considered quark-antiquark pairs. The term meson therefore is restricted to these objects.
3. This model was independently considered by E. Teller (Lawrence Radiation Laboratory), who calls it "quark chemistry" (private communication).
4. M. Cresti et al., Padova-Oxford collaboration; *Proc. of the Sienna Int'l. Conf. on Elem. Part.*, Pg. 263 (1963).
5. C. Baltay et al., Columbia-Rutgers collaboration; *Phys. Rev. Letters* 10, 62 (1963).
6. G. B. Chadwick et al., Oxford-Padova collaboration; *Phys. Rev. Letters* 15, 532 (1965).
7. C. Baltay, P. Franzini, N. Gelfand, G. Lütjens, J. C. Severiens, J. Steinberger, D. Tycho, and D. Zanello, *Phys. Rev.* 140, B 1039 (1965), and *Phys. Rev.* 140, B 1042 (1965).
8. R. Armenteros et al., CERN - College de France - Ecole Polytechnique Collaboration; *Proc. of the 1962 Int'l. Conf. on High Energy Phys. at CERN*, Pg. 351. The quoted number depends on the assumption that annihilations leading to $K^+ K^-$, $K^- K^0$, $K^+ \overline{K^0}$, and $K^0 \overline{K^0}$ and equally probable.
9. C. Baltay, P. Franzini, G. Lütjens, J. C. Severiens, D. Tycho, and D. Zanello, *Phys. Rev.* 145, 1103 (1966).
10. N. Barash, L. Kirsch, D. Miller, and T. H. Tan, *Phys. Rev.* 145, 1095 (1966).
11. C. Baltay et al., Yale-Brookhaven collaboration, *Phys. Rev.* 142, 932 (1966).

12. Another recent compilation gives

$$\frac{\pi^- p \rightarrow \phi n}{\pi^+ n \rightarrow \omega p} \approx \frac{1}{50}$$

at equal c. m. momenta for the final states. R. I. Hess,
Production of $K\bar{K}$ Pairs in $\pi^- p$ Interactions (Ph. D. Thesis)
UCRL-16832, (in preparation).

13. In Table I we list only those channels for which experimental data are available. In Ref. (1) a more complete list of predictions is given.
14. The authors of Ref. (1) also predict that η production will be strongly limited. With the suggested modification in the model this prediction no longer holds; $\eta \pi^+ \pi^-$ and $\eta \pi^0 \pi^0$ become the dominant annihilation channels in the $I^G_J P = 0^+ 0^-$ state. Experimentally the $\eta \pi^+ \pi^-$ final state is observed to account for only about 1% of all annihilations.^{4, 9}
15. J. G. Loken and M. Derrick, Phys. Letters 3, 334 (1963).

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

Faint, illegible text covering the majority of the page, possibly bleed-through from the reverse side.

