

UCSF

UC San Francisco Previously Published Works

Title

Skeletal muscle adiposity and outcomes in candidates for lung transplantation: a lung transplant body composition cohort study.

Permalink

<https://escholarship.org/uc/item/4pt8p66f>

Journal

Thorax, 75(9)

ISSN

0040-6376

Authors

Anderson, Michaela R
Easthausen, Imaani
Gallagher, Grace
[et al.](#)

Publication Date

2020-09-01

DOI

10.1136/thoraxjnl-2019-214461

Peer reviewed



Published in final edited form as:

Thorax. 2020 September ; 75(9): 801–804. doi:10.1136/thoraxjnl-2019-214461.

Skeletal muscle adiposity and outcomes in candidates for lung transplantation: a lung transplant body composition cohort study

Michaela R Anderson¹, Imaani Easthausen², Grace Gallagher¹, Jayaram Udupa³, Yubing Tong³, Drew Torigian³, Joshua Matthew Diamond⁴, Mary Katherine Porteous⁴, Scott M Palmer^{5,6}, Laurie D Snyder⁵, Luke Benvenuto¹, Meghan Aversa¹, Selim Arcasoy¹, John R Greenland^{7,8}, Steven R Hays⁸, Jasleen Kukreja⁹, Edward Cantu³, John Shinn Kim¹⁰, Dympna Gallagher¹, Matthew R Baldwin¹, R Graham Barr¹, David J Lederer^{1,11}, Jason D Christie^{3,4}, Jonathan Paul Singer⁸

¹Medicine, Columbia University Medical Center, New York, New York, USA

²Population Health Sciences, Weill Cornell Medical College, New York, New York, USA

³Radiology, University of Pennsylvania, Philadelphia, Pennsylvania, USA

⁴Pulmonary, Allergy, and Critical Care, University of Pennsylvania, Philadelphia, Pennsylvania, USA

⁵Medicine, Duke University, Durham, North Carolina, USA

⁶Duke Clinical Research Institute, Durham, North Carolina, USA

⁷Medicine, VA Medical Center, San Francisco, California, USA

⁸Medicine, University of California, San Francisco, California, USA

⁹Surgery, University of California, San Francisco, California, USA

¹⁰Medicine, University of Virginia School of Medicine, Charlottesville, Virginia, USA

¹¹Regeneron Pharmaceuticals Inc, Tarrytown, New York, USA

Abstract

CT measurement of body composition may improve lung transplant candidate selection. We assessed whether skeletal muscle adipose deposition on abdominal and thigh CT scans was associated with 6 min walk distance (6MWD) and wait-list survival in lung transplant candidates.

Correspondence to Dr Michaela R Anderson, Medicine, Columbia University, New York NY 10032, New York, USA; mdr2140@cumc.columbia.edu, **Twitter** John Shinn Kim @jskim8223 and David J Lederer @davidledererm.

MRA and IE are joint first authors.

DJL, JDC and JPS are joint senior authors.

Contributors Study design: MRA, IE, DJL, JC and JPS. Data acquisition, analysis, interpretation: MRA, IE, GG, JU, YT, DT, JMD, MKP, SMP, LDS, LB, MA, SA, JRG, SRH, JK, EC, JSK, DG, MRB, RGB, DJL, JC and JPS. First draft of the manuscript: MRA, IE, JC and JPS. Final manuscript review and approval: all authors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Each ½-SD decrease in abdominal muscle attenuation (indicating greater lipid content) was associated with 14 m decrease in 6MWD (95% CI –20 to –8) and 20% increased risk of death or delisting (95% CI 10% to 40%). Each ½-standard deviation decrease in thigh muscle attenuation was associated with 15 m decrease in 6MWD (95% CI –21 to –10). CT imaging may improve candidate risk stratification.

INTRODUCTION

In the USA, despite increased lung transplant surgeries performed, wait-list mortality has risen since 2005.¹ Refining metrics for candidate evaluation may help reverse rising wait-list mortality.

Obesity, as measured by body mass index (BMI), is associated with increased risk of death after lung transplantation.² BMI cannot discriminate adipose from muscle whereas CT imaging can.³ CT quantification of body composition may improve lung transplant candidate selection and risk stratification.

We hypothesised that greater skeletal muscle adiposity, as measured by mean muscle attenuation and intramuscular adipose tissue (IMAT) area on thigh and abdominal CT scans, would be associated with decreased 6 min walk distance (6MWD) and wait-list survival among lung transplant candidates.

METHODS

We enrolled subjects 18 years old, undergoing transplant evaluation at Columbia University, Duke University or University of Pennsylvania between 2011 and 2014.

Subjects underwent single slice (10 mm) CT of the mid-thigh and/or at the L4/L5 vertebrae.⁴ Muscle was outlined by blinded personnel. Tissue within the outlined area with Hounsfield units (HU) of –40 to 124 was quantified as muscle (online supplementary figure S1), and tissue with adipose attenuation (–40 to –170 HU) was quantified as IMAT.

The primary outcome was 6MWD measured in a standardised fashion.⁵ Wait-list survival was operationalised as time from evaluation to death or delisting for severe illness. Subjects with unknown vital status were censored on 4 September 2019.

Predictors were mean muscle attenuation (lower attenuation indicating greater lipid content) and IMAT area/height² (IMAT index).⁶ Change in outcome was reported per ½-SD change in attenuation. IMAT index was log₂-transformed to normalise its distribution with change reported per doubling.

We evaluated predictor convergent and divergent validity with traditional body composition measures by Pearson correlation. We used mixed-effects linear regression to evaluate the association between predictors and 6MWD, adjusted for age, sex, diagnosis, forced vital capacity and muscle index (muscle cross-sectional area/height),² with centre as random effect. We controlled for BMI in sensitivity analyses given potential collinearity. Subgroup analysis was performed by diagnosis using likelihood ratios.

Survival analyses used competing risks regression for competing end points of transplantation and death/delisting adjusted for covariates above. Diagnosis violated the proportional hazards assumption and was included as a multiplicative interaction with time.

Analyses were performed using STATA/IC V.15.1 (StataCorp) and R V.3.3.1 (R-Foundation for Statistical Computing).

RESULTS

Three-hundred and six subjects had thigh or abdominal CT and 6MWD (online supplementary figure S2). Fifty-five per cent were men with median(IQR) age of 60(52–67) and BMI of 25.5 kg/m²(22.3–29.3); 48% had interstitial lung disease and 28% had chronic obstructive pulmonary disease (COPD) (online supplementary table S1). Subjects in the lowest tertile of muscle attenuation (ie, more adiposity) were older, more likely to have COPD and higher BMI (online supplementary tables S1–S2).

Thigh and abdominal muscle attenuation was weakly correlated with BMI while thigh IMAT index was moderately correlated with BMI (table 1). There was no correlation between thigh muscle attenuation or IMAT and thigh muscle index. Abdominal muscle attenuation was weakly correlated with abdominal muscle index.

In adjusted models, each ½-SD decrease in thigh muscle attenuation (indicating greater lipid content) was associated with a 15 m decrease in 6MWD (95% CI –21 to –10, p<0.001, figure 1A, table 2). Each ½-SD decrease in abdominal muscle attenuation was associated with a 14 m decrease in 6MWD (95% CI –20 to –8, p<0.001, figure 1B). Associations remained after controlling for BMI.

Each doubling in IMAT index was associated with a 13 m decline in 6MWD (95% CI –20 to –6, p=0.001, figure C). BMI was not associated with 6MWD (figure D).

Among subjects with suppurative lung disease, decreased muscle attenuation may be associated with increased walk distance, though this was not statistically significant (online supplementary table S3).

In the full cohort, 41 subjects died or were delisted; 144 underwent transplantation over median 2.3 years (IQR 0.5–4.4). Every ½-SD decrease in abdominal muscle attenuation was associated with 20% increased risk of death/delisting (subdistribution HR 1.2, 95% CI 1.1 to 1.4, p=0.006, table 2). Thigh muscle attenuation was not associated with death/delisting. Among subjects with IMAT, 24 died or were delisted, 97 were transplanted over median 2.3 years (IQR 0.5–4.5). There was no association between IMAT index and death/delisting. Survival was not assessed by subgroups due to small number of events for each diagnosis.

DISCUSSION

Greater skeletal muscle adiposity was associated with decreased 6MWD and wait-list survival. While BMI remains the primary metric for evaluating body composition in lung transplant candidates, it was not associated with walk distance. This contrasts with work

demonstrating an inverse association between BMI and walk distance in COPD and may reflect transplant candidate selection criteria. Ours adds to work demonstrating the utility of cross-sectional imaging to improve assessments of frailty,⁷ posttransplant mechanical ventilation and tracheostomy.⁸

Concordantly, prior work demonstrated an association between greater muscle adiposity and decreased 6MWD in COPD⁹ and decreased liver transplant survival.³ Numerous mechanisms link muscle adiposity and 6MWD. Increased fatty acid accumulation in and around muscle drives preferential utilisation of fatty acids resulting in decreased protein synthesis and regenerative capacity.¹⁰ Obese and aged adipose contains more senescent cells that impair muscle regeneration¹¹ and are associated with decreased gait speed.¹²

In contrast, greater muscle adiposity may be associated with increased 6MWD in suppurative lung disease. This may reflect nutritional status given known associations between higher BMI and greater 6MWD in cystic fibrosis.¹³

Adipose tissue is a buffer for free fatty acids, and in states of inflammation insufficient buffering may lead to ectopic deposition.¹⁴ Frailty is associated with decreased survival and greater abdominal visceral adipose tissue (VAT) deposition in lung transplant candidates.^{7,15} Whether muscle adiposity, which moderately correlates with VAT, is a similar marker of metabolic derangement or causes frailty, is unknown.

Our study has limitations. We found no association between thigh muscle attenuation and survival despite an association with abdominal muscle attenuation potentially reflecting underpowered analyses or different mechanisms of adipose deposition. Our cross-sectional analysis does not establish causality. Thirty-four subjects were censored at the time of analysis for unknown vital status, though missing deaths are unlikely given close clinical follow-up at transplant centres.

Cross-sectional imaging provides information not captured by BMI. Future work should investigate its role in candidate selection and prioritisation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding Supported by National Institutes for Health Grants: T32 HL105323, K24 HL131937, R01 HL 114626, R01 HL 134851, R01 HL 087115, K24 HL115354, Stony Wold-Herbert Foundation, VA-ORD: IK2 CX001034.

REFERENCES

1. Valapour M, Lehr CJ, Skeans MA, et al. OPTN/SRTR 2016 annual data report: lung. *Am J Transplant* 2018;18 Suppl 1:363–433. [PubMed: 29292602]
2. Singer JP, Peterson ER, Snyder ME, et al. Body composition and mortality after adult lung transplantation in the United States. *Am J Respir Crit Care Med* 2014;190:1012–21. [PubMed: 25233138]

3. Hamaguchi Y, Kaido T, Okumura S, et al. Impact of skeletal muscle mass index, intramuscular adipose tissue content, and visceral to subcutaneous adipose tissue area ratio on early mortality of living donor liver transplantation. *Transplantation* 2017;101:565–74. [PubMed: 27926595]
4. Katzmarzyk PT, Mire E, Bouchard C. Abdominal obesity and mortality: the Pennington center longitudinal study. *Nutr Diabetes* 2012;2:e42. [PubMed: 23168527]
5. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. Ats statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 2002;166:111–7. [PubMed: 12091180]
6. Frank-Wilson AW, Chalhoub D, Figueiredo P, et al. Associations of quadriceps torque properties with muscle size, attenuation, and intramuscular adipose tissue in older adults. *J Gerontol A Biol Sci Med Sci* 2018;73:931–8. [PubMed: 29342246]
7. Anderson MR, Kolaitis NA, Gao Y, et al. A nonlinear relationship between visceral adipose tissue and frailty in adult lung transplant candidates. *Am J Transplant* 2019;19:3155–61. [PubMed: 31278829]
8. Weig T, Milger K, Langhans B, et al. Core muscle size predicts postoperative outcome in lung transplant candidates. *Ann Thorac Surg* 2016;101:1318–25. [PubMed: 26794887]
9. Maddocks M, Shrikrishna D, Vitoriano S, et al. Skeletal muscle adiposity is associated with physical activity, exercise capacity and fibre shift in COPD. *Eur Respir J* 2014;44:1188–98. [PubMed: 24993908]
10. Hyde R, Hajdich E, Powell DJ, et al. Ceramide down-regulates system A amino acid transport and protein synthesis in rat skeletal muscle cells. *Faseb J* 2005;19:1–24. [PubMed: 15629889]
11. Belizário JE, Fontes-Oliveira CC, Borges JP, et al. Skeletal muscle wasting and renewal: a pivotal role of myokine IL-6. *Springerplus* 2016;5:619. [PubMed: 27330885]
12. Justice JN, Gregory H, Tchkonja T, et al. Cellular senescence biomarker p16INK4a+ cell burden in thigh adipose is associated with poor physical function in older women. *J Gerontol A Biol Sci Med Sci* 2018;73:939–45. [PubMed: 28658942]
13. Hollander FM, van Pierre DD, de Roos NM, et al. Effects of nutritional status and dietetic interventions on survival in cystic fibrosis patients before and after lung transplantation. *J Cyst Fibros* 2014;13:212–8. [PubMed: 24041590]
14. Yang S, Zhang W, Zhen Q, et al. Impaired adipogenesis in adipose tissue associated with hepatic lipid deposition induced by chronic inflammation in mice with CheW diet. *Life Sci* 2015;137:7–13. [PubMed: 26188592]
15. Singer JP, Diamond JM, Gries CJ, et al. Frailty phenotypes, disability, and outcomes in adult candidates for lung transplantation. *Am J Respir Crit Care Med* 2015;192:1325–34. [PubMed: 26258797]




Figure 1. Association between 6 min walk distance and (A) mean thigh muscle attenuation, (B) mean abdominal muscle attenuation, (C) thigh intramuscular adipose tissue index and (D) body mass index. Black line represents the mixed-effects linear model with median values for covariates (sex, forced vital capacity, diagnosis and age) and centre as random effect. Models A–C are additionally adjusted for muscle index (area/height²). Surrounding grey areas represent 95% confidence bands. Vertical lines along the x-axis each represent a single study subjects.

Table 1
Convergent and divergent validity of measures of muscle quality with traditional measures of body composition

	Thigh MMAr (95% CI)	P value	TMIr (95% CI)	P value	Thigh IMAT Index r (95% CI)	P value	Abdominal MMAr (95% CI)	P value	AMIr (95% CI)	P value
Thigh MMA	1									
TMI	0.05 (-0.07 to 0.16)	0.44	1							
Thigh IMAT Index	-0.52 (-0.62 to -0.41)	<0.001	0.07 (-0.07 to 0.21)	0.31	1					
Abdominal MMA	0.78 (0.72 to 0.82)	<0.001	0.11 (-0.01 to 0.22)	0.08	-0.47 (-0.57 to -0.35)	<0.001	1			
AMI	0.19 (0.07 to 0.30)	0.002	0.61 (0.54 to 0.68)	<0.001	-0.003 (-0.15 to 0.14)	0.97	0.27 (0.16 to 0.37)	<0.01	1	1
Body mass index	-0.35 (-0.45 to -0.24)	<0.001	0.59 (0.50 to 0.66)	<0.001	0.52 (0.41 to 0.62)	<0.001	-0.38 (-0.48 to -0.28)	<0.001	0.28 (0.18 to 0.38)	<0.01
Height	-0.05 (-0.17 to 0.06)	0.37	0.18 (0.06 to 0.29)	0.003	0.11 (-0.03 to 0.25)	0.12	0.1 (-0.01 to 0.21)	0.09	0.06 (-0.05 to 0.17)	0.28
Weight	-0.31 (-0.42 to -0.20)	<0.001	0.57 (0.49 to 0.65)	<0.001	0.47 (0.36 to 0.58)	<0.001	-0.25 (-0.36 to -0.15)	<0.001	0.27 (0.16 to 0.37)	<0.001
Abdominal VAT area (mm ²)	-0.33 (-0.43 to -0.22)	<0.001	0.20 (0.08 to 0.31)	0.001	0.40 (0.28 to 0.52)	<0.001	-0.44 (-0.53 to -0.34)	<0.001	0.06 (-0.06 to 0.17)	0.32
Triceps skinfold thickness	-0.19 (-0.35 to -0.03)	0.03	-0.11 (-0.27 to 0.06)	0.2	0.2 (-0.006 to 0.39)	0.06	-0.24 (-0.38 to -0.08)	0.003	-0.26 (-0.40 to -0.10)	0.001

Results displayed as Pearson correlation coefficient (95% CI).

AMI, abdominal muscle index (abdominal skeletal muscle area/height²); Thigh IMAT Index, thigh intramuscular adipose tissue (intramuscular adipose tissue area/height²); MMA, mean muscle attenuation (Hounsfield unit); TMI, thigh muscle index (thigh muscle area/height²); VAT, visceral adipose tissue.

Table 2

Association between muscle attenuation, thigh IMAT index, muscle attenuation and 6 min walk distance, gait speed and survival

6MWD	Change in 6MWD (m) per ½-SD decrease in thigh muscle attenuation				Change in 6MWD (m) per ½-SD decrease in abdominal mean muscle attenuation				
	N	95% CI	P value	N	95% CI	P value	N	95% CI	P value
Unadjusted	276	-19	<0.001	206	-10	0.01	302	-18	<0.001
Model 1*	275	-15	<0.001	205	-12	0.002	301	-14	<0.001
Model 2 [†]	275	-15	<0.001	190	-13	0.001	301	-14	<0.001
Model 3 [‡]	275	-13	<0.001	190	-6	0.18	301	-14	<0.001

Survival	Sub-hazard distribution for death or delisting per ½-SD decrease in thigh mean muscle attenuation				Subhazard distribution for death or delisting per doubling in thigh IMAT index				Subhazard distribution for death or delisting per ½-SD decrease in abdominal mean muscle attenuation						
	N	95% CI	P value	N	95% CI	P value	N	95% CI	P value	N	95% CI	P value	N	95% CI	P value
Unadjusted	276	1.1	0.40	206	1.0	0.86	302	1.2	0.86	302	1.2	0.86	302	1.2	0.04
Model 1*	275	1.1	0.31	205	1.0	0.83	301	1.2	0.83	301	1.2	0.83	301	1.2	0.06
Model 2 [†]	275	1.1	0.25	190	1.0	0.47	301	1.2	0.47	301	1.2	0.47	301	1.2	0.006
Model 3 [‡]	275	1.1	0.21	190	1.0	0.89	301	1.3	0.89	301	1.3	0.89	301	1.3	<0.001

* Adjusted for sex, age, forced vital capacity and diagnosis; centre as random effect in 6MWD models; standard errors clustered by centre in survival models.

[†]Model 1=skelletal muscle index.

[‡]Model 2=body mass index.

IMAT, intramuscular adipose tissue; 6MWD, 6 min walk distance.