

Lawrence Berkeley National Laboratory

Recent Work

Title

THE DEGRADATION OF COLCHICINE TO OCTAHYDEODMETHOXYDES-OXYDESACETAMIDOCHOLCHICINE1

Permalink

<https://escholarship.org/uc/item/4q679678>

Authors

Rapoport, Henry
Williams, Arthur R.
Campion, J.E.
et al.

Publication Date

1954-03-01

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

THE DEGRADATION OF COLCHICINE TO OCTAHYDRODEMETHOXY-
DESOXYDESACETAMIDOCOLCHICINE ¹

Henry Rapoport, Arthur R. Williams, ² John E. Campion,
and Donald E. Pack

March, 1954

Berkeley, California

THE DEGRADATION OF COLCHICINE TO OCTAHYDRODEMETHOXY-
DES OXYDESACETAMIDOCOLCHICINE¹

Henry Rapoport, Arthur R. Williams,² John E. Campion,
and Donald E. Pack

Department of Chemistry and Chemical Engineering and the Radiation Laboratory,
University of California, Berkeley, California

ABSTRACT

Although a complex mixture of difficultly separable products is formed on hydrogenation of colchicine (I), replacement of the methoxyl of ring C by dimethylamino and hydrogenation of the thus formed N,N-dimethylaminocolchicide (II) results in a good yield of easily purified tetrahydrodemethoxycolchicine (III). By conversion to the mercaptole and desulfurization, the carbonyl group is transformed to methylene giving hexahydrodemethoxydesoxycolchicine (V). Removal of the acetamido group is accomplished by heating with phosphorus pentoxide in xylene and hydrogenation yields octahydrodemethoxydesoxydesacetamidocolchicine (VI). This degradative sequence very reasonably takes place without rearrangement in the colchicine carbon skeleton and thus affords a degradation product whose synthesis would establish the nature of ring C.

-
- (1) Supported (in part) by Cancer Research Funds of the University of California and (in part) by the U.S. Atomic Energy Commission.
 - (2) American Cancer Society Postdoctoral Fellow, 1951.

THE DEGRADATION OF COLCHICINE TO OCTAHYDRODEMETHOXY-
DESOXYDESACETAMIDOCOLCHICINE¹

Henry Rapoport, Arthur R. Williams,² John E. Campion,
and Donald E. Pack

Department of Chemistry and Chemical Engineering and the Radiation Laboratory,
University of California, Berkeley, California

With the synthesis of colchinol methyl ether,^{3,4} the only remaining portion of the colchicine molecule (I) for which a definitive proof of structure is lacking is ring C. The features yet to be established by decisive chemical evidence are the seven-membered, tropoloid nature of the ring⁵ and the relative positions of the carbonyl and methoxyl groups.

Much evidence⁶ has accumulated supporting a tropolone methyl ether formulation as the most feasible structural representation for ring C. This includes

-
- (1) Supported (in part) by Cancer Research Funds of the University of California and (in part) by the U. S. Atomic Energy Commission.
 - (2) American Cancer Society Postdoctoral Fellow, 1951.
 - (3) H. Rapoport, A. R. Williams, and M. E. Cisney, *J. Am. Chem. Soc.*, **73**, 1414 (1951).
 - (4) J. W. Cook, J. Jack, and J. D. Loudon, *J. Chem. Soc.*, (1951), p. 1397.
 - (5) First proposed by M. J. S. Dewar, *Nature*, **155**, 141, 479 (1945).
 - (6) (a) H. R. V. Arnstein, D. S. Tarbell, G. P. Scott, and H. T. Huang, *J. Am. Chem. Soc.*, **71**, 2448 (1949); (b) G. P. Scott and D. S. Tarbell, *ibid.*, **72**, 240 (1950); (c) W. E. Doering and L. H. Knox, *ibid.*, **73**, 828 (1951); (d) J. W. Cook, A. R. Gibb, R. A. Raphael, and A. R. Somerville, *J. Chem. Soc.*, 503 (1951); (e) D. S. Tarbell and J. C. Bill, *J. Am. Chem. Soc.*, **74**, 1234 (1952).

spectral data,^{6b} hydrogenation to a glycol,^{6a,5} and the parallelism in properties between known tropoloid compounds and colchicine,^{6c,d,e} albeit in several instances there are marked differences in degree. This similarity in properties, as has been pointed out,^{6c} is necessary but insufficient to establish the tropoloid nature of ring C.

Although the carbonyl and methoxyl groups must be confined to the two positions shown in structure I on the basis of mechanistic interpretations^{6c,7} of the rearrangements of colchicine, only one attempt⁸ has been made to indicate the specific position of each group.⁹ From the similarity of infrared spectra (in the 7μ region) and optical rotations of colchicine and several isocolchicine derivatives, it was suggested that colchicine was a single species belonging to the iso series, and this non-tautomeric nature of colchicine could be due to hydrogen-bonding to the amide carbonyl. Thus the hydroxyl group of colchicine and the methoxyl of isocolchicine occupy the position more proximate to the acetamido group, and this conclusion supports structure I for colchicine.

We have sought direct chemical evidence for these structural features of ring C by unambiguous degradation to compounds, the synthesis of which would be feasible and would establish the structure in question. The present work reports the degradation to such a compound, octahydrodemethoxydesoxy-desacetamidocolchicine (VI), containing the intact carbon skeleton of col-


-
- (7) J. Čech and F. Šantavý, Collection Czech Chem. Commun., 14, 532 (1949).
- (8) R. M. Horowitz and G. E. Ulliot, J. Am. Chem. Soc., 74, 587 (1952).
- (9) Also, an X-ray diffraction study by King and Pepinsky (Abstracts Papers Am. Chem. Soc., 119, 33C (1951) is reported⁸ to give structure I for colchicine, but we could find no conclusions in the abstract.

chicine.¹⁰

A fruitful approach to the desired compound (VI) was suggested by the studies of Bursian¹¹ who reported the hydrogenation of colchicine and the isolation of hexahydrodemethoxycolchicine (IV) and a very small amount of what was apparently hexahydrodemethoxydesoxycolchicine (V). Since these compounds arise by mild catalytic hydrogenation, it is extremely probable that no rearrangement of the original, labile carbon skeleton of ring C is involved. In addition, they still retained a double-bond, and since the most reasonable position for this resistant-to-hydrogenation double bond was at the fusion of rings B and C, the stereoisomer problem would be obviated. However, in a recent re-examination¹² of this work, the carbinol (IV) was isolated only in very small yield after a laborious separation procedure, and none of the desoxy compound (V) was detected among the hydrogenation products.

An attractive alternative which might markedly facilitate separation of the hydrogenation products was to hydrogenate the derivative in which methoxyl had been replaced by amino. Non-hydrogenolyzed material now might be easily separable by virtue of its basic properties. The derivative

-
- (10) A preliminary report of this work appeared as a Communication to the Editor, H. Rapoport and A. R. Williams, J. Am. Chem. Soc., 73, 1896 (1951).
- (11) K. Bursian, Ber., 71, 245 (1938).
- (12) A. D. Kemp and D. S. Tarbell, J. Am. Chem. Soc., 72, 243 (1950).



MU-7336

chosen for this purpose was N,N-dimethylaminocolchicide (II),¹³ formed by heating colchicine with dimethylamine.

N,N-Dimethylaminocolchicide has been reported several times in the literature^{15,16,17} with a m.p. varying from 203° to 207°. Although we have prepared this compound numerous times, following as closely as the published details permit the prescribed conditions of the other investigators and also using some variations of our own, our purified material always melted

-
- (13) Some confusion exists in the nomenclature of this type of compound in which an amino group has replaced the methoxyl of colchicine. Zeisel (Monatsh., 9, 1 (1888) who first prepared the amino compound, called it the amide of colchicine, amide of acetyltrimethylcolchicinic acid, and colchicamide, and all three names have been used by various investigators. However, we find this nomenclature (which has its origin in the mistaken belief that colchicine was an ester and colchicine a carboxylic acid) misleading, since the compounds are distinctly basic in aqueous media, a property certainly not characteristic of amides. Furthermore, the term colchiceinamide,^{8,14,15} leads to difficulty in naming the compound in which carbonyl and amino groups are interchanged. Isocolchiceinamide implies a non-existent compound, isocolchicine, and the alternative colchicineisoamide implies an "isoamide" as opposed to a normal amide, which is clearly not intended. For these reasons, we have adopted the name aminocolchicide as being more suitable. The basic properties are clearly implied, and the root name colchicide may serve quite flexibly for any number of compounds in which the methoxyl is replaced by another group, colchicide itself serving for the replacement by hydrogen. This nomenclature very conveniently accommodates the iso series also through isocolchicide.
- (14) A. Uffer, Helv. Chim. Acta, 35, 2135 (1952).
- (15) J. L. Hartwell, M. V. Nadkarni, and J. Leiter, J. Am. Chem. Soc., 74, 3180 (1952).
- (16) A. J. Ewins, J. N. Ashley, and J. O. Harris, Brit. Patent 577,606 (1945).
- (17) F. Šantavý, Chem. Listy, 46, 280 (1952).

at 178-179°. That this difference in m.p. is not due to an impurity¹⁸ in our product is demonstrated by the fact that it displayed constant properties on crystallization from any of six different solvents, chromatography on alumina, or sublimation. Also, a paper chromatogram on alumina-impregnated paper¹⁹ showed only one spot under ultraviolet light. Fortunately, an optical rotation has been reported,¹⁷ and our value is in good agreement.²⁰ On the basis of the above data, we have concluded our compound is undoubtedly a dimorphic form, and its characteristic ultraviolet absorption spectrum is given in Fig. 1.


Since N,N-dimethylaminocolchicide (II) was a key compound in the degradation scheme, its structure was of considerable importance and was established by the fact that (a) hydrolysis gave colchicine and (b) hexahydrodemethoxycolchicine (IV) could be obtained by hydrogenation. The hydrolysis to colchicine indicates the same two positions of ring C that were occupied by methoxyl and carbonyl in colchicine were now occupied by dimethylamine and carbonyl; and the hydrogenation to IV, the same compound obtained from the hydrogenation of colchicine,^{11,12} indicates the carbonyl group has retained its initial position. Therefore, N,N-dimethylaminocolchicide (II) has resulted through replacement of the methoxyl of colchicine by dimethylamino.²¹

-
- (18) A possible but highly improbable prospect that our compound was N-methylaminocolchicide for which m.p.'s of 173-174°,¹⁶ 176-178°,¹⁷ and 230-232°¹⁵ have been recorded was eliminated by the marked depression exhibited in a mixed m.p. determination with authentic N-methylaminocolchicide (m.p. 172-174°).
- (19) S. P. Datta and B. G. Overell, *Biochem. J.*, 44, xliii (1949); S. P. Datta, B. G. Overell and M. Stack-Dunne, *Nature*, 164, 673 (1949).
- (20) In this regard, it should be noted that the optical rotation is quite variable with temperature, a not uncommon behavior for colchicine derivatives. For N,N-dimethylaminocolchicide in chloroform, the specific rotation varies inversely with the temperature to the extent of 4.1°/°C.
- (21) A comparison of the specific rotation of N,N-dimethylaminocolchicide $[\alpha]_D^{27.5} + 465^\circ$ (chloroform) and N,N-dimethylaminoisocolchicide

As was expected, N,N-dimethylaminocolchicide was quite reactive toward hydrogen. After an initial, rapid absorption of three moles of hydrogen, absorption continued at a slower but still appreciable rate and ceased at a final value of five moles. Both three-mole and five-mole reaction mixtures were examined and each found to consist of a neutral and basic portion. In neither case could any crystalline material be isolated from the basic fraction. The neutral fraction, however, afforded hexahydrodemethoxycolchicine (IV) from the five-mole reaction and tetrahydrodemethoxycolchicine (III) from the three-mole reaction. Since the ketone appeared more promising for conversion to the desoxy compound (V), the three-mole hydrogenation was investigated in detail and conditions established which led to a 53% yield of tetrahydrodemethoxycolchicine (III),²² isolated via its bisulfite addition product.

[α]_D²⁷ = 520° (chloroform), unpublished work by J. B. Lavigne] is also consistent with this structural assignment according to the generalizations of Horowitz and Ulliot⁸ (see also Uffer, ref. 14) who find the iso compound of a given pair to have the more negative rotation.

- (22) In view of the strong evidence for the carbon skeleton of colchicine (I) and therefore for that of tetrahydrodemethoxycolchicine (III), we have employed the C. A. numbering method of the parent ring system, benzo[a]heptalene,



to locate substituents. However, the common names as derived from colchicine have been used for the degradation products since they are more lucid at present.

The carbonyl group of III is undoubtedly at the same position as that of colchicine since on hydrogenation both yield the carbinol (IV). This position is either 9 or 10, depending on the relative position of methoxyl and carbonyl in colchicine. The double bond that remains resistant to hydrogenation has been placed at the fusion of rings B and C since this position seems the most hindered toward catalyst adsorption. However, unsaturation between carbons 12 and 12a also might be subject to considerable resistance to hydrogenation, and although 7a-12a appears the most reasonable location for the double bond, the 12-12a alternative certainly obtains for tetrahydrodemethoxycolchicine (III) and any of the succeeding compounds. Both formulations are consistent with the ultraviolet and infrared absorption spectra (Figs. 2 and 3) which indicate the presence of a conjugate double bond and benzene ring, and an unconjugate carbonyl.

Conversion to the dimethylmercaptole²³ and desulfurization proceeded readily and in excellent yield to hexahydrodemethoxydesoxycolchicine (V). Although the olefinic double bond remained resistant to catalytic hydrogenation, its presence was clearly demonstrated by the consumption of one mole of perbenzoic acid in chloroform at 0° and isolation of a crystalline epoxide.²⁴

-
- (23) Using the general procedure of M. S. Newman and H. M. Walborsky, J. Am. Chem. Soc., 72, 4296 (1950).
- (24) Although methoxylated benzene nuclei react with perbenzoic acid [H. Fernholtz, Ber., 84, 110 (1951); S. L. Friess et al., J. Am. Chem. Soc., 72, 2611 (1950), 74, 1305 (1952)] the difference in rate as compared to the alicyclic double bond is of such great magnitude that there is no difficulty in isolating the one mole product in the case of compounds V and VI. After the initial rapid reaction, further consumption of perbenzoic acid continued but at a tremendously decreased rate.

To remove the acetamido group from hexahydrodemethoxydesocylchicine (V), it was subjected to the action of phosphorus pentoxide in refluxing xylene.²⁵ By strict adherence to reaction conditions, reproducible results could be obtained, and the reaction product on immediate hydrogenation absorbed one mole of hydrogen and gave octahydrodemethoxydesoxydesacetamidocolchicine (VI). Again, the persistence of a double bond was shown by consumption of one mole of perbenzoic acid and isolation of an epoxide.

The ultraviolet and infrared²⁶ absorption spectra of the compounds in the degradation scheme are shown in Figs. 1, 2, and 3. In every case, the spectra are consistent with the structures presented, although they do not permit a unique structure assignment (e.g., as to whether the compounds are $\Delta^{7a(12a)}$ or Δ^{12}). Considering the method of degradation employed and the avoidance of rearrangement, octahydrodemethoxydesoxydesacetamidocolchicine (VI) appears to be a compound whose synthesis would establish definitively the nature of ring C in colchicine.

(25) J. W. Cook, W. Graham, A. Cohen, R. W. Lapsley, and C. A. Lawrence, J. Chem. Soc., (1944), p. 322.

(26) We are indebted to Dr. N. K. Freeman of the Radiation Laboratory for the infrared spectra and helpful discussions concerning their interpretation.

Experimental²⁷

Colchicine (I). — Colchicine (U.S.P.) was purified by chromatography²⁸ followed by crystallization from ethyl acetate, m.p. 154-155°.

N,N-Dimethylaminocolchicide (II). — A solution of pure dimethylamine in methanol was prepared by adding saturated methanolic potassium hydroxide to dimethylamine hydrochloride, purified by several crystallizations from absolute ethanol, and absorbing the liberated amine in methanol. To 80 ml. of such a solution (3N) was added 31.1 g. (78 mmoles.) of colchicine and the resulting solution was heated in a sealed tube completely immersed in an oil bath at 105° for 18 hours. After cooling, the contents of the tube were evaporated to a thick syrup which was dissolved in 150 ml. of benzene and then extracted with three portions of 2N hydrochloric acid and two portions of water (50 ml. each). The combined aqueous extracts, after washing with benzene, were made strongly alkaline with sodium hydroxide solution and extracted with three 100 ml. portions of benzene. Evaporation of the washed (water) and dried benzene layers gave a residue which was chromatographed on alumina (Merck) using benzene-chloroform (1:1) as eluent. Crystallization of the eluted material from ethyl acetate gave 23.7 g.

(27) All melting points are corrected and those above 200° were taken in evacuated capillaries; microanalyses were performed by the Microchemical Laboratory, University of California.

(28) J. N. Ashley and J. O. Harris, J. Chem. Soc., (1944), p. 677.

(58 mmoles., 74% yield) of N,N-dimethylaminocolchicide, m.p. 175-177°.

Recrystallization gave material of m.p. 178-179°, $[\alpha]_D^{25} + 69.4^\circ$ (c, 1.03, ethanol), $[\alpha]_D^{17} + 508^\circ$ (c, 1.04, chloroform), $[\alpha]_D^{27.5} + 465^\circ$ (c, 1.00, chloroform) reported m.p. 204-206°,¹⁶ 203-205°,¹⁵ 205-207°,¹⁷ $[\alpha]_D^{21} + 510^\circ$ (c, 0.79, chloroform).¹⁷

Anal. Calcd. for $C_{23}H_{28}N_2O_5$: C, 67.0; H, 6.8; N, 6.8; OCH_3 , 22.6

Found: C, 66.9; H, 6.9; N, 7.0; OCH_3 , 22.2.

Further chromatography on alumina, sublimation, or recrystallization from ether, acetone, ethanol, methyl isobutyl ketone, or xylene had no effect on the properties of N,N-dimethylaminocolchicide, and an alumina-impregnated paper chromatogram showed only one spot. A mixed melting point determination with N-methylaminocolchicide¹⁸ showed a depression of 20°.

The picrate, prepared with saturated ethanolic picric acid, was recrystallized from absolute ethanol, m.p. 186-188°, $[\alpha]_D^{25} + 171^\circ$ (c, 1.08, chloroform).

Anal. Calcd. for $C_{29}H_{31}N_5O_{12}$: C, 54.3; H, 4.9; OCH_3 , 14.5.

Found: C, 54.3; H, 5.0; OCH_3 , 14.3.

Hydrolysis of N,N-Dimethylaminocolchicide to Colchicine. -- A mixture of 50 mg. (0.12 mmole.) of N,N-dimethylaminocolchicide, and 5 ml. of 0.1 N sodium hydroxide was heated on the steam bath for four hours, then cooled, washed with 5 ml. of benzene, and acidified with one ml. of 1 N hydrochloric acid. The mixture was extracted with three 5 ml. portions of chloroform and evaporation of the combined chloroform extracts after drying gave a residue which was digested on the steam bath with one ml. of saturated sodium bicarbonate solution and filtered. Acidification of the bicarbonate solution with 1 N hydrochloric acid and cooling several hours precipitated

material, which was crystallized from ethyl acetate-n-butyl ether to yield 27 mg. (.07 mmole., 59%) of colchicine, m.p. and mixed m.p. with an authentic sample, 178-179°.

Tetrahydrodemethoxycolchicine (III). — Hydrogenation of 4.0 g. (9.7 mmoles.) of N,N-dimethylaminocolchicine in 120 ml. of glacial acetic acid proceeded rapidly in the presence of 500 mg. of 5% palladized carbon and 250 mg. of platinum oxide at 20° and a pressure of 30 p.s.i. Three moles of hydrogen were absorbed in about 19 minutes, and an additional 0.1 mole in the next two minutes after which the hydrogenation was stopped and the reaction mixture was filtered. The filtrate was concentrated under reduced pressure, basified with 6 N sodium hydroxide, extracted with three 30 ml. portions of benzene, and the separate benzene extracts were in turn extracted with two 25 ml. portions of 2 N hydrochloric acid and two 20 ml. portions of distilled water. Evaporation of the combined, dried, benzene extracts left a yellow viscous oil which was digested for successive 30 minute periods on a steam bath with two 50 ml. portions of 20% aqueous sodium bisulfite and one 50 ml. portion of water. The combined aqueous digests were basified with potassium carbonate and extracted with four 30 ml. portions of benzene which were then washed, dried, concentrated to 25 ml., and applied to a column (30 x 1 cm.) of alumina (Merck). Elution was accomplished with 500 ml. of chloroform, from which on evaporation the ketone was obtained as a white amorphous solid, 1.91 g. (5.1 mmoles., 53%). This material was of suitable purity for use in the degradative sequence; however, crystallization from a mixture of ethyl acetate and n-butyl ether (2:1) gave a total of 1.61 g. (4.3 mmoles., 44%) of tetrahydrodemethoxycolchicine in several crops, melting

variously between 140-144°. A recrystallized sample melted at 143-144°;
 $[\alpha]_D^{25} = 174^\circ$ (c, 1.11, ethanol).

Anal. Calcd. for $C_{21}H_{27}O_5N$: C, 67.5; H, 7.3; N, 3.8; $-OCH_3$, 24.9.

Found: C, 67.5; H, 7.3; N, 3.8; $-OCH_3$, 24.8.

Hexahydrodemethoxycolchicine (IV). A. By hydrogenation of N,N-dimethylaminocolchicide (II). -- Hydrogenation of 0.5 g. (1.2 mmoles.) of N,N-dimethylaminocolchicide in 12 ml. of glacial acetic acid was allowed to proceed for 72 hours at 25° and atmospheric pressure with 50 mg. of 5% palladized carbon and 25 mg. of platinum oxide, at which time hydrogen absorption (4.9 moles) ceased. The neutral fraction (.21 g.), on crystallization from ethyl acetate gave hexahydrodemethoxycolchicine, m.p. 168-170°, $[\alpha]_D^{25} = 166^\circ$ (c, 1.01, ethanol) (reported m.p. 171⁰¹¹ and 173⁰¹²).

Anal. Calcd. for $C_{21}H_{29}O_5N$: C, 67.2; H, 7.8; N, 3.7.

Found: C, 67.2; H, 7.8; N, 3.9.

B. By hydrogenation of Tetrahydrodemethoxycolchicine (III). --

A solution of 0.25 g. (0.7 mmole.) of tetrahydrodemethoxycolchicine in 5 ml. of glacial acetic acid was hydrogenated over 10 mg. of platinum oxide and 20 mg. of 5% palladized carbon, and resulted in the absorption of 0.95 mole of hydrogen. After removal of the catalyst, the solution was made alkaline with 6 N sodium hydroxide and extracted with three 25 ml. portions of benzene. Drying and evaporating the benzene left a residue which was crystallized from ethyl acetate to give 0.16 g. (0.4 mmole., 64%) of hexahydrodemethoxycolchicine, m.p. 168-170° after drying at 100° in vacuo. This material was identical with the neutral material from the 5 mole hydrogenation of N,N-dimethylaminocolchicide.

When treated with acetic anhydride, hexahydrodemethoxycolchicine formed an acetate, m.p. 206-208° (reported¹¹ m.p. 210°).

Tetrahydrodemethoxycolchicine dimethylmercaptole. -- A sealed tube containing 0.91 g. of anhydrous fused zinc chloride, 0.91 g. of anhydrous sodium sulfate, 3.19 g. (8.5 mmoles.) of tetrahydrodemethoxycolchicine (III), and 36 ml. of methyl mercaptan was shaken until the ketone and zinc chloride dissolved and then allowed to stand at room temperature for 18 hours. The tube was opened, the methyl mercaptan evaporated, and the residue dissolved in chloroform. After being washed with water and 1 N sodium hydroxide, the chloroform was evaporated and the crude mercaptole was crystallized from aqueous methanol; yield, 3.21 g., 83%; m.p. 190-192°; $[\alpha]_D^{25} - 160^\circ$ (c, 0.96, ethanol).

Anal. Calcd. for $C_{23}H_{33}O_4NS_2$: C, 61.2; H, 7.4; S, 14.2.

Found: C, 61.4; H, 7.5; S, 14.0.

Hexahydrodemethoxydesoxycolchicine (V). -- A vigorously stirred and refluxing mixture of 3.26 g. (7.2 mmoles.) of tetrahydrodemethoxycolchicine dimethylmercaptole in 300 ml. of 90% aqueous ethanol, and 64 g. of Raney nickel²⁹ was filtered after 16 hours. The nickel was digested with two 200 ml. portions of benzene and the combined benzene digests and ethanol filtrate evaporated. Crystallization of the residue from aqueous methanol gave 2.29 g., 88%, of hexahydrodemethoxydesoxycolchicine, m.p. 183.5-184°; $[\alpha]_D^{25} - 162^\circ$ (c, 1.10, ethanol).

Anal. Calcd. for $C_{21}H_{29}O_4N$: C, 70.2; H, 8.1; OCH_3 , 25.9.

Found: C, 70.1; H, 8.2; OCH_3 , 26.0.

(29) R. Mozingo, "Organic Syntheses," Vol. 21, John Wiley and Sons, New York, N. Y., 1941, p. 15.

Hexahydrodemethoxydesoxycolchicine epoxide. — Using a 0.04 M solution of perbenzoic acid in chloroform,³⁰ the epoxidation of hexahydrodemethoxydesoxycolchicine (V) at 0° was followed by withdrawal of aliquots and determination of unreacted perbenzoic acid by titration in the usual manner.³⁰ During 3.2 hours, one mole of perbenzoic acid was consumed per mole of compound after which the rate markedly decreased and during the next three hours only an additional 0.1 mole of perbenzoic acid was consumed. A parallel reaction containing originally 100 mg. (0.28 mmole.) of hexahydrodemethoxydesoxycolchicine in 25 ml. of chloroform (0.04 M in perbenzoic acid) was allowed to continue for four hours during which 105 mole % of perbenzoic acid was consumed as determined by the removal of five 1 ml. aliquots. To quench the reaction, 10 ml. of 1 N potassium carbonate solution was then added and the chloroform phase was washed with two 10 ml. portions of water. Evaporation of the dried chloroform solution left 80 mg. of crystalline material which was applied as a benzene solution to an alumina (Merck) column (22 x 1 cm.). Benzene (150 ml.) and 25% chloroform-benzene (150 ml.) eluted 62 mg. (0.17 mmole., 74%) of colorless, crystalline epoxide. Two crystallizations from hexane-benzene gave hexahydrodemethoxydesoxycolchicine epoxide of m.p. 209.5–210° and $[\alpha]_D^{25} + 28.9^\circ$ (c, 0.85, ethanol).

Anal. Calcd. for $C_{21}H_{29}O_5N$: C, 67.2; H, 7.8; OCH_3 , 24.8.

Found: C, 67.2; H, 7.7; OCH_3 , 24.0.

(30) G. Braun, "Organic Syntheses," Coll. Vol. I, John Wiley and Sons, New York, N. Y., 1947, p. 431.

Octahydrodemethoxydesoxydesacetamidocolchicine (VI). -- To a solution of 0.5 g. (1.4 mmoles.) of hexahydrodemethoxydesoxycolchicine in 50 ml. of xylene (purified by distillation from phosphorus pentoxide) heated to 100° was added 1.0 g. of phosphorus pentoxide with vigorous stirring and in a nitrogen atmosphere. The stirred mixture was heated to reflux in 10 minutes and reflux was continued for 25 minutes. After cooling and decanting the xylene, the residue was digested with two 100 ml. portions of benzene, and the combined digests were allowed to cool, filter aid was added and the solution filtered. Evaporation of the filtrate left a viscous oil which was dissolved in two ml. of hexane (in which hexahydrodemethoxydesoxycolchicine is insoluble) and filtered from a small amount of solid. The 370 mg. of oil obtained on evaporation of the hexane was dissolved in 5 ml. of glacial acetic acid and hydrogenated over 20 mg. of platinum oxide and 40 mg. of 5% palladized carbon. Hydrogen absorption ceased at 110 mole %, after which the catalyst was removed by filtration, and the filtrate, made basic with 6 N sodium hydroxide, was extracted with two 10 ml. portions of benzene. The combined benzene extracts were washed with 10 ml. portions of water, 2 N hydrochloric acid, and 1 N potassium carbonate and were then dried and evaporated to an oil which was dissolved in 10 ml. of hexane and applied to a column (15 x 1 cm.) of alumina (Merck). The octahydrodemethoxydesoxydesacetamidocolchicine was eluted with 20% benzene-hexane (200 ml.) after hexane (50 ml.) had removed only a trace of material. Slow sublimation at 40°/64 of the white solid, m.p. 49-50°, obtained by evaporation of the benzene-hexane eluent gave 250 mg. (60% yield) of material still melting at 49-50°; $[\alpha]_D^{25}$ 0° (c, 1.01, ethanol).


Anal. Calcd. for $C_{19}H_{26}O_3$: C, 75.5; H, 8.7; $-OCH_3$, 30.8; $C-CH_3$, 0.

Found: C, 75.4; H, 8.7; $-OCH_3$, 30.9; $C-CH_3$, <0.5.

Octahydrodemethoxydesoxydesacetamidocolchicine epoxide. -- Perbenzoic acid oxidation of octahydrodemethoxydesoxydesacetamidocolchicine (VI) was carried out as described above for hexahydrodemethoxydesoxycolchicine and resulted in the consumption of one mole of perbenzoic acid per mole of compound. An oxidation reaction which consumed 82 mole % of perbenzoic acid and then became very much slower was treated as above in order to isolate epoxide. The chloroform residue was applied to the alumina column in hexane and this was followed by 10% benzene-hexane, 20% benzene-hexane, and benzene. Evaporation of the benzene gave crystalline epoxide in 42% yield and this was recrystallized from hexane; m.p. 116-117°.


Anal. Calcd. for $C_{19}H_{26}O_4$: C, 71.7; H, 8.2.

Found: C, 72.0; H, 8.2




MU-7333

Fig. 1



MU-7334

Fig. 2



MU-7335

Fig. 3