

UCSF

UC San Francisco Previously Published Works

Title

A cohort study of maternal cardiometabolic risk factors and primary cesarean delivery in an integrated health system.

Permalink

<https://escholarship.org/uc/item/4qd621zc>

Journal

PLoS ONE, 13(7)

Authors

Hedderson, Monique

Xu, Fei

Sridhar, Sneha

[et al.](#)

Publication Date

2018

DOI

10.1371/journal.pone.0199932

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

RESEARCH ARTICLE

A cohort study of maternal cardiometabolic risk factors and primary cesarean delivery in an integrated health system

Monique M. Hedderson^{1*}, Fei Xu¹, Sneha B. Sridhar¹, Emily S. Han¹, Charles P. Quesenberry¹, Yvonne Crites²

1 Division of Research, Kaiser Permanente Northern California, Oakland, California, United States of America, **2** The Kaiser Permanente Northern California Medical Group, Oakland, California, United States of America

* Monique.M.Hedderson@kp.org



OPEN ACCESS

Citation: Hedderson MM, Xu F, Sridhar SB, Han ES, Quesenberry CP, Crites Y (2018) A cohort study of maternal cardiometabolic risk factors and primary cesarean delivery in an integrated health system. *PLoS ONE* 13(7): e0199932. <https://doi.org/10.1371/journal.pone.0199932>

Editor: Kelli K. Ryckman, University of Iowa, UNITED STATES

Received: February 16, 2018

Accepted: June 15, 2018

Published: July 3, 2018

Copyright: © 2018 Hedderson et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Individual level data may not be made publicly available due to IRB and privacy concerns. The data used for this study contain protected health information (PHI) and access is protected by the Kaiser Permanente Northern California Institutional Review Board (IRB). Data are available from the Kaiser Permanente Division of Research for researchers who meet the criteria for access to confidential data. For more information about data access and criteria for access to confidential data, please contact Kaiser Permanente Division of Research:

Abstract

Background

Maternal cardiometabolic risk factors (i.e., hyperglycemia, pre-existing hypertension and high body mass index) impact fetal growth and risk of having a cesarean delivery. However, the independent and joint contribution of maternal cardiometabolic risk factors to primary cesarean section is unclear. We aimed to elucidate the degree to which maternal cardiometabolic risk factors contribute to primary cesarean deliveries and whether associations vary by infant size at birth in an integrated health system.

Methods

A cohort study of 185,045 singleton livebirths from 2001 to 2010. Poisson regression with robust standard errors provided crude and adjusted relative risks (RR) and 95% confidence intervals (CIs) for cesarean delivery risk associated with risk factors. We then estimated the proportion of cesarean sections that could be prevented if the cardiometabolic risk factor in pregnant women were eliminated (the population-attributable risk [PAR]).

Results

In a single multivariable model, maternal cardiometabolic risk factors were independently associated with cesarean delivery: RR (95% CI) abnormal glucose screening 1.04 (1.01–1.08); gestational diabetes 1.18 (1.11–1.18) and pre-existing diabetes 1.60 (1.49–1.71); pre-existing hypertension 1.16 (1.10–1.23); overweight 1.27 (1.24–1.30); obese class I 1.46 (1.42–1.51); obese class II 1.73 (1.67–1.80); and obese class III 1.97 (1.88–2.07); adjusting for established risk factors, medical facility and year. The associations between maternal cardiometabolic risk factors and primary cesarean delivery remained among infants with appropriate weights for gestational age. The PARs were 17.4% for overweight/obesity, 7.0% for maternal hyperglycemia, 2.0% for pre-existing hypertension and 20.5% for any cardiometabolic risk factor.

DOR.IRB.Submissions@kp.org. All other relevant data are within the paper.

Funding: This research was supported by a Kaiser Permanente Community Benefit grant (MMH). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Conclusions

Maternal cardiometabolic risk factors were independently associated with risk of primary cesarean delivery, even among women delivering infants born at an appropriate size for gestational age. Effective strategies to increase the proportion of women entering pregnancy at an optimal weight with normal blood pressure and glucose before pregnancy could potentially eliminate up to 20% of cesarean deliveries.

Introduction

Primary cesarean delivery rates in the United States increased from 14.6% of births in 1996[1] to 21.9% in 2012[2]. Cesarean sections are now the most common surgery performed in the United States. This rapid increase occurred despite the fact that cesarean delivery increases the risk of adverse perinatal outcomes, including surgical complications, severe hemorrhage, puerperal infection and cardiac arrest[3] as well as complications requiring admission to the neonatal intensive care unit, obstetrical care related costs and the likelihood of delivery by cesarean section in subsequent births[4, 5]. A 2011 Joint Commission report determined that there are no data supporting that higher rates of cesarean sections improve perinatal outcomes, and yet the rates continue to rise[6]. Thus, reducing primary cesarean deliveries is essential to reduce the related morbidities. Studies are needed on potentially modifiable risk factors associated with primary cesarean delivery to inform future interventions to reduce rates of cesarean deliveries[6].

A growing body of literature is showing the multifactorial reasons for the rise in cesarean sections. Increases in obesity[7–9], pre-gestational diabetes[10], gestational diabetes mellitus (GDM)[11], and hypertension, for example, have occurred in parallel to the rise in primary cesarean section deliveries in recent decades, which provides evidence that adverse maternal cardiometabolic profile in pregnancy contributes to the high rates of primary cesarean section delivery. While several studies have examined the separate effects of obesity, glucose intolerance, and hypertension in relation to cesarean section[12–17], the independent and joint effects of these cardiometabolic risk factors is unclear. In addition, while these maternal cardiometabolic risk factors impact fetal growth[18, 19], whether the impact of these risk factors on risk of cesarean delivery varies by infant size for gestational age remains unclear.

This cohort study examined the independent and joint associations between maternal cardiometabolic risk factors (BMI, pregnancy glycemia and pre-existing hypertension) and primary cesarean delivery overall and by infant size for gestational age in a racially-ethnically diverse cohort of 185,045 women delivering in Kaiser Permanente Northern California (KPNC) from 2001 to 2010. Our overall aim was to elucidate the degree to which maternal cardiometabolic risk factors contribute to primary cesarean deliveries, and whether these associations vary by infant size at birth in an integrated health care system with uniform access to care and finally to quantify the extent to which we could reduce the rates of cesarean section if we improved maternal cardiometabolic risk factors at a population level.

Materials and methods

Kaiser Permanente Northern California (KPNC) provides comprehensive medical services through 14 delivery hospitals and 23 outpatient clinics to over 3 million members located in a 14-county region of Northern California. KPNC membership is well-representative of the

population living in the geographical area served by this large, integrated health care delivery system, except that the KPNC population has slightly lower representation at the extremes of income[20]. As KPNC is an integrated health care delivery system, all pregnant women have access to care in a setting where the cost structure does not incentivize cesarean sections.

We identified 204,231 singleton live births to women aged 15–45 between 2001 and 2010 among women without a prior cesarean section. We then matched this cohort to the California birth certificate record (99% successful linkage). We excluded women whose gestational age was less than 24 weeks or greater than 45 weeks ($N = 3,274$). We also excluded women for whom we were unable to verify whether the delivery resulted in a vaginal birth or cesarean delivery ($N = 1,553$). To confirm the delivery was a primary cesarean section, we first used ICD-9 code 654.2x to identify whether women had a prior cesarean, which we confirmed by verifying that the California state birth certificate was coded as a repeat cesarean. We excluded 2,059 deliveries for whom we were unable to confirm if they had a prior cesarean delivery consistently between these two sources. We also excluded deliveries at medical facilities that were not in existence during the entire study period ($N = 12,300$), leaving us with a final analytic cohort of 185,045. Of these, there were 152,011 unique women in the cohort due to the inclusion of repeat pregnancies.

Cesarean delivery was based on ICD-9 codes available in the EMR: 74.xx (excluding 74.3 and 74.91) and ICD-9 diagnosis: 669.7. Primary cesarean deliveries identified by ICD-9 codes were also verified against the California birth certificate record and the KPNC infant cohort database[21] (~98.9% were validated based on at least two sources).

We obtained maternal age at delivery, height, pregnancy body weight (assessed on average at 17 weeks' gestation, with a range of 6–26 weeks' gestation), and gestational age at the weight measurement from electronic medical records (EMR).

We searched the KPNC Diabetes Registry[22] to identify women with pre-existing diabetes. We then searched the KPNC Pregnancy Glucose Tolerance Registry[11], which classifies women without pre-existing diabetes by glucose values obtained from a 50-g glucose challenge test (hereafter referred to as the screening test). Women with an abnormal result (≥ 140 mg/dl) receive a follow up 100-g, 3-hour oral glucose tolerance test (hereafter referred to as the diagnostic test). At KPNC, 94% of women delivering live-born singletons undergo the recommended screening test for GDM[11].

We classified women as having GDM if two or more of the four plasma glucose values obtained during the 3-hour diagnostic test were abnormal according to the Carpenter-Coustan plasma glucose thresholds for GDM (fasting, 95 mg/dl; 1-hour, 180 mg/dl; 2-hour, 155 mg/dl; 3-hour, 140 mg/dl) or if there was an outpatient diagnosis of GDM[23]. Abnormal screening was defined as women having an abnormal screening test (≥ 140 mg/dl), but a normal diagnostic test (one or fewer abnormal values on the 3-hour diagnostic test).

We calculated pregnancy BMI as the maternal weight (kilograms) divided by height (meters) squared. BMI categories were created based on the World Health Organization International Classification: underweight (< 18.5 kg/m²), normal (18.5–24.99 kg/m²), overweight (25.0–29.99 kg/m²), obese (30.00–34.99 kg/m²), obese class II (35.0 kg/m²–39.9 kg/m²), and obese class III (40.0 kg/m² or higher)[24].

Covariates

Infant size for gestational age. We defined large for gestational age (LGA) as birthweight greater than the 90th percentile and small for gestational age (SGA) as birthweight less than the 10th percentile, both according to the study population's race-ethnicity and gestational age-specific birth weight distribution.

We obtained maternal education and race-ethnicity from the California birth certificate. We identified the following potential indications for cesarean section: obstructed labor using ICD-9 codes: 660.0, 660.3 and 660.4; fetopelvic disproportion using ICD-9 codes: 653.4–653.9; breech position using ICD-9 codes 652.2 and Placenta previa using ICD-9 codes 641.0 and 641.1. Pre-existing hypertension as essential hypertension; hypertension secondary to a chronic underlying condition; pre-eclampsia or eclampsia superimposed on pre-existing hypertension; and unspecified hypertension of pregnancy (ICD-9 codes 642.0, 642.1, 642.2 and 642.7).

Statistical analysis

Univariable and multivariable Poisson regression with robust standard errors were used to provide crude and adjusted estimates of relative risks (RR) and 95% CIs for cesarean delivery risk associated with maternal cardiometabolic factors [25]. RR estimates were adjusted for maternal age at delivery, race-ethnicity, maternal education, parity, and medical facility to account for possible changes in the demographics and underlying risk factor for cesarean deliveries in the study population. We calculated partial population attributable risks (PAR) for being overweight or obese during pregnancy by the method described by Spiegelman and colleagues [26], using a publicly-available SAS Macro <http://www.hsph.harvard.edu/faculty/spiegelman/par.html>. We used the KPNC population data to estimate the prevalence of GDM for the PAR calculations. We examined effect modification between maternal cardiometabolic risk factors and infant size for gestational age and all were significant (p-interaction BMI = 0.001, maternal glucose tolerance status: 0.003 and hypertension 0.010). Therefore, we conducted a stratified analysis by infant size for gestational age to see if the associations with maternal cardiometabolic risk factors were independent of infant size. We used SAS version 9.3 (SAS Institute Inc., Cary, NC) for all analyses.

The Kaiser Permanente Northern California Institutional Review Board and the State of California Committee for the Protection of Human Subjects approved this study and waived the requirement for obtaining written informed consent from study participants. Patient records were anonymized and de-identified prior to analysis.

Results

The cohort was racially and ethnically diverse with more than 63% from non-white racial ethnic minority groups and over half of the women were overweight or obese, half of women were nulliparous and just over one third of women had a college education or higher (Table 1).

Several maternal cardiometabolic risk factors appear to independently contribute to risk of cesarean delivery above and beyond other established risk factors. Table 2 shows the relative risks (RRs) and 95% CIs from a single multivariable model containing all cardiometabolic risk factors; the risk of having a cesarean delivery increased along with maternal BMI, increasing glucose intolerance, and maternal hypertension. All maternal cardiometabolic risk factors were significantly and independently associated with having a cesarean section delivery. These associations remained significant after further adjusting for other indications for cesarean delivery: fetopelvic disproportion, obstructed labor, placenta previa and breech position (Table 2). In addition, women delivering either SGA [RR 1.12 (95% CI: 1.09–1.16)] or LGA [RR 1.44 (95% CI: 1.40–1.49)] infants also had an increased risk of cesarean delivery when compared with women delivering appropriate size for gestational age infants (Table 2). We found a trend of increasing risk of cesarean delivery with increasing number of cardiometabolic risk factor (RR: 0 risk factors = reference group, 1 risk factor 1.37 (95% CI, 1.34–1.41), 2

Table 1. Demographic characteristics by calendar year: Kaiser Permanente Northern California, 2001–2010¹.

<i>Maternal/infant factor</i>	N = 185,045
	n (%)
Primary cesarean rate	30,347 (16.4)
Age at delivery (years)	
Mean ± SD	29.0 ± 5.5
Race/Ethnicity	
Non-Hispanic White	67,922 (36.8)
Hispanic	51,533 (27.9)
Asian	41,856 (22.7)
Black	14,114 (7.7)
Other	9,079 (4.9)
Pregnancy body mass index (kg/m²)	
< 18.5 (Underweight)	2,396 (1.3)
18.5–24.9 (Normal weight)	81,117 (43.8)
25.0–29.9 (Overweight)	57,899 (31.3)
30.0–34.9 (Obese class I)	26,160 (14.1)
35.0–39.9 (Obese class II)	11,131 (6.0)
≥ 40 (Obese class III)	6,342 (3.4)
Glucose tolerance	
Normal screening	140,781 (78.5)
Abnormal screening and Normal 3-h OGTT	20,780 (11.6)
Gestational diabetes	16,546 (9.2)
Pre-existing diabetes	1,304 (0.7)
Hypertension status	
None	162,673 (89.9)
Pre-existing hypertension	4,190 (2.3)
Gestational hypertension	6,226 (3.4)
Preeclampsia/Eclampsia	7,860 (4.3)
Infant birth weight (grams)	
Mean ± SD	3,395.7 ± 545.9
Size for gestational age	
Small for gestational age	18,154 (9.8)
Appropriate for gestational age	150,311 (81.2)
Large for gestational age	16,575 (9.0)
Fetopelvic disproportion	3,590 (2.0)
Obstructed Labor	9,159 (5.1)
Placenta Previa	1,526 (0.8)
Breech Position	10,950 (6.1)

¹ Not reported, missing, and unknown are as follows: Race: n = 541, Glucose tolerance: n = 5,634, size for gestational age: n = 5 and n = 4096 women were missing all hospital discharge diagnoses codes and were missing information on hypertension, fetopelvic disproportion, obstructed labor, placenta previa, and breech position.

<https://doi.org/10.1371/journal.pone.0199932.t001>

risk factors 1.69 (95% CI, 1.64–1.74), and 3+ risk factors (2.34 (95% CI, 2.19–2.51); *P*- trend = <0.01).

Maternal overweight/obesity synergistically increases the risk of cesarean delivery above and beyond maternal glucose intolerance and pre-existing hypertension. The joint effects of maternal hypertension and maternal overweight/obesity are reported in Fig 1. Compared to

Table 2. Risk Ratios (RRs) and 95% confidence intervals (CIs) for primary cesarean delivery associated with maternal and infant factors.

Maternal/infant factor	RR (95% CI)	
	Model 1 ¹	Model 2 ²
Year of delivery		
2001	1.00 (ref)	1.00 (ref)
2002	1.04 (0.98–1.10)	1.05 (0.99–1.11)
2003	1.02 (0.96–1.08)	1.05 (0.99–1.11)
2004	1.04 (0.99–1.10)	1.10 (1.04–1.16)
2005	1.08 (1.02–1.14)	1.15 (1.09–1.21)
2006	1.08 (1.03–1.14)	1.21 (1.15–1.28)
2007	1.10 (1.04–1.16)	1.29 (1.23–1.36)
2008	1.11 (1.06–1.17)	1.29 (1.23–1.36)
2009	1.08 (1.02–1.13)	1.23 (1.17–1.30)
2010	1.00 (0.95–1.06)	1.16 (1.10–1.22)
Pregnancy body mass index (kg/m²)		
< 18.5 (underweight)	0.79 (0.70–0.89)	0.79 (0.71–0.88)
18.5–24.9 (normal weight)	1.00 (ref)	1.00 (ref)
25.0–29.9 (overweight)	1.26 (1.23–1.30)	1.26 (1.23–1.29)
30.0–34.9 (obese class I)	1.46 (1.42–1.51)	1.43 (1.38–1.47)
35.0–39.9 (obese class II)	1.72 (1.65–1.79)	1.68 (1.62–1.75)
≥ 40 (obese class III)	1.98 (1.89–2.07)	1.90 (1.81–1.99)
Glucose tolerance		
Normal screening	1.00 (ref)	1.00 (ref)
Abnormal screening and normal 3-hour OGTT	1.05 (1.02–1.08)	1.04 (1.01–1.07)
Gestational diabetes	1.16 (1.13–1.20)	1.15 (1.11–1.18)
Pre-existing diabetes	1.53 (1.43–1.63)	1.47 (1.37–1.59)
Hypertension status		
No hypertension	1.00 (ref)	1.00 (ref)
Gestational hypertension	1.18 (1.12–1.23)	1.15 (1.10–1.21)
Pre-existing hypertension	1.28 (1.22–1.35)	1.23 (1.16–1.30)
Preeclampsia / Eclampsia	1.58 (1.53–1.64)	1.53 (1.47–1.58)
Size for gestational age		
Small for gestational age	1.07 (1.04–1.11)	1.09 (1.06–1.13)
Appropriate for gestational age	1.00 (ref)	1.00 (ref)
Large for gestational age	1.68 (1.64–1.73)	1.44 (1.40–1.49)

¹ Adjusted for maternal age at delivery, race/ethnicity, education level, medical facility of delivery, and parity.

² Adjusted for maternal age at delivery, race/ethnicity, education level, medical facility of delivery, parity, obstructed labor status, fetopelvic disproportion status, placenta previa status, and breech position status.

<https://doi.org/10.1371/journal.pone.0199932.t002>

low-risk women with a BMI <25.0 kg/m² and normal blood pressure, women with a BMI < 25.0 with pre-existing hypertension had a 28% increased risk, overweight/obese women with normal blood pressure had a 45% increased risk, and overweight/obese women with pre-existing hypertension had a 2-fold increased risk of having a primary cesarean section. The independent and joint effects of maternal glucose tolerance status and maternal overweight/obesity are reported in Fig 2. For all degrees of glucose intolerance status in pregnancy, the risk of requiring a cesarean delivery was higher among overweight and obese pregnant women. For women with pre-existing diabetes, we found a 2.5 fold increased risk of requiring cesarean section that was independent of maternal BMI (Fig 2).




Fig 1. The joint effects of maternal BMI on maternal pre-existing hypertension in relation to primary cesarean delivery.

<https://doi.org/10.1371/journal.pone.0199932.g001>

The increased risk for cesarean delivery appears to occur even among infants who are appropriate for gestational age (Table 3). The relative risks for having a cesarean section associated with pre-pregnancy BMI among women whose infants were appropriate size for gestational age were higher than for women whose infants were large for gestational age. These rates were not different than the risk ratios for women whose infants were small for gestational age, suggesting cesarean sections associated with women with high pregnancy BMIs are occurring even among women whose infants are appropriate size. Similarly, even among women whose infants were appropriate for gestational age, the risk ratio associated with gestational diabetes was 1.16 (95% CI: 1.12–1.20).

The population attributable risks suggest that the percentages of cesarean deliveries that could be prevented if maternal cardiometabolic risk factors were eliminated were 17.4% for overweight/obesity, 7.0% for hyperglycemia, 2% for hypertension and 20.5% for any cardiometabolic risk factor.

Discussion

We found adverse maternal cardiometabolic risk factors including increasing degree of pregnancy glucose intolerance status, pre-existing hypertension and maternal overweight or




Fig 2. The joint effects of maternal BMI and maternal glucose tolerance status and primary cesarean delivery.

<https://doi.org/10.1371/journal.pone.0199932.g002>

Table 3. Risk Ratios (RRs)¹ and 95% confidence intervals (CIs) for primary cesarean delivery associated with maternal cardiometabolic risk factors, stratified by infant size at delivery.

	SGA	AGA	LGA
Maternal Cardiometabolic Risk Factor			
Pregnancy body mass index (kg/m²)			
< 18.5 (underweight)	0.79 (0.61–1.02)	0.80 (0.70–0.92)	0.84 (0.48–1.47)
18.5–24.9 (normal weight)	1.00	1.00	1.00
25.0–29.9 (overweight)	1.29 (1.20–1.39)	1.27 (1.23–1.30)	1.19 (1.11–1.28)
30.0–34.9 (obese class I)	1.51 (1.37–1.66)	1.47 (1.42–1.53)	1.34 (1.24–1.45)
35.0–39.9 (obese class II)	1.72 (1.52–1.95)	1.75 (1.67–1.84)	1.49 (1.36–1.64)
≥ 40 (obese class III)	2.13 (1.85–2.44)	2.07 (1.96–2.18)	1.61 (1.46–1.78)
Glucose tolerance			
Normal screening	1.00	1.00	1.00
Abnormal screening/normal 3-hour OGTT	1.08 (0.98–1.19)	1.04 (1.00–1.08)	1.09 (1.01–1.17)
Gestational diabetes	1.07 (0.96–1.18)	1.14 (1.10–1.19)	1.33 (1.25–1.42)
Pre-existing diabetes	1.42 (1.10–1.83)	1.53 (1.40–1.67)	1.70 (1.52–1.89)
Hypertension status			
No hypertension	1.00		
Gestational hypertension	1.23 (1.08–1.39)	1.19 (1.12–1.25)	1.09 (0.99–1.21)
Preeclampsia / Eclampsia	1.81 (1.67–1.96)	1.58 (1.51–1.65)	1.29 (1.18–1.41)
Pre-existing hypertension	1.52 (1.34–1.72)	1.28 (1.20–1.35)	1.08 (0.95–1.23)

¹ Adjusted for year of delivery, maternal age at delivery, race/ethnicity, education level, medical facility of delivery and parity.

<https://doi.org/10.1371/journal.pone.0199932.t003>

obesity were all independent risk factors for primary cesarean delivery, after adjusting for other established risk factors for cesarean delivery. Maternal glucose intolerance status and pre-existing hypertension both synergistically increased the risk of primary cesarean section beyond the risk associated with maternal overweight/obesity alone. The estimated population attributable risk for having any cardiometabolic risk factor was 20.5% and the population attributable risk for maternal overweight/obesity it was 17%. The high population-attributable risk for high BMI is driven by the high prevalence of overweight/obesity among pregnant women and highlights the need for strategies to decrease overweight and obesity among reproductive-aged women.

The risk of requiring a cesarean delivery increased with increasing BMI category and was highest among obese class III women. These findings are consistent with prior studies which suggested there is an increased risk of cesarean delivery with increasing BMI category [12–15]. Surprisingly, we found the relative risk of cesarean section among obese class III women was highest among women whose infants were either small or appropriate for gestational age, suggesting this effect is independent of excess fetal growth. The reason underlying the increased risk of cesarean delivery among severely obese women occurring even when the infant is normal or small size may include the practice of “defensive medicine” [27] for fear of malpractice lawsuits, biologic risk factors, or other maternal medical indications for a planned cesarean section. For example, it has been suggested that obese women may have more labor complications due to having a longer active labor phase and decreased cervical dilation rate which may prompt providers to perform a cesarean delivery [28, 29].

We found that maternal pre-existing hypertension was significantly associated with an increased risk of cesarean delivery independent of other known risk factors and indications. However, the population attributable risk associated with pre-existing hypertension was only

2.0% compared with 17.0% for overweight/obesity, suggesting that while pre-existing hypertension is risk factor for requiring a primary cesarean delivery, at the population level they are not accounting for a large a percentage of cesarean deliveries compared to maternal overweight and obesity since the overall prevalence of pre-existing hypertension among pregnant women is still low. Recently, a large study of 123 million singleton deliveries in the United States (1979–2010) found period effects in primary cesarean increasing after 1997 that were explained by a combination of trends in obesity and chronic hypertension, as well as demographic shifts over time[30].

A prior study of 42,071 singleton births in South Carolina found that both pre-existing diabetes and GDM increased the risk of cesarean delivery independently of their effects on birth weight[16]; however, they lacked information on maternal BMI and objectively measured glucose levels during pregnancy. Another study of women undergoing a labor trial found that both prepregnancy obesity and diabetes independently increased the risk for cesarean delivery [17]. We were able to look at increasing degree of objectively measured maternal glucose tolerance status and found a trend of increasing risk of primary cesarean delivery with increasing degree of maternal glucose intolerance. We observed a modest increased risk of primary cesarean section among women with GDM even when infants were appropriate for gestational age. This is similar to a finding in the landmark study by Naylor et al which found that clinicians performed cesarean sections more often among women diagnosed with GDM, even in the absence of excess fetal growth[31]. Further research is required to understand the increased primary cesarean section rates among women with GDM and mild hyperglycemia even when delivering appropriate size for gestational age infants. We also found a significant increased risk of primary cesarean section among women whose infants were born appropriate size for gestational age associated with maternal overweight and obesity and hypertension. This suggests that the associations between maternal cardiometabolic risk factors and primary cesarean section are not due exclusively to the excess or restricted fetal growth associated with the risk factors.

The strengths of this study include the large racially, ethnically, and socioeconomically diverse population of women, the integrated health care delivery system which ensures access to medical care, and the electronic health record which provides high quality data. This enabled us to look at the contribution of maternal cardiometabolic risk factors to risk of primary cesarean deliveries. In addition, in KPNC there is no cost incentive to perform additional cesarean sections. However, the study also had limitations. We were unable to determine whether the cesarean section deliveries were planned or emergency. KPNC may not be generalizable to the extremes of the income distribution which are slightly under-represented in the KPNC membership [32, 33]; the findings may also not be generalizable to other health systems with different practice guidelines or to the uninsured population. We lacked more detailed measures of quality of obstetric care, such as provider characteristics which might have helped us better understand the observed associations. Future studies are needed to determine if the increased performance of cesarean deliveries among women with cardiometabolic risk factors improves health outcomes for the mother and the baby or if the recognition of these conditions causes clinicians to have a lower threshold for performing cesarean deliveries without added benefit for mom and baby.

Maternal adverse cardiometabolic risk factors including overweight/obesity, glucose tolerance status and pre-existing hypertension were all independently associated with risk of cesarean delivery even after controlling for infant size for gestational age and other indications for having a cesarean delivery. Strategies to increase the proportion of women entering pregnancy at an optimal weight and to improve glucose and hypertension before and/or during pregnancy could potentially eliminate up to 20% of cesarean deliveries. Given that maternal

cardiometabolic risk factors are associated with cesarean deliveries even when infants are appropriate for gestational age, it is important to further clarify the patient, clinician and health system-level factors contributing to primary cesarean deliveries, especially among women with adverse cardiometabolic risk profiles.

Author Contributions

Conceptualization: Monique M. Hedderson, Yvonne Crites.

Formal analysis: Monique M. Hedderson, Fei Xu, Charles P. Quesenberry.

Funding acquisition: Monique M. Hedderson.

Investigation: Monique M. Hedderson.

Methodology: Monique M. Hedderson.

Writing – original draft: Monique M. Hedderson, Sneha B. Sridhar, Emily S. Han, Yvonne Crites.

Writing – review & editing: Monique M. Hedderson, Sneha B. Sridhar, Emily S. Han, Yvonne Crites.

References

1. Martin JA, Hamilton BE, Sutton PD, Ventura SJ, Menacker F, Kirmeyer S. Births: final data for 2004. *Natl Vital Stat Rep.* 2006; 55(1):1–101. PMID: [17051727](#)
2. Osterman MJ, Martin JA. Primary cesarean delivery rates, by state: results from the revised birth certificate, 2006–2012. *Natl Vital Stat Rep.* 2014; 63(1):1–11. PMID: [24461076](#).
3. Conroy K, Koenig AF, Yu YH, Courtney A, Lee HJ, Norwitz ER. Infectious morbidity after cesarean delivery: 10 strategies to reduce risk. *Rev Obstet Gynecol.* 2012; 5(2):69–77. PubMed Central PMCID: PMC3410505. PMID: [22866185](#)
4. Liston FA, Allen VM, O'Connell CM, Jangaard KA. Neonatal outcomes with caesarean delivery at term. *ArchDisChild Fetal Neonatal Ed.* 2008; 93(3):F176–F82.
5. Pfuntner A, Levit K, Elixhauser A. Components of Cost Increases for Inpatient Hospital Procedures, 1997–2009: Statistical Brief #133. Healthcare Cost and Utilization Project (HCUP) Statistical Briefs. Rockville (MD)2006.
6. Commission TJ. Specifications manual for Joint Commission national quality measures (v2011A): PC-02NTSV cesarean section2011.
7. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA.* 2012; 307(5):491–7. <https://doi.org/10.1001/jama.2012.39> PMID: [22253363](#).
8. Flegal KM, Carroll MD, Ogden CL, Johnson CL. Prevalence and trends in obesity among US adults, 1999–2000. *JAMA.* 2002; 288(14):1723–7. PMID: [12365955](#).
9. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA.* 2010; 303(3):235–41. <https://doi.org/10.1001/jama.2009.2014> PMID: [20071471](#).
10. Lawrence JM, Contreras R, Chen W, Sacks DA. Trends in the prevalence of preexisting diabetes and gestational diabetes mellitus among a racially/ethnically diverse population of pregnant women, 1999–2005. *Diabetes Care.* 2008; 31(5):899–904. <https://doi.org/10.2337/dc07-2345> PMID: [18223030](#)
11. Ferrara A, Kahn HS, Quesenberry C, Riley C, Hedderson MM. An increase in the incidence of gestational diabetes mellitus: Northern California, 1991–2000. *ObstetGynecol.* 2004; 103(3):526–33.
12. Crane SS, Wojtowycz MA, Dye TD, Aubry RH, Artal R. Association between pre-pregnancy obesity and the risk of cesarean delivery. *ObstetGynecol.* 1997; 89(2):213–6.
13. Vahratian A, Siega-Riz AM, Savitz DA, Zhang J. Maternal pre-pregnancy overweight and obesity and the risk of cesarean delivery in nulliparous women. *Ann Epidemiol.* 2005; 15(7):467–74. <https://doi.org/10.1016/j.annepidem.2005.02.005> PMID: [15921926](#)
14. Dietz PM, Callaghan WM, Morrow B, Cogswell ME. Population-based assessment of the risk of primary cesarean delivery due to excess prepregnancy weight among nulliparous women delivering term

- infants. *Matern Child Health J.* 2005; 9(3):237–44. <https://doi.org/10.1007/s10995-005-0003-9> PMID: 16078011
15. Dempsey JC, Ashiny Z, Qiu CF, Miller RS, Sorensen TK, Williams MA. Maternal pre-pregnancy overweight status and obesity as risk factors for cesarean delivery. *JMaternFetal Neonatal Med.* 2005; 17(3):179–85.
 16. Remsberg KE, McKeown RE, McFarland KF, Irwin LS. Diabetes in pregnancy and cesarean delivery. *Diabetes Care.* 1999; 22(9):1561–7. PMID: 10480526
 17. Clark SL, Xu W, Porter TF, Love D. Institutional influences on the primary cesarean section rate in Utah, 1992 to 1995. *AmJObstetGynecol.* 1998; 179(4):841–5.
 18. Li N, Liu E, Guo J, Pan L, Li B, Wang P, et al. Maternal prepregnancy body mass index and gestational weight gain on pregnancy outcomes. *PLoS One.* 2013; 8(12):e82310. <https://doi.org/10.1371/journal.pone.0082310> PMID: 24376527; PubMed Central PMCID: PMC3869661.
 19. Hesse BW, Nelson DE, Kreps GL, Croyle RT, Arora NK, Rimer BK, et al. Trust and sources of health information: the impact of the Internet and its implications for health care providers: findings from the first Health Information National Trends Survey. *Arch Intern Med.* 2005; 165(22):2618–24. <https://doi.org/10.1001/archinte.165.22.2618> PMID: 16344419
 20. Gordon NP. Similarity of the adult Kaiser Permanente membership in Northern California to the insured and general population in Northern California: Statistics from the 2011 California Health Interview Survey. Oakland, CA: Kaiser Permanente Division of Research, 2015.
 21. Escobar GJ, Fischer A, Kremers R, Usatin MS, Macedo AM, Gardner MN. Rapid retrieval of neonatal outcomes data: the Kaiser Permanente Neonatal Minimum Data Set. *QualManagHealth Care.* 1997; 5(4):19–33.
 22. Selby JV, Ray GT, Zhang D, Colby CJ. Excess costs of medical care for patients with diabetes in a managed care population. *Diabetes Care.* 1997; 20:1396–402. PMID: 9283786
 23. Committee opinion no. 504: screening and diagnosis of gestational diabetes mellitus. *ObstetGynecol.* 2011; 118(3):751–3.
 24. WHO. The International Classification of adult underweight, overweight and obesity according to BMI2012.
 25. Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol.* 2004; 159(7):702–6. PMID: 15033648.
 26. Spiegelman D, Hertzmark E, Wand HC. Point and interval estimates of partial population attributable risks in cohort studies: examples and software. *Cancer Causes Control.* 2007; 18(5):571–9. <https://doi.org/10.1007/s10552-006-0090-y> PMID: 17387622
 27. Dubay L, Kaestner R, Waidmann T. The impact of malpractice fears on cesarean section rates. *J Health Econ.* 1999; 18(4):491–522. PMID: 10539619.
 28. Nuthalapaty FS, Rouse DJ, Owen J. The association of maternal weight with cesarean risk, labor duration, and cervical dilation rate during labor induction. *Obstet Gynecol.* 2004; 103(3):452–6. <https://doi.org/10.1097/01.AOG.0000102706.84063.C7> PMID: 14990405.
 29. Buhimschi CS, Buhimschi IA, Malinow AM, Weiner CP. Intrauterine pressure during the second stage of labor in obese women. *Obstet Gynecol.* 2004; 103(2):225–30. <https://doi.org/10.1097/01.AOG.0000102706.84063.C7> PMID: 14754688.
 30. Ananth CV, Friedman AM, Keyes KM, Lavery JA, Hamilton A, Wright JD. Primary and Repeat Cesarean Deliveries: A Population-based Study in the United States, 1979–2010. *Epidemiology.* 2017; 28(4):567–74. <https://doi.org/10.1097/EDE.0000000000000658> PMID: 28346271; PubMed Central PMCID: PMC5501955.
 31. Naylor CD, Sermer M, Chen E, Sykora K. Cesarean delivery in relation to birth weight and gestational glucose tolerance: pathophysiology or practice style? Toronto Trihospital Gestational Diabetes Investigators [see comments]. *JAMA.* 1996; 275:1165–70. PMID: 8609683
 32. Gordon NP. Similarity of the Adult Kaiser Permanente Membership in Northern California to the Insured and General Population in Northern California: Statistics from the 2009 California Health Interview Survey. Oakland, CA: Kaiser Permanente Division of Research, 2012.
 33. Krieger N. Overcoming the absence of socioeconomic data in medical records: validation and application of a census-based methodology. *AmJPublic Health.* 1992; 82(5):703–10.