

UC San Diego

UC San Diego Previously Published Works

Title

Social Isolation and Serious Mental Illness: The Role of Context-Aware Mobile Interventions.

Permalink

<https://escholarship.org/uc/item/4rw8g3jm>

Journal

IEEE pervasive computing, 23(1)

ISSN

1536-1268

Authors

Nepal, Subigya

Pillai, Arvind

Parrish, Emma

et al.

Publication Date

2024

DOI

10.1109/mprv.2024.3377200

Peer reviewed



Published in final edited form as:

IEEE Pervasive Comput. 2024 ; 23(1): 46–56. doi:10.1109/mprv.2024.3377200.

Social Isolation and Serious Mental Illness: The Role of Context-Aware Mobile Interventions

Subigya Nepal*,

Dartmouth College Hanover, NH, USA

Arvind Pillai*,

Dartmouth College Hanover, NH, USA

Emma M. Parrish,

San Diego State University University of California San Diego San Diego, CA

Jason Holden,

University of California San Diego San Diego, CA, USA

Colin Depp,

University of California San Diego San Diego, CA, USA

Andrew T. Campbell,

Dartmouth College Hanover, NH, USA

Eric Granholm

University of California San Diego San Diego, CA, USA

Abstract

Social isolation is a common problem faced by individuals with serious mental illness (SMI), and current intervention approaches have limited effectiveness. This paper presents a blended intervention approach, called mobile Social Interaction Therapy by Exposure (mSITE), to address social isolation in individuals with serious mental illness. The approach combines brief in-person cognitive-behavioral therapy (CBT) with context-triggered mobile CBT interventions that are personalized using mobile sensing data. Our approach targets social behavior and is the first context-aware intervention for improving social outcomes in serious mental illness.

I. INTRODUCTION

Mental health is a pressing public health concern affecting a significant portion of the population in the United States. Approximately one in five adults in the country experience mental illness each year [1]. Among individuals with serious mental illness (SMI) (e.g., schizophrenia, bipolar disorder), social isolation is a pervasive issue that negatively impacts quality of life, functional ability and mental health outcomes. For example, studies have shown that over 75% of people with psychotic disorders report loneliness [2]. Studies have

sknepal@cs.dartmouth.edu .

*These authors contributed equally to this work.

shown that social isolation can lead to physical health problems, such as, an increased risk of cardiovascular disease and mortality. In fact, social isolation is a severe problem with health comparable to smoking and stroke combined [3]. Unfortunately, the worldwide rate of social isolation is increasing, and those with SMI are at particularly high risk. Studies have found that patients who are more socially isolated have more severe negative symptoms, depression, and worse social functioning [2]. Addressing these facets of social isolation and loneliness is crucial for enhancing treatment outcomes and promoting recovery. Note that while loneliness refers to a subjective feeling of being isolated that can occur even in the presence of others, social isolation is an objective lack of social contacts. Both aspects critically influence mental health and require targeted interventions.

Mobile technology, particularly through mobile sensing, has emerged as a promising tool for mental health interventions. Mobile sensing involves the use of sensors in smartphones and wearable devices to automatically collect data about an individual's behavior and environment, such as physical activity, sleep patterns, and social interactions. This capability allows for the development of personalized interventions targeting specific issues like social isolation. Moreover, mobile-based interventions, including cognitive-behavioral therapy (CBT) programs and mindfulness apps, have been effective in addressing a range of mental health issues, from depression to anxiety [4]–[7]. However, despite these advancements, there's still a significant gap in interventions specifically designed to enhance social functioning and its contribution to overall wellbeing.

Research suggests that in psychotic disorders, social isolation is determined by both reduced approach and avoidance mechanisms [8]. The approach mechanism encourages people to engage in actions that are rewarding or pleasurable, while the avoidance mechanism motivates people to avoid situations or actions that are threatening or negative. Individuals with schizophrenia, for example, may experience decreased motivation to socialize and have diminished reward learning from social interactions, which is worsened by negative symptoms. At the same time, anxiety and social threat perception may result in social avoidance. Interventions aimed at increasing motivation for socialization or reducing social anxiety have had modest success in treating negative symptoms of schizophrenia. However, most of such interventions are intensive and are not scalable. Previous studies have shown that both diminished reward and anxious avoidance play a role in determining social interactions in individuals with schizophrenia. These studies have also found that context plays an important role in social behavior, with different thoughts and emotions being present when interacting with others (negative defeatist appraisals) versus being alone at home (anxious threat about leaving) [9]–[12]. The premise of this study, therefore, is that, a context-aware CBT intervention that targets specific thoughts and emotions related to social behavior in different contexts may be more effective (than digital CBT interventions and in-person CBT interventions that are not context-aware) in improving social motivation and functioning in individuals with SMI. Our work seeks to evaluate the feasibility, acceptability and preliminary impact of such intervention.

We propose a novel blended intervention approach that combines brief in-person CBT with context-triggered mobile CBT interventions to address social isolation in SMI. The intervention is personalized, participatory, and employs a novel and flexible architecture

that facilitates intervention based on sensor-detected context. It is also the first context-aware intervention for improving social outcomes in SMI. Our blended intervention targets social behavior by adjusting perceptions (like not belonging) and encouraging social engagement, addressing both social isolation and loneliness within diverse contexts. The intervention employs brief in-person CBT to set socialization goals, introduce the cognitive model and thought challenging skills to address defeatist attitudes and social threat beliefs, and plan pleasurable social activities for the mobile component of the intervention. The mobile component consists of context-triggered CBT interventions that aim to increase social behavior through reminders, reinforcement, activity pleasure savoring, and cognitive restructuring. The mobile component is personalized based on the data collected through mobile sensing technology (specifically data on the patient's location and social interactions) as well as with the help of personalized messages entered by the therapists. Furthermore, the architecture of our intervention is flexible, enabling dynamic modification of intervention strategies based on the patient's context, paving the way for future "Just-in-time" and adaptive interventions. Note that this manuscript represents initial results and insights from an ongoing study. Full trial results, as well as a greater focus on the therapy aspect of the intervention, will be reported in a later paper. The aims of this manuscript are as follows:

1. We describe the details of our study and the design of mobile-based tools that captures novel sensing modalities (e.g., conversation-mobility sensor using gps and microphone).
2. We present a representative case study to illustrate the effectiveness of context-aware mobile interventions for improving social functioning
3. We present the implications of our work and elucidate extensions that broadly benefit the mobile sensing and mental health community.

II. RELATED WORK

Mobile-based mental health interventions have exhibited potential for providing real-time support to individuals. For instance, [13] observed that longitudinal text message-based interventions improve social isolation in individuals with schizophrenia or schizoaffective disorder. There has been some research on enhancing social functioning in SMI through mobile technology, however, as noted by [4], it is limited. Despite this, [5] reveals that a few mobile interventions, even without passive sensing, have demonstrated improvements in social functioning. Studies have also employed CBT algorithms in mobile apps to address mental health symptoms and enhance social interactions and self-efficacy beliefs [6], [7], but none of the existing context-aware systems have specifically focused on social behavior in severe mental illness (SMI). Blended interventions, which merge mobile and in-person therapies while reducing in-person contact by 50–90%, have been more effective than app-only approaches [6], [14], [15]. One recent pilot trial of a blended intervention for psychosis used coping-focused sessions for voices blended with an app and found improved coping and trend-level reduction in voices [16]. However, it focused on symptoms, and there have been limited blended interventions for social isolation tailored to SMI. We identified only one brief 3-week intervention providing feedback of ecological momentary

assessment (EMA) responses and behavior change suggestions, which showed improvement in symptoms but not social functioning [17]. Our study presents a novel contribution to the field by introducing a personalized, blended context-aware mobile intervention aimed at enhancing social functioning in individuals with SMI. This innovative approach specifically targets social behavior, expanding on previous work and addressing a gap in context-aware systems for SMI populations.

III. METHODOLOGY

A. Participant Selection

Our study aims to recruit a diverse sample of 50 participants aged 18–65 with SMI, including schizophrenia, schizoaffective, or bipolar I disorder with psychosis. In this manuscript, we report on the preliminary data on feasibility and app performance in N=5 participants with schizophrenia. 75% of the participants we have enrolled are white (N=3), 50% are male (N=2) and their mean age and mean years of education are 54.8 years (Range=47–59) and 12 respectively. Inclusion criteria of our study include a minimum level of social avoidance, defined as a score of ≥ 2 on the Scale for Assessment of Negative Symptoms asociality item, a reading level of at least 6th grade on the Wide Range Achievement Test-4 Reading subtest, and stability in medications (no hospitalizations or medication changes in the 4 months prior to enrollment). Exclusion criteria include prior receipt of CBT in the past 2 years, greater than moderate disorganization on the Positive and Negative Symptom Scale, and alcohol or substance dependence in the past 3 months based on DSM-5 criteria. In addition, participants requiring a higher level of care (e.g., hospitalization, severe medical illness) are excluded, as well as those who are unable to adequately see or manually manipulate a mobile phone, since ours is a blended intervention that uses a mobile device.

B. Study Design

The study involves a brief blended intervention that combines in-person and mobile CBT to reduce social isolation. Social isolation can be caused by a lack of reward from relationships and avoiding social situations due to anxiety. When someone is alone, they might start to have negative thoughts about social situations and then avoid them, which only reinforces their negative beliefs. Modifying negative beliefs and behaviors when alone can help someone feel more comfortable and engaged in social situations, which can lead to positive changes in their thoughts, emotions, and social behavior. Our hypothesis is that the use of our mobile-assisted CBT intervention, called mobile Social Interaction Therapy by Exposure (mSITE), will lead to an increase in social interactions among individuals with SMI. This increase in social interactions will in turn lead to a reduction in the severity of experiential negative symptoms and ultimately improve social functioning. Note that the study is approved by the Institutional Review Board of University of California, San Diego and Dartmouth College. It is also registered as a clinical trial in clinicaltrials.gov.

We are conducting a 24-week open pilot trial of mSITE. Our trial focuses on feasibility, acceptability, and initial indication of improvement in our target, i.e., social activity. The intervention involves eight weekly 1-hour in-person sessions combined with mobile app

use, followed by weekly 15-minute remote coaching calls with mobile app use for an additional 16 weeks, for a total of 24 treatment weeks. We have included weekly remote coaching, as the literature suggests that mobile interventions are more effective when coaching is included. Our assessments will be carried out at baseline, 8 weeks (end of in-person treatment), 12 weeks, 18 weeks, and 24 weeks, with one-week bursts of ecological momentary assessments (EMA) surveys at each of these weeks. Our prior research has demonstrated that improvements in outcomes can be achieved in 24 weeks or fewer. However, if we can attain our target engagement at the earliest possible assessment point, which could be at week 12 or 18, it would allow us to create a shorter yet efficient intervention. Effectively strengthening and shortening intensive psychosocial interventions could help overcome implementation barriers that arise due to the high cost and burden of treatment. In addition, it could improve access to evidence-based practices for individuals who are underserved by the current mental healthcare system.

C. In-person Intervention Component

The intervention we are using has eight sessions and is based on different evidence-based treatments for mental health issues, including CBT. During the sessions, masters-level therapists, comparable to therapists in a community mental health system, work with patients to develop skills to help them achieve their goals for recovery. We start by setting a recovery goal, then we teach patients how to use a thought challenging skill called the 3Cs (Catch-It, Check-It, Change-It; where *It* is a thought) to address defeatist beliefs and social avoidance and work towards their goals. 3Cs is a CBT technique used to help individuals identify and challenge dysfunctional thoughts and replace them with more helpful, accurate ones. We teach patients how to challenge defeatist attitudes, social threats, and avoidance behaviors using CBT skills. We also educate patients about how their thoughts, actions, and feelings are related, and use experiments and exposure to help them change dysfunctional beliefs. The intervention includes tests of expectations in social interactions to help patients achieve their goals. Essentially, this means that the intervention uses real-life social interactions as a way to help patients challenge and modify their beliefs and behaviors.

D. Mobile Intervention Component

The mobile intervention component uses a smartphone application (referred to as the mSITE app) to passively collect sensing data, e.g., activity, GPS, audio, light, phone usage, application usage, etc. The GPS sensors and microphones are used to detect if the user is home or away and around conversation, respectively. Moreover, the app alerts the users three times daily with personalized CBT scripts using information from the sensors and therapists. Note that the mSITE app does not record any raw conversation (only whether conversation was present or not and the amplitude of sound). While we collect other passive sensing data, only the GPS and conversations related information are used to drive interventions. Other passive sensing data are used only for analysis purposes.

The mSITE app is installed on the participants' phone or an Android phone provided during in-person and coaching treatment phases. Our application consists of five elements:

1) Goals: mSITE provides on-demand resources that are populated during individual sessions. Participants set a long-term goal and enter short-term goals and steps to achieve the long-term goal (see Fig. 2a). They can add their own steps on demand in and out of session.

2) Pleasurable Activities: Participants create a list of enjoyable social activities and rate their anticipated pleasure in doing them (see Fig. 2b). After doing them, they again rate their pleasure and are prompted to do something to savor pleasure (e.g., write about it or use the phone to take a photo or video), which are used as reminders that activities can be more enjoyable than they think (see Fig. 2c).

3) Conversation-Location Sensing: As mentioned earlier, our app identifies whether participants are home/away and around conversations to fire contextual EMAs, an example is shown in Fig. 3. To identify the participant's home, we first run DBScan clustering to group significant locations of a participant and then label the location where they spend most of their time from 2 am - 4 am as their home location. We use a geofence tolerance of ± 100 meters to mark whether they are home or away. In addition, the machine learning based conversation detector on the mSITE app identifies whether a participant has been around social interactions. Our application identifies speech and conversation by utilizing pre-trained classifiers that analyze accelerometer and microphone-derived audio data, respectively. To safeguard users' privacy, the application does not retain raw audio data; instead, it stores audio information when speech or conversation is detected. This detection is based on a two-stage classifier: initially, a voice activity detector employs a Hidden Markov Model and the Viterbi algorithm to ascertain the presence of speech, followed by a conversation classifier that determines whether a conversation has commenced and records its duration. We also apply a filtering process that evaluates voice presence alongside audio amplitude, where samples with lower amplitudes are more likely considered noise and thus removed. Although we cannot be certain if the subject is actively participating in the conversation, the inferred presence of conversation indicates that the subject is in proximity to it. We employ this information as an indicator of social engagement or isolation. The detector analyzes ambient audio every 10 minutes for a duration of one minute to detect the presence of nearby conversations. We tested and refined it to ensure it can distinguish conversations from other sounds such as television and background noise. Our team has previously deployed this conversation detector in numerous studies [18], [19].

The context-aware scripts using the conversation-location sensor are triggered in afternoon and evening. For example, when the participant is home and alone, the app challenges their negative beliefs about social situations (i.e., anxiety and social threat appraisal) and reminds them of positive experiences they have had in the past. Similarly, when the individual has recently had a social interaction, mSITE helps them savor the positive aspects of that interaction and challenges any negative thoughts they have about future interactions (i.e., defeatist appraisal). This is based on findings from our prior research [9]–[12]; we found that, when home alone, individuals with schizophrenia report threat beliefs and anxiety about going out which contribute to social avoidance, so we challenge threat beliefs. When out around others, they report defeatist beliefs about being judged negatively, so we challenge those beliefs in that context. The app automatically detects whether they are home/

away or have been around conversations to fire the personalized interventions. However, we first prompt the participants to confirm that we have indeed correctly detected their social context (e.g., “I think you have been home and around other people in the afternoon; am I right?”). If they answer with “no”, the app shows “Sorry that we got it wrong! Were you around others this morning/afternoon?”. The challenges that follow are then based on their input to this response rather than on the automatically detected context (i.e., home/away and alone/with others).

4) Therapist-in-the-loop & personalization: Here, the therapist can enter unlimited personalized messages that challenge the individuals’ beliefs using our online therapist dashboard. An example prompt challenging a belief is, “But it was fun when you played cards with Joe at the clubhouse”. For each participant, the therapists craft personalized evidence post-session, tailored to the session’s content and the individual’s specific therapeutic needs. This customization introduces variability in message content and frequency, reflecting the unique therapeutic journey of each participant. Such variability, aligned with traditional CBT practices, is intended to enhance engagement by ensuring relevance to the participant’s personal goals and challenges. We view this individualized approach not as a limitation but as a strength, fostering a more personalized and engaging therapeutic experience. In addition, the database is composed of several generic messages—a list of pre-composed messages that are not unique to a participant’s situation (e.g., “The best way to find out if someone will like you is to talk to them and see.”). The mSITE app randomly selects personalized and generic messages 60% and 40% of the times, respectively. The purpose of generic messages are to accommodate new participants who join the study and may have only one or two personalized message. In such cases, using the 60–40 threshold avoid repeated personalized messages. The message is displayed when the context-aware EMAs are fired as shown in Fig. 3e.

5) Gamifying social interactions: mSITE provides users with an “Awards” screen that provides information about their intervention progress on demand (see Fig. 2d). After successful completion of goal steps and activities, an “awards wheel” randomly gives users “diamonds”. Thus, mildly gamifies the social interaction progress through non-monetary “achievements”.

IV. APPLICATION ACCEPTABILITY

Our study participants have shown sustained adherence to our mobile app, which is a promising indication of the app’s robustness. On average, participants answered 77% of the EMAs presented to them on their 8 weeks of in-person use. It is worth noting that participants have the option to respond to up to three EMAs daily, including an action plan EMA in the morning and a context-aware EMA in the afternoon and evening. Of the EMAs that participants have answered so far, 46% are context-aware EMAs triggered by our app’s detection of whether the participant is home or away and around conversations. The remaining 54% of the EMAs are daily action plan EMAs that encourage participants to set goals for their day, such as engaging in a fun activity outside their home, interacting with someone or achieving a custom goal that they can set for themselves.

Our application demonstrated an initial contextual sensing accuracy of 75%. To confirm the contextual Ecological Momentary Assessment (EMA) prompt, we have a validation mechanism that solicits participant verification before providing a response. For example, if the application detects that the participant is at home and engaged in conversation, the contextual EMA's initial prompt states, "I believe you have been at home and in the presence of others this morning; is this correct?" The accuracy of this detection was initially confirmed by participants 75% of the time. There is a delay between data collection from the application and its integration into the system. For instance, our first contextual EMA triggers at 12 PM, using data collected between 6 AM and 12 PM to determine whether participants have been at home, away, or engaged in conversations. We observed that by adjusting the frequency of data uploads to our server and altering server-side processing intervals, we could enhance accuracy. Given that many of our participants reside in care homes, it was essential to adapt our voice activity detection model to function effectively in noisy environments. To achieve this, we calibrated the model using a Samsung Galaxy S10 across various settings, including areas with heavy traffic, quiet rooms, and spaces where televisions or music were playing. This process ensured that our system could reliably perform in diverse acoustic conditions. The calibration involved fine-tuning the model's parameters to accurately differentiate between human speech and other types of background noise. In addition, we fine-tuned the geo-fence settings, adjusting the DBSCAN algorithm's epsilon (which determines the radius of neighborhood) and minimum samples (which affects the density required to form a cluster) parameters and the data upload frequency to improve location and interaction accuracy. The improved calibration of our application has achieved an accuracy exceeding 85% in internal tests, specifically in detecting participants' presence at home and their involvement in conversations, validated by their self-reports.

In addition to the high adherence rates to EMAs, our mobile sensing data collection has also been successful. Each participant has at least 18 hours of location and audio data available, including information on the presence and amplitude of conversations on 85% of their study days. Note that we do not record any raw audio. Overall, the acceptability and feasibility of our mobile app-based intervention have been promising, as evidenced by the high adherence rates to both EMAs and mobile sensing data collection. We are continually evaluating the app's effectiveness in improving mental health outcomes and hope to provide additional insights as the study progresses.

V. PARTICIPANT CASE STUDY

We have enrolled five participants so far for a duration of a little over eight weeks. While it is premature to have any compelling indication about behavior change in this short period, here we highlight two case studies which provides a representative example of the effectiveness of context-aware mobile interventions for improving social functioning.

Participant A, a 47-year-old White male with 12 years of schooling, aimed to form a close friendship, despite being content with his acquaintances and social life. He found the app useful, using it daily to work towards his goal or related tasks. Over 8 weeks, he responded to all daily action plan EMAs, focusing mainly on initiating interactions (54 out of 56 times). He also completed 47 morning and 40 evening contextual EMAs, revealing he was in

social situations 64% of the time when prompted and spent most days away from home. The app's personalized messages from the therapist, aimed at overcoming negative self-beliefs and encouraging social engagement, made up a significant part of his experience. With 46% of messages addressing defeatist thoughts and 44% promoting social interaction, these communications supported his active efforts to meet his goal. A appreciated tracking his progress within the app, undertaking various activities to meet new people, such as visiting potential friend-meeting locations, introducing himself to new individuals, and practicing conversation skills. He set 10 long-term goals and listed 25 activities ranging from doctor visits to attending social events. While he hasn't finished the remote coaching sessions yet, A reports increased happiness in his social endeavors and enhanced satisfaction with his existing relationships.

Participant B, a 59-year-old Asian male with a 10-year education background, initially aimed to make three new friends but shifted his focus to volunteering at a local animal shelter. He felt comfortable conversing but wanted to be more open and involved in his community. Variable app engagement was noted, as he often left his phone at home to avoid losing it, leading to 89% completion of daily action plan EMAs and only 21% of contextual EMAs in the first eight weeks. Coaching encouraged him to carry his phone, boosting his engagement and EMA response rates. Despite struggling with some app features and technology, targeted messages from the therapist helped him confront social fears and negative self-perceptions. B's active session involvement and steps toward his goals, like initiating conversations, visiting the animal shelter, and submitting a volunteer application, demonstrated his progress. Regular outings with a friend from his board and care increased his social interactions. By the end of the remote coaching sessions, B had made significant strides towards his goal of volunteering at the animal shelter.

Fig. 4 shows weekly trends of Participant A's conversations and home time via the mSITE app, with conversations increasing post-week four and less time at home. Table I indicates improved interaction enjoyment for participants A and B, with A showing more interest in socializing. Both participants spent less time at home. Social defeatist attitudes rose for A but fell for B. While eight weeks is brief for conclusive outcomes, early signs suggest the intervention is fostering more social engagement and less isolation. We anticipate further positive changes in social activity for both participants as the trial progresses.

VI. DISCUSSION

A. Preliminary Insights and Implications

Our preliminary findings demonstrate the feasibility and acceptability of our blended intervention approach that combines in-person CBT with context-triggered mobile CBT interventions to address social isolation in individuals with SMI. Participants have shown sustained adherence to the mobile app, and the app's robustness is evident through its successful data collection and contextual sensing capabilities. As the study progresses, we will continue to evaluate the app's effectiveness in improving social functioning and mental health outcomes in our target population. The context-aware aspect of our intervention is particularly noteworthy, as it sets our approach apart from existing interventions in the field. Most context-aware interventions in the literature have focused on physical health,

with limited exploration of their potential in addressing social isolation or mental health outcomes [20]. However, our focus on social activity and functioning is relatively novel in the mental health domain. In fact, it is the very first study to use socially mediated contextual sensing intervention.

The high adherence rates and the case study provided highlight the importance of personalization and therapist involvement in designing mobile interventions. In our study, the mSITE app allows therapists to input personalized messages that challenge participants' beliefs and tailored interventions. In addition, our mobile intervention system enables users to set their long-term goals and short-term steps, providing them with the tools to work towards achieving those goals. This approach empowers users to take an active role in their recovery and fosters a sense of agency and control, which is crucial for individuals with SMI. Another important implication of our study is the potential for context-aware mobile interventions to be applied in other areas of mental health research. For example, our approach could be used to study social anxiety or agoraphobia, where individuals may be hesitant to leave their homes or interact with others in social situations. In addition, the approach could be used to study healthy aging, where social isolation and loneliness can have negative effects on physical and mental health.

B. Privacy Measures

Privacy is a key focus in our study and we have implemented rigorous safeguards to protect it. Participants give written informed consent, fully briefed on what data will be collected and who will have access. They also undergo a decisional capacity assessment to ensure they grasp the study's procedures. We monitor for any adverse effects, such as participant dropouts or concerns about data collection, and have observed none to date. This absence of concerns may stem from the trust built during individual coaching sessions where the rationale behind data-driven prompts is explained. Furthermore, participants are motivated to enhance their social interactions and understand that the prompts are constructive feedback for their goals, rather than unexpected surveillance. This understanding, coupled with the personal support from coaches, likely contributes to the high acceptability of our intervention.

C. Limitations and Future Work

Our study on social isolation in individuals with SMI has limitations: it's ongoing, requiring more data for conclusions; conducted over a short period; lacks a control group; and focuses on a specific SMI population. In addition, the personalized intervention using mobile sensing technology raises privacy concerns, although no conversation content is recorded. Future research should address these concerns and ensure that the interventions are implemented in a way that respects patient privacy and security.

Overall, our study represents an important first step in addressing social isolation in individuals with SMI. However, future research is needed to further investigate the effectiveness of context-aware mobile interventions and to address the limitations of our study.

VII. CONCLUSION

Our study describes a novel blended intervention approach that combines brief in-person CBT with context-triggered mobile CBT interventions to address social isolation in SMI. Our mobile intervention system provides personalized interventions at the right time in the right context to improve social functioning, addressing a critical need in the mental health field. The intervention was acceptable and feasible in this small sample of individuals with SMI. Future research can expand on our work to develop “Just-in-time” adaptive interventions, paving the way for further advancements in the field of mobile sensing and mental health.

ACKNOWLEDGMENTS

This work is supported by National Institute of Mental Health (NIMH), grant number 5R61MH126094.

REFERENCES

- [1]. SAMHSA, “Key substance use and mental health indicators in the united states: Results from the 2020 national survey on drug use and health (hhs publication no. pep21–07-01–003, nsduh series h-56),” 2021.
- [2]. Michalska da Rocha B, Rhodes S, Vasilopoulou E, and Hutton P, “Loneliness in psychosis: a meta-analytical review,” *Schizophrenia bulletin*, vol. 44, no. 1, pp. 114–125, 2018. [PubMed: 28369646]
- [3]. Valtorta NK, Kanaan M, Gilbody S, Ronzi S, and Hanratty B, “Loneliness and social isolation as risk factors for coronary heart disease and stroke: systematic review and meta-analysis of longitudinal observational studies,” *Heart*, 2016.
- [4]. Cornet VP and Holden RJ, “Systematic review of smartphone-based passive sensing for health and wellbeing,” *Journal of Biomedical Informatics*, vol. 77, p. 120–132, Jan. 2018. [PubMed: 29248628]
- [5]. Bell IH, Lim MH, Rossell SL, and Thomas N, “Ecological momentary assessment and intervention in the treatment of psychotic disorders: A systematic review,” *Psychiatric Services*, vol. 68, no. 11, p. 1172–1181, Nov. 2017. [PubMed: 28669284]
- [6]. Ben-Zeev D, Brenner CJ, Begale M, Duffecy J, Mohr DC, and Mueser KT, “Feasibility, acceptability, and preliminary efficacy of a smartphone intervention for schizophrenia,” *Schizophrenia Bulletin*, vol. 40, no. 6, pp. 1244–1253, Mar. 2014. [PubMed: 24609454]
- [7]. Bucci S, Barrowclough C, Ainsworth J, Machin M, Morris R, Berry K, Emsley R, Lewis S, Edge D, Buchan I, and Haddock G, “Actissist: Proof-of-concept trial of a theory-driven digital intervention for psychosis,” *Schizophrenia Bulletin*, vol. 44, no. 5, pp. 1070–1080, Mar. 2018. [PubMed: 29566206]
- [8]. Fulford D, Campellone T, and Gard DE, “Social motivation in schizophrenia: How research on basic reward processes informs and limits our understanding,” *Clinical Psychology Review*, vol. 63, p. 12–24, Jul. 2018. [PubMed: 29870953]
- [9]. Depp CA, Bashem J, Moore RC, Holden JL, Mikhael T, Swendsen J, Harvey PD, and Granholm EL, “GPS mobility as a digital biomarker of negative symptoms in schizophrenia: a case control study,” *npj Digital Medicine*, vol. 2, no. 1, Nov. 2019.
- [10]. Parrish EM, Depp CA, Moore RC, Harvey PD, Mikhael T, Holden J, Swendsen J, and Granholm E, “Emotional determinants of life-space through GPS and ecological momentary assessment in schizophrenia: What gets people out of the house?” *Schizophrenia Research*, vol. 224, pp. 67–73, Oct. 2020. [PubMed: 33289659]
- [11]. Depp CA, Moore RC, Perivoliotis D, Holden JL, Swendsen J, and Granholm EL, “Social behavior, interaction appraisals, and suicidal ideation in schizophrenia: The dangers of being alone,” *Schizophrenia Research*, vol. 172, no. 1–3, pp. 195–200, Apr. 2016. [PubMed: 26948502]

- [12]. Granholm E, Ben-Zeev D, Fulford D, and Swendsen J, "Ecological momentary assessment of social functioning in schizophrenia: Impact of performance appraisals and affect on social interactions," *Schizophrenia Research*, vol. 145, no. 1–3, pp. 120–124, Apr. 2013. [PubMed: 23402693]
- [13]. Granholm E, Ben-Zeev D, Link PC, Bradshaw KR, and Holden JL, "Mobile assessment and treatment for schizophrenia (mats): a pilot trial of an interactive text-messaging intervention for medication adherence, socialization, and auditory hallucinations," *Schizophrenia bulletin*, vol. 38, no. 3, pp. 414–425, 2012. [PubMed: 22080492]
- [14]. Alvarez-Jimenez M, Alcazar-Corcoles M, González-Blanch C, Bendall S, McGorry P, and Gleeson J, "Online, social media and mobile technologies for psychosis treatment: A systematic review on novel user-led interventions," *Schizophrenia Research*, vol. 156, no. 1, pp. 96–106, Jun. 2014. [PubMed: 24746468]
- [15]. Ben-Zeev D, Kaiser SM, Brenner CJ, Begale M, Duffecy J, and Mohr DC, "Development and usability testing of FOCUS: A smartphone system for self-management of schizophrenia." *Psychiatric Rehabilitation Journal*, vol. 36, no. 4, pp. 289–296, Dec. 2013. [PubMed: 24015913]
- [16]. Bell IH, Rossell SL, Farhall J, Hayward M, Lim MH, Fielding-Smith SF, and Thomas N, "Pilot randomised controlled trial of a brief coping-focused intervention for hearing voices blended with smartphone-based ecological momentary assessment and intervention (SAVVy): Feasibility, acceptability and preliminary clinical outcomes," *Schizophrenia Research*, vol. 216, pp. 479–487, Feb. 2020. [PubMed: 31812327]
- [17]. Hanssen E, Balvert S, Oorschot M, Borkelmans K, van Os J, Delespaul P, and Fett A-K, "An ecological momentary intervention incorporating personalised feedback to improve symptoms and social functioning in schizophrenia spectrum disorders," *Psychiatry Research*, vol. 284, p. 112695, Feb. 2020.
- [18]. Wang R, Chen F, Chen Z, Li T, Harari G, Tignor S, Zhou X, Ben-Zeev D, and Campbell AT, "Studentlife: Assessing mental health, academic performance and behavioral trends of college students using smartphones," in *Proceedings of the 2014 ACM International Joint Conference on Pervasive and Ubiquitous Computing*, ser. UbiComp '14. New York, NY, USA: Association for Computing Machinery, 2014, p. 3–14.
- [19]. Wang W, Mirjafari S, Harari G, Ben-Zeev D, Brian R, Choudhury T, Hauser M, Kane J, Masaba K, Nepal S, Sano A, Scherer E, Tseng V, Wang R, Wen H, Wu J, and Campbell A, "Social sensing: Assessing social functioning of patients living with schizophrenia using mobile phone sensing," in *Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems*, ser. CHI '20. New York, NY, USA: Association for Computing Machinery, 2020, p. 1–15.
- [20]. Craig KJT, Morgan LC, Chen C-H, Michie S, Fusco N, Snowdon JL, Scheufele E, Gagliardi T, and Sill S, "Systematic review of context-aware digital behavior change interventions to improve health," *Translational Behavioral Medicine*, vol. 11, no. 5, pp. 1037–1048, Oct. 2020.




Fig. 1: mSITE's blended CBT approach using context-triggered mobile interventions and therapist-in-the-loop personalization.




Fig. 2: Important mSITE screens: (a) User entered goal steps in a hierarchical view, i.e., Top-level long term goals, contains short-term goals, which contains goal steps, (b) User enters a pleasurable social activity for the day, (c) User describes how their pleasurable activity went using freeform text, sliders, and can add photos, (d) Gamification using awards the user has accumulated through the study.




Fig. 3: An example of a context-aware EMA from mSITE. (a) The conversation-location sensor detects that the user is away and alone in the morning. (b)-(d) EMA questions appraising the user's current defeatist or threat beliefs, (e) Challenging the user's defeatist belief through personalization and their conversations with the therapist, (f) A prompt helping the user savor positive actions.




Fig. 4: Objectively sensed number of conversations (left) and time spent at home (right) of Participant A. We can see that as the study progressed, their number of conversations increased while time spent at home decreased. The x-axis is the week number and the y-axis is the sensed data.

8-week study summary for participants A and B evaluating pleasure and interest during interactions, motivation for future interactions, social defeatist attitudes, and time spent at home. The values presented are mean of surveys.

TABLE I:

Participant	Time Point	How much pleasure or enjoyment did you feel in the interactions?	In the past hour, how much interest or motivation did you have for interacting with others? [Had 0 interactions]	How much interest or motivation do you have for engaging in interactions later today? [Had 1+ interactions]	Social Defeatist Attitudes	In the past hour, about how much time did you spend at home?
A	Week 0	3.84	2.67	3.16	3.88	45.98
	Week 8	4.08	2.80	3.68	3.96	42.00
B	Week 0	5.39	2.32	3.61	2.48	54.69
	Week 8	5.45	1.52	3.55	2.36	53.27