

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

HIGH-ENERGY TOTAL CROSS SECTIONS FOR POSITIVE PIONS AND PROTONS ON HYDROGEN

### Permalink

<https://escholarship.org/uc/item/4sc6d60f>

### Authors

Longo, Michael J.  
Helland, Jerome A.  
Hess, Wilmot N.  
et al.

### Publication Date

1959-11-10

0 0 1 0 0 3 0 5 3 3 4

UCRL 8955

UNIVERSITY OF  
CALIFORNIA

*Ernest O. Lawrence*

*Radiation  
Laboratory*

**For Reference**  
Not to be taken from this room  
(Bldg. 50)

BERKELEY, CALIFORNIA

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

0 0 1 0 0 3 0 5 3 3 5

For pub. Phys. Rev. Letters

UCRL-8955

UNIVERSITY OF CALIFORNIA  
Lawrence Radiation Laboratory  
Berkeley, California

Contract No. W-7405-eng-48

HIGH-ENERGY TOTAL CROSS SECTIONS  
FOR POSITIVE PIONS AND PROTONS ON HYDROGEN

Michael J. Longe, Jerome A. Helland, Wilmet N. Hess,  
Burton J. Moyer, and Victor Perez-Mendez

November 10, 1959

**HIGH-ENERGY TOTAL CROSS SECTIONS  
FOR POSITIVE PIONS AND PROTONS ON HYDROGEN\***

Michael J. Longo, Jerome A. Helland, Wilmot N. Hess,  
Burton J. Moyer, and Victor Perez-Mendez

Lawrence Radiation Laboratory  
University of California  
Berkeley, California

November 10, 1959

On the basis of dispersion relations, Pomeranchuk has shown that if the  $(\pi^+, p)$  and  $(\pi^-, p)$  total cross sections approach constant values at high energies, these limits must be equal.<sup>1</sup> Considerable information is already available on the  $(\pi^-, p)$  total cross sections in the Bev range.<sup>2, 3, 4</sup> We present the results of an experiment at the Berkeley Bevatron in which  $(\pi^+, p)$  cross sections between 1.4 and 4.0 Bev/c were measured. Data on  $(p, p)$  total cross sections are also presented.

The experimental arrangement is shown in Fig. 1. Positive particles produced when the circulating proton beam strikes a target are bent inwards towards the center of the Bevatron by the magnetic field. In order to obtain positive pions with the highest possible momentum, the pion beam was brought out of the machine through a window on the inside radius. The beam is further deflected by bending magnets to keep it clear of the Bevatron structure and is focused by an 8-in. -bore doublet quadrupole.

Scintillation counters  $M_1$ ,  $M_2$ , and  $M_3$ , each 1-1/2 in. diam, define the beam. Pions are distinguished from protons by requiring a coincidence with C, a gas Cerenkov counter<sup>5</sup> filled with sulfur hexafluoride to give an index of refraction of approximately 1.008. By putting C in anticoincidence with  $M_1 M_2 M_3$ , we obtained data on p-p scattering simultaneously with the  $(\pi^+, p)$  data.

\* Work done under auspices of U. S. Atomic Energy Commission.

After passing through the defining telescope, the beam impinged on a 4-ft-long liquid hydrogen target. The number of pions (and protons) transmitted was measured simultaneously at three different solid angles by means of counters  $S_1$ ,  $S_2$ , and  $S_3$ . An extra coincidence in  $S_0$  was added to reduce accidentals. The solid angles defined by  $S_1$ ,  $S_2$ , and  $S_3$  ranged from 0.6 to 4.3 milliradians.

The coincidence circuits used were of the type described by Wenzel.<sup>6</sup> With the clipping lines employed the resolving time was about  $5 \times 10^{-9}$  sec. The output of the monitor coincidence circuit was used as an input to a second circuit where a coincidence with  $S_0$  and  $S_1$  (for example) was required. Several species of accidentals were monitored throughout the experiment, and corrections to the data were made where necessary. Accidentals involving C were always less than 2% of the number of pions.

The momentum spread accepted by the counter telescope was approximately 2.5%, and the uncertainty in the momentum determination is estimated to be  $\pm 3\%$ . Extensive magnetic shielding to reduce the Bevatron's leakage field was required along most of the beam line.

Data collection at each momentum involved series of runs with the  $H_2$  target filled and empty; each series consisted of a full run preceded and followed by an empty run. The number of counts recorded in each series was usually sufficient to give a cross section with a statistical uncertainty of less than 1%. An analysis of the data from different measurements at the same momentum indicated that the probable error assigned to each measurement had to be increased to about 3%. The results given below are averages of two or more measurements at each momentum. The uncertainty in the extrapolation to the zero solid angle subtended by the last counter was considered in assigning the errors. Where necessary the data at the smaller solid angles were corrected for multiple coulomb scattering by a method similar to that of Sternheimer.<sup>7</sup> These

corrections were assigned a probable error of  $\pm 20\%$  and had little effect on the extrapolated cross sections. Other possible sources of error are discussed below.

A. Sources of error in the  $(\pi, p)$  measurements

A curve of the ratio of  $M_1 M_2 C M_3$  counts to  $M_1 M_2 M_3$  counts as a function of gas pressure in the Čerenkov counter indicated less than 2% electron contamination and less than 1% muon contamination at 1.8 Bev/c. Such a curve gives only the fraction of muons with momentum equal to or greater than that of the pions. The total muon contamination for the geometries and momenta used was determined by calculation, and corrections were made. The maximum was 2%. Practically all of the muons came from pions decaying after  $M_2$ , where the calculation is relatively simple.

B. Sources of error in the  $(p, p)$  measurements

The errors assigned to the  $(p, p)$  cross sections have been increased to allow for the uncertainty in the efficiency of the anticoincidence circuits and the Čerenkov counter. If these were not 100% efficient in eliminating pion counts, the measured  $(p, p)$  cross sections would be low. The assigned errors are based on an efficiency between 90% and 100%.

K-meson contamination in the "proton beam" is estimated to be less than 1% for the momenta and take-off angles used in this experiment.

Our cross sections are shown in Table I and Fig. 2 which also shows data of other experimenters. Our data for  $\sigma(\pi^+, p)$  indicate a maximum near 1.5 Bev/c and an essentially constant cross section above 2.4 Bev/c. Our value at 4.0 Bev/c is  $29.3 \pm 0.4$  mb. The best high-energy values for  $\sigma(\pi^-, p)$  to date are  $28.7 \pm 2.6$  mb at 4.3 Bev/c<sup>3</sup> and  $29.1 \pm 2.9$  mb at 5.2 Bev/c.<sup>4</sup>

We wish to acknowledge our indebtedness to Ken Lou for handling the many engineering problems encountered in the experiment, to Thomas Devlin for his help in conducting the experiment, and to the Bevatron staff and crew.



Table I

Total ( $\pi^+$ , p) and (p, p) cross sections

<u>Momentum</u> <u>(Bev/c)</u>	<u><math>\sigma(\pi^+, p)</math></u> <u>(mb)</u>	<u><math>\sigma(p, p)</math></u> <u>(mb)</u>
1.40	$39.4 \pm .6$	$46.9^{+1.0}_{-0.6}$
1.46	$39.1 \pm .8$	$47.5^{+2.2}_{-1.1}$
1.60	$35.8 \pm .9$	$47.7^{+3.0}_{-1.1}$
1.73	$30.1 \pm .5$	$46.5^{+2.0}_{-0.6}$
1.89	$28.4 \pm .6$	$46.3^{+3.2}_{-0.8}$
2.05	$27.8 \pm .6$	$45.0^{+3.0}_{-1.0}$
2.47	$29.0 \pm .6$	$45.6^{+1.9}_{-0.7}$
2.97	$29.2 \pm .5$	$45.1^{+0.9}_{-0.5}$
3.58	$29.2 \pm .4$	$43.3^{+0.6}_{-0.5}$
4.00	$29.3 \pm .4$	$42.4 \pm 0.6$


## References

1. Ia. Pomeranchuk, Soviet Physics JETP 34(7), 499 (1958).
2. R. Cool, O. Piccioni, and D. Clark, Phys. Rev. 103, 1082 (1956).
3. Fredrick Wikner, Nuclear Cross Sections for 4.2-Bev Negative Pions (thesis), UCRL-3639, January 1957.
4. Richard G. Thomas, The Elastic Scattering of 5-Bev  $\pi^-$  Mesons of Hydrogen, UCRL-8965, November 3, 1959.
5. J. H. Atkinson and V. Perez-Mendez, Rev. Sci. Instr. 30, 864 (1959).
6. William A. Wenzel, Millimicrosecond Coincidence Circuit for High Speed Counting, UCRL-8000, October 1957.
7. R. M. Sternheimer, Rev. Sci. Instr. 25, 1070 (1954).
8. Devlin, Perez-Mendez, Hess, Barish, and Solomon, Lawrence Radiation Laboratory, University of California, private communication.
9. Chen, Leavitt, and Shapiro, Phys. Rev. 103, 211 (1956).

Figure legends

Fig. 1. Experimental arrangement.

Fig. 2. Total ( $\pi^+$ , p) and (p, p) cross sections vs. momentum.



1.8.7

