

# UCSF

## UC San Francisco Previously Published Works

### Title

Barriers to a cure for HIV: new ways to target and eradicate HIV-1 reservoirs

### Permalink

<https://escholarship.org/uc/item/4tb179pp>

### Journal

The Lancet, 381(9883)

### ISSN

0140-6736

### Authors

Katlama, Christine  
Deeks, Steven G  
Autran, Brigitte  
[et al.](#)

### Publication Date

2013-06-01

### DOI

10.1016/s0140-6736(13)60104-x

Peer reviewed

Published in final edited form as:

*Lancet*. 2013 June 15; 381(9883): . doi:10.1016/S0140-6736(13)60104-X.

## Barriers to a Cure: New concepts in targeting and eradicating HIV-1 reservoirs

Christine Katlama<sup>1</sup>, Steven G. Deeks<sup>2</sup>, Brigitte Autran<sup>3</sup>, Javier Martinez-Picado<sup>4</sup>, Jan van Lunzen<sup>5</sup>, Christine Rouzioux<sup>6</sup>, Michael Miller<sup>7</sup>, Stefano Vella<sup>8</sup>, Joern E. Schmitz<sup>9</sup>, Jeffrey Ahlers<sup>11</sup>, Douglas D. Richman<sup>10</sup>, and Rafick P. Sekaly<sup>11</sup>

<sup>1</sup>Department of Infectious Diseases, Pierre et Marie Curie University, Pitié-Salpêtrière Hospital, Paris, France

<sup>2</sup>Department of Medicine, University of California, San Francisco, CA, United States

<sup>3</sup>Laboratory Immunity and Infection, UMR-S 945, Pierre et Marie Curie University, Hospital Pitié-Salpêtrière, Paris, France

<sup>4</sup>AIDS Research Institute Irsi Caixa, ICREA and Universitat Autònoma de Barcelona, Badalona, Barcelona, Spain

<sup>5</sup>University Medical Center Eppendorf, Infectious Diseases Unit, Hamburg, Germany

<sup>6</sup>Paris-Descartes University Necker Hospital, Department of Virology, Paris, France

<sup>7</sup>Department of West Point Discovery Chemistry, Merck Research Labs, West Point, USA

<sup>8</sup>Department of Pharmacology and Therapeutic Research, Istituto Superiore di Sanità, Rome, Italy

<sup>9</sup>Center for Virology and Vaccine Research, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA

<sup>10</sup>VA San Diego Healthcare System and Departments of Pathology and Medicine, Center for AIDS Research, University of California, San Diego, California, USA

<sup>11</sup>Vaccine and Gene Therapy Institute of Florida, Port Saint Lucie, Florida, USA

### Abstract

Antiretroviral therapy for HIV infection requires life-long access and strict adherence to regimens that are both expensive and associated with toxicities. There is growing recognition that a curative intervention will be needed to fully stop the epidemic. The failure to eradicate HIV infection during long-term antiretroviral therapy reflects the intrinsic stability of the viral genome in latently infected CD4<sup>+</sup> T cells and other cells and perhaps ongoing low-level viral replication.

© 2013 Elsevier Ltd. All rights reserved.

Corresponding authors: Christine Katlama, Department of Infectious Diseases, Hospital Pitié-Salpêtrière, 83 Bd de l'Hôpital, Paris, 75013, France, christine.katlama@psl.aphp.frss. Rafick Pierre Sekaly, Vaccine and Gene Therapy Institute of Florida, Port Saint Lucie, Florida, USA, rpssekaly@vgtifl.org.

Authors Contributions: All authors contributed to the ideas and interpretation of data in this manuscript. CK, SGD, BA, JMP, DDR, JA and RPS were primarily responsible for the writing and editing of the manuscript and inclusion of data provided in the Tables. Dr. Miller is an employee of Merck Research Laboratories which holds U.S. Patent Nos. RE 38,506 E, 6,087,367 ZOLINZA ® (vorinostat).

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Heterogeneity in latently infected cell populations and homeostatic proliferation of infected cells may influence the dynamics of virus production and persistence. Chronic immune activation, inflammation and immune dysfunction persist despite potent antiretroviral therapy and likely have important effects on the size and distribution of the viral reservoir. The inability of the immune system to recognize cells harboring latent virus and to eliminate cells actively producing virus represents the biggest challenge to finding a cure. In this perspective, we highlight new approaches toward unraveling the complex virus-host interactions that lead to persistent infection and latency and discuss the rationale for combining novel therapeutic strategies with current antiretroviral treatment options with the goal of curing HIV disease.

## Keywords

Histone Deacetylases; Epigenetics; Gene Silencing; HIV-1 /immunology

---

## Finding a cure for HIV: The magnitude of the problem

Antiretroviral therapy (ART) can be considered one of the major medical successes of the late 20<sup>th</sup> century. When used effectively, it results in indefinite viral suppression, restoration of immune function, an improved quality of life, the near normalization of expected life span and a reduction in viral transmission. Despite the inherent potency of ART to suppress virus replication, current therapeutic approaches have limitations. Antiretroviral therapy does not eliminate viral reservoirs, requiring life-long adherence to regimens that are expensive and which have potential short-term and long-term toxicities. There were over 34 million people estimated to be living with HIV in 2010, which represents an increase of 17% over the last ten years, and the numbers are expected to continue to increase. It is unclear if there will be sufficient resources on a global level to provide treatment and monitoring to all in need. Even for those with access to ART, significant challenges remain in terms of individual adherence. In addition, despite virus control, HIV-associated complications persist, including higher than normal risk for developing cardiovascular disease, cancer, osteoporosis and other end-organ diseases. This risk might be due to treatment toxicities and/or the consequences of persistent HIV-associated inflammation and immune dysfunction. Thus, there is a very strong need for novel therapeutic approaches that could eliminate persistent virus and would not require life-long adherence to expensive and potentially toxic antiretroviral drugs. For all these reasons, there is now widespread interest in potentially curative interventions.

## Potential outcomes of the cure approach

There are two broadly defined categories of a cure for HIV infection. A *functional cure* can be defined as host-mediated control of HIV replication, in the absence of ART, during which there is (i) suppression of viral replication for a pre-defined period (*e.g.*, five years) in absence of therapy, (ii) restoration and stabilization of effective immune function, (iii) limited HIV induced inflammation that could lead to an increased risk for AIDS or non-AIDS morbidity, and (iv) limited risk of transmitting virus to others with stable low level plasma viral load. A *sterilizing cure* for HIV infection requires the complete elimination of replication-competent virus.

There are now examples for both types of cure. A functional cure is achieved spontaneously by a rare group of HIV-infected individuals who naturally control HIV replication in the absence of therapy (“elite controllers”) These patients are characterized by a favourable HLA profile and potent HIV-specific CD8<sup>+</sup> T cell responses that are associated with a low viral DNA reservoir. A second group of patients was recently identified; they initiated ART

during acute infection and were found to control HIV for several years after interruption of ART<sup>1</sup>. These “post-treatment controllers” are exceedingly rare; in contrast to elite controllers, they do not exhibit strong HIV-specific CD8<sup>+</sup> T cell responses or possess protective HLA alleles<sup>2, 3</sup>.

A sterilizing cure was likely achieved following myeloablative chemotherapy, whole body irradiation, and subsequent successful transplantation of hematopoietic stems cells from a CCR5  $\Delta 32$  homozygous donor into an HIV infected individual who had developed acute myelogenous leukemia (the “Berlin Patient”)<sup>4, 5</sup>. In a recent report from Boston, two antiretroviral-treated subjects with relapsed Hodgkin’s lymphoma that received a CCR5<sup>+/+</sup> hematopoietic stem cell transplant, proviral DNA and replication competent HIV were undetectable 8–17 months after transplantation. These observations suggest that ablative conditioning, immunosuppressive therapy, and/or post-transplant graft-versus-host disease (GVHD) — all of which were common to the Berlin Patient and the Boston cases— may cause dramatic and perhaps curative reductions in the size of the reservoir<sup>6</sup>. Whether the Boston-based individuals were truly cured will require interruption of antiretroviral therapy.

Efforts to pursue both *functional* and *sterilizing* cures are on-going. It is possible that an effective cure will likely require combinatorial approaches. For example, attempts at eradicating the reservoir may not work unless the capacity of the immune system to clear and control the virus are enhanced.

## HIV Reservoirs: Obstacles to a Cure

### Establishment and maintenance of HIV latency

The HIV reservoir is established during primary infection. Administration of antiretroviral therapy in very early acute infection appears to result in a lower post-treatment total and integrated DNA and HIV-RNA levels, suggesting aggressive treatment can limit the size of the viral reservoir<sup>1, 7–9</sup>. Although early treatment can substantially reduce the total reservoir size, a stable population of latently infected CD4<sup>+</sup> cells exists that transits in to the long-lived latent reservoir and is relatively unaffected by early cART<sup>10</sup>.

The vast majority of HIV proviral DNA is detected in CD4<sup>+</sup> T lymphocytes in lymphoid tissue<sup>11, 12</sup>. In blood, most HIV DNA can be found in central memory (TCM) and in transitional memory T cells (TTM); these cells maintain the reservoir because of their intrinsic capacity to persist through homeostatic proliferation and renewal<sup>13</sup>. Other cellular reservoirs may exist, including naïve CD4<sup>+</sup> T cells, monocytes/macrophages, astrocytes, and microglial cells<sup>14</sup>. During long-term effective antiretroviral therapy, a steady state low-level plasma HIV RNA, typically from less than one to three copies RNA/ml, is eventually achieved<sup>15</sup>. The source for this persistent HIV is not fully known. Chronic production of HIV from a stable reservoir of long-lived infected cells (the “latent reservoir”) is likely the dominant source, but some have argued that persistent low-level replication is also involved particularly in tissues where it has been suggested that ongoing persistent viral replication despite ART is due to cell to cell spread and lack of drug penetrance in tissues<sup>16, 17</sup>.

A prerequisite for the establishment of HIV latency is the integration of viral DNA into the host chromatin and epigenetic silencing of active viral transcription. Two models have been proposed to explain latent infection in memory cells. The “pre-activation latency” model relies on the supposition that HIV can directly infect a subset of resting CD4<sup>+</sup> T cells<sup>18, 19</sup> while the post-activation latency model proposes that activated antigen-specific CD4<sup>+</sup> memory T cells become preferentially infected yet avoid cell death and then revert to a resting state<sup>20</sup>. The frequency of resting CD4<sup>+</sup> T cells that become latently infected varies

widely between individuals and has been estimated to be as few as one per million CD4<sup>+</sup> T cells.

The molecular mechanisms contributing to HIV-DNA silencing are complex<sup>21</sup>. Infected cells containing replication-competent provirus are transcriptionally silenced by corepressor complexes containing histone deacetylases (HDAC), histone methyltransferases, and heterochromatin proteins; active methylation of the LTR may also be involved<sup>22, 23</sup>. Epigenetic silencing of provirus can be reversed by agents that mobilize chromatin remodeling complexes<sup>24</sup> to replace repressive complexes poised at the viral LTR<sup>24</sup>. Signals delivered through the T cell receptor (TCR/CD3 complex) and CD28 costimulation can drive productive transcription suggesting that physiologic activation of memory CD4<sup>+</sup> T cells can lead to virus production *in vivo*<sup>25</sup>.

Activated CD4<sup>+</sup> T cells are the most permissive target for HIV infection. How recently infected activated cells become long-lived latently infected resting memory cells is not fully understood. Multiple regulatory pathways are turned on that are designed to blunt the impact of cell activation. These pathways include upregulation of the “negative regulators of T cell activation” such as the cell-surface receptors PD-1, CTLA-4, TIM-3, LAG3, CD160, and 2B4. Theoretically, cells expressing these receptors may be preferentially latently infected with HIV. In a cross-sectional study of long-term treated individuals, PD-1 expressing cells were enriched for HIV DNA<sup>13</sup>. Agents aimed at interrupting the molecular pathways associated with these negative regulators might help clear the reservoir.

## Novel strategies towards an HIV cure

### ART treatment intensification

If persistent low-level replication of HIV (or *de novo* infection of new target cells) persists during therapy, then addition of potent agents (“intensification”) to a stable regimen may help reduce reservoir size. To date, treatment intensification studies with new classes of drugs, such as the HIV integrase inhibitor raltegravir or the CCR5 inhibitor maraviroc, have had no effect in reducing residual viremia<sup>26</sup>. A few reports have shown that intensification with an integrase inhibitor leads to transient accumulation of pre-integration episomal DNA (2-LTR circles) and simultaneous reductions in CD8 T cell activation markers, suggesting inhibition of low-levels of viral replication<sup>26,27</sup>. Another study showed that intensification with raltegravir reduced HIV RNA levels in ileum, which is rich in infected target cells, and may be an ideal environment to support cell-to-cell transfer of virus during ART; however, there was no evidence of a reduction in HIV-1 DNA or residual viremia<sup>28</sup>. Antiretroviral therapy may not be able to fully penetrate all tissues. Persistence of HIV replication in tissues justifies efforts to optimize drug delivery to enhance penetration into cells in tissues where current drugs fail to reach, including the central nervous system and possibly lymphoid tissues<sup>29</sup>. Measurements of virus activity in lymphoid tissue and other anatomical compartments such as the gut, reproductive organs, and the brain are needed to better understand the current limitations of ART intensification therapy to eradicate residual virus production.

### The promise of gene therapy approaches

The “Berlin Patient” case has generated widespread interest in cell-based curative interventions. As performing allogeneic transplants using stem cells from those very rare donors who are naturally resistant to HIV is not a feasible strategy, much of the interest is now focused on using gene therapy to delete virus from infected cells or to produce cells resistant to HIV infection<sup>30</sup>. Three classes of DNA-editing enzymes are currently being evaluated for safety and efficacy to target HIV coreceptors and proviral sequences: zinc finger nucleases (ZFNs), transcription activator-like (TAL) effector nucleases (TALENs),

and homing endonucleases (HEs)<sup>31, 32</sup>. The targeted gene therapy approach to block CCR5 (CCR5–ZFN knockout T cells) is currently being evaluated in phase I clinical trials of adoptive transfer of ex vivo expanded autologous T cells (Table 1)<sup>33</sup>. Phase I studies have shown the safety and feasibility of such approaches.

Other gene-based therapies that target HIV viral proteins using either an anti-HIV ribozyme or antisense RNA oligonucleotide constructs have demonstrated safety and feasibility in reducing viral load in phase I/IIa clinical trials<sup>34</sup>, while CD4 aptamer-CCR5-siRNA chimera have proven safe and efficacious in the humanized mouse model<sup>35</sup>. Interventions that interfere with pre-integration steps in the viral life cycle are promising and currently being evaluated in phase I trials<sup>36</sup>. The safety and, efficacy of gene delivery to specific cells and tissues, and ultimately access to these treatments are major hurdles for these approaches in eradicating HIV infection<sup>37</sup>.

### A Three Tiered Approach to Reverse Latency and Eliminate the Reservoir

Most investigators believe that a safe and scalable cure will require a three-tiered strategy: 1) reactivation of HIV proviral genomes to purge HIV from all latently infected cells, 2) suppression of residual (cryptic) viral replication, and 3) enhancement of host immune function to recognize virus-infected cells and eliminate rebound virus replication. Potential approaches to meet this goal are summarized below.

**Step 1. Reactivation of latent infectious proviral genomes**—The goal of HIV-1 reactivation therapy is to purge cellular reservoirs by activating the transcriptional activity of latent HIV-1 provirus in infected CD4<sup>+</sup> T cells resulting in the generation of virus-producing cells that either die from the direct cytopathic effects of the virus or are cleared by host mechanisms. Although complex molecular mechanisms regulate expression of provirus in latently infected cells, one possible strategy is to trigger NF- $\kappa$ B activity, a major host transcription factor for HIV-1 replication<sup>38, 39</sup>. Initial attempts to reactivate virus in patients on ART using two known inducers of NF- $\kappa$ B, IL-2 or anti-CD3 MAb (OKT3), were unsuccessful in reactivating virus and were associated with dose-dependent toxicity<sup>40–42</sup>. Other candidate NF- $\kappa$ B activating agents effective at inducing virus transcription in *in vitro* latency models include prostratin, TNF- $\alpha$ , and the transient NF- $\kappa$ B activating agent HIV-reactivating factor (HRF)<sup>43</sup>. Pharmacological agents that stimulate autophagy, protein kinase C, or the generation of reactive oxygen species (ROS) can also activate the NF- $\kappa$ B pathway to induce virus replication. Bryostatin 1 induce the PKC–NF- $\kappa$ B pathway and can synergize with other drugs to remodel chromatin transcriptional coactivator complexes in the cell, reactivate virus, and promote cellular apoptosis<sup>44, 45</sup>. Enthusiasm for these approaches is tempered by the recognition that generalized activation of CD4<sup>+</sup> T cells has significant risk associated with systemic induction of proinflammatory cytokines.

Using small molecule drugs to more specifically reverse those pathways involved in chromatin silencing was advocated at about the time the concept of latency was first introduced<sup>46</sup>. There is now intense interest in inhibiting histone deacetylases (HDAC) as these molecules block HIV DNA transcription by preventing transcription factors from accessing the HIV promoter. Two candidate HDAC inhibitors (HDACi) have already entered clinical trial evaluation (Table I). Valproic acid, a relatively weak non-toxic HDACi, was the first drug studied. Despite initial promising results in a pilot study, subsequent studies failed to find any benefit and the field has moved on to more potent drugs<sup>47–49</sup>. Vorinostat (SAHA) has shown promise in reactivating virus in *in vitro* models of latency. In a recent pilot study, 11 of 16 antiretroviral-treated subjects exhibited potential susceptibility to vorinostat *ex vivo*<sup>50</sup>. Eight of these subjects were then administered a single dose of vorinostat and showed a significant increase in their expression of RNA levels compared to

baseline. Other clinical trials of vorinostat are ongoing (Table 1). A recent report showed that the HDAC inhibitor FK288 (romidepsin) was 1000 times more potent than vorinostat at inducing latent HIV and induced HIV RNA expression *ex vivo* in twelve of thirteen HIV infected subjects on cART<sup>51</sup>. Other HDACi, such as LBH589 (panobinostat), ITF2357 (givinostat), PXD101 (belinostat) also showed greater potency than vorinostat in reactivating virus in latently infected cell lines (Table 2)<sup>52</sup>.

Coordinated action of multiple epigenetic modifications might be required to initiate viral transcription. Optimum reactivation of epigenetically silenced provirus DNA might require multi-drug combinations that include HDACi, PKC activators, and/or specific HMTase inhibitors. The combination of a HDACi with the H3K9 histone methyltransferase (HMT) inhibitor, BIX01294, reactivated latent virus better than either agent alone<sup>53, 54</sup>. The discovery of new molecules that are safe and can act alone or synergize with other drugs to reactivate virus transcription with high efficiency is a major priority. For example, three FDA-approved drugs (dactinomycin, aclacinomycin, and cytarabine) were recently described to act as priming agents when combined with other reactivating molecules to increase the frequency of cells producing virus<sup>55</sup>. Disulfiram, used for the treatment of alcoholism, was shown through its active metabolite diethyldithiocarbamate (DDTC) to reactivate latent virus in an *in vitro* latency model<sup>56</sup>. In a small pilot study, a single dose of disulfiram given daily for 2 weeks to cART treated subjects was well tolerated and led to a transient increase in plasma RNA in a subset of subjects<sup>57</sup> (Table 1).

There are other potential drug targets that maintain latency including the above described immune checkpoint blockers such as PD-1, which is expressed on latently infected CD4<sup>+</sup> cells in HIV infection. These molecules present a major barrier to T-cell activation by inhibiting signals transmitted through the TCR<sup>58-60</sup>. In ART treated patients, PD-1 expression remains elevated and correlates with the size of the latent reservoir and high levels of cell-associated RNA and proviral DNA<sup>13, 61, 62</sup>. High levels of PD-1 expression on CD4 memory T cells in HIV-1 infection prevents antigen-specific TCR activation and downstream signals that are needed for proliferation and effector function and HIV replication. PD-1 or PDL-1 blocking antibodies can relieve inhibitory signals on TCR activation and increase the sensitivity of the TCR to antigen. High levels of PD-1 expression found on antigen-specific CD8 T<sub>TM</sub> and T<sub>EM</sub> cells in HIV infection even in patients successfully treated with ART could prevent recognition and killing of CD4 cells in which virus has been reactivated. Blocking PD-1 on antigen-specific CD8 T cells can prevent negative signals delivered through PDL-1 and PDL-2 ligand binding. Thus, blocking PD-1 or combinations of co-inhibitory molecules might serve a dual role in reactivating latent virus in CD4 cells, as well as relieving a functional block on virus-specific CD8<sup>+</sup> memory T cells by enhancing antigen-specific proliferation and effector cell differentiation<sup>63, 64</sup>.

The cytokine IL-7 is required for homeostatic proliferation and survival of naïve and memory CD4<sup>+</sup> T cells that express the IL-7R<sup>65</sup>. In ART treated patients with low CD4<sup>+</sup> T cells, IL-7 was shown to reconstitute naïve and TCM cells<sup>66, 67</sup> and in one study induced transient viremia in 6 of 26 subjects<sup>68</sup>. Exogenous IL-7 could reactivate HIV replication in CD4 memory T cells from some HIV-infected donors *in vitro* when cells were stimulated in PBMC cocultures<sup>69</sup>. IL-7 induces the proliferation of CD4 memory T cells that carry integrated viral DNA including TCM and TTM, yet the ability to reactivate virus replication with IL-7 in *ex vivo* cultures is highly variable<sup>70</sup>. Based on these findings, IL-7 therapy as a reactivating intervention is currently tested in combination with ART intensification with raltegravir plus maraviroc to prevent proliferation or reseeding of the latent reservoir (ERAMUNE 01 study) (Table I).

**Step 2. Targeting residual virus replication**—Chronic inflammation is a major factor contributing to HIV immune activation and disease pathogenesis<sup>71–73</sup>. The inflammatory process appears to be most destructive in lymphoid tissues, leading to collagen deposition, fibrosis and irreversible damage to the lymphoid infrastructure as a result of early recruitment of Tregs and production of TGF- $\beta$ <sup>74, 75</sup>. Abundant amounts of HIV proteins (e.g. p24, p17, gp120) persist on the surface of the follicular dendritic cell network (FDC) despite suppressive ART and may contribute to chronic immune activation<sup>41, 76–78</sup>. Proinflammatory cytokines and IFN- $\gamma$  production results in a higher frequency of activated CD4<sup>+</sup> T cells that might contribute to persistent low level viremia and prevent healing damaged gut and lymph nodes<sup>79, 80</sup>. Chronic inflammation results in the persistent upregulation of co-inhibitory receptors and impaired immune function on antigen-specific T cells, including those that are needed to clear HIV reservoirs. Drugs aimed at preventing the negative aspects of inflammation on functional immune responses (e.g., blocking PD-1-PDL interactions; blocking Type I Interferons and IL-6) and enabling the positive aspects (e.g., enhancing innate and HIV-specific T-cell immunity) may prove to be an important combination for an effective cure strategy.

A number of anti-inflammatory drugs are now being advanced into hypothesis-testing phase I and phase II clinical trials to establish a role for attenuating chronic inflammatory responses in balance with immune responses needed for eradication of latent reservoirs. For example, a recent study found that the expression of CD38 and PD-1 on total CD8<sup>+</sup> and Gag-specific CD8<sup>+</sup> T cells was reduced in untreated HIV infected patients administered a COX-2 inhibitor<sup>81, 82</sup>. The Peroxisome Proliferator-Activated Receptor (PPAR- $\gamma$ ) agonists represent a promising class of anti-inflammatory molecules. Among these, pioglitazone and leflunomide are relatively well tolerated and effective for treatment of chronic inflammation<sup>83</sup> and might reduce comorbidities (metabolic syndrome, dyslipidemias, and glucose intolerance) that have been associated with long-term ART in HIV infected patients<sup>84</sup>. Chloroquine or hydroxy chloroquine treatment can significantly reduce HLA-DR<sup>+</sup>CD38<sup>+</sup> CD8<sup>+</sup> T cells, as well as Ki-67 expression in CD4<sup>+</sup> and CD8<sup>+</sup> T cells<sup>85, 86</sup>, but failed to show efficacy in meeting these endpoints in HIV asymptomatic ART-naïve patients<sup>87</sup>. Other drugs currently being evaluated include methotrexate, mesalamine, which should reduce inflammation in the gut mucosa, where much of the virus resides, and anti-fibrotic agents such as ACE inhibitors (which may restore immune function) and a variety of drugs aimed at other microbial infections which may cause chronic immune activation (including drugs aimed at reducing translocation of gut microbes across the damaged mucosal surfaces of the gut). Whether these anti-inflammatory agents will help decrease virus production and ultimately decrease viral reservoirs remains to be demonstrated.

**Step 3: Enhancing host-mediated clearance of residual virus**—The importance of cellular immunity in controlling the size of the HIV reservoir has been suggested by several studies. HIV-1 Gag-specific CD8 T cells isolated from elite controllers but not subjects treated with ART were shown to effectively kill autologous resting CD4 T cells in which virus was reactivated with SAHA<sup>88</sup>. Moreover, functional anti-viral CD8<sup>+</sup> T cells are associated with limiting the size of the central memory CD4<sup>+</sup> T cell reservoir in patients controlling their virus in the absence of ART<sup>89</sup>. Indeed, high-avidity multifunctional CD8<sup>+</sup> CTL that target vulnerable regions in Gag are particularly important in limiting virus diversity and reservoirs in HIV-infected individuals possessing “protective” HLA-class I alleles<sup>89–91</sup>. Therapeutic vaccines have been proposed to restimulate CD8 CTL in order to prevent or control virus relapses and the reestablishment of latent infection in CD4 T cells after treatment interruptions<sup>92–94</sup>. A few therapeutic vaccine trials such as the Ad5-HIV-1 gag vaccine (ACTG A5197), or infusions of dendritic cells pulsed with inactivated HIV particles have shown transient viral suppression following treatment interruption<sup>95, 96</sup>. The Eramune-02 study is currently testing whether the VRC HIV-Gag, Pol, Nef, and Env



vaccine in a DNA prime-recombinant replication-defective adenovirus serotype-5 (rAd5) boost regimen can reduce the viral reservoir in subjects undergoing an ARV intensification regimen (Table 1).

The Type I IFN response is an essential innate mediator of the acute antiviral response and for the induction of effector CD8<sup>+</sup> CTL. A recent study found that pegylated interferon (IFN)- $\alpha$ 2A treatment could suppress virus replication in subjects whose immune function was partially restored by cART<sup>97</sup>. Adjuvants inducing the production of type I IFNs by conventional dendritic cells, plasmacytoid dendritic cells, and nonhematopoietic cells can potentially drive activation and differentiation of quiescent memory CD4<sup>+</sup> T cells and enhance *de novo* CD8<sup>+</sup> CTL responses and viral clearance. Push–Pull strategies combining synergistic combinations of molecular adjuvants such as Toll-like receptor (TLR) ligands and costimulatory molecules, while blocking of negative regulatory molecules (PD-1, CTLA-4, TIM-3, CD160, 2B4), immunosuppressive cytokines, and T regulatory function might further optimize induction of CD8<sup>+</sup>T effector and effector memory cells<sup>96, 98–100</sup>.

High dose IL-15 can skew differentiation of high levels of CD8<sup>+</sup> T effector memory cells<sup>101</sup>. IL-15 was also shown to be 4 fold more potent than IL-7 for inducing virus production in latently infected CD4 cells<sup>70</sup>. Therapeutic vaccines adjuvanted with IL-15 in combination with blockage of co-inhibitory molecules such as PD1 might enhance killing of infected cells and prevent reactivated virus from replenishing latent reservoirs.

## The clinical challenges ahead: a “new look” at the design of clinical studies

To achieve a cure for HIV, the efforts of the global research community must be coordinated and transparent in sharing both positive as well as negative clinical results. Specifically we must:

1. Identify virus producing cells and the location of viral sanctuaries and determine the mechanisms by which virus establishes and maintains latent infection.
2. Develop standardized *in vitro* latency models in primary memory CD4 cells to facilitate the screening and prioritization of reactivating compounds and combinations.
3. Harmonize on-going efforts of different laboratories to develop a high through put primary T-cell model of HIV latency to screen for novel anti-latency compounds.
4. Develop and implement robust monitoring tools to quantitate single copy cell-associated HIV RNA and HIV DNA as a measure of the HIV reservoir and the proportion of cells producing virus in response to treatment. Since the majority of proviral DNA is replication-incompetent, assays that can directly measure replication competent HIV DNA and quantitate HIV DNA/RNA double positive cells are needed.
5. Develop non-human primate (NHP) models of antiretroviral-treated SIV infection and conduct proof of concept studies testing the safety and efficacy of reactivation strategies and immune-based therapies
6. Develop humanized mouse models of HIV infection to assess novel eradication approaches and combinatorial strategies.
7. Develop non-comparative phase I/II clinical studies to rapidly screen multiple clinical strategies as in the oncology field to determine safety and efficacy and a regulatory framework that allows fast implementation of small scale (*e.g.* 10 to 20 subjects per arm) proof-of-concept hypothesis driven trials.

8. Develop guidelines for selection of patient populations according to the risk/benefit ratio associated with the primary objective of the study, *i.e.* sterilizing or functional cure.
9. Develop consensus protocols for defining time to primary endpoints and the follow-up period for secondary endpoints that quantify (i) reduction in the HIV reservoir in blood and tissues, (ii) levels of inflammatory biomarkers, and (iii) functional antigen-specific immune responses for advancing to phase II studies. A phase IIb/III approach will ultimately require analytical treatment interruption of ART in responders and evaluation of HIV reservoirs in blood and tissues after withdrawing therapy
10. Develop a universal web-based database for meta-analyses and hierarchical modelling of multi-center trials and protocols that can facilitate the identification of best approaches.
11. Involve the HIV community, funding agencies, and regulatory authorities prospectively to discuss the complex ethical considerations with special regard to the design and financial support of clinical studies in HIV cure research.
12. Prepare the public opinion for this new phase of research and discovery, preventing high expectations in a short-term and readdressing the emphasis on prevention as the main available strategy to fight the HIV/AIDS epidemic.

## Conclusions

Given the burden of HIV infection throughout the world, the goal of finding a cure for HIV is one of the major medical challenges of our time. This goal has rallied all sectors of the biomedical community (public, private, academic and community groups). A consultation has led to elaboration of a document that describes a strategy that will lead to a cure<sup>14</sup>. There are still numerous scientific challenges ahead to identify combinatorial approaches that can effectively induce reactivation of the latent reservoir(s) and enhance specific immune responses to control virus replication and eliminate virus-infected cells. Ethical issues are also of great concern in strategies that reactivate virus infection in individuals that are being successfully treated on ART. Recent studies that immediate/early antiretroviral treatment following infection might minimize the size and complexity of the latent reservoir and could lead to post-treatment control provide us with the optimism for moving forward with therapeutic interventions aimed at eradicating HIV infection. We should all keep in mind that prevention and early initiation of ART is currently the best way to control the HIV pandemic.

## Acknowledgments

NIH funded Human clinical trial identifiers were obtained from a search of the [www.clinicaltrials.gov](http://www.clinicaltrials.gov) website.

Support for CK was provided through ORVACS by Fondation Bettencourt Schuller. JMP was supported by the European Union's Seventh Framework Programme (CHAIN project; grant agreement 223131), the Spanish Ministry of Science and Innovation (grant SAF2010-21224), and the Catalan HIV Vaccine Development Program (HIVACAT). DDR was supported by The Department of Veterans Affairs, and by awards AI080193, AI69432, AI047745, AI74621, AI306214 (CFAR), and AI096113 (Collaboratory for AIDS Research on Eradication [CARE]) from the National Institutes of Health. SGD was supported by NIH grant award K24 AI069994 and SGD and RPS were supported by DARE: Delaney AIDS Research Enterprise (DARE; U19AI096109). We thank Dr. Chomont for helpful discussion and valuable insights.

## References

1. Hocqueloux L, Prazuck T, Avettand-Fenoel V, Lafeuillade A, Cardon B, Viard J-P, et al. Long-term immunovirologic control following antiretroviral therapy interruption in patients treated at the time of primary HIV-1 infection. *AIDS*. 2010; 24(10):1598–601. [PubMed: 20549847]
2. Lodi S, Meyer L, Kelleher A, Rosinska M, Ghosn J, Sannes M, et al. Immunovirologic control 24 months after interruption of antiretroviral therapy initiated close to HIV seroconversion. *Archives of Internal Medicine*. 2012; 172(16):1–4.
3. Goujard C, Girault I, Rouzioux C, Lécuroux C, Deveau C, Chaix M, et al. HIV-1 control after transient antiretroviral treatment initiated in primary infection: role of patient characteristics and effect of therapy. *Antivir Ther*. 2012; 17(6):1001–9. [PubMed: 22865544]
4. Hutter G, Nowak D, Mossner M, Ganepola S, Müssig A, Allers K, et al. Long-Term Control of HIV by CCR5 Delta32/Delta32 Stem-Cell Transplantation. *New England Journal of Medicine*. 2009; 360(7):692–8. [PubMed: 19213682]
5. Allers K, Hütter G, Hofmann J, Loddenkemper C, Rieger K, Thiel E, et al. Evidence for the cure of HIV infection by CCR5 32/ 32 stem cell transplantation. *Blood*. 2011; 117(10):2791–9. [PubMed: 21148083]
6. Henrich, TJ.; Sciaranghella, G.; Li, J.; Gallien, S.; Ho, V.; LaCasce, A., et al. Long-term reduction in peripheral blood HIV-1 reservoirs following reduced-intensity conditioning allogeneic stem cell transplantation in two HIV-positive individuals. XIX International AIDS Conference; Washington, DC. July 22–27, 2012; p. Abstract THAA0101
7. Chun T-W, Justement JS, Moir S, Hallahan CW, Maenza J, Mullins JJ, et al. Decay of the HIV Reservoir in Patients Receiving Antiretroviral Therapy for Extended Periods: Implications for Eradication of Virus. *Journal of Infectious Diseases*. 2007; 195(12):1762–4. [PubMed: 17492591]
8. Ananworanich J, Schuetz A, Vandergeeten C, Sereti I, de Souza M, Rerknimitr R, et al. Impact of Multi-Targeted Antiretroviral Treatment on Gut T Cell Depletion and HIV Reservoir Seeding during Acute HIV Infection. *PLoS ONE*. 2012; 7(3):e33948. [PubMed: 22479485]
9. Archin NM, Vaidya NK, Kuruc JD, Liberty AL, Wiegand A, Kearney MF, et al. Immediate antiviral therapy appears to restrict resting CD4+ cell HIV-1 infection without accelerating the decay of latent infection. *Proceedings of the National Academy of Sciences*. 2012; 109(24):9523–8.
10. von Wyl V, Gianella S, Fischer M, Niederöst B, Kuster H, Battegay M, et al. Early Antiretroviral Therapy During Primary HIV-1 Infection Results in a Transient Reduction of the Viral Setpoint upon Treatment Interruption. *PLoS ONE*. 2011; 6(11):e27463. [PubMed: 22102898]
11. Hufert F, van Lunzen J, Janossy G, Bertram S, Schmitz J, Haller O, et al. Germinal centre CD4+ T cells are an important site of HIV replication in vivo. *AIDS*. 1997; 11(7):849–57. [PubMed: 9189209]
12. Stellbrink H, van Lunzen J, Hufert F, Fröschele G, Wolf-Vorbeck G, Zöllner B, et al. Asymptomatic HIV infection is characterized by rapid turnover of HIV RNA in plasma and lymph nodes but not of latently infected lymph-node CD4+ T cells. *AIDS*. 1997; 11(9):1103–10. [PubMed: 9233456]
13. Chomont N, El-Far M, Ancuta P, Trautmann L, Procopio FA, Yassine-Diab B, et al. HIV reservoir size and persistence are driven by T cell survival and homeostatic proliferation. *Nat Med*. 2009; 15(8):893–900. [PubMed: 19543283]
14. Deeks SG, Autran B, Berkhout B, Benkirane M, Cairns S, Chomont N, et al. Towards an HIV cure: a global scientific strategy. *Nat Rev Immunol*. 2012; 12(8):607–14. [PubMed: 22814509]
15. Palmer S, Maldarelli F, Wiegand A, Bernstein B, Hanna GJ, Brun SC, et al. Low-level viremia persists for at least 7 years in patients on suppressive antiretroviral therapy. *Proceedings of the National Academy of Sciences*. 2008; 105(10):3879–84.
16. Chun T-W, Murray D, Justement JS, Hallahan CW, Moir S, Kovacs C, et al. Relationship Between Residual Plasma Viremia and the Size of HIV Proviral DNA Reservoirs in Infected Individuals Receiving Effective Antiretroviral Therapy. *Journal of Infectious Diseases*. 2011; 204(1):135–8. [PubMed: 21628667]
17. Sigal A, Kim JT, Balazs AB, Dekel E, Mayo A, Milo R, et al. Cell-to-cell spread of HIV permits ongoing replication despite antiretroviral therapy. *Nature*. 2011; 477(7362):95–8. [PubMed: 21849975]

18. Swiggard WJ, Baytop C, Yu JJ, Dai J, Li C, Schretzenmair R, et al. Human Immunodeficiency Virus Type 1 Can Establish Latent Infection in Resting CD4+ T Cells in the Absence of Activating Stimuli. *Journal of Virology*. 2005; 79(22):14179–88. [PubMed: 16254353]
19. Cameron PU, Saleh S, Sallmann G, Solomon A, Wightman F, Evans VA, et al. Establishment of HIV-1 latency in resting CD4+ T cells depends on chemokine-induced changes in the actin cytoskeleton. *Proceedings of the National Academy of Sciences*. 2010; 107(39):16934–9.
20. Bosque A, Planelles V. Induction of HIV-1 latency and reactivation in primary memory CD4+ T cells. *Blood*. 2009; 113(1):58–65. [PubMed: 18849485]
21. Karn J, Stoltzfus C. Transcriptional and Posttranscriptional Regulation of HIV-1 Gene Expression. *Cold Spring Harb Perspect Med*. 2012 Feb 2.(2):a006916. [PubMed: 22355797]
22. Van Duynne R, Guendel I, Narayanan A, Gregg E, Shafagati N, Tyagi M, et al. Varying Modulation of HIV-1 LTR Activity by BAF Complexes. *Journal of Molecular Biology*. 2011; 411(3):581–96. [PubMed: 21699904]
23. Friedman J, Cho WK, Chu CK, Keedy KS, Archin NM, Margolis DM, et al. Epigenetic Silencing of HIV-1 by the Histone H3 Lysine 27 Methyltransferase Enhancer of Zeste 2. *Journal of Virology*. 2011; 85(17):9078–89. [PubMed: 21715480]
24. Hakre S, Chavez L, Shirakawa K, Verdin E. HIV latency: experimental systems and molecular models. *FEMS Microbiology Reviews*. 2012; 36(3):706–16. [PubMed: 22372374]
25. Rong L, Perelson AS. Modeling Latently Infected Cell Activation: Viral and Latent Reservoir Persistence, and Viral Blips in HIV-infected Patients on Potent Therapy. *PLoS Comput Biol*. 2009; 5(10):e1000533. [PubMed: 19834532]
26. Llibre J, Buzón MJ, Massanella M, Esteve A, Dahl V, Puertas M, et al. Treatment intensification with raltegravir in subjects with sustained HIV-1 viraemia suppression: a randomized 48-week study. *Antivir Ther*. 2012; 17(2):355–64. [PubMed: 22290239]
27. Buzon MJ, Massanella M, Llibre JM, Esteve A, Dahl V, Puertas MC, et al. HIV-1 replication and immune dynamics are affected by raltegravir intensification of HAART-suppressed subjects. *Nat Med*. 2010; 16(4):460–5. [PubMed: 20228817]
28. Yukl SA, Shergill AK, McQuaid K, Gianella S, Lampiris H, Hare CB, et al. Effect of raltegravir-containing intensification on HIV burden and T-cell activation in multiple gut sites of HIV-positive adults on suppressive antiretroviral therapy. *AIDS*. 2010; 24(16):2451–60. [PubMed: 20827162]
29. Cohen J. Tissue Says Blood Is Misleading, Confusing HIV Cure Efforts. *Science*. 2011; 334(6063):1614. [PubMed: 22194536]
30. Cannon P, June C. Chemokine receptor 5 knockout strategies. *Current Opinion in HIV and AIDS*. 2011; 6(1):74–9. [PubMed: 21242897]
31. Mussolino C, Morbitzer R, Lütge F, Dannemann N, Lahaye T, Cathomen T. A novel TALE nuclease scaffold enables high genome editing activity in combination with low toxicity. *Nucleic Acids Research*. 2011; 39(21):9283–93. [PubMed: 21813459]
32. Schiffer JT, Aubert M, Weber ND, Mintzer E, Stone D, Jerome KR. Targeted DNA Mutagenesis for the Cure of Chronic Viral Infections. *Journal of Virology*. 2012; 86(17):8920–36. [PubMed: 22718830]
33. Tebas, P. Disruption of CCR5 in zinc finger nuclease-treated CD4 T cells: phase I trials. 18th Conference on Retroviruses and Opportunistic Infections; 2011. p. Paper 165
34. Mitsuyasu RT, Merigan TC, Carr A, Zack JA, Winters MA, Workman C, et al. Phase 2 gene therapy trial of an anti-HIV ribozyme in autologous CD34+ cells. *Nat Med*. 2009; 15(3):285–92. [PubMed: 19219022]
35. Wheeler LA, Trifonova R, Vrbancac V, Basar E, McKernan S, Xu Z, et al. Inhibition of HIV transmission in human cervicovaginal explants and humanized mice using CD4 aptamer-siRNA chimeras. *The Journal of Clinical Investigation*. 2011; 121(6):2401–12. [PubMed: 21576818]
36. van Lunzen J, Glaunsinger T, Stahmer I, Baehr Vv, Baum C, Schilz A, et al. Transfer of Autologous Gene-modified T Cells in HIV-infected Patients with Advanced Immunodeficiency and Drug-resistant Virus. *Mol Ther*. 2007; 15(5):1024–33. [PubMed: 17356541]

37. von Laer D, Baum C, Schambach A, Kuhlcke K, Zahn R, Newrzela S, van Lunzen J, Johnson RP, Schmitz JE. Gene Therapeutic Approaches for Immune Modulation in AIDS. *Anti-Inflammatory & Anti-Allergy Agents in Medicinal Chemistry*. 2007; 6:121–40.
38. Chan JK, Greene WC. Dynamic roles for NF- $\kappa$ B in HTLV-I and HIV-1 retroviral pathogenesis. *Immunological Reviews*. 2012; 246(1):286–310. [PubMed: 22435562]
39. Wolschendorf F, Bosque A, Shishido T, Duverger A, Jones J, Planelles V, et al. Kinase control prevents HIV-1 reactivation in spite of high levels of induced NF- $\kappa$ B activity. *Journal of Virology*. 2012; 86(8):4548–58. [PubMed: 22345467]
40. Mitsuyasu R, Gelman R, Cherng D, Landay A, Fahey J, Reichman R, et al. The virologic, immunologic, and clinical effects of interleukin 2 with potent antiretroviral therapy in patients with moderately advanced human immunodeficiency virus infection: a randomized controlled clinical trial-AIDS Clinical Trials Group 328. *Arch Intern Med*. 2007; 167(6):597–605. [PubMed: 17389292]
41. Stellbrink H, Hufert F, Tenner-Racz K, Lauer J, Schneider C, Albrecht H, et al. Kinetics of productive and latent HIV infection in lymphatic tissue and peripheral blood during triple-drug combination therapy with or without additional interleukin-2. *Antivir Ther*. 1998; 3(4):209–14. [PubMed: 10682140]
42. Kulkosky J, Nunnari G, Otero M, Calarota S, Dornadula G, Zhang H, et al. Intensification and Stimulation Therapy for Human Immunodeficiency Virus Type 1 Reservoirs in Infected Persons Receiving Virally Suppressive Highly Active Antiretroviral Therapy. *Journal of Infectious Diseases*. 2002; 186(10):1403–11. [PubMed: 12404155]
43. Wolschendorf F, Duverger A, Jones J, Wagner FH, Huff J, Benjamin WH, et al. Hit-and-Run Stimulation: a Novel Concept To Reactivate Latent HIV-1 Infection without Cytokine Gene Induction. *Journal of Virology*. 2010; 84(17):8712–20. [PubMed: 20538859]
44. Perez M, de Vinuesa AG, Sanchez-Duffhues G, Marquez N, Bellido ML, Munoz-Fernandez A, et al. Bryostatin-1 Synergizes with Histone Deacetylase Inhibitors to Reactivate HIV-1 from Latency. *Current HIV Research*. 2010 Sep; 8(6):418–29. [PubMed: 20636281]
45. DeChristopher BA, Loy BA, Marsden MD, Schrier AJ, Zack JA, Wender PA. Designed, synthetically accessible bryostatin analogues potently induce activation of latent HIV reservoirs in vitro. *Nat Chem*. 2012; 4(9):705–10. [PubMed: 22914190]
46. Van Lint C, Emiliani S, Ott M, Verdin E. Transcriptional activation and chromatin remodeling of the HIV-1 promoter in response to histone acetylation. *EMBO J*. 1996; 15(5):1112–20. [PubMed: 8605881]
47. Siliciano JD, Lai J, Callender M, Pitt E, Zhang H, Margolick JB, et al. Stability of the Latent Reservoir for HIV-1 in Patients Receiving Valproic Acid. *Journal of Infectious Diseases*. 2007; 195(6):833–6. [PubMed: 17299713]
48. Archin NM, Cheema M, Parker D, Wiegand A, Bosch RJ, Coffin JM, et al. Antiretroviral Intensification and Valproic Acid Lack Sustained Effect on Residual HIV-1 Viremia or Resting CD4+ Cell Infection. *PLoS ONE*. 2010; 5(2):e9390. [PubMed: 20186346]
49. Routy JP, Tremblay CL, Angel JB, Trottier B, Rouleau D, Baril JG, et al. Valproic acid in association with highly active antiretroviral therapy for reducing systemic HIV-1 reservoirs: results from a multicentre randomized clinical study. *HIV Medicine*. 2012; 13(5):291–6. [PubMed: 22276680]
50. Archin NM, Liberty AL, Kashuba AD, Choudhary SK, Kuruc JD, Crooks AM, et al. Administration of vorinostat disrupts HIV-1 latency in patients on antiretroviral therapy. *Nature*. 2012; 487(7408):482–5. [PubMed: 22837004]
51. Geleziunas, R.; Stephan, G.; Wei, G.; Yu, H.; Graupe, M.; Pagratis, N., et al. Search for Small Molecule Activators of Latent HIV. *Retrovirology; International Symposium on HIV and Emerging Infectious Diseases; Marseille, France. May 24, 2012; 2012. p. 02*
52. Sahu G, Cloyd M. Latent HIV in primary T lymphocytes is unresponsive to histone deacetylase inhibitors. *Virology*. 2011; 8 (400):1–6. [PubMed: 21205327]
53. Sakane N, Kwon HS, Pagans S, Kaehlcke K, Mizusawa Y, Kamada M, et al. Activation of HIV Transcription by the Viral Tat Protein Requires a Demethylation Step Mediated by Lysine-specific Demethylase 1 (LSD1/KDM1). *PLoS Pathog*. 2011; 7(8):e1002184. [PubMed: 21876670]

54. Imai K, Togami H, Okamoto T. Involvement of Histone H3 Lysine 9 (H3K9) Methyltransferase G9a in the Maintenance of HIV-1 Latency and Its Reactivation by BIX01294. *Journal of Biological Chemistry*. 2010; 285(22):16538–45. [PubMed: 20335163]
55. Shishido T, Wolschendorf F, Duverger A, Wagner F, Kappes J, Jones J, et al. Selected Drugs with Reported Secondary Cell-Differentiating Capacity Prime Latent HIV-1 Infection for Reactivation. *Journal of Virology*. 2012; 86(17):9055–69. [PubMed: 22696646]
56. Xing S, Bullen CK, Shroff NS, Shan L, Yang HC, Manucci JL, et al. Disulfiram Reactivates Latent HIV-1 in a Bcl-2-Transduced Primary CD4+ T Cell Model without Inducing Global T Cell Activation. *Journal of Virology*. 2011; 85(12):6060–4. [PubMed: 21471244]
57. Spivak, A.; Andrade, A.; Hoh, R.; Bacchetti, P.; Eisele, E.; Buckheit, R., III, et al. Safety and Feasibility of Using Disulfiram to Enhance HIV Transcription among Long-term ARV-treated Adults: Preliminary Results from a Pilot Study. 19th Conference on Retroviruses and Opportunistic Infections; March 5–8, 2012; Seattle, WA. p. Session 48
58. Patsoukis N, Brown J, Petkova V, Liu F, Li L, Boussiotis VA. Selective Effects of PD-1 on Akt and Ras Pathways Regulate Molecular Components of the Cell Cycle and Inhibit T Cell Proliferation. *Sci Signal*. 2012; 5(230):ra46. [PubMed: 22740686]
59. Yokosuka T, Takamatsu M, Kobayashi-Imanishi W, Hashimoto-Tane A, Azuma M, Saito T. Programmed cell death 1 forms negative costimulatory microclusters that directly inhibit T cell receptor signaling by recruiting phosphatase SHP2. *The Journal of Experimental Medicine*. 2012; 209(6):1201–17. [PubMed: 22641383]
60. Che K, Shankar E, Muthu S, Zandi S, Sigvardsson M, Hinkula J, et al. P38MAPK/STAT3 Pathway Signaling Regulates Expression of Inhibitory Molecules in T-cells Activated by HIV-1 Exposed Dendritic Cells. *Mol Med*. 2012; 18:1169–82. [PubMed: 22777388]
61. Said EA, Dupuy FP, Trautmann L, Zhang Y, Shi Y, El-Far M, et al. Programmed death-1-induced interleukin-10 production by monocytes impairs CD4+ T cell activation during HIV infection. *Nat Med*. 2010; 16(4):452–9. [PubMed: 20208540]
62. Hatano H, Jain V, Hunt PW, Lee TH, Sinclair E, Do TD, et al. Cell-based Measures of Viral Persistence Are Associated with Immune Activation and PD-1 Expressing CD4+ T cells. *Journal of Infectious Diseases*. 2012; 10:1093.
63. Trautmann L, Janbazian L, Chomont N, Said EA, Gimmig S, Bessette B, et al. Upregulation of PD-1 expression on HIV-specific CD8+ T cells leads to reversible immune dysfunction. *Nat Med*. 2006; 12(10):1198–202. [PubMed: 16917489]
64. Peretz Y, He Z, Shi Y, Yassine-Diab B, Goulet JP, Bordi R, et al. CD160 and PD-1 Co-Expression on HIV-Specific CD8 T Cells Defines a Subset with Advanced Dysfunction. *PLoS Pathogens*. 2012; 8(8):e1002840. [PubMed: 22916009]
65. Mackall CL, Fry TJ, Gress RE. Harnessing the biology of IL-7 for therapeutic application. *Nat Rev Immunol*. 2011; 11(5):330–42. [PubMed: 21508983]
66. Levy Y, Lacabaratz C, Weiss L, Viard J, Goujard C, Lelièvre J, et al. Enhanced T cell recovery in HIV-1-infected adults through IL-7 treatment. *The Journal of Clinical Investigation*. 2009; 119(4):997–1007. [PubMed: 19287090]
67. Sereti I, Dunham RM, Spritzler J, Aga E, Proschan MA, Medvik K, et al. IL-7 administration drives T cell cycle entry and expansion in HIV-1 infection. *Blood*. 2009; 113(25):6304–14. [PubMed: 19380868]
68. Levy Y, Sereti I, Tambussi G, Routy JP, Lelievre JD, Delfraissy JF, et al. Effects of Recombinant Human Interleukin 7 on T-Cell Recovery and Thymic Output in HIV-Infected Patients Receiving Antiretroviral Therapy: Results of a Phase I/IIa Randomized, Placebo-Controlled, Multicenter Study. *Clinical Infectious Diseases*. 2012; 55(2):291–300. [PubMed: 22550117]
69. Wang F-X, Xu Y, Sullivan J, Souder E, Argyris EG, Acheampong EA, et al. IL-7 is a potent and proviral strain-specific inducer of latent HIV-1 cellular reservoirs of infected individuals on virally suppressive HAART. *The Journal of Clinical Investigation*. 2005; 115(1):128–37. [PubMed: 15630452]
70. Vandergeeten C, DaFonseca S, Fromentin R, Sekaly R, Chomont N. Differential Impact of IL-7 and IL-15 on HIV Reservoir Persistence. In Fianl Program and Abstracts of the 5th International

Workshop on HIV Persistence during Therapy, St Maarten, West Indies, December 6–9, 2011, oral presentation 29. *Global Antiviral J.* 2011; 7(suppl 2):23.

71. Sandler NG, Wand H, Roque A, Law M, Nason MC, Nixon DE, et al. Plasma Levels of Soluble CD14 Independently Predict Mortality in HIV Infection. *Journal of Infectious Diseases.* 2011; 203(6):780–90. [PubMed: 21252259]
72. Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, et al. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med.* 2006; 12(12):1365–71. [PubMed: 17115046]
73. Marchetti G, Cozzi-Lepri A, Merlini E, Bellistri GM, Castagna A, Galli M, et al. Microbial translocation predicts disease progression of HIV-infected antiretroviral-naïve patients with high CD4+ cell count. *AIDS.* 2011; 25(11):1385–94. [PubMed: 21505312]
74. van Grevenynghe J, Halwani R, Chomont N, Ancuta P, Peretz Y, Tanel A, et al. Lymph node architecture collapse and consequent modulation of FOXO3a pathway on memory T- and B-cells during HIV infection. *Seminars in Immunology.* 2008; 20(3):196–203. [PubMed: 18757210]
75. Zeng M, Southern PJ, Reilly CS, Beilman GJ, Chipman JG, Schacker TW, et al. Lymphoid Tissue Damage in HIV-1 Infection Depletes Naïve T Cells and Limits T Cell Reconstitution after Antiretroviral Therapy. *PLoS Pathog.* 2012; 8(1):e1002437. [PubMed: 22241988]
76. Tenner-Racz K, Stellbrink H, van Lunzen J, Schneider C, Jacobs J, Raschdorff B, et al. The Unenlarged Lymph Nodes of HIV-1-Infected, Asymptomatic Patients with High CD4 T Cell Counts Are Sites for Virus Replication and CD4 T Cell Proliferation. The Impact of Highly Active Antiretroviral Therapy. *The Journal of Experimental Medicine.* 1998; 187(6):949–59. [PubMed: 9500797]
77. Stellbrink H, van Lunzen J. Lymph nodes during antiretroviral therapy. *Current Opinion in Infectious Diseases.* 2001; 14(1):17–22. [PubMed: 11979110]
78. Popovic M, Tenner-Racz K, Pelsler C, Stellbrink HJ, van Lunzen J, Lewis G, et al. Persistence of HIV-1 structural proteins and glycoproteins in lymph nodes of patients under highly active antiretroviral therapy. *Proceedings of the National Academy of Sciences of the United States of America.* 2005; 102(41):14807–12. [PubMed: 16199516]
79. Catalfamo M, Wilhelm C, Tcheung L, Proschan M, Friesen T, Park J, et al. CD4 and CD8 T cell immune activation during chronic HIV infection: roles of homeostasis, HIV, type I IFN, and IL-7. *J Immunol.* 2011; 186(4):2106–16. [PubMed: 21257970]
80. Chen H, Li C, Huang J, Cung T, Seiss K, Beamon J, et al. CD4+ T cells from elite controllers resist HIV-1 infection by selective upregulation of p21. *The Journal of Clinical Investigation.* 2011; 121(4):1549–60. [PubMed: 21403397]
81. Kvale D, Ormaasen V, Kran A, Johansson C, Aukrust P, Aandahl E, et al. Immune modulatory effects of cyclooxygenase type 2 inhibitors in HIV patients on combination antiretroviral treatment. *AIDS.* 2006; 20(6):813–20. [PubMed: 16549964]
82. Pettersen FO, Torheim EA, Dahm AE, Aaberge IS, Lind A, Holm M, et al. An Exploratory Trial of Cyclooxygenase Type 2 Inhibitor in HIV-1 Infection: Downregulated Immune Activation and Improved T Cell-Dependent Vaccine Responses. *Journal of Virology.* 2011; 85(13):6557–66. [PubMed: 21490090]
83. Peters JM, Shah YM, Gonzalez FJ. The role of peroxisome proliferator-activated receptors in carcinogenesis and chemoprevention. *Nat Rev Cancer.* 2012; 12(3):181–95. [PubMed: 22318237]
84. Mencarelli A, Francisci D, Renga B, D'Amore C, Cipriani S, Basile F, et al. Ritonavir-induced lipodystrophy and dyslipidaemia is reversed by the anti-inflammatory drug leflunomide in a PPAR- $\alpha$ -dependent manner. *Antivir Ther.* 2012; 17(4):669–78. [PubMed: 22297608]
85. Murray SM, Down CM, Boulware DR, Stauffer WM, Cavert WP, Schacker TW, et al. Reduction of Immune Activation with Chloroquine Therapy during Chronic HIV Infection. *Journal of Virology.* 2010; 84(22):12082–6. [PubMed: 20844049]
86. Piconi S, Parisotto S, Rizzardini G, Passerini S, Terzi R, Argentero B, et al. Hydroxychloroquine drastically reduces immune activation in HIV-infected, antiretroviral therapy-treated immunologic nonresponders. *Blood.* 2011; 118(12):3263–72. [PubMed: 21576701]
87. Paton N, Goodall R, Dunn D, Franzen S, Collaco-Moraes Y, Gazzard B, et al. Effects of hydroxychloroquine on immune activation and disease progression among HIV-infected patients

- not receiving antiretroviral therapy: a randomized controlled trial. *The Journal of the American Medical Association*. 2012; 308(4):353–61.
88. Shan L, Deng K, Shroff N, Durand C, Rabi S, Yang H, et al. Stimulation of HIV-1-Specific Cytolytic T Lymphocytes Facilitates Elimination of Latent Viral Reservoir after Virus Reactivation. *Immunity*. 2012; 36(3):491–501. [PubMed: 22406268]
  89. Descours B, Avettand-Fenoel V, Blanc C, Samri A, Mélard A, Supervie V, et al. Immune Responses Driven by Protective Human Leukocyte Antigen Alleles From Long-term Nonprogressors Are Associated With Low HIV Reservoir in Central Memory CD4 T Cells. *Clinical Infectious Diseases*. 2012; 54(10):1495–1503. [PubMed: 22441653]
  90. Almeida J, Price D, Papagno L, Arkoub Z, Sauce D, Bornstein E, et al. Superior control of HIV-1 replication by CD8+ T cells is reflected by their avidity, polyfunctionality, and clonal turnover. *The Journal of Experimental Medicine*. 2007; 204(10):2473–85. [PubMed: 17893201]
  91. Berger CT, Frahm N, Price DA, Mothe B, Ghebremichael M, Hartman KL, et al. High-Functional-Avidity Cytotoxic T Lymphocyte Responses to HLA-B-Restricted Gag-Derived Epitopes Associated with Relative HIV Control. *Journal of Virology*. 2011; 85(18):9334–45. [PubMed: 21752903]
  92. Autran B, Carcelain G, Combadiere B, Debre P. Therapeutic Vaccines for Chronic Infections. *Science*. 2004; 305(5681):205–8. [PubMed: 15247470]
  93. Lichterfeld M, Gandhi R, Simmons R, Flynn T, Sbrolla A, Yu X, et al. Induction of strong HIV-1-specific CD4+ T-cell responses using an HIV-1 gp120/NefTat vaccine adjuvanted with AS02A in antiretroviral-treated HIV-1-infected individuals. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2012; 59(1):1–9.
  94. Vanham G, Van Gulck E. Can immunotherapy be useful as a “functional cure” for infection with Human Immunodeficiency Virus-1? *Retrovirology*. 2012; 9(1):72. [PubMed: 22958464]
  95. Garcia F, Climent N, Assoumou L, Gil C, Gonzalez N, Alcami J, et al. A Therapeutic Dendritic Cell-Based Vaccine for HIV-1 Infection. *Journal of Infectious Diseases*. 2011; 203(4):473–8. [PubMed: 21233310]
  96. Li JZ, Brumme CJ, Lederman MM, Brumme ZL, Wang H, Spritzler J, et al. Characteristics and Outcomes of Initial Virologic Suppressors during Analytic Treatment Interruption in a Therapeutic HIV-1 gag Vaccine Trial. *PLoS ONE*. 2012; 7(3):e34134. [PubMed: 22479542]
  97. Azzoni L, Foulkes AS, Pappasavvas E, Mexas AM, Lynn KM, Mounzer K, et al. Pegylated Interferon-alpha-2A mono-therapy results in suppression of HIV-1 replication and decreased cell-associated HIV DNA integration. *Journal of Infectious Diseases*. 2013; 207(2):213–22. [PubMed: 23105144]
  98. Ahlers JD, Belyakov IM. Lessons learned from natural infection: focusing on the design of protective T cell vaccines for HIV/AIDS. *Trends in Immunology*. 2010; 31(3):120–30. [PubMed: 20089450]
  99. Vezys V, Penaloza-MacMaster P, Barber DL, Ha S-J, Konieczny B, Freeman GJ, et al. 4-1BB Signaling Synergizes with Programmed Death Ligand 1 Blockade To Augment CD8 T Cell Responses during Chronic Viral Infection. *The Journal of Immunology*. 2011; 187(4):1634–42. [PubMed: 21742975]
  100. Aubert RD, Kamphorst AO, Sarkar S, Vezys V, Ha S-J, Barber DL, et al. Antigen-specific CD4 T-cell help rescues exhausted CD8 T cells during chronic viral infection. *Proceedings of the National Academy of Sciences*. 2011; 108(52):21182–7.
  101. Sneller MC, Kopp WC, Engelke KJ, Yovandich JL, Creekmore SP, Waldmann TA, et al. IL-15 administered by continuous infusion to rhesus macaques induces massive expansion of CD8+ T effector memory population in peripheral blood. *Blood*. 2011; 118(26):6845–8. [PubMed: 22067383]
  102. Kim YK, Mbonye U, Hokello J, Karn J. T-Cell Receptor Signaling Enhances Transcriptional Elongation from Latent HIV Proviruses by Activating P-TEFb through an ERK-Dependent Pathway. *Journal of Molecular Biology*. 2011; 410(5):896–916. [PubMed: 21763495]
  103. Gallastegui E, Marshall B, Vidal D, Sanchez-Duffhues G, Collado J, Alvarez-Fernández C, et al. Combination of Biological Screening in a Cellular Model of Viral Latency and Virtual Screening



- Identifies Novel Compounds That Reactivate HIV-1. *Journal of Virology*. 2012; 86(7):3795–808. [PubMed: 22258251]
104. Kovoichich M, Marsden MD, Zack JA. Activation of Latent HIV Using Drug-Loaded Nanoparticles. *PLoS ONE*. 2011; 6(4):e18270. [PubMed: 21483687]
105. Wightman F, Ellenberg P, Churchill M, Lewin SR. HDAC inhibitors in HIV. *Immunol Cell Biol*. 2012; 90(1):47–54. [PubMed: 22083528]
106. Rasmussen, T.; Sjøgaard, O.; Melchjorsen, J.; Brinkmann, C.; Østergaard, L.; Dinarello, C., et al. The Histone Deacetylase Inhibitor (HDACi) panobinostat (LBH589) Stimulates HIV-1 Expression More Potently than Other HDACi in Clinical Use and Disrupts HIV Latency at Clinically Achievable Concentrations. 19th Conference on Retrovirus and Opportunistic Infections; March 5–8, 2012; Seattle, Washington, USA. p. 370
107. Contreras X, Mzoughi O, Gaston F, Peterlin BM, Bahraoui E. Protein kinase C-delta regulates HIV-1 replication at an early post-entry step in macrophages. *Retrovirology*. 2012; 9(1):37. [PubMed: 22554282]
108. Bouchat S, Gatot J-S, Kabeya K, Cardona C, Colin L, Herbein G, et al. Histone methyltransferase inhibitors induce HIV-1 recovery in resting CD4+ T cells from HIV-1-infected HAART-treated patients. *AIDS*. 2012; 26(12):1473–82. [PubMed: 22555163]
109. Banerjee C, Archin N, Michaels D, Belkina AC, Denis GV, Bradner J, et al. BET bromodomain inhibition as a novel strategy for reactivation of HIV-1. *Journal of Leukocyte Biology*. 2012:92. 10.1189.
110. Lisziewicz J, Bakare N, Calarota S, Bánhegyi D, Szlávik J, Ujhelyi E, et al. Single DermaVir Immunization: Dose-Dependent Expansion of Precursor/Memory T Cells against All HIV Antigens in HIV-1 Infected Individuals. *PLoS ONE*. 2012; 7(5):e35416. [PubMed: 22590502]
111. Reuse S, Calao M, Kabeya K, Guiguen A, Gatot JS, Quivy V, et al. Synergistic Activation of HIV-1 Expression by Deacetylase Inhibitors and Prostratin: Implications for Treatment of Latent Infection. *PLoS ONE*. 2009; 4(6):e6093. [PubMed: 19564922]
112. Desimie BA, Humbert M, Lescrinier E, Hendrix J, Vets S, Gijsbers R, et al. Phage Display-directed Discovery of LEDGF/p75 Binding Cyclic Peptide Inhibitors of HIV Replication. *Mol Ther*. 2012; 20(11):2064–2075. [PubMed: 22828501]

Table 1

CURE-related clinical pilot trials ongoing in 2012

Study/Principal investigator	Intervention	Study Population	Primary End -point	Clinicaltrials.gov
<b>ERAMUNE 01C Katlama</b> Objectif Recherche Vaccins SIDA (ORVACS) University PARIS VI	IL-7 plus ART intensification with RAL and MVC	29 chronically HIV infected patients with HIV RNA < 50 cp/ml on ART -HIVDNA :10–1000 cp/10 <sup>6</sup> PBMC	Decrease of 0.5 log <sub>10</sub> HIV DNA from baseline at 56 weeks	NCT01019551
<b>ERAMUNE02 R. Murphy</b> ORVACS Northwestern University	HIV rAD5 vaccine and ART intensification with RAL and MVC	28 chronically HIV infected patient with HIV RNA < 50 cp/ml on ART-HIV DNA 10+1000 cp/10 <sup>6</sup> PBMC -Negative AD5 Ab	Decrease of -0.5 log <sub>10</sub> HIV DNA from baseline at 56 weeks	NCT00976404
<b>Disulfiram S. Deeks, A. Andrade</b> University of California, San Francisco. Johns Hopkins University	Disulfiram: 500mg/d x 14 days	20 chronically HIV-1 patients with HIV RNA < 50 cp/ml on ART	2 weeks frequency of replication, competent HIV-1 in resting CD4+ T cells	NCT01286259
<b>Vorinostat D. Margolis</b> University of North Carolina, Chapel Hill	Vorinostat (87) 200 mg to 600 mg	20 chronically HIV-1 patients with HIV RNA < 50 cp/ml	HIV expression in resting CD4+ T cells	NCT01319383
<b>Vorinostat S. Lewin</b> Monash University and The Alfred Hospital	Vorinostat 400 mg/d	20 chronically HIV-1 patients with HIV RNA < 50 cp/ml - CD4 >500/mm <sup>3</sup>	HIV unspliced RNA in resting CD4+ T cells at 28 days	NCT01365065
<b>Panobinostat T. Rasmussen</b> Aarhus University Denemark	Panobinostat 20mg Three times a week Every other week for 8 weeks	16 chronic HIV+ patients with suppressed viremia and CD4>	HIV unspliced RNA in resting CD4+ T cells Change from baseline	NCT01680094).
<b>CD4+ T cells modified at CCR5 by ZFN</b> Winson TangSangamo Biosciences	Autologous CD4+ T cells modified at CCR5 gene by Zinc-finger nuclease SB-728-T	Chronic HIV+ patients with suppressed viremia and CD4 < 300/mm <sup>3</sup> or ART failing	Safety Persistence and activation of CCR5+ZFN modified autologous T cells	NCT01252641
<b>CD4+ T cells modified at CCR5 by ZFN</b> P. Tebas, D. Stein, C. June Pennsylvania University Jacobi Medical Center	Autologous CD4+ T cells modified at CCR5 gene by Zinc-finger nuclease	18 patients: 3 cohorts of HIV+ patients, either failing ART or with suppressed viremia	Safety	NCT00842634
<b>Lentivirus vector rHIV7-shI-TARCCR5RZ-transduced hematopoietic progenitor cells</b> A. Krishnan City of Hope Medical Center	Autologous CD34+ hematopoietic cells modified by lentivirus transduced nonfunctional CCR5RZ gene	Hematopoietic stem cell transplant AIDS related lymphoma	Safety and durability of transduced cells	NCT00569985
<b>Interferon 2b F. Maldarelli</b> National Cancer Institute (NCI)	INF 2b intensification	Recruiting, Non-randomized, single group assignment	Efficacy: viral RNA levels in blood and sequence diversification	NCT01295515

**Table 2**

## Experimental Approaches For Reactivating Latent HIV infection

Approach	Mechanism of Action	Activation	Refs.
<b><i>NF-κB Inducing Agents</i></b>			
Anti-CD3/CD28	T-cell receptor activation	++++	102,103
TNF- <sup>*</sup>	TNFR	++	38
Prostratin PMA/ionomycin Bryostatin-1, Picolog	PKC activation	+++	44, 104
<b><i>HDAC inhibitors</i></b>			
SAHA (vorinostat)	Class I HDAC1, HDAC3 inhibitor; HDAC4 and HDAC6 Cell cycle arrest: Myc •p21 Reactive oxygen species– apoptosis nonhomologous DSB repair Angiogenesis: HIF <sup>*</sup> , VEGF Innate immunity, proinflammatory cytokines STAT 5 hyperacetylation		
Depsiptide/FK228 (romidepsin)	HDACs 1, 2, 4, and 6 inhibitor	+++	51
BH589 (panobinostat) ITF2357 (givinostat) PXD101 (belinostat)	Potency in latently infected cell lines: LBH589 >ITF2357 and PXD101 >SAHA >VPA	+++	105, 106
Sirtuin inhibitors	Acetylation of nonhistone proteins: HIV Tat, p65 RelA, p53, FOXO3a	untested	
<b><i>HMT inhibitors</i></b>			
BIX-01294	G9a (H3K9me2) inhibitor	++	54, 107
chaetocin	SUV39H (H3K9me3), inhibitor	++	108
<b><i>Proapoptotic and cell differentiating molecules</i></b>			
JQ1	Blocks TAT binding to P-TEFb Inhibits bromodomain reader BRD4 recruitment to H3K9ac, H4K16ac myc transcription SIRT1	+ priming	109
Nutlin3	MDM2 blocker, p53 transcription, caspase activation	untested	
Disulfiram, aphidicolin	Oxidative stress, apoptosis	++	56
HMBA Dactinomycin, aclacinomycin Cytarabine	P-TEFb release associated with RNAP II triggers transcriptional elongation	+	55
Wnt small molecule inhibitors, Notch ( -secretase) inhibitors	myc, CtBP1, Cyclin D, cJun HES target genes	untested	
<b><i>Immune-modulators</i></b>			
Anti-PD-1, CTLA-4, TIM-3 mAb	Block negative regulatory receptors	anti-PD-1 (ACTG 5301)	
Anti-PD1mAb + IL-15	STAT5, PI3K signaling, CD4 <sup>+</sup> T cell effector memory differentiation	untested	
Therapeutic CD4 <sup>+</sup> T cell vaccine + raltegravir	Reactivate virus in latently infected antigen-specific CD4 cells To limit reseeding viral reservoirs	Not determined	93, 110
<b><i>Synergistic combinations</i></b>			
SAHA + chateocin + prostratin	HDAC inhibitor + HMTI HDAC inhibitor + NF <sup>*</sup> B inducer	+++	108, 111

Approach	Mechanism of Action	Activation	Refs.
HDACi+ anti-PD-1	HDAC inhibitor + anti-PD-1mAb.	preclinical studies	
Anti-PD1mAb + IL-15 + Raltegravir, peptidomimetics	CD4 <sup>+</sup> T effector memory cell differentiation, Inhibit Integrase, LEDGF/p75	untested	112

\* limited effect in primary cell models

phorbol ester 13-phorbol-12-myristate acetate (PMA), suberoylanilidehydroxamic acid (SAHA); Hexamethylenebisacetamide (HMBA); positive transcription elongation factor (p-TEFb)